


- c) $2r$ d) $2\sqrt{3}r$
- (n) If the ratio of mode and median of a certain data is $6 : 5$, then the ratio of its mean and median is [1]
- a) $10 : 9$ b) $9 : 10$
- c) $10 : 8$ d) $8 : 10$
- (o) **Assertion (A):** $a_n - a_{n-1}$ is not independent of n then the given sequence is an AP. [1]
- Reason (R):** Common difference $d = a_n - a_{n-1}$ is constant or independent of n .
- a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.
- c) A is true but R is false. d) A is false but R is true.

2. **Question 2** [12]

- (a) Mr. Gupta opened a recurring deposit account in a bank. He deposited ₹ 2,500 per month for 2 years. [4]
At the time of maturity, he got ₹67,500. Find:
- i. the total interest earned by Mr. Gupta
- ii. the rate of interest per annum.
- (b) Find the third proportional to [4]
- i. 16 and 36
- ii. $(x^2 + y^2 + xy)^2$ and $(x^3 - y^3)$
- (c) Prove that: $\frac{(1 + \cot \theta + \tan \theta)(\sin \theta - \cos \theta)}{\sec^3 \theta - \operatorname{cosec}^3 \theta} = \sin^2 \theta \cos^2 \theta$ [4]

3. **Question 3** [13]

- (a) The internal and external diameters of a hollow hemispherical vessel are 7cm and 14 cm, respectively. [4]
The cost of silver plating of 1 sq cm surface is ₹ 0.60. Find the total cost of silver plating the vessel all over.
- (b) The side AB of a square ABCD is parallel to the Y-axis as shown in the given figure. [4]



Calculate

- i. the slope of AD.
- ii. the slope of BD.
- iii. the slope of AC. [Given, $\tan(90^\circ + \theta) = -\cot\theta$]
- (c) Use graph paper to answer this question: [5]
- i. The point $P(2, -4)$ is reflected about the line $x = 0$ to get the image Q. Find the coordinates of Q.
- ii. Point Q is reflected about the line $y = 0$ to get the image R. Find the coordinates of R.
- iii. Name the figure PQR.
- iv. Find the area of figure PQR.

Section B

Attempt any 4 questions

4. **Question 4** [10]

(a) A shopkeeper bought an article with market price ₹1200 from the wholesaler at a discount of 10%. [3]

The shopkeeper sells this article to the customer on the market price printed on it. If the rate of GST is 6%, then find:

- i. GST paid by the wholesaler.
- ii. Amount paid by the customer to buy the item.

(b) The sum of the squares of two consecutive odd positive integers is 290. Find them. [3]

(c) Draw a Histogram for the given data, using a graph paper: [4]

Weekly Wages (in ₹)	No. of People
3000-4000	4
4000-5000	9
5000-6000	18
6000-7000	6
7000-8000	7
8000-9000	2
9000-10000	4

Estimate the mode from the graph.

5. **Question 5** [10]

(a) Evaluate, $\begin{bmatrix} 4 \sin 30^\circ & 2 \cos 60^\circ \\ \sin 90^\circ & 2 \cos 0^\circ \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 5 & 4 \end{bmatrix}$. [3]

(b) O is the circumcentre of the $\triangle ABC$ and D is mid-point of the base BC. Prove that $\angle BOD = \angle A$. [3]

(c) Use factor theorem to factorise $6x^3 + 17x^2 + 4x - 12$ completely. [4]

6. **Question 6** [10]

(a) Calculate the ratio in which the line joining A (-4, 2) and B(3, 6) is divided by P(x, 3). Also, find [3]

- i. x
- ii. length of AP

(b) Prove that: $1 - \frac{\cos^2 \theta}{1 + \sin \theta} = \sin \theta$ [3]

(c) Sum of the first n terms of an AP is $5n^2 - 3n$. Find the AP and also find its 16th term. [4]

7. **Question 7** [10]

(a) A grassy land is in the shape of a right triangle. The hypotenuse of the land is 1 m more than twice the shortest side. If the third side is 7 m more than the shortest side, find the sides of the grassy land. [5]

(b) The marks obtained by 100 students in a Mathematics test are given below [5]

Marks	Number of students
0-10	3
10-20	7
20-30	12

30-40	17
40-50	23
50-60	14
60-70	9
70-80	6
80-90	5
90-100	4


Draw an ogive for the given distribution on a graph sheet, (use a scale of 2 cm = 10 units on both axes). Use the ogive to estimate the

- median.
- lower quartile.
- number of students who obtained more than 85% marks in the test.
- number of students who did not pass in the test, if the pass percentage was 35.

8. **Question 8**

[10]

- A number is selected at random from first 50 natural numbers. Find the probability that it is a multiple of 3 and 4. [3]
- How many solid spheres of diameter 6 cm are required to be melted to form a cylindrical solid of height 45 cm and diameter 4 cm? [3]
- In the given figure, $AC = AE$. [4]



Show that


- $CP = EP$
- $BP = DP$

9. **Question 9**

[10]

- Solve the following inequation and represent the solution set on the number line. [3]

$$\frac{3x}{5} + 2 < x + 4 \leq \frac{x}{2} + 5, x \in \mathbb{R}$$
- Mode and mean of a data are $12k$ and $15k$ respectively. Find the median of the data. [3]
- In the given figure, if $DE \parallel BC$, find the ratio of ar ($\triangle ADE$) and ar (DECB). [4]



10. **Question 10**

[10]

- Find the fourth proportional to $(a^3 + 8)$, $(a^4 - 2a^3 + 4a^2)$ and $(a^2 - 4)$. [3]

- (b) Use a ruler and a pair of compasses to construct a $\triangle ABC$, in which $BC = 4.2$ cm, $\angle ABC = 60^\circ$ and $AB = 5$ cm. Construct a circle of radius 2 cm to touch both the arms of $\angle ABC$. [3]
- (c) A man observes the angle of elevation of the top of a building to 30° . He walks towards it in a horizontal line through its base. On covering 60 m, the angle of elevation changes to 60° . Find the height of the building correct to the nearest metre. [4]

Solution

Section A

1. Question 1 Choose the correct answers to the questions from the given options:

(i) (a) 18%

Explanation: {

C.P. = ₹ 25,000, CGST = ₹ 2250

∴ GST = 2 × ₹ 2250 = ₹ 4500

let rate of GST = r%

∴ r% of ₹ 25,000 = ₹ 4500

⇒ r = (4500 × 100) ÷ 25000 = 18%

(ii) (c) 576

Explanation: {

Let the total number of Saras birds be x.

Then, number of Saras birds moving in lotus plants = $\frac{x}{4}$

Number of Saras birds moving on a hill = $\frac{x}{9} + \frac{x}{4} + 7\sqrt{x}$

Number of Saras birds sitting on the Bakula trees = 56

According to the question,

$$\frac{x}{4} + \frac{x}{9} + \frac{x}{4} + 7\sqrt{x} + 56 = x$$

$$\Rightarrow 7\sqrt{x} = x - \frac{x}{4} - \frac{x}{9} - \frac{x}{4} - 56$$

$$\Rightarrow 7\sqrt{x} = \frac{36x - 9x - 4x - 9x}{36} = 56$$

$$\Rightarrow 7\sqrt{x} = \frac{7x}{18} - 56 \Rightarrow \sqrt{x} = \frac{x}{18} - 8$$

$$\Rightarrow x = \frac{x^2}{324} + 64 - \frac{8x}{9} \text{ [squaring on both sides]}$$

$$\Rightarrow x = \frac{x^2 + 20736 - 288x}{324}$$

$$\Rightarrow 324x = x^2 + 20736 - 288x$$

$$\Rightarrow x^2 - 612x + 20736 = 0$$

$$\Rightarrow x^2 - 36x - 576x + 20736 = 0 \text{ [splitting the middle term]}$$

$$\Rightarrow x(x - 36) - 576(x - 36) = 0$$

$$\Rightarrow (x - 36)(x - 576) = 0$$

$$\Rightarrow x - 36 = 0 \text{ or } x - 576 = 0$$

$$\Rightarrow x = 576 \text{ or } x = 36$$

Here, x = 36 is not possible, because if there are only 36 birds, then 56 cannot be on the trees.

Thus, total number of Saras birds is 576.

(iii) (c) 6

Explanation: {

Let f(x) = 3x³ + kx² + 7x + 4

As x + 1 is a factor of f(x), f(-1) = 0

$$\Rightarrow 3(-1)^3 + k(-1)^2 + 7(-1) + 4 = 0$$

$$\Rightarrow -3 + k - 7 + 4 = 0$$

$$\Rightarrow k = 6$$

(iv) (a) 100

Explanation: {

$$\text{We have, } A = \begin{bmatrix} 5 & 5 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 5 & 5 \end{bmatrix} = \begin{bmatrix} 25 & 25 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 5^2 & 5^2 \\ 0 & 0 \end{bmatrix}$$

$$\therefore A^2 = A \cdot A = \begin{bmatrix} 5^2 & 5^2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 5^2 & 5^2 \\ 0 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 5^4 & 5^4 \\ 0 & 0 \end{bmatrix}$$

$$A^n = \begin{bmatrix} 5^{2n} & 5^{2n} \\ 0 & 0 \end{bmatrix}$$

$$\text{Thus, } \begin{bmatrix} 5^{2n} & 5^{2n} \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 5^{200} & 5^{200} \\ 0 & 0 \end{bmatrix}$$

$$\Rightarrow 5^{2n} = 5^{200}$$

$$\Rightarrow 2n = 200$$

$$\Rightarrow n = 100$$

(v) (d) 0

Explanation: {

Let A be the first term and R be the common ratio of the given GP.

Then, a = pth term $\Rightarrow a = AR^{p-1}$

$\Rightarrow \log a = \log A + (p-1)\log R$... (i)

b = qth term $\Rightarrow b = AR^{q-1}$

$\Rightarrow \log b = \log A + (q-1)\log R$... (ii)

c = rth term $\Rightarrow c = AR^{r-1}$

$\Rightarrow \log c = \log A + (r-1)\log R$... (iii)

\Rightarrow Now, consider $\{q-r\}\log a + (r-p)\log b + (p-q)\log c$

$= (q-r)\{\log A + (p-1)\log R\} + (r-p)\{\log A + (q-1)\log R\}$ [from Eqs. (i), (ii) and (iii)]

$= \log A\{(q-r) + (r-p) + (p-q)\} + \log R\{(p-1)(q-r) + (q-1)(r-p) + (r-1)(p-q)\}$

$= (\log A)0 + \{p(q-r) + q(r-p) + r(p-q) - (q-r) - (r-p) - (p-q)\}\log R$

$= (\log A)0 + (\log R)0 = 0$

(vi) (d) (4, -5)

Explanation: {

Since, the image of any point (x, y) under X-axis is (x, -y).

\therefore Coordinate of M \equiv (4, 5)

Since, the image of any point (x, y) under Y-axis is (-x, y).

\therefore Coordinate of M'' \equiv (4, -5)


Since, the image of any point (x, y) under origin is (-x, -y).

\therefore Coordinate of M''' = (4, -5)

(vii) (a) $3\sqrt{17}$ cm

Explanation: {

Given AB = 8 cm and BC = 6 cm



$\therefore AC = \sqrt{8^2 + 6^2} = 10$ cm

Also, given AC : CE = 2 : 1

Now, produce BC to meet DE at the point P as CP is parallel to AD,

$\triangle ECP \sim \triangle EAD$... (i)

$\Rightarrow \frac{CP}{AD} = \frac{CE}{AE} \Rightarrow \frac{CP}{6} = \frac{1}{3}$... (ii)

$\Rightarrow CP = 2$ cm

Also, $\triangle CPD$ is right triangle.

$\therefore DP = \sqrt{CD^2 + CP^2}$

$= \sqrt{6^2 + 2^2} = 2\sqrt{17}$ cm

But DP = PE = 2 : 1 [from Eq.(i)]


$\therefore PE = \sqrt{17}$ cm

Thus, DE = DP + PE = $2\sqrt{17} + \sqrt{17} = 3\sqrt{17}$ cm

(viii) (c) 489.84 cm^2

Explanation: {

According to the given information, a shape of figure is shown below



When the hanging pipes touches the surface paper, a circular shape ABCD is formed on the graph paper. The size of circle ABCD is equal to the size of circular base of the cone.

∴ Radius of the circle ABCD is 6 cm.

Hence, the coordinates of A, B, C and D are (6, 0), (0, 6), (-6, 0) and (0, -6), respectively.

The figure formed in the given information is cylindrical in outer surface and conical in the inner surface. Now, total surface area of the figure

= Curved surface area of the cylinder + Curved surface area of the cone

$$= 2\pi rh + \pi rl = \pi r(2h + l)$$

$$= \pi r(2h + \sqrt{r^2 + h^2})$$

$$= 3.14 \times 6(2 \times 8 + \sqrt{6^2 + 8^2})$$

$$= 18.84(16 + \sqrt{36 + 64})$$

$$= 18.84(16 + \sqrt{100}) = 18.84(16 + 10)$$

$$= 18.84 \times 26 = 489.84 \text{ cm}^2$$

(ix) (a) $(-\infty, \frac{3}{2})$

Explanation: {

We have, $(x + 1)^2 - (x - 1)^2 < 6$

$$\Rightarrow (x^2 + 1 + 2x) - (x^2 + 1 - 2x) < 6 \quad [\because (a \pm b)^2 = a^2 + b^2 \pm 2ab]$$

$$\Rightarrow 4x < 6$$

$$\Rightarrow x < \frac{6}{4}$$

$$\Rightarrow x < \frac{3}{2}$$

$$\Rightarrow x \in (-\infty, \frac{3}{2})$$

(x) (a) $\frac{1}{6}$

Explanation: {

In a wall clock, the minute hand cover the 60 min in on complete round.

∴ Total number of possible outcomes = 60

The minute hand cover the time from 5 to 15 min,

Number of outcomes favourable to E

= Distance from 5 to 15 min = 10

∴ Required probability = $\frac{10}{60} = \frac{1}{6}$

(xi) (c) $2^{-\frac{3}{2}}$

Explanation: {

$$\text{We have, } \begin{bmatrix} a^x \\ a^{-x} \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix} = \begin{bmatrix} p & a^{-2} \\ q & \log_2 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} a^x & 2a^x \\ a^{-x} & 2a^{-x} \end{bmatrix} = \begin{bmatrix} p & a^{-2} \\ q & 1 \end{bmatrix} \quad \left[\because \log_2 2 = \frac{\log 2}{\log 2} = 1 \right]$$

On comparing the corresponding elements both sides, we get

$$\Rightarrow a^x = p \dots(i)$$

$$\Rightarrow 2a^x = a^{-2} \dots(ii)$$

$$\Rightarrow a^{-x} = q \dots(iii)$$

$$\text{and } 2a^{-x} = 1 \dots(iv)$$

On multiplying Eqs. (ii) and (iv), we get

$$4a^{x-x} = a^{-2}$$

$$\Rightarrow 4a^0 = a^{-2} \Rightarrow 4 = a^{-2} \Rightarrow 4 = \frac{1}{a^2}$$

$$\Rightarrow a^2 = \frac{1}{4} \Rightarrow a = \frac{1}{2} [\because a > 0]$$

$$\text{Now, } a^{p-q} = a^{a^x - a^{-x}} \text{ [from Eqs. (i) and (iii)]}$$

$$= a^{\frac{1}{2}a^{-2} - \frac{1}{2}} = a^{2 - \frac{1}{2}} \text{ [from Eqs. (ii) and (iv)]}$$

$$= a^{\frac{1}{2} \cdot 4 - \frac{1}{2}} [\because a^{-2} = 4]$$


$$= \left(\frac{1}{2}\right)^{\frac{3}{2}} = 2^{-\frac{3}{2}}$$

(xii) (d) $45^\circ, 45^\circ$

Explanation: {

Given, coordinates of pole be $P(1, 3)$ and $Q(3, 3)$ and $A(1, 1)$ be the position of man

$$\begin{aligned} \text{i. Now, } AP &= \sqrt{(1-1)^2 + (3-1)^2} [\because \text{distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}] \\ &= \sqrt{0^2 + 2^2} = 2 \text{ units} \end{aligned}$$



$$\text{and } PO = \sqrt{(3-1)^2 + (3-3)^2} = \sqrt{2^2 + 0^2} = 2 \text{ units}$$

Now, in $\triangle APQ$, we have

$$\tan \theta = \frac{PQ}{AP} \Rightarrow \tan \theta = \frac{2}{2} = 1$$

$$\Rightarrow \theta = 45^\circ [\because \tan 45^\circ = 1]$$

ii. When we shift the origin at $(1, 1)$, then the angle will remain same, i.e. $\theta = 45^\circ$.

(xiii) (a) $2(\sqrt{3} + 1)r$


Explanation: {

As PQR is an equilateral triangle, hence PS will be perpendicular to QR and will divide it into 2 equal parts.

Since, $\angle P$ and $\angle S$ will be supplementary, so

$$\angle S = 120^\circ \text{ and } \angle QSA = \angle RSA = 60^\circ$$

$$\text{Now, } PA = PQ \cos 30^\circ \text{ and } OA = OQ \sin 30^\circ = \frac{r}{2}$$



$$\Rightarrow AS = OA = \frac{r}{2} \text{ and } PA = PO + OA = r + \frac{r}{2}$$

$$\text{Hence, } PQ = \frac{PA}{\cos 30^\circ} = \frac{r + \frac{r}{2}}{\frac{\sqrt{3}}{2}} = \sqrt{3}r$$

$$\text{In } \triangle QAS, AS = QS \cos 60^\circ \Rightarrow QS = \frac{\frac{r}{2}}{\frac{1}{2}} = r$$

Since, $AQ = AR$, AS is common and $\angle QAS = \angle RAS = 90^\circ$

So, $QS = RS$.

$$\therefore \text{Perimeter of } PQSP = 2(PQ + QS) = 2(\sqrt{3} + 1)r$$

(xiv) (b) 9 : 10

Explanation: {

We know that,

$$\text{Mode} = 3 \text{ Median} - 2 \text{ Mean}$$

On dividing both sides by median, we get

$$\begin{aligned} \frac{\text{Mode}}{\text{Median}} &= 3 - 2 \frac{\text{Mean}}{\text{Median}} \\ \Rightarrow \frac{6}{5} &= 3 - 2 \frac{\text{Mean}}{\text{Median}} \quad [\because \frac{\text{mode}}{\text{median}} = \frac{6}{5}, \text{ given}] \\ \Rightarrow \frac{6}{5} - 3 &= -2 \frac{\text{Mean}}{\text{Median}} \\ \Rightarrow \frac{6-15}{5} &= -2 \frac{\text{Mean}}{\text{Median}} \\ \Rightarrow \frac{-9}{5} &= -2 \frac{\text{Mean}}{\text{Median}} \\ \Rightarrow \frac{\text{Mean}}{\text{Median}} &= \frac{9}{10} \end{aligned}$$

(xv) (d) A is false but R is true.

Explanation: {

We have, common difference of an AP

$$d = a_n - a_{n-1} \text{ is independent of } n \text{ or constant.}$$

So, A is false but R is true.

2. Question 2

(i) i. $P = ₹ 2,500$

$$n = 24 \text{ months}$$

$$r = ?$$

$$\text{M.V.} = ₹ 67,500$$

Total money deposited in 2 years

$$= P \times n$$

$$= 2,500 \times 24$$

$$= 60,000$$

Total interest earned by Mr. Gupta = Maturity value - Money deposited

$$= 67,500 - 60,000$$

$$= ₹ 7,500$$

$$\text{ii. M.V.} = P \times n + \frac{P \times n(n+1) \times r}{2400}$$

$$67,500 = 2500 \times 24 + \frac{2500 \times 24 \times 25 \times r}{2400}$$

$$67,500 = 60,000 + 625r$$

$$67,500 - 60,000 = 625r$$

$$7,500 = 625r$$

$$\frac{7500}{625} = r$$

$$r = 12\% \text{ p.a.}$$

(ii) i. 16 and 36

Let the third proportional to 16 and 36 be x .

$$\Rightarrow 16, 36 \text{ and } x \text{ in continuous proportion.}$$

$$\Rightarrow 16 : 36 = 36 : x$$

$$\Rightarrow 16 \times x = 36 \times 36$$

$$\Rightarrow x = \frac{36 \times 36}{16}$$

$$\Rightarrow x = 81$$

ii. $(x^2 + y^2 + xy)^2$ and $x^3 - y^3$

Let third proportional to $(x^2 + y^2 + xy)^2$ and $x^3 - y^3$ be x .

$\Rightarrow (x^2 + y^2 + xy)^2, x^3 - y^3$ and x are in continuous proportion.

$$\Rightarrow (x^2 + y^2 + xy)^2 : x^3 - y^3 = x^3 - y^3 : x$$

$$x = \frac{(x^3 - y^3)^2}{(x^2 + y^2 + xy)^2}$$

$$x = \frac{(x-y)^2 (x^2 + y^2 + xy)^2}{(x^2 + y^2 + xy)^2} [\because x^3 - y^3 = (x-y)(x^2 + y^2 + xy)]$$

$$x = (x-y)^2$$

$$(iii) L.H.S. = \frac{(1 + \cot \theta + \tan \theta)(\sin \theta - \cos \theta)}{\sec^3 \theta - \operatorname{cosec}^3 \theta}$$

$$= \frac{\left(1 + \frac{\cos \theta}{\sin \theta} + \frac{\sin \theta}{\cos \theta}\right)(\sin \theta - \cos \theta)}{\frac{1}{\cos^3 \theta} - \frac{1}{\sin^3 \theta}}$$

$$= \frac{\left(\frac{1 + \cos^2 \theta + \sin^2 \theta}{\sin \theta \cos \theta}\right)(\sin \theta - \cos \theta)}{\frac{\sin^3 \theta - \cos^3 \theta}{\sin^3 \theta \cos^3 \theta}} \{\sin^2 \theta + \cos^2 \theta = 1\}$$

$$= \frac{(\sin \theta \cos \theta + 1)(\sin \theta - \cos \theta)(\sin^3 \theta \cos^3 \theta)}{(\sin^3 \theta - \cos^3 \theta) \sin \theta \cos \theta}$$

Since, we know,

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$

$$= \frac{(\sin \theta \cos \theta + 1)(\sin \theta - \cos \theta)(\sin^3 \theta \cos^3 \theta)}{(\sin \theta - \cos \theta)(\sin^2 \theta + \cos^2 \theta + \sin \theta \cos \theta(\sin \theta \cos \theta))}$$

$$= \frac{(\sin \theta \cos \theta + 1)(\sin^2 \theta \cos^2 \theta)}{(1 + \sin \theta \cos \theta)} \{\because \sin^2 \theta + \cos^2 \theta = 1\}$$

$$= \sin^2 \theta \cos^2 \theta = \text{RHS proved}$$

3. Question 3

(i) Given, internal diameter of hollow hemispherical vessel = 7 cm

external diameter of a hollow hemispherical vessel = 14 cm

$$r_1 = \frac{7}{2} \text{ cm}$$

$$r_2 = \frac{14}{2} = 7 \text{ cm}$$

$$\text{Area of Ring} = \pi r_2^2 - \pi r_1^2$$

Total area to be painted

$$= 2\pi r_2^2 + 2\pi r_1^2 + (\pi r_2^2 - \pi r_1^2)$$

$$= 3\pi r_2^2 + \pi r_1^2$$

$$= \pi (3\pi r_2^2 + r_1^2)$$


$$= \pi [3 \times 49 + (3.5)^2] = 3.14 \times (147 + 12.25)$$

$$= 500.045 \text{ cm}^2$$

Hence, the total cost of silver painting the vessel = 500.045×0.6

$$= ₹ 300.027$$

(ii) i. The slope of AD



We know that the slope of any line parallel to x-axis is 0.

\therefore The slope of AD = 0

ii. The slope of BD.

As ABCD in a square

∴ the diagonal BD makes an angle of 45° with +ve direction of x-axis

∴ Slope of BD = $\tan 45^\circ = 1$

iii. The diagonal AC make angle of 135° with positive direction of x-axis.

∴ Slope of AC = $\tan 135^\circ$


$$= \tan (90 \times 2 - 45)$$

$$= \tan (-45)$$

$$= -\tan 45$$

$$= -1$$

(iii)



i. The coordinates of Q are $(-2, -4)$

ii. The coordinates of R are $(-2, 4)$

iii. PQR is Right angle Triangle

iv. Area of $\triangle PQR = \frac{1}{2} \times \text{Base} \times \text{height}$

$$= \frac{1}{2} \times 4 \times 8$$

$$= 2 \times 8$$

$$= 16 \text{ sq. unit}$$

Section B

4. Question 4

(i) i. C.P for the shopkeeper

$$= 1200 \times \frac{90}{100} = ₹1080$$

GST paid by the wholesaler

$$= 1080 \times \frac{60}{100} = ₹64.80$$

ii. S.P of the article = ₹1200

GST paid by the customer

$$= 1200 \times \frac{6}{100} = ₹72$$

Amount paid by the customer

$$= \text{S.P.} + \text{GST} = 1200 + 72 = ₹1272$$

(ii) Let the two consecutive odd no. be x and $x + 2$.

A/c question

$$x^2 + (x + 2)^2 = 290$$

$$\Rightarrow x^2 + x^2 + 4x + 4 = 290$$

$$\Rightarrow 2x^2 + 4x - 286 = 0$$


$$\Rightarrow 2(x^2 + 2x - 143) = 0$$

$$\Rightarrow x^2 + 2x - 143 = 0$$

$$\Rightarrow x^2 + 13x - 11x - 143 = 0$$

$$\Rightarrow x(x + 13) - 11(x + 13) = 0$$

$$x = 11 \text{ or } x = -13 \text{ rejected}$$



In the given histogram, inside the highest rectangle, which represents the maximum frequency.

∴ Modal class = 5000 - 6000

Then, mode = 5500.

5. Question 5

(i)


$$\begin{bmatrix} 4 \sin 30^\circ & 2 \cos 60^\circ \\ \sin 90^\circ & 2 \cos 0^\circ \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 5 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} 4 \times \frac{1}{2} & 2 \times \frac{1}{2} \\ 1 & 2 \times 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 5 & 4 \end{bmatrix} \quad [\because \sin 30^\circ = \cos 60^\circ = \frac{1}{2} \text{ and } \sin 90^\circ = \cos 0^\circ = 1]$$

$$= \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 5 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} 2 \times 4 + 1 \times 5 & 2 \times 5 + 1 \times 4 \\ 1 \times 4 + 2 \times 5 & 1 \times 5 + 2 \times 4 \end{bmatrix}$$

$$= \begin{bmatrix} 8 + 5 & 10 + 4 \\ 4 + 10 & 5 + 8 \end{bmatrix} = \begin{bmatrix} 13 & 14 \\ 14 & 13 \end{bmatrix}.$$



To prove $\angle BOD = \angle A$

In $\triangle BDO$ and $\triangle CDO$

$BD = DC$ (given)

$OD = OD$ (common)

$BO = OC$ (Radius)

By SSS

$\triangle BDO \cong \triangle CDO$

$\angle BOD = \angle COD = x$

Now,

$\angle BAC = \frac{1}{2} \angle BOC$ (angle made by same arc)

$\angle BAC = \frac{1}{2} \times (2x)$

$\angle BAC = x$

$\angle BAC = \angle BOD$

$\angle A = \angle BOD$

Hence proved.

(iii) Let $p(x) = 6x^3 - 17x^2 + 4x - 12$

Remainder $p(-2) = 6(-2)^3 + 17(-2)^2 + 4(-2) - 12$

$$= -48 + 68 - 8 - 12$$

$$= 68 - 68 = 0$$

∴ (x + 2) is a factor of given polynomial p(x)

$$\begin{array}{r} x+2 \overline{) 6x^3 + 17x^2 + 4x - 12} \\ \underline{6x^3 + 12x^2} \\ (-) 5x^2 + 4x - 12 \\ \underline{(-) 5x^2 + 10x} \\ (-) - 6x - 12 \\ \underline{(-) - 6x - 12} \\ (+) 0 \end{array}$$

$$\therefore 6x^3 + 17x^2 + 4x - 12$$

$$= (x + 2)(6x^2 + 5x - 6)$$

$$= (x + 2)\{6x^2 + 9x - 4x - 6\}$$

$$= (x + 2)\{3x(2x + 3) - 2(2x + 3)\}$$

$$= (x + 2)(2x + 3)(3x - 2)$$

6. Question 6

(i) i. By using section formula,

$$\begin{array}{ccc} A(-4, 2) & P(x, 3) & B(3, 6) \\ \leftarrow & \bullet & \rightarrow \end{array}$$

Let the ratio be k : 1

$$y = \frac{m_2 y_1 + m_1 y_2}{m_1 + m_2}$$

$$3 = \frac{1 \times 2 + k \times 6}{k + 1}$$

$$3k + 3 = 2 + 6k$$

$$3k - 6k = 2 - 3$$

$$-3k = -1$$

$$k = \frac{1}{3}$$

Ratio = 1 : 3

$$x = \frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}$$

$$x = \frac{1 \times 3 + 3 \times (-4)}{1 + 3}$$

$$x = \frac{3 - 12}{4}$$

$$x = \frac{-9}{4}$$

ii. Here coordinate of P is $\left(\frac{-9}{4}, 3\right)$

$$\text{Length of, AP} = \sqrt{(y_2 - y_1)^2 + (x_2 - x_1)^2}$$

$$= \sqrt{(3 - 2)^2 + \left(-\frac{9}{4} - (-4)\right)^2}$$

$$= \sqrt{(1)^2 + \left(\frac{-9}{4} + 4\right)^2}$$

$$= \sqrt{(1)^2 + \left(\frac{-9 + 16}{4}\right)^2}$$

$$= \sqrt{(1)^2 + \left(\frac{7}{4}\right)^2}$$

$$= \sqrt{1 + \frac{49}{16}}$$

$$= \sqrt{\frac{65}{16}}$$

$$= \frac{\sqrt{65}}{4}$$

$$(ii) \text{ LHS} = \frac{1}{1} - \frac{\cos^2 \theta}{1 + \sin \theta} = \frac{1 + \sin \theta - \cos^2 \theta}{1 + \sin \theta}$$

$$= \frac{1 + \sin \theta - (1 - \sin^2 \theta)}{1 + \sin \theta}$$

$$= \frac{1 + \sin \theta - 1 + \sin^2 \theta}{1 + \sin \theta}$$

$$= \frac{1 + \sin \theta}{1 + \sin \theta}$$

$$= \sin \theta$$

$$= \text{RHS}$$

Hence Proved

$$(iii) S_n = 5n^2 - 3n$$

$$S_1 = a_1$$

$$= 5(1)^2 - 3(1)$$

$$= 5 - 3$$

$$= 2$$

$$a_1 = 2$$

$$S_2 = a_1 + a_2$$

$$a_1 + a_2 = 5(2)^2 - 3(2)$$

$$= 20 - 6$$

$$= 14$$

$$\therefore a_1 + a_2 = 14$$

$$2 + a_2 = 14$$

$$a_2 = 14 - 2$$

$$a_2 = 12$$

$$\text{Again, } S_3 = a_1 + a_2 + a_3$$

$$a_1 + a_2 + a_3 = 5(3)^2 - 3(3)$$

$$= 45 - 9$$

$$= 36$$

$$a_1 + a_2 + a_3 = 36$$

$$14 + a_3 = 36$$

$$a_3 = 36 - 14$$

$$a_3 = 22$$

$$a_1 = 2, a_2 = 12, a_3 = 22$$

Hence square becomes 2, 12, 22, ...

$$a_1 = a = 2$$

$$d = 12 - 2 = 10$$

$$a_{16} = a + (16 - 1)d$$

$$= 2 + 15 \times 10$$

$$= 2 + 150$$


$$a_{16} = 152$$

7. Question 7

(i) Let the shortest side be x m.

$$\text{hypotenuse} = (2x + 1)\text{m}$$

$$\text{3rd side} = (x + 7)\text{m}$$



$$(2x + 1)^2 = (x + 7)^2 + x^2 \text{ \{by Pythagoras theorem\}}$$

$$\Rightarrow 4x^2 + 1 + 4x = x^2 + 49 + 14x + x^2$$

$$\Rightarrow 2x^2 - 10x - 48 = 0$$

$$\Rightarrow x^2 - 5x - 24 = 0$$

$$\Rightarrow x^2 - 8x + 3x - 24 = 0$$

$$\Rightarrow x(x - 8) + 3(x - 8) = 0$$

$$\Rightarrow (x + 3)(x - 8) = 0$$

$$x = -3, 8$$

$x = -3$ rejected (\because length can never be -ve)

$$\therefore x = 8$$

hypotenuse i.e. AC

$$= 2x + 1$$

$$= 2 \times 8 + 1 = 17$$

$$BC = x = 8\text{m}$$

$$AB = x + 7$$


$$= 8 + 7$$

$$AB = 15\text{ m}$$

(ii) The cumulative frequency table for the given continuous distribution is given below

Marks	Number of students	Cumulative frequency (cf)
0-10	3	3
10-20	7	10
20-30	12	22
30-40	17	39
40-50	23	62
50-60	14	76
60-70	9	85
70-80	6	91
80-90	5	96
90-100	4	100

On the graph paper, we plot the following points A (10, 3), B (20, 10), C (30, 22), D (40, 39), E (50, 62), F (60, 76), G (70, 85), H (80, 91), I (90, 96) and V(100, 100). Join all these points by a free hand drawing. The required ogive is shown on the graph paper given below



Here, number of students (n) = 100, which is even.

i. Let P be the point on Y-axis representing frequency

$$= \frac{n}{2} = \frac{100}{2} = 50$$

Through P, draw a horizontal line to meet the ogive at point Q. Through Q, draw a vertical line to meet the X-axis at

T. The abscissa of the point T represents 43 marks. Hence, the median marks is 43.

ii. Let R be the point on Y-axis representing frequency

$$= \frac{n}{4} = \frac{100}{4} = 25.$$

Through R, draw a horizontal line to meet the ogive at point S. Through S, draw a vertical line to meet the X-axis at N. The abscissa of the point N represents 31 marks. Hence, the lower quartile = 31 marks.

iii. 85% marks = 85% of 100 = 85 marks.

Let the point M on X-axis represents 85 marks. Through M, draw a vertical line to meet the ogive at the point O.

Through O draw a horizontal line to meet the Y-axis at point J. The ordinate of point J represents 95 students.

\therefore Number of students who obtained more than 85% in the test = 100 - 95 = 5

iv. 35% marks = 35% of 100 = 35

Let the point K on X-axis represents 35 marks. Through K, draw a vertical line to meet the ogive at the point L.

Through L, draw a horizontal line to meet the Y-axis at point U. The ordinate of point U represents 30 students on Y-axis. Hence, the number of students, who did not pass in the test is 30.

8. Question 8

(i) $n(s) = 50$

$n(\text{multiple of 3 and 4}) = \{12, 24, 36, 48\}$

i.e multiple of 12

$$(\text{multiple of 3 and 4}) = \frac{4}{50} = \frac{2}{25}$$

(ii) Given, diameter of solid sphere, $d_1 = 6$ cm

$$\therefore \text{Radius of sphere, } r_1 = \frac{6}{2} = 3 \text{ cm}$$

Also, given diameter of cylinder, $d_2 = 4$ cm

$$\therefore \text{Radius of cylinder, } r_2 = \frac{4}{2} = 2 \text{ cm}$$

\therefore Height of cylinder, $h = 45$ cm [given]

Let the required number of spheres be N.

$$\therefore N \times \text{Volume of sphere} = \text{Volume of cylinder}$$


$$\Rightarrow N \times \frac{4}{3} \pi r_1^3 = \pi r_2^2 h$$

$$\Rightarrow N \times \frac{4}{3} \pi \times (3)^3 = \pi \times (2)^2 \times 45$$

$$\therefore N = \frac{2 \times 2 \times 45}{4 \times 3 \times 3} = 5$$

Hence, the required number of solid spheres is 5.

(iii)



In $\triangle ADC$ and $\triangle ABE$

$\angle ACD = \angle AEB$ (angle in the same segment BD)

$AC = AE$ (given)

$\angle A = \angle A$ (common)

$\therefore \triangle ADC \cong \triangle ABE$ (ASA Cong rule)

$\Rightarrow AB = AD$ (CPCT)

But $AC = AE$

$\therefore AC - AB = AE - AD$

$\Rightarrow BC = DE$

In $\triangle BPC$ and $\triangle DPE$

$\angle C = \angle E$ (angle in the same segment)

$BC = DE$

$\angle CBP = \angle CDE$ (angle on the same segment)

$\therefore \triangle BPC \cong \triangle DPE$ (ASA cong rule)

$\Rightarrow BP = DP$ and $CP = PE$ (CPCT)

9. Question 9

(i) $\frac{3x}{5} + 2 < x + 4$

$$\Rightarrow \frac{3x+10}{5} < x + 4$$


$$\Rightarrow 3x + 10 < 5(x + 4)$$

$$\Rightarrow 3x + 10 < 5x + 20$$

$$\Rightarrow 10 - 20 < 5x - 3x$$

$$\Rightarrow -10 < 2x$$

$$\Rightarrow -5 < x$$




And

$$x + 4 \leq \frac{x}{2} + 5$$

$$\Rightarrow x + 4 \leq \frac{x+10}{2}$$

$$\Rightarrow 2x + 8 \leq x + 10$$

$$\Rightarrow x \leq 2$$



Solution Set = $\{x: -5 < x \leq 2, x \in \mathbb{R}\}$

(ii) Given:

mode = 12 k, mean = 15 k, median = ?

Using Empirical relation;

$$\text{Mode} = 3 \text{ median} - 2 \text{ mean}$$

$$12k = 3 \text{ median} - 2(15k)$$

$$3 \text{ median} = 12k + 30k$$

$$3 \text{ median} = 42k$$

$$\text{median} = \frac{42k}{3}$$

$$\text{median} = 14k$$

(iii) Given, $DE \parallel BC$, $DE = 6$ cm and $BC = 12$ cm.

In $\triangle ABC$ and $\triangle ADE$,

$$\angle ABC = \angle ADE \text{ [corresponding angles]}$$

$$\angle ACB = \angle AED \text{ [corresponding angles]}$$

$$\text{and } \angle A = \angle A \text{ [common angle]}$$

$$\therefore \triangle ABC \sim \triangle ADE \text{ [by AAA similarity criterion]}$$

We know that the ratio of areas of two similar triangles is equal to the ratio of squares of their corresponding sides.

$$\therefore \frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle ABC)} = \frac{(DE)^2}{(BC)^2} = \frac{(6)^2}{(12)^2} = \left(\frac{1}{2}\right)^2$$

$$\Rightarrow \frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle ABC)} = \frac{1}{4}$$

$$\text{Let ar}(\triangle ADE) = k, \text{ then ar}(\triangle ABC) = 4k$$

$$\text{Now, ar}(\text{DECB}) = \text{ar}(\triangle ABC) - \text{ar}(\triangle ADE)$$

$$= 4k - k = 3k$$

$$\therefore \text{Required ratio} = \text{ar}(\triangle ADE) : \text{ar}(\text{DECB})$$

$$= k : 3k = 1 : 3$$

10. Question 10

(i) Let fourth proportional be x.

$$\text{Then, } (a^3 + 8) : (a^4 - 2a^3 + 4a^2) :: (a^2 - 4) : x$$

$$\Rightarrow \frac{a^3 + 8}{a^4 - 2a^3 + 4a^2} = \frac{a^2 - 4}{x}$$

$$\Rightarrow x(a^3 + 8) = (a^2 - 4)(a^4 - 2a^3 + 4a^2) \text{ [by cross-multiplication]}$$

$$\therefore x = \frac{(a^2 - 2^2) \times a^2(a^2 - 2a + 4)}{(a^3 + 2^3)}$$

$$= \frac{a^2(a-2)(a+2)(a^2-2a+4)}{(a+2)(a^2-2a+4)} = a^2(a-2)$$

Hence, the required value of fourth proportional is $a^2(a-2)$.

(ii) i. Draw a line $BC = 4.2$ cm.


ii. At B, draw an angle of 60° using compass.

iii. Cut an arc of 5 cm on the angle arm at B and name this point as A.

iv. Draw angle bisector of angle ABC.


v. Draw a line EFIBC at a distance of 2 cm which cuts the angle bisector at O.

vi. Take O as centre and 2 cm as radius, draw a circle which touches both the arms of the angle.



(iii) Let the height of the building be h m and D be the position of a man.

Here, $BC = x$ m



Now, In $\triangle ABC$,

$$\tan 60^\circ = \frac{AB}{BC}$$

$$\sqrt{3} = \frac{h}{x}$$

$$x = \frac{h}{\sqrt{3}} \dots (i)$$

In $\triangle ABD$,

$$\tan 30^\circ = \frac{AB}{BD}$$

$$\frac{1}{\sqrt{3}} = \frac{h}{60+x}$$

$$60 + x = h\sqrt{3}$$

$$\sqrt{3}h = 60 + \frac{h}{\sqrt{3}} \text{ [From eq. (i)]}$$

$$\frac{\sqrt{3}h}{1} - \frac{h}{\sqrt{3}} = 60$$

$$\frac{3h-h}{\sqrt{3}} = 60$$

$$2h = 60\sqrt{3}$$

$$h = \frac{60\sqrt{3}}{2}$$

$$\Rightarrow h = 30\sqrt{3}$$

$$= 30 \times 1.732$$

Height of building = 51.96 m = 52 m (Approx).