

- Please check that this question paper contains 23 printed pages.
- (II) कृपया जाँच कर लें कि इस प्रश्न-पत्र में 33 प्रश्न हैं।

Please check that this question paper contains 33 questions.

- (III) प्रश्न-पत्र में दाहिने हाथ की ओर दिए गए प्रश्न-पत्र कोड को परीक्षार्थी उत्तर-पुस्तिका के मुख-पृष्ठ पर लिखें । Q.P. Code given on the right hand side of the question paper should be written on the title page of the answer-book by the candidate.
- (IV) कृपया प्रश्न का उत्तर लिखना शुरू करने से पहले, उत्तर-पुस्तिका में यथा स्थान पर प्रश्न का क्रमांक अवश्य लिखें।

Please write down the serial number of the question in the answerbook at the given place before attempting it.

(V) इस प्रश्न-पत्र को पढ़ने के लिए 15 मिनट का समय दिया गया है। प्रश्न-पत्र का वितरण पूर्वाह्न में 10.15 बजे किया जाएगा। 10.15 बजे से 10.30 बजे तक परीक्षार्थी केवल प्रश्न-पत्र को पढ़ेंगे और इस अवधि के दौरान वे उत्तर-पुस्तिका पर कोई उत्तर नहीं लिखेंगे।

15 minute time has been allotted to read this question paper. The question paper will be distributed at 10.15 a.m. From 10.15 a.m. to 10.30 a.m., the candidates will read the question paper only and will not write any answer on the answer-book during this period.



रसायन विज्ञान (सैद्धांतिक) CHEMISTRY (Theory)



निर्धारित समय : 3 घण्टे Time allowed : 3 hours







अधिकतम अंक : 70 Maximum Marks : 70 ~~~~~~

P.T.O.

सामान्य निर्देश : निम्नलिखित निर्देशों को ध्यानपूर्वक पढ़िए और उनका पालन कीजिए :

- (i) इस प्रश्न-पत्र में 33 प्रश्न हैं। सभी प्रश्न अनिवार्य हैं।
- (ii) यह प्रश्न-पत्र **पाँच** खण्डों में विभाजित है खण्ड क, ख, ग, घ तथा ङ ।
- (iii) खण्ड क प्रश्न संख्या 1 से 16 तक बहुविकल्पीय प्रकार के प्रश्न हैं । प्रत्येक प्रश्न 1 अंक का है ।
- (iv) खण्ड ख प्रश्न संख्या 17 से 21 तक अति लघु-उत्तरीय प्रकार के प्रश्न हैं । प्रत्येक प्रश्न 2 अंकों का है ।
- (v) खण्ड ग प्रश्न संख्या 22 से 28 तक लघु-उत्तरीय प्रकार के प्रश्न हैं। प्रत्येक प्रश्न 3 अंकों का है।
- (vi) खण्ड घ प्रश्न संख्या 29 तथा 30 प्रकरण आधारित प्रश्न हैं। प्रत्येक प्रश्न 4 अंकों का है।
- (vii) खण्ड ङ प्रश्न संख्या 31 से 33 दीर्घ-उत्तरीय प्रकार के प्रश्न हैं । प्रत्येक प्रश्न 5 अंकों का है ।
- (viii) प्रश्न-पत्र में समग्र विकल्प नहीं दिया गया है। यद्यपि, खण्ड क के अतिरिक्त अन्य सभी खण्डों के कुछ प्रश्नों में आंतरिक विकल्प का चयन दिया गया है।
- (ix) ध्यान दें कि दृष्टिबाधित परीक्षार्थियों के लिए अलग प्रश्न-पत्र है।
- (x) कैल्कुलेटर का उपयोग वर्जित है।

जहाँ आवश्यक हो, आप निम्नलिखित भौतिक नियतांकों के मानों का उपयोग कर सकते हैं :

· 2

$$\begin{split} \mathbf{c} &= 3 \times 10^8 \text{ m/s} \\ \mathbf{h} &= 6.63 \times 10^{-34} \text{ Js} \\ \mathbf{e} &= 1.6 \times 10^{-19} \text{ C} \\ \mu_0 &= 4\pi \times 10^{-7} \text{ T m A}^{-1} \\ \epsilon_0 &= 8.854 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2} \\ \frac{1}{4\pi\epsilon_0} &= 9 \times 10^9 \text{ N m}^2 \text{ C}^{-2} \\ \mathbf{s} \\ \mathbf{c} \\ \mathbf{r} \\ \mathbf{r} \\ \mathbf{z} \\ \mathbf{z} \\ \mathbf{r} \\ \mathbf{r} \\ \mathbf{r} \\ \mathbf{z} \\ \mathbf{z} \\ \mathbf{r} \\ \mathbf{r} \\ \mathbf{r} \\ \mathbf{z} \\ \mathbf{r} \\ \mathbf{r}$$

56/1/1

同時期	

General Instructions :

Read the following instructions carefully and follow them :

- (i) This question paper contains **33** questions. **All** questions are **compulsory**.
- (ii) This question paper is divided into five sections Section A, B, C, D and E.
- (iii) Section A questions number 1 to 16 are multiple choice type questions. Each question carries 1 mark.
- (iv) Section B questions number 17 to 21 are very short answer type questions. Each question carries 2 marks.
- (v) Section C questions number 22 to 28 are short answer type questions. Each question carries 3 marks.
- (vi) Section D questions number 29 and 30 are case-based questions. Each question carries 4 marks.
- (vii) Section E questions number 31 to 33 are long answer type questions. Each question carries 5 marks.
- (viii) There is no overall choice given in the question paper. However, an internal choice has been provided in few questions in all the sections except Section –A.
- (ix) Kindly note that there is a separate question paper for Visually Impaired candidates.

(x) Use of calculator is NOT allowed. You may use the following values of physical constants wherever necessary : $c = 3 \times 10^8 \text{ m/s}$ $h = 6.63 \times 10^{-34} \text{ Js}$ $e = 1.6 \times 10^{-19} \text{ C}$ $\mu_0 = 4\pi \times 10^{-7} \text{ T m A}^{-1}$ $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$ $\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ N m}^2 \text{ C}^{-2}$ Mass of electron (m_e) = 9.1 × 10⁻³¹ kg. Mass of neutron = 1.675×10^{-27} kg. Mass of proton = 1.673×10^{-27} kg. Avogadro's number = 6.023×10^{23} per gram mole Boltzmann's constant = $1.38 \times 10^{-23} \text{ JK}^{-1}$

P.T.O.

:通服:
agente.
日空神感

खण्ड – क

ख्या 1	. से 16 तक बहुविकल्पीय प्रकार के 1 अंक के	प्रश्न हैं	1
संगुणन	। में विलेय का अपसामान्य मोलर द्रव्यमान		
(A)	बढ़ेगा	(B)	घटेगा
(C)	अपरिवर्तित रहेगा	(D)	पहले बढ़ेगा और फिर घटेगा
${ m Sn}^{4+}$ -0.7	-/Sn ²⁺ युग्म के लिए मानक इलेक्ट्रोड विभ 4 V है। दो युग्म अपनी मानक अवस्थाओं में	व +0. संयोजि	15 V और ${ m Cr}^{3+}\!/{ m Cr}$ युग्म के लिए यह त होकर एक सेल का निर्माण करते हैं। सेल
विभव	होगा		
(A)	+1.19 V	(B)	+0.89 V
(C)	+0.18 V	(D)	+1.83 V

चुंबकीय आघूर्ण इसके प्रचक्रणी कोणीय संवेग और कक्षीय कोणीय संवेग से जुड़ा होता है । ${
m Cr}^{3+}$ आयन 3. (परमाणु क्रमांक : Cr = 24) का प्रचक्रण मात्र चुंबकीय आधूर्ण मान _____ है।

(A)	2.87 B.M.	(B)	3.87 B.M.
(C)	3.47 B.M.	(D)	3.57 B.M.

अम्लीय KMnO_4 सल्फाइट को ऑक्सीकृत कर देता है 4.

प्रश्न संख्या 1

1.

2.

- (A) $S_2 O_3^{2-} \dot{H}$ (B) $S_2O_8^{2-}\dot{H}$ (D) SO_4^{2-} \ddot{H} (C) SO₂(g) में
- $[\mathrm{Pt}(\mathrm{NH}_3)_2\mathrm{C}l_2]^{2+}$ का सही IUPAC नाम है 5.
 - (A) डाइऐम्मीनडाइक्लोरिडोप्लैटिनम (II)
 - (B) डाइऐम्मीनडाइक्लोरिडोप्लैटिनम (IV)
 - (C) डाइऐम्मीनडाइक्लोरिडोप्लैटिनम (O)
 - (D) डाइऐम्मीनडाइक्लोरिडोप्लैटिनेट (IV)
- निम्नलिखित यौगिकों को उनके क्वथनांक के बढ़ते क्रम में व्यवस्थित करें : 6.

CH_{3}	CH_3
(i) $CH - CH_2Br$	(ii) $CH_3CH_2CH_2CH_2Br$ (iii) $H_3C - CH_3$
CH ₃ सही कम है	Br
(A) (ii) $<$ (i) $<$ (iii)	(B) (i) < (ii) < (iii)
(C) (iii) $<$ (i) $<$ (ii)	(D) (iii) $<$ (ii) $<$ (i)
56/1/1	~ 4 ~

 $(16 \times 1 = 16)$

SECTION – A

Questions No. 1 to 16 are Multiple Choice type questions carrying 1 mark each.

- 1. In case of association, abnormal molar mass of solute will
 - (A) increase (B) decrease
 - (C) remain same (D) first increase and then decrease
- 2. Standard electrode potential for $\text{Sn}^{4+}/\text{Sn}^{2+}$ couple is +0.15 V and that for the Cr^{3+}/Cr couple is -0.74 V. The two couples in their standard states are connected to make a cell. The cell potential will be

(A)	+1.19 V	(B)	+0.89 V
(C)	+0.18 V	(D)	+1.83 V

3. The magnetic moment is associated with its spin angular momentum and orbital angular momentum. Spin only magnetic moment value of Cr^{3+} ion (Atomic no. : Cr = 24) is _____.

(A)	2.87 B.M.	(B)	3.87 B.M.
(C)	3.47 B.M.	(D)	3.57 B.M.

4. Acidified $KMnO_4$ oxidises sulphite to

(A)
$$S_2O_3^{2-}$$
 (B) $S_2O_8^{2-}$

(C)
$$SO_2(g)$$
 (D) SO_4^2

- 5. The correct IUPAC name of $[Pt(NH_3)_2Cl_2]^{2+}$ is
 - (A) Diamminedichloridoplatinum (II)
 - (B) Diamminedichloridoplatinum (IV)
 - (C) Diamminedichloridoplatinum (O)
 - (D) Diamminedichloridoplatinate (IV)

6. Arrange the following compounds in increasing order of their boiling points :

 $(16 \times 1 = 16)$

- 7. नाभिकरागी द्विआणविक प्रतिस्थापन अभिक्रिया से गुजरने वाले ऐल्किल हैलाइड में सम्मिलित है
 - (A) विन्यास का धारण (B) रेसिमिक मिश्रण का बनना
 - (C) विन्यास का प्रतिलोमन (D) कार्बोधनायन का निर्माण
- 8. निम्नलिखित में अम्ल सामर्थ्य का सही क्रम कौन सा है ?
 - (A) $C_6H_5OH > H_2O > ROH$ (B) $C_6H_5OH > ROH > H_2O$
 - (C) $\operatorname{ROH} > \operatorname{C_6H_5OH} > \operatorname{H_2O}$ (D) $\operatorname{H_2O} > \operatorname{C_6H_5OH} > \operatorname{ROH}$
- 9. प्रोपिल मैग्नीशियम ब्रोमाइड को ${
 m CO}_2$ के साथ अभिक्रियित करने के पश्चात् अम्ल जलअपघटन करने पर बनने वाला अम्ल है
 - (A) C_3H_7COOH (B) C_2H_5COOH
 - (C) CH_3COOH (D) C_3H_7OH
- 10. प्रोपेनेमाइड को प्रोपेनेमीन में परिवर्तित करने के लिए सबसे उत्तम अभिकर्मक ______ है।
 - (A) H₂ का आधिक्य
 - (B) जलीय NaOH में ${\rm Br}_2$
 - (C) लाल फॉस्फोरस की उपस्थिति में आयोडीन
 - (D) ईथर में $LiAlH_4$
- 11. निम्नलिखित में से कौन सा कथन ग्लूकोज़ के विषय में सत्य नहीं है ?
 - (A) यह एक एल्डोहेक्सोस है।
 - (B) HI के साथ गर्म करने पर यह n-हेक्सेन बनाता है।
 - (C) यह फ्यूरेनोस रूप में उपस्थित होता है।
 - (D) यह शिफ-परीक्षण नहीं देता है।
- 12. अचार बनाने के लिए नमक के सांद्र विलयन में रखा गया कच्चा आम सिकुड़ जाता है, क्योंकि _____
 - (A) यह परासरण के कारण जल प्राप्त करता है।
 - (B) यह उत्क्रम परासरण के कारण जल खो देता है।
 - (C) यह उत्क्रम परासरण के कारण जल प्राप्त करता है।
 - (D) यह परासरण के कारण जल खो देता है।

56/1/1



- 7. Alkyl halides undergoing nucleophilic bimolecular substitution reaction involve
 - (A) retention of configuration
 - (C) inversion of configuration
- (B) formation of racemic mixture
- (D) formation of carbocation
- 8. Which is the correct order of acid strength from the following ?
 - (A) $C_6H_5OH > H_2O > ROH$ (B) $C_6H_5OH > ROH > H_2O$
 - (C) $\operatorname{ROH} > \operatorname{C}_6\operatorname{H}_5\operatorname{OH} > \operatorname{H}_2\operatorname{O}$ (D) $\operatorname{H}_2\operatorname{O} > \operatorname{C}_6\operatorname{H}_5\operatorname{OH} > \operatorname{ROH}$
- 9. The acid formed when propyl magnesium bromide is treated with CO_2 followed by acid hydrolysis is :
 - (A) C_3H_7COOH (B) C_2H_5COOH (C) CH_3COOH (D) C_3H_7OH
- 10. The best reagent for converting propanamide into propanamine is
 - (A) excess H_2
 - (B) Br_2 in aqueous NaOH
 - (C) iodine in the presence of red phosphorus
 - (D) $LiAlH_4$ in ether
- 11. Which of the following statements is not true about glucose ?
 - (A) It is an aldohexose.
 - (B) On heating with HI it forms n-hexane.
 - (C) It exists in furanose form.
 - (D) It does not give Schiff's test.
- 12. An unripe mango placed in a concentrated salt solution to prepare pickle, shrivels because _____.
 - (A) it gains water due to osmosis
 - (B) it loses water due to reverse osmosis
 - (C) it gains water due to reverse osmosis
 - (D) it loses water due to osmosis
 - ~ 7 ~

56/1/1



प्रश्न संख्या 13 से 16 के लिए, दो कथन दिए गए हैं – जिनमें एक को अभिकथन (A) तथा दूसरे को कारण (R) द्वारा अंकित किया गया है। इन प्रश्नों के सही उत्तर नीचे दिए गए कोडों (A), (B), (C) और (D) में से चुनकर दीजिए :

- (A) अभिकथन (A) और कारण (R) दोनों सही हैं और कारण (R), अभिकथन (A) की सही व्याख्या करता है।
- (B) अभिकथन (A) और कारण (R) दोनों सही हैं, परन्तु कारण (R), अभिकथन (A) की सही व्याख्या नहीं करता है।
- (C) अभिकथन (A) सही है, परन्तु कारण (R) गलत है।
- (D) अभिकथन (A) गलत है, परन्तु कारण (R) सही है।
- 13. अभिकथन (A) : [Cr(H₂O)₆]Cl₂ तथा [Fe(H₂O)₆]Cl₂ होमोलेप्टिक संकुल के उदाहरण हैं ।
 कारण (R) : धातु के साथ जुड़े सभी लिगैन्ड एक प्रकार के हैं ।
- 14. अभिकथन (A) : ऐल्किल हैलाइडों के क्वथनांक निम्न क्रम में घटते हैं : RI > RBr > RCl > RF ।
 - **कारण (R)** : ऐल्किल क्लोराइड, ब्रोमाइड और आयोडाइड के क्वथनांक तुलनीय आण्विक द्रव्यमान वाले हाइड्रोकार्बन की तुलना में काफी अधिक होते हैं।
- 15. अभिकथन (A) : एथेनॉल का क्वथनांक मेथॉक्सीमेथेन से अधिक होता है।
 - कारण (R) : एथेनॉल में अंतः अणुक हाइड्रोजन बंध होता है।
- 16. अभिकथन (A) : ग्लाइसीन के अतिरिक्त सभी प्राकृतिक रूप से पाए जाने वाले α-ऐमीनो अम्ल ध्रुवण घूर्णक होते हैं।
 - **कारण (R) :** अधिकांश प्राकृतिक रूप से पाए जाने वाले ऐमीनो अम्लों में L-विन्यास होता है।

56/1/1 ~ 8



For questions number 13 to 16, two statements are given – one labelled as Assertion (A) and the other labelled as Reason (R). Select the correct answer to these questions from the codes (A), (B), (C) and (D) as given below :

- (A) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A).
- (B) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).
- (C) Assertion (A) is true, but Reason (R) is false.
- (D) Assertion (A) is false, but Reason (R) is true.
- 13. Assertion (A) : $[Cr(H_2O)_6]Cl_2$ and $[Fe(H_2O)_6]Cl_2$ are examples of homoleptic complexes.

Reason (R) : All the ligands attached to the metal are the same.

- 14. Assertion (A) : The boiling points of alkyl halides decrease in the order : RI > RBr > RCl > RF.
 - Reason (R) : The boiling points of alkyl chlorides, bromides and iodides are considerably higher than that of the hydrocarbon of comparable molecular mass.
- 15. Assertion (A) : The boiling point of ethanol is higher than that of methoxymethane.
 - **Reason (R)** : There is intramolecular hydrogen bonding in ethanol.
- 16. Assertion (A) : All naturally occurring α -amino acids except glycine are optically active.
 - **Reason (R)** : Most naturally occurring amino acids have L-configuration.

P.T.O.



(A) कारण बताइए : 17. (1 + 1 = 2)प्रेशर-कुकर में खाना पकाना, खुले बर्तन (कड़ाही) की तुलना में तेज़ होता है। (a) द्रव X और द्रव Y को मिलाने पर, परिणामी विलयन का आयतन कम हो जाता है। (b) परिणामी विलयन राउल्ट के नियम से किस प्रकार का विचलन दर्शाता है ? द्रव X और Y को मिलाने के बाद आप ताप में क्या परिवर्तन प्रेक्षित करेंगे ? अथवा (B) स्थिरक्वाथी को परिभाषित करें । राउल्ट के नियम में ऋणात्मक विचलन से किस प्रकार का स्थिरक्वाथी बनता है ? एक उदाहरण दीजिए। 2 18. निम्नलिखित रासायनिक समीकरण को पूर्ण एवं संतुलित कीजिए : 2 $2MnO_{4}(aq) + 10I(aq) + 16H(aq) \longrightarrow$ (a) $\operatorname{Cr}_2O_7^{2-}(\operatorname{aq}) + 6\operatorname{Fe}^{2+}(\operatorname{aq}) + 14\operatorname{H}^+(\operatorname{aq}) \longrightarrow$ (b) क्या आप अपेक्षा करते हैं कि प्रोपेनल की तुलना में बेंजैल्डिहाइड नाभिकरागी योगज अभिक्रियाओं में 19. अधिक अभिक्रियाशील या कम अभिक्रियाशील होगा ? अपना उत्तर स्पष्ट कीजिए। 2 निम्नलिखित प्रत्येक अनुक्रम अभिक्रिया में A और B की पहचान कीजिए : 20.(1 + 1 = 2) $CH_3CH_2Cl \xrightarrow{NaCN} A \xrightarrow{H_2/Ni} B$ (a) (b) $C_6H_5NH_2 \xrightarrow{NaNO_2/HCl} A \xrightarrow{C_6H_5NH_2} B$ D-ग्लूकोज़ को निम्नलिखित अभिकर्मकों के साथ अभिक्रियित करने पर होने वाली अभिक्रियाएँ लिखिए : 21.(1 + 1 = 2)(a) **HCN**

~~~~~~~~~~~~~~~~~

 $\operatorname{Br}_2$  जल

(b)



# SECTION – B

# 17. (A) Give reasons :

- (a) Cooking is faster in pressure cooker than in an open pan.
- (b) On mixing liquid X and liquid Y, volume of the resulting solution decreases. What type of deviation from Raoult's law is shown by the resulting solution ? What change in temperature would you observe after mixing liquids X and Y ?

#### OR

- (B) Define Azeotrope. What type of Azeotrope is formed by negative deviation from Raoult's law ? Give an example.2
- 18. Complete and balance the following chemical equations :
  - (a)  $2MnO_4(aq) + 10I(aq) + 16H(aq) \longrightarrow$
  - (b)  $\operatorname{Cr}_2O_7^{2-}(\operatorname{aq}) + 6\operatorname{Fe}^{2+}(\operatorname{aq}) + 14\operatorname{H}^+(\operatorname{aq}) \longrightarrow$
- 19. Would you expect benzaldehyde to be more reactive or less reactive in nucleophilic addition reactions than propanal ? Justify your answer.2
- 20. Identify A and B in each of the following reaction sequence : (1 + 1 = 2)

(a) 
$$CH_3CH_2Cl \xrightarrow{NaCN} A \xrightarrow{H_2/Ni} B$$
  
(b)  $C_6H_5NH_2 \xrightarrow{NaNO_2/HCl} A \xrightarrow{C_6H_5NH_2} H^+ B$ 

- 21. Write the reactions involved when D-glucose is treated with the following reagents : (1 + 1 = 2)
  - (a) HCN
  - (b) Br<sub>2</sub> water

(1 + 1 = 2)

 $\mathbf{2}$ 

खण्ड – ग

- 22. जल में ग्लूकोस (मोलर द्रव्यमान = 180 g mol<sup>-1</sup>) के विलयन का क्वथनांक 100.20 °C है । उसी विलयन के हिमांक की गणना करें । पानी के लिए, K<sub>f</sub> और K<sub>b</sub> मोलल स्थिरांक क्रमशः 1.86 K kg mol<sup>-1</sup> और 0.512 K kg mol<sup>-1</sup> हैं ।
- 23. (a) निम्न को समझाइए :
  - (i) आयनों के स्वतंत्र अभिगमन का कोलराऊश नियम
  - (ii) फैराडे के विद्युत्-अपघटन का प्रथम नियम
  - (b) नीचे दिए गए E° मानों का उपयोग करते हुए, प्रागुक्ति कीजिए कि संक्षारण रोकने के लिए लोहे की सतह पर लेपन के लिए कौन–सा बेहतर है और क्यों।

दिया गया है :  $E_{X^{2+}/X}^{\circ} = -2.36 \text{ V},$  $E_{Y^{2+}/Y}^{\circ} = -0.14 \text{ V},$  $E_{Fe^{2+}/Fe}^{\circ} = -0.44 \text{ V}$ 

24. कोई अभिक्रिया 300 K पर 20 मिनट में 50% पूर्ण हो जाती है और वही अभिक्रिया 350 K पर 5 मिनट में 50% पूर्ण हो जाती है। यदि यह अभिक्रिया प्रथम कोटि की है तो सक्रियण ऊर्जा की गणना कीजिए।

 $[R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}; \log 4 = 0.602]$ 

- 25. 3d संक्रमण श्रेणी के तत्व इस प्रकार दिए गए हैं : Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn निम्नलिखित का उत्तर दीजिए :
  - (a) कॉपर का  $E^{\circ}_{M^{2+}/M}$  मान असाधारण रूप से धनात्मक है, क्यों ?
  - (b) कौन सा तत्व +2 ऑक्सीकरण अवस्था में एक प्रबल अपचायक है और क्यों ?
  - (c) Zn<sup>2+</sup> लवण रंगहीन होते हैं, क्यों ?

3

(1 + 1 + 1 = 3)

~~~~~~~~~~~~~~~

3



SECTION – C

- 22. A solution of glucose (molar mass = 180 g mol⁻¹) in water has a boiling point of 100.20 °C. Calculate the freezing point of the same solution. Molal constants for water K_f and K_b are 1.86 K kg mol⁻¹ and 0.512 K kg mol⁻¹ respectively.
- 23. (a) State the following :
 - (i) Kohlrausch law of independent migration of ions and
 - (ii) Faraday's first law of electrolysis.
 - (b) Using E_{values}° of X and Y given below, predict which is better for coating the surface of iron to prevent corrosion and why?

Given $E_{X^{2+}/X}^{\circ} = -2.36 \text{ V},$ $E_{Y^{2+}/Y}^{\circ} = -0.14 \text{ V},$ $E_{Fe^{2+}/Fe}^{\circ} = -0.44 \text{ V}$

- 24. A certain reaction is 50% complete in 20 minutes at 300 K and the same reaction is 50% complete in 5 minutes at 350 K. Calculate the activation energy if it is a first order reaction.
 3 [R = 8.314 J K⁻¹ mol⁻¹; log 4 = 0.602]
- 25. The elements of 3d transition series are given as : (1 + 1 + 1 = 3)
 Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn
 Answer the following :
 - (a) Copper has exceptionally positive $E^{\circ}_{M^{2^+}/M}$ value, why ?
 - (b) Which element is a strong reducing agent in +2 oxidation state and why?
 - (c) Zn^{2+} salts are colourless. Why ?

56/1/1



3





27.

28.



26.Draw the structure of the major monohalo product for each of the (A) following reaction : (1+1+1=3)



How do you convert : (B)

 $(3 \times 1 = 3)$

- Chlorobenzene to biphenyl (a)
- (b) Propene to 1-Iodopropane
- (c) 2-bromobutane to but-2-ene.
- 27.Arrange the following compounds in increasing order of their boiling (a) point : (1+2=3)(CH₃)₂NH, CH₃CH₂NH₂, CH₃CH₂OH.
 - Give plausible explanation for each of the following: (b)
 - Aromatic primary amines cannot be prepared by Gabriel (i) Phthalimide synthesis.
 - (ii) Amides are less basic than amines.
- 28.What is the difference between native protein and denatured protein ? (a)

(1+1+1=3)

P.T.O.

- Which one of the following is a disaccharide? (b) Glucose, Lactose, Amylose, Fructose
- Which vitamin is responsible for the coagulation of blood ? (c)



खण्ड – ঘ

निम्नलिखित प्रश्न प्रकरण आधारित प्रश्न है । परिच्छेद को ध्यानपूर्वक पढ़िए और दिए गए प्रश्नों के उत्तर दीजिए ।

29. रासायनिक अभिक्रिया का वेग या तो प्रति इकाई समय में अभिकारक की सांद्रता में कमी या उत्पाद की सांद्रता में वृद्धि के रूप में व्यक्त किया जाता है । अभिक्रिया का वेग अभिकारकों की प्रकृति, अभिकारकों की सांद्रता, तापमान, उत्प्रेरक की उपस्थिति, अभिकारकों के पृष्ठीय क्षेत्रफल और प्रकाश की उपस्थिति पर निर्भर करता है । अभिक्रिया का वेग अभिकारकों के पृष्ठीय क्षेत्रफल और प्रकाश की उपस्थिति पर निर्भर करता है । अभिक्रिया का वेग अभिकारक की सांद्रता से सीधे संबंधित होता है । वेग नियम बताता है कि अभिक्रिया का वेग सांद्रता पदों पर निर्भर करता है जिस पर अभिक्रिया का वेग वास्तव में निर्भर करता है, जैसा कि प्रयोगात्मक रूप से देखा गया है । वेग नियम अभिव्यक्ति में अभिकारकों की सांद्रता की घातों के योग को अभिक्रिया की कोटि कहा जाता है जबकि एक प्राथमिक अभिक्रिया में भाग लेने वाली स्पीशीज़ की संख्या जो रासायनिक अभिक्रिया सम्पन्न करने के लिए एक साथ संघट्ट करती है, अभिक्रिया की आण्विकता कहलाती है ।

निम्न प्रश्नों के उत्तर दीजिए :

(a) (i) वेग निर्धारक पद क्या है ? (1 + 1 = 2)

1

- (ii) जटिल अभिक्रिया को परिभाषित कीजिए।
- (b) वेग स्थिरांक पर ताप का क्या प्रभाव पड़ता है ?

अथवा

- (b) आण्विकता केवल प्राथमिक अभिक्रियाओं के लिए ही क्यों लागू होती है जबकि कोटि प्राथमिक और जटिल अभिक्रिया दोनों के लिए लागू होती हैं ?
- (c) अणु X का Y में रूपांतरण द्वितीय कोटि की बलगतिकी के अनुरूप होता है । यदि X की सांद्रता तीन गुनी कर दी जाए तो Y के निर्माण होने के वेग पर क्या प्रभाव पड़ेगा ?



SECTION – D

The following questions are case based questions. Read the passage carefully and answer the questions that follow.

29. The rate of a chemical reaction is expressed either in terms of decrease in the concentration of reactants or increase in the concentration of a product per unit time. Rate of the reaction depends upon the nature of reactants, concentration of reactants, temperature, presence of catalyst, surface area of the reactants and presence of light. Rate of reaction is directly related to the concentration of reactant. Rate law states that the rate of reaction depends upon the concentration terms on which the rate of reaction actually depends, as observed experimentally. The sum of powers of the concentration of the reactants in the Rate law expression is called order of reaction while the number of reacting species taking part in an elementary reaction which must collide simultaneously in order to bring about a chemical reaction is called molecularity of the reaction.

Answer the following questions :

- (a) (i) What is a rate determining step ? (1 + 1 = 2)
 - (ii) Define complex reaction.
- (b) What is the effect of temperature on the rate constant of a reaction ? 1

OR

- (b) Why is molecularity applicable only for elementary reactions whereas order is applicable for elementary as well as complex reactions?
- (c) The conversion of molecule X to Y follows second order kinetics. If concentration of X is increased 3 times, how will it affect the rate of formation of Y ?

P.T.O.

^^^^^

1



30. बेन्जीन वलय से जुड़े OH समूह के प्रबल सक्रियण प्रभाव के कारण फ़ीनॉल आसानी से इलेक्ट्रॉनरागी प्रतिस्थापन अभिक्रियाएँ देते हैं । चूँकि OH समूह o– और p– स्थितियों पर इलेक्ट्रॉन घनत्व को अधिक बढ़ाता है, इसलिए OH समूह ऑर्थो और पैरा निर्देशक होता है । राइमर–टीमन इलेक्ट्रॉनरागी प्रतिस्थापन अभिक्रिया के उदाहरण में से एक है । जिसके द्वारा फ़ीनॉल की एरोमैटिक वलय में हाइड्रॉक्सिल समूह की ऑर्थो स्थिति पर एल्डिहाइड समूह प्रवेश कर जाता है । यह एक सामान्य विधि है जो फ़ीनॉलों के ऑर्थो फार्मिलन के लिए प्रयुक्त की जाती है ।

निम्नलिखित प्रश्नों के उत्तर दीजिए :

| (a) | क्या होता है जब फ़ीनॉल निम्न से अभिक्रिया करता है ? | 2 |
|-----|---|---|
|-----|---|---|

- (i) Br_2/CS_2
- (ii) सांद्र HNO_3
- (b) फ़ीनॉल का प्रोटॉनन आसानी से क्यों नहीं होता है ? 1
- (c) कौन सा एक प्रबल अम्ल है : फ़ीनॉल अथवा क्रीसॉल ? कारण दीजिए। 1

अथवा

(c) राइमर-टीमन अभिक्रिया में बनने वाले उत्पाद का IUPAC नाम लिखें। 1

खण्ड – ङ

31. (A) (a) निम्नलिखित के लिए सेल अभिक्रिया लिखिए तथा 298 K पर सेल का e.m.f.
 परिकलित कीजिए : (3 + 2 = 5)

 $Sn(s) | Sn^{2+} (0.004 \text{ M}) | | H^{+} (0.02 \text{ M}) | H_{2}(g) (1 \text{ Bar}) | Pt (s)$

(दिया गया है : $E_{Sn^{2+}/Sn}^{\circ} = -0.14 \text{ V}, E_{H+|H_2(g), Pt}^{\circ} = 0.00 \text{V}$)

56/1/1 ~ 18 ~



30. Phenols undergo electrophilic substitution reactions readily due to the strong activating effect of OH group attached to the benzene ring. Since, the OH group increases the electron density more to o- and p- positions therefore OH group is ortho, para-directing. Reimer-Tiemann reaction is one of the examples of aldehyde group being introduced on the aromatic ring of phenol, ortho to the hydroxyl group. This is a general method used for the ortho-formylation of phenols.

Answer the following questions :

| (a) | What happens when phenol reacts with | 2 |
|-----|--------------------------------------|---|
|-----|--------------------------------------|---|

- (i) Br_2/CS_2
- (ii) Conc. HNO₃
- (b) Why phenol does not undergo protonation readily ? 1
- (c) Which is a stronger acid phenol or cresol ? Give reason. 1

OR

(c) Write the IUPAC name of the product formed in the Reimer-Tiemann reaction.

SECTION – E

31. (A) (a) Write the cell reaction and calculate the e.m.f. of the following cell at 298 K: (3 + 2 = 5)

 $Sn(s)\,|\,Sn^{2+}\,(0.004~M)\,\big|\,\big|\,H^{+}\,(0.02~M)\,|\,H_{2}(g)\,(1~Bar)\,|\,Pt\,(s)$

(Given : $E_{Sn^{2+}/Sn}^{\circ} = -0.14 \text{ V}, E_{H+|H_2(g), Pt}^{\circ} = 0.00 \text{V}$)

| 通過 |
|-----------------|
| 部の語 |
| 田 王 李秋 秋 |

- (b) निम्नलिखित के कारण बताइए :
 - (i) E° मानों के आधार पर, O_2 गैस एनोड पर मुक्त होनी चाहिए, लेकिन जलीय NaCl के विद्युत अपघटन में Cl_2 गैस मुक्त होती है ।
 - (ii) तनुकरण पर CH₃COOH की चालकता कम हो जाती है।

अथवा

- 31. (B) (a) लेड स्टोरेज बैटरी के उपयोग के दौरान होने वाली ऐनोड और कैथोड अभिक्रियाएँ और समग्र
 सेल अभिक्रिया लिखें ।
 (2 + 3 = 5)
 - (b) 0.01 M K₂Cr₂O₇(aq), 0.01 M Cr³⁺(aq) और 1.0 × 10⁻⁴ M H⁺(aq) युक्त अर्ध सेल के लिए विभव की गणना करें । अर्ध-सेल अभिक्रिया है : Cr₂O₇²⁻(aq) + 14H⁺(aq) + 6e⁻ → 2Cr³⁺(aq) + 7H₂O(*l*) और मानक इलेक्ट्रोड विभव E[°] = 1.33 V दिया गया है ।

[दिया गया है : $\log\,10$ = 1]

- 32. (A) निम्नलिखित के उत्तर दीजिए :
 - (a) निम्न प्रचक्रण चतुष्फलकीय संकुल क्यों नहीं बनते ?
 - (b) प्रबल लिगैंड की उपस्थिति में Co²⁺ आसानी से Co³⁺ में ऑक्सीकृत हो जाता है। [परमाणु संख्या Co = 27]
 - (c) $[Co(NH_3)_6] [Cr(CN)_6]$ संकुल द्वारा किस प्रकार की समावयवता दर्शाई जाती है?
 - (d) $[Ni(H_2O)_6]^{2+}$ का विलयन हरा क्यों होता है जबकि $[Ni(CN)_4]^{2-}$ का विलयन रंगहीन होता है ? (Ni की परमाणु संख्या = 28)
 - (e) निम्नलिखित संकुल का IUPAC नाम लिखिए : $[Co(NH_3)_5(CO_3)]Cl$

अथवा



5



- (b) Account for the following ;
 - On the basis of E° values, O_2 gas should be liberated at (i) anode but it is Cl_2 gas which is liberated in the electrolysis of aqueous NaCl.
 - Conductivity of CH₃COOH decreases on dilution. (ii)

OR

- 31. (B) Write the anode and cathode reactions and the overall cell (a) reaction occurring in a lead storage battery during its use. (2 + 3 = 5)
 - Calculate the potential for half-cell containing 0.01 M (b) $\rm K_2Cr_2O_7(aq),~0.01~M~Cr^{3+}$ (aq) and $\rm 1.0\times10^{-4}~M~H^+(aq).$

The half cell reaction is

$$\mathrm{Cr}_{2}\mathrm{O}_{7}^{2-}(\mathrm{aq}) + 14\mathrm{H}^{+}(\mathrm{aq}) + 6\mathrm{e}^{-} \longrightarrow 2\mathrm{Cr}^{3+}(\mathrm{aq}) + 7\mathrm{H}_{2}\mathrm{O}(l)$$

and the standard electrode potential is given as $E^{\circ} = 1.33 \text{ V}$. [Given : $\log 10 = 1$]

5

32.(A) Answer the following :

56/1/1

- (a) Low spin tetrahedral complexes are not known.
- Co^{2+} is easily oxidised to Co^{3+} in the presence of a strong ligand (b) [At. No. of Co = 27]
- (c) What type of isomerism is shown by the complex $[Co(NH_3)_6]$ $[Cr(CN)_6]$?
- Why a solution of $[Ni(H_2O)_6]^{2+}$ is green while a solution of (d) $[Ni(CN)_4]^{2-}$ is colourless. (At. No. of Ni = 28)
- Write the IUPAC name of the following complex : $[Co(NH_3)_5(CO_3)]Cl$ (e)

| | OR |
|---|----------------|
| ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~ | |
| P.T.O. | $\sim 21 \sim$ |
| | |



32. (B) (a) 'कीलेट प्रभाव' से क्या तात्पर्य है ? एक उदाहरण दीजिए। (2 + 2 + 1 = 5)

- (b) [Fe(CN)₆]^{4–} का संकरण एवं चुम्बकीय व्यवहार लिखिए ।
 (परमाणु संख्या : Fe = 26)
- (c) यदि $PtCl_2 \cdot 2NH_3$; $AgNO_3$ के साथ अभिक्रिया नहीं करता है, तो इसका सूत्र क्या होगा ?
- 33. (A) (a) निम्नलिखित रूपांतरण करें : (2 + 3 = 5)
 - (i) एथेनेल से ब्यूट-2-ईनैल
 - (ii) प्रोपेनोइक अम्ल से एथेन
 - (b) आण्विक सूत्र C₅H₁₀ वाला एक एल्कीन A ओज़ोनी अपघटन पर दो यौगिकों B और C का मिश्रण देता है । यौगिक B धनात्मक फेलिंग परीक्षण देता है और आयोडीन और NaOH विलयन के साथ भी अभिक्रिया करता है । यौगिक C फेलिंग परीक्षण नहीं देता है लेकिन आयोडोफार्म बनाता है । यौगिक A, B और C की पहचान कीजिए ।

अथवा

33. (B) एक कार्बनिक यौगिक (A) (आण्विक सूत्र C₈H₁₆O₂) को तनु सल्फ्यूरिक अम्ल के साथ जल अपघटन करके कार्बोक्सिलिक अम्ल (B) और एल्कोहॉल (C) प्राप्त हुआ । (C) का क्रोमिक अम्ल के साथ ऑक्सीकरण करने पर (B) बनता है । निर्जलीकरण पर (C) ब्यूट-1-ईन देता है । (A), (B) और (C) को पहचानिए तथा सम्मिलिति अभिक्रियाओं के लिए रासायनिक समीकरण लिखिए ।

5

^^^^^

56/1/1



- 32. (B) (a) What is meant by 'Chelate effect' ? Give an example. (2 + 2 + 1 = 5)
 - (b) Write the hybridization and magnetic behaviour of $[Fe(CN)_6]^{4-}$. (Atomic number : Fe = 26)
 - (c) If $PtCl_2 \cdot 2NH_3$ does not react with $AgNO_3$, what will be its formula ?

33. (A) (a) Carry out the following conversions : (2 + 3 = 5)

- (i) Ethanal to But-2-enal
- (ii) Propanoic acid to ethane
- (b) An alkene A with molecular formula C_5H_{10} on ozonolysis gives a mixture of two compounds B and C. Compound B gives positive Fehling test and also reacts with iodine and NaOH solution. Compound C does not give Fehling solution test but forms iodoform. Identify the compounds A, B and C.

OR

33. (B) An organic compound (A) (molecular formula $C_8H_{16}O_2$) was hydrolysed with dilute sulphuric acid to get a carboxylic acid (B) and an alcohol (C). Oxidation of (C) with chromic acid produced (B). (C) on dehydration gives But-l-ene. Identify (A), (B) and (C) and write chemical equations for the reactions involved.

 $\mathbf{5}$



56/1/1

730-1

Marking Scheme Strictly Confidential (For Internal and Restricted use only) Senior School Certificate Examination, 2024-25 SUBJECT NAME CHEMISTRY (Theory) -043 (Q.P.CODE 56/1/1)- MM: 70

General Instructions: -

You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.

"Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its' leakage to public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in News Paper/Website etc may invite action under various rules of the Board and IPC."

Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one's own interpretation or any other consideration. Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-X, while evaluating two competency-based questions, please try to understand given answer and even if reply is not from marking scheme but correct competency is enumerated by the candidate, due marks should be awarded.

The Marking scheme carries only suggested value points for the answers

These are in the nature of Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly.

The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after delibration and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.

Evaluators will mark($\sqrt{}$) wherever answer is correct. For wrong answer CROSS 'X" be marked. Evaluators will not put right (\checkmark) while evaluating which gives an impression that answer is correct and no marks are awarded. This is most common mistake which evaluators are committing.

If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly.

If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.

If a student has attempted an extra question, answer of the question deserving more marks should be retained and the other answer scored out with a note "**Extra Question**".

No marks to be deducted for the cumulative effect of an error. It should be penalized only once.

A full scale of marks ______(example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it.

Every examiner has to necessarily do evaluation work for full working hours i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper.

Ensure that you do not make the following common types of errors committed by the Examiner in the past:-

- Leaving answer or part thereof unassessed in an answer book.
- Giving more marks for an answer than assigned to it.
- Wrong totaling of marks awarded on an answer.
- Wrong transfer of marks from the inside pages of the answer book to the title page.
- Wrong question wise totaling on the title page.
- Wrong totaling of marks of the two columns on the title page.
- Wrong grand total.
- Marks in words and figures not tallying/not same.
- Wrong transfer of marks from the answer book to online award list.
- Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.)

Half or a part of answer marked correct and the rest as wrong, but no marks awarded.

While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0)Marks.

Any unassessed portion, non-carrying over of marks to the title page, or totaling error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, in order to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.

The Examiners should acquaint themselves with the guidelines given in the "**Guidelines for Spot Evaluation**" before starting the actual evaluation.

Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words.

The candidates are entitled to obtain photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme.

MARKING SCHEME 2024-25

CHEMISTRY (Theory)- 043

QP CODE 56/1/1

MM: 70

| Q. No | Value points | Mark |
|-------|--|-------|
| | SECTION A | |
| 1 | (A) | 1 |
| 2 | (B) | 1 |
| 3 | (B) | 1 |
| 4 | (D) | 1 |
| 5 | (B) | 1 |
| 6 | (C) | 1 |
| 7 | (C) | 1 |
| 8 | (A) | 1 |
| 9 | (A) | 1 |
| 10 | (D) | 1 |
| 11 | (C) | 1 |
| 12 | (D) | 1 |
| 13 | (A) | 1 |
| 14 | (B) | 1 |
| 15 | (C) | 1 |
| 16 | (B) | 1 |
| | SECTION B | |
| 17 | (A) (a) Due to high pressure inside the pressure cooker, higher is the boiling point and | 1 |
| | faster is the cooking. | |
| | (b) | |
| | Negative deviation | 1/2 |
| | Temperature increases. | 1/2 |
| | OR | |
| 17 | (B) | |
| | Same composition in liquid and in vapour phase and boil at a constant temperature. | 1 |
| | Maximum Boiling Azeotrope | 1/2 |
| | 68% HNO ₃ + 32% H ₂ O (Or any other correct example) (Percentage can be ignored) | 1/2 |
| 18 | (a) $10I^{-} + 2MnO_{4}^{-} + 16H^{+} \rightarrow 2Mn^{2+} + 8H_{2}O + 5I_{2}$ | 1 |
| | (b) $\operatorname{Cr}_2\operatorname{O}_7^{2-} + 14 \operatorname{H}^+ + 6 \operatorname{Fe}^{2+} \to 2 \operatorname{Cr}^{3+} + 6 \operatorname{Fe}^{3+} + 7 \operatorname{H}_2\operatorname{O}$ | 1 |
| 19 | Less reactive, | 1 |
| | • The carbon atom of the carbonyl group of benzaldehyde is less electrophilic than carbon | |
| | atom of the carbonyl group present in propanal. / The polarity of the carbonyl group is | 1 |
| | reduced in benzaldehyde due to resonance. | |
| 20 | (a) $A = CH_3CH_2CN$; $B = CH_3CH_2 CH_2NH_2$ | ½ x 4 |
| | (b) $A = C_6 H_5 N_2^+ C I^-$; | |
| | | |

| 21 | (a) CHO HC CN | 1 |
|----|--|-----------|
| | | |
| | $(CHOH)_4 \longrightarrow (CHOH)_4$ | |
| | сн ₂ он сн ₂ он | |
| | | |
| | (b) CHO COOH | |
| | $ Br_2/H_2O $ | |
| | $(CHOH)_4 \longrightarrow (CHOH)_4$ | 1 |
| | сн2он сн2он | T |
| | SECTION C | |
| 22 | T_b of glucose solution = 100.20°C | |
| | $\Delta T_{b} = T_{b} - T_{b}^{o}$ | 1/2 |
| | = 100.20 $= 100$ C $= 0.20$ C 0 0.20 K | /2 |
| | $\Delta T_{b} = K_{b} .m$ | 1/_ |
| | m = <u>0.20</u> = 0.390 mol/kg | 72 |
| | 0.512 | |
| | $\Delta T_f = K_f \cdot m$ | 1/2 |
| | $\Delta T_f = 1.86 \text{ K kg/mol} \times 0.390 \text{ mol/kg}$ | 72 |
| | $\Delta T_f = 0.725 \text{ K}$ | |
| | Freezing point of solution | 1 |
| | = 273.15 - 0.725 | |
| | = 272.425 K | 1/2 |
| | Or -0.725 °C | |
| 23 | (a) (i) The limiting molar conductivity of an electrolyte can be represented as the sum of the | 1 |
| | individual contributions of the anion and cation of the electrolyte. | 1 |
| | (ii) The amount of chemical reaction which occurs at any electrode during electrolysis by a current is proportional to the quantity of electricity passed through the electrolyte. | L |
| | (b) 'X' is better, as X has more negative electrode potential than Fe / X has more oxidation | 1/2 , 1/2 |
| | potential than Fe. | |
| 24 | $t_{1/2} = \frac{0.693}{k}$ | |
| | $k_1 = \frac{0.693}{20} = 0.03465 / 3.465 \times 10^{-2} \text{ min}^{-1}$ | 1/2 |
| | $k_2 = \frac{0.693}{7} = 0.1386 / 1.386 \times 10^{-1} \text{ min}^{-1}$ | 1/2 |
| | 5 | |
| | | |
| | $\log \frac{\kappa_2}{\kappa_1} = \frac{E_a}{2.303 \text{R}} \left[\frac{I_2 - I_1}{T_1 T_2} \right]$ | 1/2 |
| | | |
| | 0.1386 Fa [350-300] | |
| | $\log \frac{0.1350}{0.03465} = \frac{2.0}{2.303 \times 8.314} \frac{1350 \times 300}{[350 \times 300]}$ | 1/2 |
| | $\log 4 = \frac{Ea}{19.15} \frac{[50]}{[350 \times 300]}$ | |
| | Ea = 24209 J mol ⁻¹ or 24.209 kJ mol ⁻¹ (Deduct $\frac{1}{2}$ mark for no or incorrect unit) | 1 |
| | | |



| 28 | (a) | | | |
|----|---|--|--|-----|
| | | Native protein | Denatured protein | |
| | | Three-dimensional structure is intact.
Biologically active | Three-dimensional structure is destroyed.
Biologically inactive | 1 |
| | | | (Or any other one correct difference) | |
| | (b) Lactose | | | |
| | (C) | Vitamin K | | 1 |
| | | SECT | | |
| 29 | (a) | (i) Slowest step. | | 1 |
| | (, | ii) Series of elementary reactions / Reactions | involving two or more steps. | 1 |
| | (b) | Increases with increase in temperature. | | 1 |
| | | (|)R | |
| | (b) Molecularity is defined only for elementary reactions whereas order is experimentally | | | |
| | det | ermined hence applicable for both / Because | molecularity of each elementary reaction in | 1 |
| | con | nplex reaction may be different and hence m | eaningless for overall complex reaction whereas | |
| | and | is therefore applicable for both | termined by the slowest step in its mechanism | |
| | (c) | 9 times | | 1 |
| 30 | (a) | | | |
| | (i) | | | |
| | | OH OH | OH | |
| | | Br | <u>_</u> | |
| | | $\left \frac{\operatorname{Br}_2 \operatorname{in} \operatorname{CS}_2}{273 \operatorname{K}} \right + \left \right $ | | 1 |
| | | | \mathbf{Y} | |
| | | | Br | |
| | | / 2-Bromophenol and 4-Bromophenol is | formed. | |
| | (ii) | | | |
| | | | NO | 1 |
| | | Conc. HNO ₃ | | |
| | | $ \begin{bmatrix} \\ \end{bmatrix} \\ \begin{bmatrix} \\ \\ \end{bmatrix} \\ \begin{bmatrix} \\ \\ \end{bmatrix} $ | | |
| | | \sim | | |
| | | NO | 2 | |
| | | / 2,4,6-Trinitrophenol / Picric acid is for | ned. | 1 |
| | b)D | ue to resonance, the lone pair of electrons o | n oxygen is not easily available for | 1 |
| | pro | tonation. | | |
| | c) | | | |
| | Phe | nol | staul group / phonoxido ion formed is loss stable | 1/2 |
| | inc | resol | thy group/ phenoxide for formed is less stable | 1/2 |
| | | OR | | |
| | c) 2 | -Hvdroxybenzaldehvde / 2- Hvdroxybenzene | carbaldehvde. | |
| | -, | , | | 1 |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

| 31 | | |
|----|--|----------|
| | (A) (a) The cell reaction is | 1 |
| | $Sn(s)+2H'(aq) \rightarrow Sn^{2}(aq)+H_2(g)$ | |
| | | |
| | $E_{Coll} = (E^{o}_{c} - E^{o}_{c}) - \frac{0.059}{100} \log \frac{[Sn^{2+}]}{100}$ | 1 |
| | $-Cen - c - u^{2} - u$ | |
| | 0.059 0.004 | |
| | $= [(0) - (-0 \cdot 14)] - \frac{0.037}{2} \log \frac{0.004}{(0.02)^2}$ | |
| | $= 0.14 - 0.0295 \log 10$
= 0.1105 V | |
| | - 0.1105 V | 1 |
| | | |
| | b) (i) overagterial of Qa | 1 |
| | | |
| | (ii) Number of ions carrying current per unit volume decreases on dilution | 1 |
| 31 | B) a) At anode: | |
| | Pb+SO ₄ ⁻² →PbSO ₄ +2e− | 1/2 |
| | At cathode: | |
| | $PbO_2 + SO_4^{-2} + 4H^+ + 2e - \rightarrow PbSO_4 + 2H_2O$ | 1/2 |
| | Overall reaction: | |
| | $Pb+PbO_{2}+2 SO_{4}^{-2}+4H^{+} \rightarrow 2PbSO_{4}+2H_{2}O$ | 1 |
| | b) | |
| | $E_{Cell} = E_{Cell}^{\circ} - \frac{0.059}{log} log \left[\frac{[Cr^{3+}]^2}{2} \right]$ | |
| | $\begin{aligned} & = 1 \\ \text{a} \\ \text{a} \\ \text{b} \\ \text{b} \\ \text{cell} = 1 \\ \text{cell} = 1 \\ \text{cell} = 1 \\ \text{cell} = \frac{0.059}{100} \log (10^{-2})^2 / (10^{-2}) (1 \\ \text{cell} + 1^{14}) \end{aligned}$ | 1 |
| | $= 4.22 - \frac{0.059}{6} (54) \log (10^{-7}) (17 \times 10^{-7})$ | |
| | $= 1.33 - \frac{1}{6} (54) \log 10$ | |
| | $= 1.33 - 0.039 \times 9$
= 1.33 - 0.531 | |
| | = 0.799 V | 1 |
| 32 | A)a) The orbital splitting energies are not sufficiently large for forcing paring of electrons. | 1 |
| | configuration / Strong field effect stabilises higher oxidation state. | 1 |
| | c) Co-ordination isomerism. | 1 |
| | a) נואו(ד2O) ₆ j ⁻¹ nas unpaired electrons whereas נאו(CN) ₄ j ⁻¹ has no unpaired electron.
e) Pentaamminecarbonatocobalt(III) chloride | 1
 1 |
| | OR | |
| 32 | B)(a) The higher stability of complexes involving chelating ligands as compare to complexes having non-chelating ligand | 1 |
| | Example: $[Co(en)_3]^{3+}$ (or any other correct example) | 1 |
| | (b) d ² sp ³ diamagnetic | 1+1 |
| | (c) $[Pt (NH_3)_2 Cl_2]$ | 1 |

| 33 | (A) (a) (i) OH | 1 |
|----|---|-----|
| | $2CH_3 - C - H \xrightarrow{OH^-} CH_3 - CH_2 - CHO \xrightarrow{\Delta} CH_3 - CH = CH - CHO$ | 1 |
| | $CH_3CH_2COOH + NaOH + CaO + heat \longrightarrow CH_3-CH_3$ (ii) | I |
| | (b) $A = (CH_3)_2CH = CHCH_3 / 2$ -Methylbut-2-ene | 1 |
| | B = CH₃CHO / Ethanal | 1 |
| | C = CH ₃ COCH ₃ / Acetone/ Propanone | 1 |
| | OR | |
| 33 | $A = C_3 H_7 COOC_4 H_9 / Butyl butanoate$ | 1 |
| | $B = C_3 H_7 COOH / Butanoic acid$ | 1/2 |
| | C= C ₄ H ₉ OH / Butan-1-ol | 1/2 |
| | $C_{3}H_{7}COOC_{4}H_{9}+ dil.H_{2}SO_{4} \rightarrow C_{3}H_{7}COOH + C_{4}H_{9}OH$ | 1 |
| | $C_4H_9OH + Conc.$ Sulphuric acid + Heat \rightarrow CH ₃ CH ₂ CH=CH ₂ | 1 |
| | $C_4H_9OH \xrightarrow{\text{CrO}_3/\text{CH}_3COOH} C_3H_7COOH$ | 1 |

यंकन यो.रना

MARKING SCHEME 2024-25

रसायनावनाम् (मेंदु)तिक CHEMISTRY (Theory)- 043 QP CODE 56/1/1

MM : 70



| 18 | $101^{-} + 2MnO_4^{-} + 16H^{+} \rightarrow 2Mn^{2+} + 8H_2O + 5I_2$ | 1 |
|----|--|-------|
| | (a) $\operatorname{Cr}_2 \operatorname{O}_7^{2-} + 14 \operatorname{H}^+ + 6 \operatorname{Fe}^{2+} \to 2 \operatorname{Cr}^{3+} + 6 \operatorname{Fe}^{3+} + 7 \operatorname{H}_2 \operatorname{O}$ | 1 |
| 19 | • कम अभिनियाशील | 3 |
| | 2-9- (-=== में कार्बानल समह का कार्बन | |
| | ग्राण गापनेन के कार्बोर्गल समूह की | |
| | त्लाना में क्य इलेक्ट्रानरागी होता है।/ | |
| | बन्दा राइडाइड में अनुमाद के कारण | Ĩ |
| | कार्बातल समूह की भुवता केम हा जाती ह | |
| | | |
| | | |
| 20 | (a) $A = CH_3CH_2CN$; $B = CH_3CH_2 CH_2NH_2$ | ½ x 4 |
| | (b) $A = C_6 H_5 N_2^+ C_1^-$; | |
| | | |
| 21 | B = CN | 1 |
| | $ \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad$ | |
| | СH ₂ OH CH ₂ OH | |
| | | |
| | $(0) CHO \qquad (COOH) \qquad (CHOH) \qquad (CHOH).$ | |
| | | 1 |
| | <u> </u> | |
| 22 | T_b of glucose solution = 100.20°C | |
| | $\Delta T_b = T_b - T_b$
= 100.20 ° - 100 °C = 0.20 °C or 0.20 K | 1/2 |
| | $\Delta T_{b} = K_{b} .m$ | 1/2 |



$$\begin{array}{c} \begin{array}{c} m = 0.20 = 0.390 \mol/kg \\ 0.512 \\ \Delta T_1 = K_1, m \\ \Delta T_1 = 1.86 \mbox{ Kymol $\times 0.390 \mol/kg } \\ \Delta T_1 = 0.725 \mbox{ K} \\ \hline \Box = 0.725$$

3 | P a g e

$$E_{a} = 24209 \text{ J mol}^{4} \text{ or } 24209 \text{ J mol}^{4} \text{ or } 24209 \text{ J mol}^{4} \text{ or } 23407 \text{ or } 757 \text{ f} \text{$$



.

खंड 'घ' 29 (a) (i) सबसे जेद पद 1 (i) प्रावनिक अगिनिया की मेरवला / देा या आणिक 1 पदों वाली आफि क्रिमा ७) ताप में वृद्धि के साथ बढ़ जाता है। 1 ७ आछितनता केवल प्राथमिक अभिक्रिया के लिए ही परिभाषित होती है जनकी आणक्रिया की कोटि एक प्राघोगिक जाता है आतः दोनों पर लग्ग होतीहें/ क्यों कि जारत आ ह आ क्रिया के लिए ज त्येक पार्षामक, 1 अभिक्रिया की आणिकता भिन्त्र होती है अत' डारिल अभिक्रिया के लिए उगाण्वकता का कोई अर्घ नहीं होता जबकि जारल को भोद्रिया के लिय को टि क्रियाविश्व का सबसे मेद पद की बाती है अत' होने' पर आग्र होती है ! (C) -110[-11 30 (i) (ii) OH 22 1 2, 4, 6- द्राइनाइ रोफा ताला)) पिक्रिक आस्त Conc. HNO₃ NO₂ वनता है। 6 | Page

े अनुनाद के कारण, ऑक्सीजान पर असह भारति इलेक्ट्रोन खुगल पोटोनन के लिए आसामी से । उपलब्ध- नहीं है! 1/2 (c) • फ़ीनॉल • माचिला समह के इलेक्टॉन विज्ञोचन प्रभाव (+ I प्रभाव) के कारण / क्रीसॉल के निर्मित फीनॉक्साइड आग्र काम स्वायी होता है। 1 31 81 1 1 c) 2- हाइड्रा क्यी केन्जी लिड हाइड / 2- हाइड्रा क्यी केनी न का कील्डल खंड 'ङ 31 (A) (a) सेल आग किया है $Sn(s)+2H^{+}(aq)\rightarrow Sn^{2+}(aq)+H_{2}(g)$ $E_{Cell} = (E^{o}_{c} - E^{o}_{a}) - \frac{0.059}{2} \log \frac{[Sn^{2+1}]}{[H^{+}]^{2}}$ $= [(0) - (-0 \cdot 14)] - \frac{0.059}{2} \log \frac{0.004}{(0.02)^2}$ = 0.14 - 0.0295 log 10 = 0.1105 V 02 के अग्रियों के मत के कारण b) (i)

तनुकरण करने पर प्रति इकाई आयतन में (ii) विद्युतणामा लेजाने वाले आयतां की संग्राम 1 बाट जाती है। 😤 সদ্মা 31 1/2 ी क्योर्ड 'पर PbO₂+ SO₄⁻²+4H⁺+2e-→PbSO₄+2H₂O 2-77777 3-1777 3-1777 3-1777 Pb+PbO2+2 SO4-2+4H⁺ → 2PbSO4+2H₂O 1/2 1 b) (大); $E_{Cell} = E_{Cell}^{\circ} - \frac{0.059}{n} log \left[\frac{[Cr3^+]2}{[Cr207^{2-}][H^+]^{14}} \right]$ 1 Ecell = $1.33 - \frac{0.059}{6} \log (10^{-2})^2 / (10^{-2}) (1 \times 10^{-4})^{14}$ 1 $= 1.33 - \frac{0.059}{6} (54) \log 10$ = 1.33 - 0.059 x 9 = 1.33 - 0.5311 = 0.799 V (A) (a) मार्शनों की विपाटन आजी इतनी अग्य के 32 नहीं होती जो इलेक्ट्रांतों की युग्मत के लिए वावम करे (b) प्रवल सेत्र लिंगन्ड की उपास्थात में, dt अधिक रूपायी de विन्यास में परिवर्तित हा जाता है। प्रवल को प्रगाव उत्पतर डॉक्सीकरण अवस्था को स्पापित्व प्रधान करता है। (с) उपसहसंगोलन स्मावयवता a) [Ni(H20)] 2+ हें अयुगलित इतेक्ट्रॉन हैं GIERA [Ni (CN]] - n' ant 31 31 31 month a maxing 1 -18' 3 e) पेन्टा रेम्सीत कार्वी ने टो को बाल्ट (11) क्लोबाइड 1

3-1 21 21 (B) (a) जिना कीलेट लिगान्ड युक्त संकुली 32 47) तुलग में भोलेट लिगन्ड युक्त से कुल अधिक स्पापी होते हैं! 1 3218-241 ;: [Co(en)3]3+ 1+1 1 (A) (a) (i) 33 он 1 $2CH_3 - C - H \longrightarrow CH_3 - CH_2 - CHO \longrightarrow CH_3 - CH = CH - CHO$ 1 CH₃CH₂COOH + NaOH + CaO + 3 (3) → CH₃-CH₃ (ii) 1 (b) A = (CH3)2CH=CHCH3 (유 · 커 @ v @ 2 - 2 ~ 동न B = CH3CHO / 관환구여 C = CH3COCH3/ 관련 고려 / 며 여 1 1 अंघवा A= C3H7COOC4H9/ 04712 M 0272 -1122 1 33 1/2 B=C3H7COOH/ 0121130 37701 C=C4HOH/ @22-7-1-31/M 1/2 $C_{3}H_{7}COOC_{4}H_{9} + (\overline{J}H_{2}SO_{4} \rightarrow C_{3}H_{7}COOH + C_{4}H_{9}OH \\C_{4}H_{9}OH + (\overline{H})G + (\overline{$ 1 1 C₄H₉OH C₄H₉OH C₃H₇COOH 1