

**CBSE Board**  
**Class XI Mathematics**  
**Sample Paper – 1**

**Time: 3 hrs**

**Total Marks: 100**

---

**General Instructions:**

1. All questions are compulsory.
  2. The question paper consist of 29 questions.
  3. Questions 1 – 4 in Section A are very short answer type questions carrying 1 mark each.
  4. Questions 5 – 12 in Section B are short-answer type questions carrying 2 mark each.
  5. Questions 13 – 23 in Section C are long-answer I type questions carrying 4 mark each.
  6. Questions 24 – 29 in Section D are long-answer type II questions carrying 6 mark each.
- 

**SECTION – A**

1. Find the derivative of  $\sin(x + 1)$ .
2. Find the truth value of p: 'Every real number is either prime or composite.'
3. Simplify:  $\frac{1+3i}{1-2i}$

**OR**

Show that  $i^n + i^{n+1} + i^{n+2} + i^{n+3} = 0$ , for all  $i \in \mathbb{N}$ .

4. A coin is tossed twice. Find the probability of getting at least one head.

**SECTION – B**

5. A and B are two sets such that  $n(A - B) = 14 + x$ ,  $n(B - A) = 3x$  and  $n(A \cap B) = x$ , draw a Venn diagram to illustrate the information. If  $n(A) = n(B)$ , then find the value of  $x$ .
6. If the power sets of two sets are equal, then show that the sets are also equal.

**OR**

If  $a \in \mathbb{N}$  such that  $a\mathbb{N} = \{ax : x \in \mathbb{N}\}$ . Describe the set  $3\mathbb{N} \cap 7\mathbb{N}$ ?

7. There are 11 teachers who teach mathematics or physics in school. Of these, 7 teach mathematics and 3 teach both subjects. How may teach physics?

**OR**

Let A and B be two sets such that :  $n(A) = 20$ ,  $n(A \cup B) = 42$  and  $n(A \cap B) = 4$ . Find  $n(B)$  and  $n(A - B)$ .

8. Let  $A = \{1, 2\}$  and  $B = \{3, 4\}$ . Write  $A \times B$ . How many subsets will  $A \times B$  have? Also list them

9. Prove that:  $\left( \frac{\cos A}{1 - \tan A} \right) + \left( \frac{\sin A}{1 - \cot A} \right) = \cos A + \sin A$

**OR**

Prove that  $\cot^4 \theta + \cot^2 \theta = \operatorname{cosec}^4 \theta - \operatorname{cosec}^2 \theta$

10. Write contrapositive of the following statements :

1. A number is divisible by 9, then it is divisible by 3.
2. If you are born in India, then you are citizen of India.

11. Find sum :  $10^3 + 11^3 + 12^3 + \dots + 20^3$

12. Three consecutive vertices of a parallelogram ABCD are A (4, -11), B (5, 3) & C (2, 15). Find D.

### SECTION - C

13. If  $f$  and  $g$  are two functions:  $R \rightarrow R$ ;  $f(x) = 2x - 1$ ,  $g(x) = 2x + 3$ , then evaluate

(i)  $(f + g)(x)$     (ii)  $(f - g)(x)$     (iii)  $(fg)(x)$     (iv)  $\left( \frac{f}{g} \right)(x)$

14. Let  $R$  be a relation from  $N$  to  $N$  defined by  $R = \{(a, b) \in N \text{ and } a = b^4\}$ . Determine if the relation is

- (i) Reflexive    (ii) Symmetric    (iii) Transitive    (iv) Equivalence

15. In a  $\Delta ABC$ , if  $a = 3$ ,  $b = 5$ ,  $c = 7$ , find  $\cos A$ ,  $\cos B$  and  $\cos C$ .

16. Find the square root of the complex number  $5 - 12i$ .

17. Find the probability such that when 7 cards are drawn from a well shuffled deck of 52 cards, all the aces are obtained.

18. Find the sum to infinity of the series:  $\frac{1}{3} + \frac{1}{5^2} + \frac{1}{3^3} + \frac{1}{5^4} + \frac{1}{3^5} + \frac{1}{5^6} + \dots$

19. In how many ways can the letters of the word 'Mathematics' be arranged so that the (i) vowels are together (ii) vowels are not together

**OR**

In how many ways can 5 girls and 3 boys be seated in a row with 11 chairs so that no two boys sit together?

20. A point M with x-coordinate 4 lies on the line segment joining the points P(2, -3, 4) and Q (8, 0, 10). Find the co-ordinates of the point M.

**OR**

Find the equation of the set of points such that the sum of the square of its distance from the points (3, 4, 5) and (-1, 3, -7) is a constant.

21. Solve for x:  $\tan 2x + \sec^2 2x - 1 = 0$

**OR**

Solve for x:  $\sin x + \sin 2x + \sin 3x = 0$

22. Evaluate:  $\lim_{x \rightarrow 0} \frac{\log 10 + \log \left( x + \frac{1}{10} \right)}{x}$

23. Write down the binomial expression  $(1 + x)^{n+1}$ , when  $x = 8$ . Deduce that  $9^{n+1} - 8n - 9$  is divisible by 64, when n is an integer.

### **SECTION - D**

24. If  $\frac{\pi}{2} \leq x \leq \pi$  and  $\tan x = -\frac{4}{3}$ , find  $\sin \frac{x}{2}$ ,  $\cos \frac{x}{2}$ ,  $\tan \frac{x}{2}$ .

**OR**

Show that  $\sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ$ .

25. Find the mean deviation about the median for the following data:

Marks	No. of students
0-10	5
10-20	10
20-30	20
30-40	5
40-50	10

26. Prove by the principle of Mathematical Induction that every even power of every odd integer greater than one when divided by 8 leaves one as the remainder.

27. Solve the following system of inequalities graphically:

$$x + 2y \leq 10; x + y \geq 1; x - y \leq 0; x \geq 0; y \geq 0$$

**OR**

For the purpose of an experiment an acid solution between 4% and 6% is required. 640 liters of 8% acid solution and a 2% acid solution are available in a laboratory. How many liters of the 2% solution needs to be added to the 8% solution?

28. The first three terms in the binomial expansion of  $(a + b)^n$  are given to be 729, 7290 and 30375 respectively. Find a, b and n.
29. A student wants to buy a computer for Rs. 12,000. He has saved up to Rs. 6000 which he pays as cash. He is to pay the balance in annual installments of Rs. 500 plus an interest of 12% on the unpaid amount. How much will the computer cost him?

**OR**

Find the value of  $\frac{1 \times 2^2 + 2 \times 3^2 + 3 \times 4^2 + \dots \text{uptill the } n\text{th term}}{1^2 \times 2 + 2^2 \times 3 + 3^2 \times 4 + \dots \text{uptill the } n\text{th term}}$

**CBSE Board**  
**Class XI Mathematics**  
**Sample Paper – 1 Solution**

---

**SECTION – A**

1.  $[\sin(x+1)]' = \cos(x+1) \cdot 1 = \cos(x+1)$
2. Giving one counter example is enough to prove the falsehood of a statement. Here counter example is: The real number 1 is neither prime nor composite. So the statement is false.

3.

$$\frac{1+3i}{1-2i} \times \frac{1+2i}{1+2i} = \frac{1-6+3i+2i}{(1)^2 - (2i)^2} = \frac{-5+5i}{1-4i^2} = \frac{-5+5i}{1+4} = \frac{-5+5i}{5} = -1+i$$

**OR**


$$i^n + i^{n+1} + i^{n+2} + i^{n+3} = i^n (1 + i + i^2 + i^3) = i^n (1 + i - 1 - i) = 0 \quad \because i^2 = -1 \text{ and } i^3 = -i$$

4. Sample space  $S = \{HH, HT, TH, TT\}$  i.e. total number of cases = 4  
Favourable cases for atleast one head are  $\{HH, HT, TH\}$ .

$$\text{Required probability} = \frac{3}{4}$$

**SECTION – B**

5.  $n(A - B) = 14 + x$ ,  $n(B - A) = 3x$  and  $n(A \cap B) = x$



$$n(A) = n(B)$$

$$n(A) = n(A - B) + n(A \cap B);$$

$$n(B) = n(B - A) + n(A \cap B)$$

$$\Rightarrow n(A - B) + n(A \cap B) = n(B - A) + n(A \cap B)$$

$$\Rightarrow 14 + x + x = 3x + x$$

$$\Rightarrow 14 = 2x \Rightarrow x = 7$$

6. Let  $a$  be any element which belongs to set  $A$ , i.e.  $a \in A$   
 $P(A)$  is the set of all subsets of the set  $A$ . Therefore  $\{a\}$  belongs to  $P(A)$   
 i.e.  $\{a\} \in P(A)$   
 But  $P(A) = P(B)$  [ Given ]  
 $\therefore \{a\} \in P(B)$   
 $\Rightarrow a \in B$   
 So  $a \in A \Rightarrow a \in B$ , Hence  $A \subseteq B$   
 Similarly, we can prove that  $A \subseteq B$   
 $\Rightarrow A = B$

**OR**

$$a\mathbb{N} = \{ax : x \in \mathbb{N}\}$$

$$3\mathbb{N} = \{3x : x \in \mathbb{N}\} = \{3, 6, 9, \dots\}$$

$$\text{And } 7\mathbb{N} = \{7x : x \in \mathbb{N}\} = \{7, 14, 21, 28, \dots\}$$

$$3\mathbb{N} \cap 7\mathbb{N} = \{21, 42, \dots\} = \{21x : x \in \mathbb{N}\} = 21\mathbb{N}$$

7. Let  $A$  be the set of teachers who teach math and  $B$  be the set of teachers who teach physics. Then,  
 $n(A \cup B) = 11$ ,  $n(A) = 7$  &  $n(A \cap B) = 3$   
 $n(A \cup B) = n(A) + n(B) - n(A \cap B)$   
 $11 = 7 + n(B) - 3$   
 $n(B) = 7$

**OR**

$$n(A) = 20, n(A \cup B) = 42 \text{ and } n(A \cap B) = 4$$

$$n(A \cup B) = n(A) + n(B) - n(A \cap B)$$

$$42 = 20 + n(B) - 4$$

$$n(B) = 42 - 20 + 4 = 26$$

$$n(A - B) = n(A) - n(A \cap B)$$

$$= 20 - 4 = 16$$

8.  $A \times B = \{(1, 3), (1, 4), (2, 3), (2, 4)\}$   
 Since  $A \times B$  contains 4 elements therefore it has  $2^4 = 16$  subsets  
 Subsets are  
 $\{\}$ ,  
 $\{(1, 3)\}, \{(1, 4)\}, \{(2, 3)\}, \{(2, 4)\},$   
 $\{(1, 3), (1, 4)\}, \{(1, 3), (2, 3)\}, \{(1, 3), (2, 4)\}, \{(1, 4), (2, 3)\}, \{(1, 4), (2, 4)\},$   
 $\{(1, 3), (1, 4), (2, 3)\}, \{(1, 3), (1, 4), (2, 4)\}, \{(1, 4), (2, 3), (2, 4)\},$   
 $\{(1, 3), (1, 4), (2, 3), (2, 4)\}.$

$$\begin{aligned}
9. \quad & \left( \frac{\cos A}{1 - \tan A} \right) + \left( \frac{\sin A}{1 - \cot A} \right) = \left( \frac{\cos A}{1 - \frac{\sin A}{\cos A}} \right) + \left( \frac{\sin A}{1 - \frac{\cos A}{\sin A}} \right) \\
& \left( \frac{\cos^2 A}{\cos A - \sin A} \right) + \left( \frac{\sin^2 A}{\sin A - \cos A} \right) \\
& \left( \frac{\cos^2 A - \sin^2 A}{\cos A - \sin A} \right) = \left( \frac{(\cos A - \sin A)(\cos A + \sin A)}{\cos A - \sin A} \right) \\
& = (\cos A + \sin A)
\end{aligned}$$

**OR**

$$\begin{aligned}
\cot^4 \theta + \cot^2 \theta &= (\cot^2 \theta)^2 + \cot^2 \theta \\
&= (\operatorname{cosec}^2 \theta - 1)^2 + \operatorname{cosec}^2 \theta - 1 \\
&= \operatorname{cosec}^4 \theta - 2\operatorname{cosec}^2 \theta + 1 + \operatorname{cosec}^2 \theta - 1 \\
&= \operatorname{cosec}^4 \theta - \operatorname{cosec}^2 \theta
\end{aligned}$$

**10.**

1. If a number is not divisible by 3, it is not divisible by 9.
2. If you are not citizen of India, then you were not born in India.

**11.** The given series :  $10^3 + 11^3 + 12^3 + \dots + 20^3$

Can be written as

$$\begin{aligned}
& (1^3 + 2^3 + 3^3 + \dots + 20^3) - (1^3 + 2^3 + 3^3 + \dots + 9^3) \\
&= \sum_{r=1}^{20} r^3 - \sum_{r=1}^{10} r^3 \\
&= \left( \frac{20(20+1)}{2} \right)^2 - \left( \frac{9(9+1)}{2} \right)^2 \\
&= 42075
\end{aligned}$$

**12.** Vertices of a parallelogram ABCD are A (4, -11), B (5, 3) & C (2, 15)

let D(x,y)

then mid point of diagonal AC =  $\left( \frac{4+2}{2}, \frac{-11+15}{2} \right)$

also mid point of diagonal BD =  $\left( \frac{5+x}{2}, \frac{3+y}{2} \right)$

but midpoint of AC BD coincides

so,

$$\left(\frac{4+2}{2}, \frac{-11+15}{2}\right) = \left(\frac{5+x}{2}, \frac{3+y}{2}\right)$$

$$5+x=4+2 \dots\dots\dots x=1$$

$$3+y=-11+15 \dots\dots\dots y=1$$

### SECTION - C

13.  $f(x) = 2x - 1, g(x) = 2x + 3; x \in \mathbb{R}$

$$(f + g)(x) = f(x) + g(x) = (2x - 1) + (2x + 3) = 4x + 2; x \in \mathbb{R}$$

$$(f - g)(x) = f(x) - g(x) = (2x - 1) - (2x + 3) = -4$$

$$(fg)(x) = f(x)g(x) = (2x - 1)(2x + 3) = 4x^2 - 2x + 6x - 3 = 4x^2 + 4x - 3$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} = \frac{2x-1}{2x+3}; x \in \mathbb{R} - \left\{-\frac{3}{2}\right\}$$

14.

$$\{(a, b), a = b^4, a, b \in \mathbb{N}\}$$

(i)  $(a, a) \in R \Rightarrow a = a^4,$

which is true for  $a = 1$  only, not for other values of  $a \in \mathbb{N}$

$\therefore$  Relation is not reflexive

(ii)  $\{(a, b), a = b^4, a, b \in \mathbb{N}\}$  and  $\{(b, a), b = a^4, a, b \in \mathbb{N}\}$

$a = b^4$  and  $b = a^4$  cannot be true simultaneously

$\therefore$  Relation is not symmetric.

(iii)  $\{(a, b), a = b^4, a, b \in \mathbb{N}\}; \{(b, c), b = c^4, b, c \in \mathbb{N}\}$

$$\Rightarrow a = b^4 = c^{16}$$

$$\text{So } a \neq c^4$$

$$\therefore (a, c) \notin R$$

$\therefore$  Relation is not transitive.

(iv) Since the relation is not reflexive, not symmetric, and also not transitive,

$\Rightarrow$  Relation is not an equivalence relation

15. We know,

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc} = \frac{25 + 49 - 9}{2(5)(7)} = \frac{65}{70} = \frac{13}{14}$$

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac} = \frac{9 + 49 - 25}{2(3)(7)} = \frac{33}{42}$$



16. Let  $\sqrt{5-12i} = x + yi$

$$\Rightarrow 5 - 12i = (x^2 - y^2) + 2xy i$$

Equating real and imaginary parts, we get

$$x^2 - y^2 = 5 \dots\dots(i) \quad \text{and}$$

$$2xy = -12 \dots\dots(ii)$$

$$\text{Now } (x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2 = 5^2 + 12^2 = 169$$

$$\therefore x^2 + y^2 = 13 \dots\dots\dots(iii)$$

From (i) and (iii), we get

$$2x^2 = 18 \Rightarrow x = \pm 3$$

$$\text{and } y = \pm 2$$

From equation (ii) we can say that  $xy$  is negative.

$$\text{As } xy \text{ is negative } \Rightarrow \text{when } x = 3, y = -2$$

$$\text{and when } x = -3, y = 2$$

The required square roots are  $(3 - 2i)$  and  $(-3 + 2i)$

$$\text{or } \pm (3 - 2i)$$

17. Total number of possible sets of 7 cards =  ${}^{52}C_7$

$$\text{Number of sets of 7 with all 4 aces} = {}^4C_4 \times {}^{48}C_3$$

(4 aces from among 4 aces and other 3 cards must be chosen from the rest 48 cards)

$$\begin{aligned} \text{Hence the probability that the 7 cards drawn contain 4 aces} &= \frac{{}^4C_4 \times {}^{48}C_3}{{}^{52}C_7} \\ &= \frac{1}{7735} \end{aligned}$$

18. We have

$$\begin{aligned} \frac{1}{3} + \frac{1}{5^2} + \frac{1}{3^3} + \frac{1}{5^4} + \frac{1}{3^5} + \frac{1}{5^6} + \dots &= \left[ \frac{1}{3} + \frac{1}{3^3} + \frac{1}{3^5} + \dots \right] + \left[ \frac{1}{5^2} + \frac{1}{5^4} + \frac{1}{5^6} + \dots \right] \\ &= \frac{\frac{1}{3}}{1 - \frac{1}{3^2}} + \frac{\frac{1}{5^2}}{1 - \frac{1}{5^2}} = \frac{1}{3} \times \frac{9}{8} + \frac{1}{25} \times \frac{25}{24} \\ &= \frac{3}{8} + \frac{1}{24} = \frac{10}{24} = \frac{5}{12} \end{aligned}$$

19. In MATHEMATICS, there are 11 letters of which there are 2 Ms, 2 As, and 2 Ts, so the total arrangements are  $\frac{11!}{2!.2!.2!} = 4,989,600$

(i) In MATHEMATICS, there are 4 vowels: 2 As, 1 E and 1 I.  
Since they must be together so 'AAEI' is treated as a single unit.  
So there are 8 objects which include 2 Ms, 2 As, and 2 Ts.

So the required number of arrangements are  $\frac{8!}{2!.2!} \times \frac{4!}{2!} = 120960$

(ii) Number of arrangements with vowels never together =  
Total arrangements – arrangements in which vowels are always together  
=  $4,989,600 - 120960 = 4868640$

**OR**

First the 5 girls are arranged in  $5!$  ways as shown below:

$(G_1 - G_2 - G_3 - G_4 - G_5)$

Now there are 6 places in which the boys can be arranged.

This can be done in  ${}^6P_3$  ways

$\Rightarrow$  Total ways =  $5! \times {}^6P_3$

=  $120 \times 6 \times 5 \times 4 = 14,400$

20. Let M divide PQ in the ratio  $k : 1$ .

The co-ordinates of the point M are given by

$$\left( \frac{8k+2}{k+1}, \frac{-3}{k+1}, \frac{10k+4}{k+1} \right)$$

here the x-coordinate =  $\frac{8k+2}{k+1} = 4$ .

$$\Rightarrow 8k + 2 = 4k + 4 \Rightarrow 4k = 2 \Rightarrow k = \frac{2}{4} = \frac{1}{2}$$

Putting  $k$  back in the x, y and z co-ordinate of the point M, we have  $(4, -2, 6)$

**OR**

Let the given points be A(3, 4, 5) and B (-1, 3, -7).

Let the required point be P: P(x, y, z)

Given:  $PA^2 + PB^2 = k^2$  ( $k^2$  is a constant)

$$\Rightarrow (x-3)^2 + (y-4)^2 + (z-5)^2 + (x+1)^2 + (y-3)^2 + (z+7)^2 = k^2$$

$$\Rightarrow x^2 - 6x + 9 + y^2 - 8y + 16 + z^2 - 10z + 25 + x^2 + 2x + 1 + y^2 - 6y + 9 + z^2 + 14z + 49 = k^2$$

$$\Rightarrow 2x^2 + 2y^2 + 2z^2 - 4x - 14y + 4z + 109 - k^2 = 0$$

This is the equation of the set of points P that satisfy the condition.

$$21. \tan 2x + \sec^2 2x - 1 = 0$$

$$\Rightarrow \tan 2x + 1 + \tan^2 2x - 1 = 0$$

$$\Rightarrow \tan 2x + \tan^2 2x = 0$$

$$\Rightarrow \tan 2x [1 + \tan 2x] = 0$$

$$\Rightarrow \tan 2x = 0 \text{ or } 1 + \tan 2x = 0$$

$$\Rightarrow \tan 2x = 0 \text{ or } \tan 2x = -1$$

$$\tan 2x = 0 \Rightarrow 2x = n\pi \Rightarrow x = \frac{n\pi}{2}$$

$$\tan 2x = -1 \Rightarrow \tan 2x = -\tan \frac{\pi}{4} = \tan \left( \pi - \frac{\pi}{4} \right) = \tan \frac{3\pi}{4}$$

$$\Rightarrow 2x = n\pi + \frac{3\pi}{4}, n \in \mathbb{Z}$$

$$\Rightarrow x = \frac{n\pi}{2} + \frac{3\pi}{8}, n \in \mathbb{Z}$$

**OR**

$$\sin x + \sin 2x + \sin 3x = 0$$

$$\sin x + \sin 3x + \sin 2x = 0$$

$$2\sin \left( \frac{x+3x}{2} \right) \cos \left( \frac{x-3x}{2} \right) + \sin 2x = 0$$

$$2\sin 2x \cos(-x) + \sin 2x = 0$$

$$\sin 2x (2\cos x + 1) = 0$$

$$\sin 2x = 0 \text{ or } \cos x = \frac{-1}{2}$$

$$\sin 2x = 0 \Rightarrow 2x = n\pi$$

$$\text{or } \cos x = \frac{-1}{2} \Rightarrow x = 2n\pi \pm \left( \pi - \frac{\pi}{3} \right)$$

$$x = \frac{n\pi}{2} \text{ or } x = 2n\pi \pm \left( \pi - \frac{\pi}{3} \right)$$

22.

$$\begin{aligned}
 & \lim_{x \rightarrow 0} \frac{\log 10 + \log \left( x + \frac{1}{10} \right)}{x} \\
 &= \lim_{x \rightarrow 0} \frac{\log \left[ 10 \left( x + \frac{1}{10} \right) \right]}{x} \\
 &= \lim_{x \rightarrow 0} \frac{\log (10x + 1)}{x} \\
 &= 10 \lim_{x \rightarrow 0} \frac{\log (10x + 1)}{10x} \\
 &= 10 \times 1 \quad \left( \because \lim_{x \rightarrow 0} \frac{\log (x + 1)}{x} = 1 \right)
 \end{aligned}$$

23. We have,

$$(1 + x)^{n+1} = {}^{n+1}C_0 + {}^{n+1}C_1x + {}^{n+1}C_2x^2 + {}^{n+1}C_3x^3 + \dots + {}^{n+1}C_{n+1}x^{n+1}$$

Putting  $x = 8$ , we get

$$(1 + 8)^{n+1} = {}^{n+1}C_0 + {}^{n+1}C_1(8)^1 + {}^{n+1}C_2(8)^2 + {}^{n+1}C_3(8)^3 + \dots + {}^{n+1}C_{n+1}(8)^{n+1} \dots (i)$$

$$9^{n+1} = 1 + (n+1) \times 8 + {}^{n+1}C_2(8)^2 + {}^{n+1}C_3(8)^3 + \dots + {}^{n+1}C_{n+1}(8)^{n+1}$$

$$9^{n+1} = 1 + 8n + 8 + 8^2 \{ {}^{n+1}C_2 + {}^{n+1}C_3(8)^1 + \dots + {}^{n+1}C_{n+1}(8)^{n-1} \}$$

$$9^{n+1} = 8n + 9 + 8^2 \{ {}^{n+1}C_2 + {}^{n+1}C_3(8)^1 + \dots + {}^{n+1}C_{n+1}(8)^{n-1} \}$$

$$9^{n+1} - 8n + 9 = 64 \{ {}^{n+1}C_2 + {}^{n+1}C_3(8)^1 + \dots + {}^{n+1}C_{n+1}(8)^{n-1} \}$$

$$9^{n+1} - 8n + 9 = 64 \times \text{an integer}$$

$$\therefore 9^{n+1} - 8n + 9 \text{ is divisible by } 64.$$

## SECTION - D

24.  $\tan x = -\frac{4}{3}$ ;  $\frac{\pi}{2} \leq x \leq \pi$

$$\tan x = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}} \quad \left( \because \tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta} \right)$$

$$\Rightarrow -\frac{4}{3} = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}} \Rightarrow 4 \left( 1 - \tan^2 \frac{x}{2} \right) = -6 \tan \frac{x}{2}$$

$$\Rightarrow 4 \tan^2 \frac{x}{2} - 6 \tan \frac{x}{2} - 4 = 0$$

$$\Rightarrow 2 \tan^2 \frac{x}{2} - 3 \tan \frac{x}{2} - 2 = 0$$

The equation is quadratic in  $\tan \frac{x}{2}$

$$\Rightarrow \tan \frac{x}{2} = \frac{-(-3) \pm \sqrt{9+16}}{2 \cdot 2} = \frac{3 \pm 5}{4} = 2, -\frac{1}{2}$$

$$\text{Given } \frac{\pi}{2} \leq x \leq \pi \Rightarrow \frac{\pi}{4} \leq \frac{x}{2} \leq \frac{\pi}{2} \Rightarrow \frac{x}{2} \in \text{I}^{\text{st}} \text{ quadrant}$$

$$\text{In I}^{\text{st}} \text{ quadrant, } \tan \frac{x}{2} \geq 0 \Rightarrow \tan \frac{x}{2} = 2$$

$$\text{We know, } 1 + \tan^2 \theta = \sec^2 \theta$$

$$\Rightarrow 1 + \tan^2 \frac{x}{2} = \sec^2 \frac{x}{2} \Rightarrow 1 + (2)^2 = \sec^2 \frac{x}{2}$$

$$\Rightarrow \sec^2 \frac{x}{2} = 5 \Rightarrow \sec \frac{x}{2} = \pm \sqrt{5} \Rightarrow \cos \frac{x}{2} = \pm \frac{1}{\sqrt{5}}$$

$$\text{In I}^{\text{st}} \text{ quadrant, } \cos \frac{x}{2} \geq 0 \Rightarrow \cos \frac{x}{2} = \frac{1}{\sqrt{5}}$$

$$\text{We know } \sin \theta = \pm \sqrt{1 - \cos^2 \theta}$$

$$\sin \frac{x}{2} = \pm \sqrt{1 - \cos^2 \frac{x}{2}} = \pm \sqrt{1 - \frac{1}{5}} = \pm \sqrt{\frac{4}{5}} = \pm \frac{2}{\sqrt{5}}$$

$$\text{In I}^{\text{st}} \text{ quadrant, } \sin \frac{x}{2} \geq 0 \Rightarrow \sin \frac{x}{2} = \frac{2}{\sqrt{5}}$$

$$\therefore \text{(i) } \sin \frac{x}{2} = \frac{2}{\sqrt{5}} \quad \text{(ii) } \cos \frac{x}{2} = \frac{1}{\sqrt{5}} \quad \text{(iii) } \tan \frac{x}{2} = 2$$

**OR**

$$\sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ$$

$$= \frac{\sqrt{3}}{\sin 20^\circ} - \frac{1}{\cos 20^\circ}$$

$$= \frac{\sqrt{3} \cos 20^\circ - \sin 20^\circ}{\sin 20^\circ \cos 20^\circ}$$

$$= \frac{2 \left( \frac{\sqrt{3}}{2} \cos 20^\circ - \frac{1}{2} \sin 20^\circ \right)}{\sin 20^\circ \cos 20^\circ}$$

$$= \frac{2(\sin 60^\circ \cos 20^\circ - \cos 60^\circ \sin 20^\circ)}{\sin 20^\circ \cos 20^\circ}$$

$$= \frac{2 \sin(60^\circ - 20^\circ)}{\sin 20^\circ \cos 20^\circ}$$

$$\begin{aligned}
&= \frac{4\sin 40^\circ}{2\sin 20^\circ \cos 20^\circ} \\
&= \frac{4\sin 40^\circ}{\sin 40^\circ} \\
&= 4
\end{aligned}$$

25.

Marks	Frequency ( $f_i$ )	Cumulative frequency (cf)
0-10	5	5
10-20	10	15
20-30	20	35
30-40	5	40
40-50	10	50
Total	N= 50	

$$\text{Median (M)} = l + \frac{\frac{n}{2} - cf}{f} \times h$$

$l$  = lower limit of the median class,  $n$  = number of observations,

cf = cumulative frequency of the class preceding the median class,  $h$  = class size and

$f$  = frequency of the median class

Substituting the values we get

$$\text{Median (M)} = 20 + \frac{(25 - 15) \times 10}{20}$$

$$\text{Median (M)} = 25$$

$x_i$	$f_i$	$ d_i  =  x_i - M $	$f_i  d_i $
5	5	20	100
15	10	10	100
25	20	0	0
35	5	10	50
45	10	20	200
	50		450

$$\therefore \sum_{i=1}^n |d_i| f_i = 450, \quad n = \sum_{i=1}^n f_i = 50$$

$$\therefore \text{M.D (M)} = \frac{\sum_{i=1}^n |d_i| f_i}{n} = \frac{450}{50} = 9$$

26. The first odd integer  $> 1$ , is 3 .

The general term for odd number  $> 1$  is  $(2r + 1)$

$P(n)$ :  $(2r + 1)^{2n} = 8m + 1$  where  $m, n$  are natural numbers

i.e.  $P(n)$ :  $(2r + 1)^{2n} - 1$  is divisible by 8

Here  $P(1)$ :  $(2r + 1)^{2 \cdot 1} - 1$  is divisible by 8.

Consider  $(2r + 1)^2 - 1 = 4r^2 + 4r = 4r(r + 1)$

$r(r + 1)$  being the product of consecutive natural numbers is even,

So,  $4r(r + 1)$  is divisible by 8.

Therefore,  $P(1)$  is true.

Let us assume  $P(k)$  to be true

$P(k)$ :  $(2r + 1)^{2k} - 1$  is divisible by 8.

Using this assumption, we will prove  $P(k + 1)$  to be true

$P(k + 1)$ :  $(2r + 1)^{2(k + 1)} - 1$  is divisible by 8.

Consider  $(2r + 1)^{2(k + 1)} - 1 = (2r + 1)^{2k} (2r + 1)^2 - 1 = (8m + 1)(8p + 1) - 1$  [using  $P(1)$  and  $P(k)$ , where  $m$  and  $p$  are integers ]

$(2r + 1)^{2(k + 1)} - 1 = 64 mp + 8(m + p) + 1 - 1 = 64 mp + 8(m + p)$ , which is divisible by 8.

Thus  $P(k + 1)$  is true whenever  $P(k)$  is true, also  $P(1)$  is true.

$\Rightarrow P(n)$  is true for every natural number  $n$ .

27.

$x + 2y = 10$  or  $x = 10 - 2y$


x	14	10	6
y	-2	0	2

$x + y = 1$  or  $y = 1 - x$

x	-2	0	3
y	3	1	-2

$x - y = 0$  or  $y = x$

x	-2	0	2
y	-2	0	2



OR

The amount of acid in 640 litres of the 8% solution =  $8\%$  of  $640 = \frac{8 \times 640}{100}$

Let  $x$  litres of the 2% solution be added to obtain a solution between 4% and 6%

The amount of acid in  $x$  litres of the 2% solution =  $\frac{2 \times x}{100}$

The resultant amount =  $640 + x$

The amount of acid in  $(640 + x)$  litres solution is =  $\frac{8 \times 640}{100} + \frac{2 \times x}{100}$

Acid percentage of the solution now =  $\frac{\frac{8 \times 640}{100} + \frac{2 \times x}{100}}{640 + x} \times 100$

$$\Rightarrow 4 < \frac{\frac{8 \times 640}{100} + \frac{2 \times x}{100}}{640 + x} \times 100 < 6$$

$$\Rightarrow \frac{4(640 + x)}{100} < \frac{8 \times 640}{100} + \frac{2 \times x}{100} < \frac{6(640 + x)}{100}$$

$$\Rightarrow 4(640 + x) < 5120 + 2x < 6(640 + x)$$

$$\Rightarrow 2(640 + x) < 2560 + x < 3(640 + x)$$

$$\Rightarrow 2(640 + x) < 2560 + x \text{ and } 2560 + x < 3(640 + x)$$



$$\begin{aligned}
&\Rightarrow 2(640+x) < 2560+x < 3(640+x) \\
&\Rightarrow 2(640+x) < 2560+x \text{ and } 2560+x < 3(640+x) \\
&\Rightarrow 1280+2x < 2560+x \text{ and } 2560+x < 1920+3x \\
&\Rightarrow x < 1280 \text{ and } 320 < x \\
&\Rightarrow 320 < x < 1280
\end{aligned}$$

Hence, the number of liters of 2% of acid which must be added should be more than 320 but less than 1280.

**28.** The first three terms in the binomial expansion  $(a+b)^n$ , ie  $t_1, t_2, t_3$  are given.

$$\Rightarrow t_1 = {}^nC_0 a^n b^0 = 729 \dots (i);$$

$$t_2 = {}^nC_1 a^{n-1} b^1 = 7290 \dots (ii);$$

$$t_3 = {}^nC_2 a^{n-2} b^2 = 30375 \dots (iii)$$

$$\text{Now, } t_1 = {}^nC_0 a^n b^0 = 729 \Rightarrow 1 \times a^n \times 1 = 729 \Rightarrow a^n = 729 \dots (iv)$$

Dividing (ii) by (i), we have

$$\frac{t_2}{t_1} = \frac{{}^nC_1 a^{n-1} b^1}{{}^nC_0 a^n b^0} = \frac{7290}{729} = 10 \Rightarrow \frac{n a^{n-1} b}{a^n} = 10 \Rightarrow \frac{nb}{a} = 10 \dots (v)$$

Multiplying (iii) by (i), we have

$$t_3 \times t_1 = {}^nC_2 a^{n-2} b^2 \times {}^nC_0 a^n b^0 = \frac{n(n-1)}{2} a^{2n-2} b^2 = 729 \times 30375 \dots (vi)$$

Squaring (ii), we have

$$\left[ {}^nC_1 a^{n-1} b^1 \right]^2 = [7290]^2 \Rightarrow n^2 a^{2n-2} b^2 = 7290 \times 7290 \dots (vii)$$

Dividing (vi) by (vii), we have

$$\frac{\frac{n(n-1)}{2} a^{2n-2} b^2}{n^2 a^{2n-2} b^2} = \frac{729 \times 30375}{7290 \times 7290}$$

$$\Rightarrow \frac{(n-1)}{2n} = \frac{30375}{7290 \times 10}$$

$$\Rightarrow \frac{(n-1)}{2n} = \frac{5}{12}$$

$$\Rightarrow 12n - 12 = 10n \Rightarrow 2n = 12 \Rightarrow n = 6$$

Putting  $n = 6$  in (iv), we have

$$a^6 = 729 \Rightarrow a = 3$$

Putting  $n = 6, a = 3$  in (v), we have

$$\frac{6b}{3} = 10 \Rightarrow b = 5$$

Hence,  $a = 3, b = 5, n = 6$

29. Interest to be paid with Installment 1 (S.I. on Rs. 6000 for 1 year) =  $\frac{6000 \times 12 \times 1}{100} = 720$

Interest to be paid with Installment 2 (S.I. on Rs. 5500 for 1 year) =  $\frac{5500 \times 12 \times 1}{100} = 660$

Interest to be paid with Installment 3 (S.I. on Rs. 5000 for 1 year) =  $\frac{5000 \times 12 \times 1}{100} = 600$

Interest to be paid with 12th Installment (S.I. On Rs 500 for 1 year) =  $\frac{500 \times 12 \times 1}{100} = 60$

Total interest paid =  $720 + 660 + 600 + \dots + 60$

This forms an A.P., with  $a = 720$  and  $d = -60$

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

$$\Rightarrow S_n = \frac{12}{2}[2 \times 720 + (12-1)(-60)] = 4680$$

The computer costed the student =  $12000 + 4680 = 16680$

**OR**

$\frac{1 \times 2^2 + 2 \times 3^2 + 3 \times 4^2 + \dots \text{uptill the } n\text{th term}}{1^2 \times 2 + 2^2 \times 3 + 3^2 \times 4 + \dots \text{uptill the } n\text{th term}}$

$1^2 \times 2 + 2^2 \times 3 + 3^2 \times 4 + \dots \text{uptill the } n\text{th term}$

Consider Numerator =  $1 \times 2^2 + 2 \times 3^2 + 3 \times 4^2 + \dots + \text{uptill the } n\text{th term}$

The  $n\text{th}$  term is  $n(n+1)^2$

$\therefore 1 \times 2^2 + 2 \times 3^2 + 3 \times 4^2 + \dots \text{uptill the } n\text{th term}$

$$= \sum n(n+1)^2$$

$$= \sum n(n^2 + 1 + 2n)$$

$$= \sum (n^3 + n + 2n^2) = \sum n^3 + 2\sum n^2 + \sum n$$

$$= \left[ \frac{n(n+1)}{2} \right]^2 + 2 \frac{n(n+1)(2n+1)}{6} + \frac{n(n+1)}{2}$$

$$= \frac{n(n+1)}{2} \left[ \frac{n(n+1)}{2} + 2 \frac{(2n+1)}{3} + 1 \right]$$

$$= \frac{n(n+1)}{12} [3n^2 + 3n + 8n + 4 + 6]$$

$$= \frac{n(n+1)}{12} [3n^2 + 11n + 10]$$

$$= \frac{n(n+1)}{12} [(3n+5)(n+2)]$$

Consider Denominator  $= 1^2 \times 2 + 2^2 \times 3 + 3^2 \times 4 + \dots$  upto the  $n$ th term

The  $n$ th term is  $n^2 (n+1)$

$\therefore 1^2 \times 2 + 2^2 \times 3 + 3^2 \times 4 + \dots$  upto the  $n$ th term

$$= \sum n^2 (n+1)$$

$$= \sum (n^3 + n^2) = \sum n^3 + \sum n^2$$

$$= \left[ \frac{n(n+1)}{2} \right]^2 + \frac{n(n+1)(2n+1)}{6}$$

$$= \frac{n(n+1)}{2} \left[ \frac{n(n+1)}{2} + \frac{(2n+1)}{3} \right]$$

$$= \frac{n(n+1)}{12} [3n^2 + 3n + 4n + 2]$$

$$= \frac{n(n+1)}{12} [(3n+1)(n+2)]$$

$$\text{The given expression} = \frac{\frac{n(n+1)}{12} [(3n+5)(n+2)]}{\frac{n(n+1)}{12} [(3n+1)(n+2)]} = \frac{3n+5}{3n+1}$$