

p-BLOCK ELEMENT

GROUP 13 ELEMENTS: THE BORON FAMILY :

Boron is a typical non-metal, aluminium is a metal but shows many chemical similarities to boron, and gallium, indium and thallium are almost exclusively metallic in character.

OCCURANCE

Boron : Boron is a fairly rare element, mainly occurs as orthoboric acid, (H_3BO_3) , borax, $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$, and kernite, $\text{Na}_2\text{B}_4\text{O}_7 \cdot 4\text{H}_2\text{O}$. There are two isotopic forms of boron ^{10}B (19%) and ^{11}B (81%).

ALUMINIUM :

Aluminium is the most abundant metal and the third most abundant element in the earth's crust (8.3% by mass) after oxygen (45.5%) and Si (27.7%). Bauxite, $\text{Al}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$ and cryolite, Na_3AlF_6 are the important minerals of aluminium.

ELECTRONIC CONFIGURATION :

The outer electronic configuration of these elements is ns^2np^1 . A close look at the electronic configuration suggests that while boron and aluminium have noble gas core, gallium and indium have noble gas plus 10 d -electrons, and thallium has noble gas plus 14 f -electrons plus 10 d -electron cores. Thus, the electronic structures of these elements are more complex than for the first two groups of elements discussed in unit 10. This difference in electronic structures affects the other properties and consequently the chemistry of all the elements of this group.

ATOMIC RADII :

On moving down the group, for each successive member one extra shell of electrons is added and, therefore, atomic radius is expected to increase. However, a deviation can be seen.

Atomic radii order

$\text{B} < \text{Ga} < \text{Al} < \text{In} < \text{Tl}$

Ionic Radii order (+3 OS)

$\text{B} < \text{Al} < \text{Ga} < \text{In} < \text{Tl}$

Atomic radius of Ga is less than that of Al. This can be understood from the variation in the inner core of the electronic configuration. The presence of additional 10 d -electrons offer only poor screening effect for the outer electrons from the increased nuclear charge in gallium. Consequently, the atomic radius of gallium (135 pm) is less than that of aluminium (143 pm).

Ionization Enthalpy

The ionisation enthalpy values as expected from the general trends do not decrease smoothly down the group.

Ionization Enthalpies order

$\text{B} > \text{Tl} > \text{Ga} > \text{Al} > \text{In}$

The decrease from B to Al is associated with increase in size. The observed discontinuity in the ionisation enthalpy values between Al and Ga, and between In and Tl are due to inability of d - and f -electrons, which have low screening effect, to compensate the increase in nuclear charge.

The order of ionisation enthalpies, as expected, is $\Delta_1 H_1 < \Delta_1 H_2 < \Delta_1 H_3$. The sum of the first three ionisation enthalpies for each of the elements is very high. Effect of this will be apparent during study their chemical properties.

Electronegativity

Down the group, electronegativity first decreases from B to Al and then increases marginally. This is because of the discrepancies in atomic size of the elements.

Physical Properties

- (i) Boron is non-metallic in nature. It is extremely hard and black coloured solid. It exists in many allotropic forms. Due to very strong crystalline lattice, boron has unusually high melting point.
- (ii) Rest of the members are soft metals with low melting point and high electrical conductivity.
- (iii) It is worth while to note that gallium with unusually low melting point (303K), could exist in liquid state during summer. Its high boiling point (2676 K) makes it a useful material for measuring high temperatures.
- (iv) Density of the elements increases down the group from boron to thallium.

Melting and Boiling points order

M.P. $B > Al > Tl > In > Ga$


B.P. $B > Al > Ga > In > Tl$

Electropositive Character

Due to high IE they are less electropositive on moving down the group metallic character increases due to decrease in IE [$\therefore$ B is nonmetals and other elements are metals.]

$B <$	$Al >$	$Ga >$	$In >$	Tl
Non metal		Metals		

Note : Boron exists in many allotropic forms. All the allotropes have basic building B_{12} icosahedral units made up of polyhedron having 20 faces and 12 corners. For example one is the simplest form α - rhombohedral boron.




But Al, In & Tl all have close packed metal structure.

Chemical Properties


Oxidation state and trends in chemical reactivity

- (i) Due to small size of boron, the sum of its first three ionization enthalpies is very high. This prevents it to form +3 ions and forces it to form only covalent compounds. But as we move from B to Al, the sum of the first three ionisation enthalpies of Al considerably decreases, and is therefore able to form Al^{3+} ions. In fact, aluminium is a highly electropositive metal.

- (ii) However, down the group, due to poor shielding effect of intervening d and f orbitals, the increased effective nuclear charge holds ns electrons tightly (responsible for inert pair effect) and thereby, restricting their participation in bonding. As a result of this, only p-orbital electrons may be involved in bonding. In fact in Ga, In and Tl, both +1 and +3 oxidation states are observed. The relative stability of +1 oxidation state progressively increases for heavier elements: $\text{Al} < \text{Ga} < \text{In} < \text{Tl}$. In thallium +1 oxidation state is predominant whereas the +3 oxidation state is highly oxidising in character.
- (iii) The compounds in +1 oxidation state, as expected from energy considerations, are more ionic than those in +3 oxidation state.
- (iv) In trivalent state, the number of electrons around the central atom in a molecule of the compounds of these elements (e.g., boron in BF_3) will be only six. Such **electron deficient** molecules have tendency to accept a pair of electrons to achieve stable electronic configuration and thus, behave as Lewis acids. The tendency to behave as Lewis acid decreases with the increase in the size down the group. BCl_3 easily accepts a lone pair of electrons from ammonia to form $\text{BCl}_3 \cdot \text{NH}_3$.





AlCl_3 achieves stability by forming a dimer



- (v) In trivalent state most of the compounds being covalent are hydrolysed in water. For example, the trichlorides on hydrolysis in water form tetrahedral $[\text{M}(\text{OH})_4]^-$ species; the hybridisation state of element M is sp^3 . Aluminium chloride in acidified aqueous solution forms octahedral $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$ ion. In this complex ion, the 3d orbitals of Al are involved and the hybridisation state of Al is sp^3d^2 .

REACTIVITY TOWARDS AIR :

- (i) Boron is unreactive in crystalline form.
- (ii) Aluminium forms a very thin oxide layer on the surface which protects the metal from further attack.
- (iii) Amorphous boron and aluminium metal on heating in air form B_2O_3 and Al_2O_3 respectively. With dinitrogen at high temperature they form nitrides.



(E = element)

The nature of these oxides varies down the group. Boron trioxide is acidic and reacts with basic (metallic) oxides forming metal borates. Aluminium and gallium oxides are amphoteric and those of indium and thallium are basic in their properties.

Reaction with Air at Water

Al should react air to form a very thin oxide film (10^{-4} to 10^{-6} mm thick) on the surface and protects the metal from further attack




Ga and In are attacked neither by cold water nor hot water unless oxygen is present. Tl form an oxide on surface.


REACTIVITY TOWARDS ACIDS AND ALKALIES :

Boron does not react with acids and alkalies even at moderate temperature; but aluminium dissolves in mineral acids and aqueous alkalies and thus shows amphoteric character.

Aluminium dissolves in dilute HCl and liberates dihydrogen.




However, concentrated nitric acid renders aluminium passive by forming a protective oxide layer on the surface. Aluminium also reacts with aqueous alkali and liberates dihydrogen.



Sodium tetrahydroxaluminate(III)

or



Ga, In, Tl dissolve in dilute acids liberating H_2 . Ga is amphoteric like Al and it dissolves in aq. NaOH liberating H_2 and forming gallates.


Reactivity towards halogens :

These elements react with halogens to form trihalides (except TlI_3).

**IMPORTANT TRENDS AND ANOMALOUS PROPERTIES OF BORON**

Certain important trends can be observed in the chemical behaviour of group 13 elements. The tri-chlorides, bromides and iodides of all these elements being covalent in nature are hydrolysed in water. Species like tetrahedral $[\text{M}(\text{OH})_4]^{-}$ and octahedral $[\text{M}(\text{H}_2\text{O})_6]^{3+}$, except in boron, exist in aqueous medium.



The monomeric trihalides, being electron deficient, are strong Lewis acids. Boron trifluoride easily reacts with Lewis bases such as NH_3 to complete octet around boron.




It is due to the absence of d orbitals that the maximum covalence of B is 4. Since the d orbitals are available with Al and other elements, the maximum covalence can be expected beyond 4. Most of the other metal halides (*e.g.*, AlCl_3) are dimerised through halogen bridging (*e.g.*, Al_2Cl_6). The metal species completes its octet by accepting electrons from halogen in these halogen bridged molecules.

Preparation of Boron :


- (i) Preparation of
- B_2O_3
- from Borax or Colemanite






- (ii) Reduction of
- B_2O_3

**Chemical Properties :**

- (i) Burning in air:
- $4B + 3O_2 \longrightarrow 2B_2O_3$








- (ii) Reaction with water




(red hot)

- (iii)
- $B + HCl \longrightarrow$
- no reaction



- (iv)
- $2B + 2NaOH + 2H_2O \xrightarrow{\Delta} 2NaBO_2 + 3H_2$




- (v)
- $2B + N_2 \longrightarrow 2BN$
- $(2Al + N_2 \longrightarrow 2AlN)$





- (vi)
- $3Mg + 2B \longrightarrow Mg_3B_2$

SOME IMPORTANT COMPOUNDS OF BORON


Some useful compounds of boron are borax, orthoboric acid and diborane. We will briefly study their chemistry.

Preparation of Borax :**Borax**

Filtered
– CaCO_3 (as residue)


**Properties :**

- (i) It is a white crystalline solid of formula $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$. In fact it contains the tetranuclear units $[\text{B}_4\text{O}_5(\text{OH})_4]^{2-}$ and correct formula, therefore, is $\text{Na}_2[\text{B}_4\text{O}_5(\text{OH})_4] \cdot 8\text{H}_2\text{O}$.
- (ii) Borax dissolves in water to give an alkaline solution.



Orthoboric acid

- (iii) On heating, borax first loses water molecules and swells up. On further heating it turns into a transparent liquid, which solidifies into glass like material known as borax bead.



The metaborates of many transition metals have characteristic colours and, therefore, borax bead test can be used to identify them in the laboratory. For example, when borax is heated in a Bunsen burner flame with CoO on a loop of platinum wire, a blue coloured $\text{Co}(\text{BO}_2)_2$ bead is formed.


Orthoboric acid :**Preparation :**

- (i) It can be prepared by acidifying an aqueous solution of borax.

$$\text{Na}_2\text{B}_4\text{O}_7 + 2\text{HCl} + 5\text{H}_2\text{O} \rightarrow 2\text{NaCl} + 4\text{B}(\text{OH})_3$$
- (ii) It is also formed by the hydrolysis (reaction with water or dilute acid) of most boron compounds (halides, hydrides, etc.)

Property :


- (i) Orthoboric acid, H_3BO_3 is a white crystalline solid, with soapy touch.
- (ii) It is sparingly soluble in water but highly soluble in hot water.
- (iii) H_3BO_3 is soluble in water and behaves as weak monobasic acid. It does not donate protons like most the acids, but rather it accepts OH^- . It therefore is Lewis acid ($\text{B}(\text{OH})_3$)




or



Since B(OH)_3 only partially reacts with water to form H_3O^+ and $[\text{B(OH)}_4]^-$ it behaves as a weak acid. Thus it cannot be titrated satisfactorily with NaOH as a sharp end point is not obtained. If certain polyhydroxy compounds such as glycerol, mannitol or sugar are added to the titration mixture then B(OH)_3 behaves as a strong monobasic acid, and hence can now be titrated with NaOH and end point is diluted using phenolphthalein as indicator.





The added compound must be a cis diol to enhance the acidic properties in this way the cis-diol forms very stable complexes with $[\text{B(OH)}_4]^-$ formed in forward direction above, thus effectively removing it from solution. Hence reaction proceeds in forward direction (Le-Chatelier principle.)



On heating, orthoboric acid above 370K forms metaboric acid, HBO_2 which on further heating yields boric oxide, B_2O_3 .


□ Heating of boric acid :



Sodium peroxy borate used in washing powder as brightner

STRUCTURE


It has a layer structure in which planar BO_3 units are joined by hydrogen bonds as shown in figure.



Structure of boric acid; the dotted lines represent hydrogen bonds

Structure & bonding in diborane :

The structure of diborane is shown in figure. The four terminal hydrogen atoms and the two boron atoms lie in one plane. Above and below this plane, there are two bridging hydrogen atoms. The four terminal B–H bonds are regular two centre-two electron bonds while the two bridge (B–H–B) bonds are different and can be described in terms of three centre–two electron bonds shown in figure.



The structure of diborane, B_2H_6




Figure: Bonding in diborane.




Each B atom uses sp^3 hybrid orbitals for bonding. Out of the four sp^3 hybrid orbital on each B atom, one is without an electron shown in broken lines. The terminal B–H bonds are normal 2-centre-2-electron bonds but the two bridge bonds are 3-centre-2-electron bonds. The 3-centre-2-electron bridge bonds are also referred to as banana bonds.

Note: Metal hydrido borates : Boron also forms a series of hydridoborates; the most important one is the tetrahedral $[BH_4]^-$ ion. Tetrahydridoborates of several metals are known. Lithium and sodium tetrahydridoborates, also known as borohydrides, are prepared by the reaction of metal hydrides with B_2H_6 in diethyl ether.




Both $LiBH_4$ and $NaBH_4$ are used as reducing agents in organic synthesis. They are useful starting materials for preparing other metal borohydrides.


❑ **Al_2O_3 preparation :**




- Uses:** (i) In making refractory bricks
 (ii) as an abrasive
 (iii) To make high alumina cement

❑ **$AlCl_3$ preparation :**



**Properties :**

- (i) Its anhydrous form is deliquescent and fumes in air.
- (ii) It sublimes at $180^\circ C$.
- (iii) It is covalent and exists in the form of dimer even if in non polar solvents e.g. alcohol, ether, benzene, where it is soluble in fair extent.



Uses: (i) Friedel-Craft reaction

(ii) Dyeing, drug & perfumes etc.

□ **Alums :** $M_2SO_4 \cdot M'_2(SO_4)_3 \cdot 24 H_2O$

Properties: Swelling characteristics

where $M = Na^+, K^+, Rb^+, Cs^+, Ag^+, Tl^+, NH_4^+$ (except Li^+)

$M' = Al^{+3}, Cr^{+3}, Fe^{+3}, Mn^{+3}, Co^{+3}$

$K_2SO_4 \cdot Al_2(SO_4)_3 \cdot 24H_2O$ Potash alum

$(NH_4)_2SO_4 \cdot Al_2(SO_4)_3 \cdot 24H_2O$ Ammonium alum

$K_2SO_4 \cdot Cr_2(SO_4)_3 \cdot 24H_2O$ Chrome alum

$(NH_4)_2SO_4 \cdot Fe_2(SO_4)_3 \cdot 24H_2O$ Ferric alum

Preparation: $Al_2O_3 + 3H_2SO_4 \longrightarrow Al_2(SO_4)_3 + 3H_2O$

$Al_2(SO_4)_3 + K_2SO_4 + aq. sol^n \longrightarrow$ crystallise

Uses: (i) Act as coagulant

(ii) Purification of water

(iii) Tanning of leather

(iv) Mordant in dyeing

(v) Antiseptic

USES OF BORON AND ALUMINIUM AND THEIR COMPOUNDS

Boron :

- (i) Boron being extremely hard refractory solid of high melting point, low density and very low electrical conductivity, finds many applications.
- (ii) Boron fibres are used in making bullet-proof vest and light composite material for aircraft.
- (iii) The boron-10 (^{10}B) isotope has high ability to absorb neutrons and, therefore, metal borides are used in nuclear industry as protective shields and control rods.
- (iv) The main industrial application of borax and boric acid is in the manufacture of heat resistant glasses(e.g., Pyrex), glass-wool and fibreglass.
- (v) Borax is also used as a flux for soldering metals, for heat, scratch and stain resistant glazed coating to earthenwares and as constituent of medicinal soaps.
- (vi) An aqueous solution of orthoboric acid is generally used as a mild antiseptic.

Aluminium :

- (i) Aluminium is a bright silvery-white metal, with high tensile strength.
- (ii) It has a high electrical and thermal conductivity.
- (iii) On a weight-to-weight basis, the electrical conductivity of aluminium is twice that of copper.
- (iv) Aluminium is used extensively in industry and every day life.
- (v) It forms alloys with Cu, Mn, Mg, Si and Zn.
- (vi) Aluminium and its alloys can be given shapes of pipe, tubes, rods, wires, plates or foils and, therefore, find uses in packing, utensil making, construction, aeroplane and transportation industry.
- (vii) The use of aluminium and its compounds for domestic purposes is now reduced considerably because of their toxic nature.

GROUP 14 ELEMENTS :**The carbon family**

Carbon (C), silicon (Si), germanium (Ge), tin (Sn) and lead (Pb) are the members of group 14.

❑ Occurrence of element

- (i) **Carbon** : Carbon is the seventeenth most abundant element by mass in the earth's crust. Naturally occurring carbon contains two stable isotopes: ^{12}C and ^{13}C . In addition to these, third isotope, ^{14}C is also present. It is a radioactive isotope with half-life 5770 years and used for radiocarbon dating.
- (ii) **Silicon** : Silicon is the second (27.7 % by mass) most abundant element on the earth's crust and is present in nature in the form of silica and silicates. Silicon is a very important component of ceramics, glass and cement.
- (iii) **Germanium** : Germanium exists only in traces.
- (iv) **Tin** : Tin occurs mainly as cassiterite, SnO_2
- (v) **Lead** : Lead as galena, PbS .

Note : Ultrapure form of germanium and silicon are used to make transistors and semiconductor devices.

❑ Electronic Configuration

The valence shell electronic configuration of these elements is ns^2np^2 . The inner core of the electronic configuration of elements in this group also differs.

❑ Covalent Radius

There is a considerable increase in covalent radius from C to Si, thereafter from Si to Pb a small increase in radius is observed. This is due to the presence of completely filled d and f orbitals in heavier members.

Covalent radii : $\text{C} < \text{Si} < \text{Ge} < \text{Sn} < \text{Pb}$

❑ Ionization Enthalpy

The first ionization enthalpy of group 14 members is higher than the corresponding members of group 13. The influence of inner core electrons is visible here also. In general the ionisation enthalpy decreases down the group. Small decrease in $\Delta_i H$ from Si to Ge, Ge to Sn and slight increase in $\Delta_i H$ from Sn to Pb is the consequence of poor shielding effect of intervening d and f orbitals and increase in size of the atom.

$\text{C} > \text{Si} > \text{Ge} > \text{Pb} > \text{Sn}$ (IE_1 values)

❑ Melting and Boiling Points

M.P. : $C > Si > Ge > Pb > Sn$

B.P. : $Si > Ge > Sn > Pb$

❑ Electronegativity

Due to small size, the elements of this group are slightly more electronegative than group 13 elements. The electronegativity values for elements from Si to Pb are almost the same.

❑ Physical Properties

All group 14 members are solids. Carbon and silicon are non-metals, germanium is a metalloid, whereas tin and lead are soft metals with low melting points. Melting points and boiling points of group 14 elements are much higher than those of corresponding elements of group 13.

❑ Chemical Properties***Oxidation states and trends in chemical reactivity***


- (i) The group 14 elements have four electrons in outermost shell.
- (ii) The common oxidation states exhibited by these elements are +4 and +2. Carbon also exhibits negative oxidation states. Since the sum of the first four ionization enthalpies is very high, compounds in +4 oxidation state are generally covalent in nature.
- (iii) In heavier members the tendency to show +2 oxidation state increases in the sequence $Ge < Sn < Pb$. It is due to the inability of ns^2 electrons of valence shell to participate in bonding. The relative stabilities of these two oxidation states vary down the group.
- (iv) Carbon and silicon mostly show +4 oxidation state.
- (v) Germanium forms stable compounds in +4 state and only few compounds in +2 state.
- (vi) Tin forms compounds in both oxidation states (Sn in +2 state is a reducing agent).
- (vii) Lead compounds in +2 state are stable and in +4 state are strong oxidising agents.
- (viii) In tetravalent state the number of electrons around the central atom in a molecule (*e.g.*, carbon in CCl_4) is eight. Being *electron precise* molecules, they are normally not expected to act as electron acceptor or electron donor species. Although carbon cannot exceed its covalence more than 4, other elements of the group can do so. It is because of the presence of *d* orbital in them. Due to this, their halides undergo hydrolysis and have tendency to form complexes by accepting electron pairs from donor species. For example, the species like, SiF_6^{2-} , $[GeCl_6]^{2-}$, $[Sn(OH)_6]^{2-}$ exist where the hybridisation of the central atom is sp^3d^2 .

❑ Reactivity towards oxygen

All members when heated in oxygen form oxides. There are mainly two types of oxides, *i.e.*, monoxide and dioxide of formula MO and MO_2 respectively. SiO only exists at high temperature. Oxides in higher oxidation states of elements are generally more acidic than those in lower oxidation states. The dioxides — CO_2 , SiO_2 and GeO_2 are acidic, whereas SnO_2 and PbO_2 are amphoteric in nature. Among monoxides, CO is neutral, GeO is distinctly acidic whereas SnO and PbO are amphoteric.

❑ Reactivity towards water

Carbon, silicon and germanium are not affected by water. Tin decomposes steam to form dioxide and dihydrogen gas.




Lead is unaffected by water, probably because of a protective oxide film formation.

❑ Reactivity towards halogen

- (i) These elements can form halides of formula MX_2 and MX_4 (where $\text{X} = \text{F}, \text{Cl}, \text{Br}, \text{I}$). Except carbon, all other members react directly with halogen under suitable condition to make halides.
- (ii) Most of the MX_4 are covalent in nature. The central metal atom in these halides undergoes sp^3 hybridisation and the molecule is tetrahedral in shape. Exceptions are SnF_4 and PbF_4 , which are ionic in nature.
- (iii) PbI_4 does not exist because Pb—I bond initially formed during the reaction does not release enough energy to unpair $6s^2$ electrons and excite one of them to higher orbital to have four unpaired electrons around lead atom.
- (iv) Heavier members Ge to Pb are able to make halides of formula MX_2 .
- (v) Stability of dihalides increases down the group. Considering the thermal and chemical stability, GeX_4 is more stable than GeX_2 , whereas PbX_2 is more than PbX_4 .
- (vi) Except CCl_4 , other tetrachlorides are easily hydrolysed by water because the central atom can accommodate the lone pair of electrons from oxygen atom of water molecule in d orbital.

Hydrolysis can be understood by taking the example of SiCl_4 . It undergoes hydrolysis by initially accepting lone pair of electrons from water molecule in d orbitals of Si, finally leading to the formation of Si(OH)_4 as shown below :



IMPORTANT TRENDS AND ANOMALOUS BEHAVIOUR OF CARBON

Like first member of other groups, carbon also differs from rest of the members of its group. It is due to its smaller size, higher electronegativity, higher ionisation enthalpy unavailability of d orbitals. In carbon, only s and p orbitals are available for bonding and, therefore, it can accommodate only four pairs of electrons around it. This would limit the maximum covalence to four whereas other members can expand their covalence due to the presence of d orbitals.

Carbon also has unique ability to form $\text{p}_\pi-\text{p}_\pi$ multiple bonds with itself and with other atoms of small size and high electronegativity. Few examples of multiple bonding are: $\text{C}=\text{C}$, $\text{C}\equiv\text{C}$, $\text{C}=\text{O}$, $\text{C}=\text{S}$ and $\text{C}\equiv\text{N}$. Heavier elements do not form $\text{p}_\pi-\text{p}_\pi$ bonds because their atomic orbitals are too large and diffuse to have effective overlapping.

Catenation Property

Carbon atoms have the tendency to link with one another through covalent bonds to form chains and rings. This property is called **catenation**. This is because C—C bonds are very strong. Down the group the size increases and electronegativity decreases, and, thereby, tendency to show catenation decreases. This can be clearly seen from bond enthalpies values. The order of catenation is $C \gg Si > Ge \approx Sn$. Lead does not show catenation.

Bond	Bond enthalpy / kJ mol^{-1}
C—C	348
Si—Si	297
Ge—Ge	260
Sn—Sn	240

Due to property of catenation and $p_\pi - p_\pi$ bond formation, carbon is able to show allotropic forms.

ALLOTROPES OF CARBON

Carbon exhibits many allotropic forms; both crystalline as well as amorphous. Diamond and graphite are two well-known crystalline forms of carbon. In 1985, third form of carbon known as **fullerenes** was discovered by H.W. Kroto, E. Smalley and R.F. Curl. For this discovery they were awarded the Nobel Prize in 1996.

SOME IMPORTANT COMPOUNDS OF CARBON AND SILICON




Types of Carbide

(i) Ionic and salt like:



Classification on basis of
no. of carbon atoms
present in hydrocarbon
found on their hydrolysis

{	(a) C_1 unit (b) C_2 unit (c) C_3 unit
---	--


C_1 unit:



C_2 unit:



C_3 unit:





(ii) Covalent carbide : SiC & B_4C

(iii) Interstitial carbide :

(Transition element or inner transitional elements forms this kind of carbide)

Interstitial carbide formation doesn't affect the metallic lustre and electrical conductivity. ($\because$ no chemical bond is present, no change in property)

SiC (Carborundum)**Preparation**

Note :

(i) SiC has diamond like or wurtzite structure

(ii) SiC is often dark purple, black or dark green due to traces of Fe and other impurities but pure sample are pale yellow to colourless.

Properties


(i) It is very hard and is used in cutting tools and abrasive powder (polishing material)

(ii) It is very much inert

(iii) It is not being affected by any acid except H_3PO_4

Carbon Monoxide**Preparation :**


(i) Direct oxidation of C in limited supply of oxygen or air yields carbon monoxide.




(ii) On small scale pure CO is prepared by dehydration of formic acid with concentrated H_2SO_4 at 373 K



(iii) On commercial scale it is prepared by the passage of steam over hot coke. The mixture of CO and H_2 thus produced is known as **water gas** or **synthesis gas**.








When air is used instead of steam, a mixture of CO and N_2 is produced, which is called **producer gas**.





Water gas and producer gas are very important industrial fuels. Carbon monoxide in water gas or producer gas can undergo further combustion forming carbon dioxide with the liberation of heat.

(iv) By heating potassium ferrocyanide with conc. H_2SO_4 : When potassium ferrocyanide in powdered state is heated with concentrated H_2SO_4 , CO is evolved. Dilute H_2SO_4 should never be used because it shall evolve highly poisonous gas HCN.







Properties :

- (i) Carbon monoxide is a colourless, odourless and almost water insoluble gas.
- (ii) It is a powerful reducing agent and reduces almost all metal oxides other than those of the alkali and alkaline earth metals, aluminium and a few transition metals. This property of CO is used in the extraction of many metals from their oxides ores.

**DETECTION**

(a) burns with blue flame

(b) CO is passed through PdCl_2 solution giving rise to black ppt.

**ESTIMATION****ABSORBERS****❑ Bonding in CO mole**



In CO molecule, there are one sigma and two π bonds between carbon and oxygen. Because of the presence of a lone pair on carbon, CO molecule acts as a donor and reacts with certain metals when heated to form **metal carbonyls**.

❑ Poisonous nature of CO


The highly poisonous nature of CO arises because of its ability to form a **complex with haemoglobin**, which is about 300 times more stable than the oxygen-haemoglobin complex. This prevents haemoglobin in the red blood corpuscles from carrying oxygen round the body and ultimately resulting in death.

❑ Carbon Dioxide**Preparation :**

- (i) It is prepared by complete combustion of carbon and carbon containing fuels in excess of air.





- (ii) **Laboratory** by the action of dilute HCl on calcium carbonate.



- (iii) Commercial scale by heating limestone.

Properties :


- (i) It is a colourless and odourless gas.
- (ii) Its low solubility in water makes it of immense biochemical and geo-chemical importance.
- (iii) With water, it forms carbonic acid, H_2CO_3 which is a weak dibasic acid and dissociates in two steps:



$\text{H}_2\text{CO}_3/\text{HCO}_3^-$ buffer system helps to maintain pH of blood between 7.26 to 7.42. Being acidic in nature, it combines with alkalies to form metal carbonates.

Use of CO_2

Carbon dioxide, which is normally present to the extent of $\sim 0.03\%$ by volume in the atmosphere, is removed from it by the process known as **photosynthesis**. It is the process by which green plants convert atmospheric CO_2 into carbohydrates such as glucose. The overall chemical change can be expressed as:




By this process plants make food for themselves as well as for animals and human beings.

Harmful effect of CO_2

Unlike CO, it is not poisonous. But the increase in combustion of fossil fuels and decomposition of limestone for cement manufacture in recent years seem to increase the CO_2 content of the atmosphere. This may lead to increase in **green house effect** and thus, raise the temperature of the atmosphere which might have serious consequences.


- (i) Carbon dioxide can be obtained as a solid in the form of **dry ice** by allowing the liquified CO_2 to expand rapidly and dry ice is used as a refrigerant for ice-cream and frozen food.
- (ii) Gaseous CO_2 is extensively used to carbonate soft drinks. Being heavy and non-supporter of combustion it is used as fire extinguisher.
- (iii) A substantial amount of CO_2 is used to manufacture urea. In CO_2 molecule carbon atom undergoes *sp* hybridisation. Two *sp* hybridised orbitals of carbon atom overlap with two *p* orbitals of oxygen atoms to make two sigma bonds while other two electrons of carbon atom are involved in $p_\pi-p_\pi$ bonding with oxygen atom. This results in its linear shape [with both C–O bonds of equal length (115 pm)] with no dipole moment. The resonance structures are shown below:



Resonating structures of carbon dioxide

Note : Carbongene has 95% O₂ and 5% CO₂ and is used as an antidote for poisoning of CO.

Teflon $-(\text{CF}_2 - \text{CF}_2)_n$



Purpose

Temperature with standing capacity upto 300°C (1st organic compound withstand this kind of high temperature)

SILICON (Si)

Occurrence


Silicon is the second most abundant (27.2%) element after oxygen (45.5%) in the earth's crust. It does not occur free in nature but in the combined state, it occurs widely in form of silica and silicates. All mineral rocks, clays and soils are built of silicates of magnesium, aluminium, potassium or iron. Aluminium silicate is however the most common constituent of rocks and clays.

Silica is found in the free state in sand, flint and quartz and in the combined state as silicates like



- (i) Feldspar – K₂O. Al₂O₃. 6SiO₂
- (ii) Kaolinite – Al₂O₃. 2SiO₂. 2H₂O
- (iii) Asbestos – CaO. 3MgO. 4SiO₂

Preparation

- (i) From silica (sand): Elemental silicon is obtained by the reduction of silica (SiO₂) with high purity coke in an electric furnace.



- (ii) From silicon tetrachloride (SiCl₄) or silicon chloroform (SiHCl₃) : Silicon of very high purity required for making semiconductors is obtained by reduction of highly purified silicon tetrachloride or silicon chloroform with dihydrogen followed by purification by zone refining.





Physical Properties :

- (i) Elemental silicon is very hard having diamond like structure.
- (ii) It has shining luster with a melting point of 1793 K and boiling point of about 3550 K.
- (iii) Silicon exists in three isotopes, i.e. $^{28}_{14}\text{Si}$, $^{29}_{14}\text{Si}$ and $^{30}_{14}\text{Si}$ but $^{28}_{14}\text{Si}$ is the most common isotope.


Chemical Properties :

Silicon is particularly unreactive at room temperature towards most of the elements except fluorine. Some important chemical reactions of silicon are discussed below.


- (i) **Action of air :** Silicon reacts with oxygen of air at 1173 K to form silicon dioxide and with nitrogen of air at 1673 K to form silicon nitride,.



(ii) **Action of steam** : It is slowly attacked by steam when heated to redness liberating dihydrogen gas.




(iii) **Reaction with halogens**: It burns spontaneously in fluorine gas at room temperature to form silicon tetrafluoride (SiF_4).



However, with other halogens, it combines at high temperatures forming tetrahalides.

(iv) **Reaction with carbon** : Silicon combines with carbon at 2500°C forming silicon carbide (SiC) known as carborundum.




Carborundum is an extremely hard substance next only to diamond. It is mainly used as an abrasive and as a refractory material.

Uses :

- (i) Silicon is added to steel as such or more usually in form of ferrosilicon (an alloy of Fe and Si) to make it acid-resistant.
- (ii) High purity silicon is used as semiconductors in electronic devices such as transistors.
- (iii) It is used in the preparation of alloys such as silicon-bronze, magnesium silicon bronze and ferrosilicon.


Compounds of Silicon :

Silane :



Only these two are found

Higher molecules are not formed. $\therefore$ Si can't show catenation property




Silicones


It is an organosilicon polymer

TYPES OF SILICONES :

(i) Linear silicones




Linear silicone




(ii) Cyclic silicones


Silicones may have the cyclic structure also having 3, 4, 5 and 6 nos. of silicon atoms within the ring. Alcohol analogue of silicon is known as silanol



cyclic silicone not planar


(iii) Dimer silicones

Note



This end of the chain can't be extended hence R_3SiCl is called as chain stopping unit

* Using R_3SiCl in a certain proportion we can control the chain length of the polymer

(iv) Crossed linked silicones

cross linked silicone

3 dimensional network

It provides the crosslinking among the chain making the polymer more hard and hence controlling the proportion of RSiCl_3 we can control the hardness of polymer.

Uses

- (1) It can be used as electrical insulator (due to inertness of Si-O-Si bonds)
- (2) It is used as water repellent ($\because$ surface is covered) eg. car polish, shoe polish, masonry works in buildings
- (3) It is used as antifoaming agent in sewage disposal, beer making and in cooking oil used to prepare potato chips.
- (4) As a lubricant in the gear boxes and light weight machinery

SILICA (SiO₂)


Occurrence :

Silica or silicon dioxide occurs in nature in the free state as sand, quartz and flint and in the combined state as silicates like, Feldspar : K₂O.Al₂O₃.6SiO₂, Kaolinite : Al₂O₃. 2SiO₂. 2H₂O etc.




Properties :

- (i) Pure silica is colourless, but sand is usually coloured yellow or brown due to the presence of ferric oxide as an impurity.
- (ii) Silicon dioxide is insoluble in water and all acids except hydrofluoric acid.

$$\text{SiO}_2(\text{s}) + 4\text{HF}(\text{l}) \longrightarrow \text{SiF}_4(\text{l}) + 2\text{H}_2\text{O}(\text{l})$$
- (iii) It also combines with metallic oxides at high temperature giving silicates e.g.




- (iv) When silica is heated strongly with metallic salts, silicates are formed and the volatile oxides are driven off as vapours.



The first two examples quoted here are important in glass making.

Structures of Silica :

Silica has a three-dimensional network structure. In silica, silicon is sp³-hybridized and is thus linked to four oxygen atoms and each oxygen atom is linked to two silicon atoms forming a three-dimensional giant molecule as shown in figure. This three-dimensional network structure imparts stability to SiO₂ crystal and hence a large amount of energy is required to break the crystal resulting in high melting point.



Uses :

- (i) Sand is used in large quantities to make mortar and cement.
- (ii) Being transparent to ultraviolet light, large crystal of quartz are used for making lenses for optical instruments and for controlling the frequency of radio-transmitters.
- (iii) Powdered quartz is used for making silica bricks.
- (iv) Silica gel (SiO₂.xH₂O) is used as a desiccant (for absorbing moisture) and as an adsorbent in chromatography.

Quartz



Quartz is extensively used as a piezoelectric material; it has made possible to develop extremely accurate clocks, modern radio and television broadcasting and mobile radio communications. Silica gel is used as a drying agent and as a support for chromatographic materials and catalysts. Kieselghur, an amorphous form of silica is used in filtration plants.

Silicates



A large number of silicates minerals exist in nature. Some of the examples are feldspar, zeolites, mica and asbestos. Two important man-made silicates are glass and cement.

Zeolites

If aluminium atoms replace few silicon atoms in three-dimensional network of silicon dioxide, overall structure known as aluminosilicate, acquires a negative charge. Cations such as Na^+ , K^+ or Ca^{2+} balance the negative charge. Examples are feldspar and zeolites. Zeolites are widely used as a catalyst in petrochemical industries for cracking of hydrocarbons and isomerisation, e.g., ZSM-5 (A type of zeolite) used to convert alcohols directly into gasoline. Hydrated zeolites are used as ion exchangers in softening of “hard” water.


TIN & ITS COMPOUND

(iii) Reaction with acid.






or




KOH [In absence of air K_2SnO_2 forms and in contact with air it readily converts into K_2SnO_3]




Both are amphoteric in nature :







But conc. hot alkali behaves differently.








(Soluble only in hot conc. H_2SO_4)








SnCl_2 & SnCl_4 :






(2) A piece of Sn is always added to preserve a solution of SnCl_2 . Explain.



(4) Reducing Properties of SnCl_2 :




(5) Readily combines with $\text{I}_2 \Rightarrow \text{SnCl}_2\text{I}_2 \Rightarrow$ This reaction is used to estimate tin.

Formation of SnCl_4 :









* $\text{SnCl}_4 \cdot 5\text{H}_2\text{O}$ is known as butter of tin $\Rightarrow$ used as mordant.

$(\text{NH}_4)_2\text{SnCl}_6$ is known as 'pink's salt' $\Rightarrow$ used in calico printing.


Mosaic gold : SnS_2 yellow crystalline substance :







Note : Mosaic gold used for filling purpose (in joining gold pieces)

Distinction of Sn^{+2} / Sn^{+4} :**COMPOUNDS OF LEAD****Oxides of lead :****Laboratory Prepⁿ. :**

$\left\{ \begin{array}{l} \text{PbO, hot oxide} \\ \text{easily reduced to Pb by} \\ \text{H}_2 \text{ or C.} \end{array} \right.$



**Preparation of Pb_2O_3 :**

This reaction suggests that Pb_2O_3 contains PbO_2 .






(3) **PbO₂** : Insoluble in water :

HNO₃, But reacts with HCl and H₂SO₄(hot conc.) but does not react with HNO₃ and soluble in hot NaOH / KOH.






↓


Excess bleaching powder
is being removed by stirring with
HNO₃




PbO₂ : Powerful oxidising agent :




PbCl₄ : Exists as H₂[PbCl₆]



{ice cold conc. saturated with Cl₂}



TetraEthyl lead :



It is antiknocking agent.

NITROGEN FAMILY

GROUP-15 ELEMENTS (N, P, As, Sb, Bi)

- (i) As we go down the group, there is a shift from non-metallic to metallic through metalloidic character.
- (ii) Nitrogen and phosphorus are **non-metals**, arsenic and antimony **metalloids** and bismuth is a **typical metal**.

Occurrence :

Nitrogen : Molecular nitrogen comprises 78% by volume of the atmosphere. It is 33rd most abundant element in the earth's crust. In the earth's crust, it occurs as sodium nitrate, NaNO_3 (called Chile saltpetre) and potassium nitrate (Indian saltpetre). It is found in the form of proteins and nucleic acid in plants and animals.

Phosphorus :

- (i) It is eleventh most abundant element in earth's crust occurs in minerals of the apatite family, $\text{Ca}_9(\text{PO}_4)_6 \cdot \text{CaX}_2$ ($\text{X} = \text{F}, \text{Cl}$ or OH) (e.g., fluorapatite $\text{Ca}_9(\text{PO}_4)_6 \cdot \text{CaF}_2$) and also found as chlorapatite $\text{Ca}_9(\text{PO}_4)_6 \cdot \text{CaCl}_2$.
- (ii) It is also present in nucleic acid (in DNA and RNA) which are the main components of phosphate rocks.
- (iii) Arsenic, antimony and bismuth are found mainly as sulphide minerals.

Electronic Configuration :

The valence shell electronic configuration of these elements is ns^2np^3 .

Atomic and Ionic Radii :

Covalent radius : $\text{N} < \text{P} < \text{As} < \text{Sb} < \text{Bi}$

Explanation :

Covalent and ionic (in a particular state) radii increase in size down the group. There is a considerable increase in covalent radius from N to P. However, from As to Bi only a small increase in covalent radius is observed. This is due to the presence of completely filled d and/or f orbitals in heavier members.

Ionisation Enthalpy :

$\text{N} > \text{P} > \text{As} > \text{Sb} > \text{Bi}$ (IE_1 values)

Explanation :

Ionisation enthalpy decreases down the group due to gradual increase in atomic size. Because of the extra stable half-filled p orbitals electronic configuration and smaller size, the ionisation enthalpy of the group 15 elements is much greater than that of group 14 elements in the corresponding periods. The order of successive ionisation enthalpies, as expected is $\Delta_1 H_1 < \Delta_1 H_2 < \Delta_1 H_3$ (See above table).

Electronegativity :

$\text{N} > \text{P} > \text{As} > \text{Sb} = \text{Bi}$
(1.9) (1.9)

Explanation :

The electronegativity value, in general, decreases down the group with increasing atomic size. However, amongst the heavier elements, the difference is not that much pronounced.

Metallic Character

$\frac{\text{N} < \text{P}}{\text{Non metal}}$	$\frac{\text{As} <}{\text{Metalloid}}$	$\frac{\text{Sb} < \text{Bi}}{\text{Metals}}$
--	--	---

Physical Properties :

- All the elements of this group are polyatomic. Dinitrogen is a diatomic gas while all others are solids.
- Metallic character increases down the group.
- Nitrogen and phosphorus are non-metals, arsenic and antimony metalloids and bismuth is a metal. This is due to decrease in ionisation enthalpy and increase in atomic size.
- The boiling points, in general, increase from top to bottom in the group but the melting point increases upto arsenic and then decreases upto bismuth.
- Except nitrogen all the elements show allotropy.

P → exists in three allotropic form as white, red and black

As, Sb → exist as yellow and grey

Bi → exist as α , β , γ , δ allotropic form

Catenation

- * The group 15 elements also show catenation property but to much smaller extent than carbon. For example hydrazine (H_2NNH_2) has two N atoms bonded together HN_3 has three N atoms.



- * Among group 15 elements P has the maximum tendency for catenation forming cyclic as well as open chain compounds consisting of many phosphorous atoms.

P_2H_4 has two P atoms bonded together the lesser tendency of elements of group 15 to show catenation in comparison to carbon is their low dissociation enthalpies.

C – C 353.3 kJ /mole

N – N 160.8 kJ / mole

P – P 201.6 kJ / mole

As – As 147.4 kJ / mole


Chemical Properties :**Oxidation states and trends in chemical reactivity**

- The common oxidation states of these elements are –3, +3 and +5.
- The tendency to exhibit –3 oxidation state decreases down the group due to increase in size and metallic character. Bismuth hardly forms any compound in –3 oxidation state.
- The stability of +5 oxidation state decreases down the group. The only well characterised Bi (V) compound is BiF_5 .

- (iv) The stability of +5 oxidation state decreases and that of +3 state increases (due to inert pair effect) down the group.
- (v) Nitrogen exhibits + 1, + 2, + 4 oxidation states also when it reacts with oxygen. Phosphorus also shows +1 and +4 oxidation states in some oxoacids.
- (vi) In the case of nitrogen, all oxidation states from +1 to +4 tend to disproportionate in acid solution. For example,



- (vii) Similarly, in case of phosphorus nearly all intermediate oxidation states disproportionate into +5 and -3 both in alkali and acid.



- (viii) +3 oxidation state in case of arsenic, antimony and bismuth becomes increasingly stable with respect to disproportionation.
- (ix) Nitrogen is restricted to a maximum covalency of 4 since only four (one s and three p) orbitals are available for bonding.
- (x) The heavier elements have vacant d orbitals in the outermost shell which can be used for bonding (covalency) and hence, expand their covalency as in PF_6^- .

Anomalous properties of nitrogen

- (i) Nitrogen has unique ability to form $p_\pi - p_\pi$ multiple bonds with itself and with other elements having small size and high electronegativity (e.g., C, O).
- (ii) Heavier elements of this group do not form $p_\pi - p_\pi$ bonds as their atomic orbitals are so large and diffuse that they cannot have effective overlapping.
- (iii) Nitrogen exists as a diatomic molecule with a triple bond (one σ and two π) between the two atoms. N_2 bond enthalpy ($941.4 \text{ kJ mol}^{-1}$) is very high.
- (iv) Phosphorus, arsenic and antimony form single bonds as P–P, As–As and Sb–Sb while bismuth forms metallic bonds in elemental state.
- (v) The single N–N bond is weaker than the single P–P bond because of high interelectronic repulsion of the non-bonding electrons, owing to the small bond length. As a result the catenation tendency is weaker in nitrogen.
- (vi) Another factor which affects the chemistry of nitrogen is the absence of d orbitals in its valence shell. Besides restricting its covalency to four, nitrogen cannot form $d_\pi - p_\pi$ bond as the heavier elements can e.g., $\text{R}_3\text{P}=\text{O}$ or $\text{R}_3\text{P}=\text{CH}_2$ (R=alkyl group).
- (vii) Phosphorus and arsenic can form $d_\pi - d_\pi$ bond also with transition metals when their compounds like $\text{P}(\text{C}_2\text{H}_5)_3$ and $\text{As}(\text{C}_6\text{H}_5)_3$ act as ligands.

(i) Reactivity towards hydrogen:

All the elements of Group 15 form hydrides of the type EH_3 where $\text{E} = \text{N}, \text{P}, \text{As}, \text{Sb}$ or Bi . Some of the properties of these hydrides are shown in Table. The hydrides show regular gradation in their properties. The stability of hydrides decreases from NH_3 to BiH_3 which can be observed from their bond dissociation enthalpy. Consequently, the reducing character of the hydrides increases. Ammonia is only a mild reducing agent while BiH_3 is the strongest reducing agent amongst all the hydrides. Basicity also decreases in the order $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3 \geq \text{BiH}_3$.

Table : Properties of Hydrides of Group 15 Elements

Property	NH_3	PH_3	AsH_3	SbH_3	BiH_3
Melting point/K	195.2	139.5	156.7	185	—
Boiling point/K	238.5	185.5	210.6	254.6	290
(E–H) distance/pm	101.7	141.9	151.9	170.7	—
HEH angle ($^\circ$)	107.8	93.6	91.8	91.3	—
$\Delta_f H^\ominus / \text{kJ mol}^{-1}$	– 46.1	13.4	66.4	145.1	278
$\Delta_{\text{diss}} H^\ominus (\text{E} - \text{H}) / \text{kJ mol}^{-1}$	389	322	297	255	—

(ii) Reactivity towards oxygen: All these elements form two types of oxides: E_2O_3 and E_2O_5 . The oxide in the higher oxidation state of the element is more acidic than that of lower oxidation state. Their acidic character decreases down the group. The oxides of the type E_2O_3 of nitrogen and phosphorus are purely acidic, that of arsenic and antimony amphoteric and those of bismuth predominantly basic.

(iii) Reactivity towards halogens: These elements react to form two series of halides: EX_3 and EX_5 . Nitrogen does not form pentahalide due to non-availability of the d orbitals in its valence shell. Pentahalides are more covalent than trihalides. All the trihalides of these elements except those of nitrogen are stable. In case of nitrogen, only NF_3 is known to be stable. Trihalides except BiF_3 are predominantly covalent in nature.

(iv) Reactivity towards metals: All these elements react with metals to form their binary compounds exhibiting –3 oxidation state, such as, Ca_3N_2 (calcium nitride) Ca_3P_2 (calcium phosphide), Na_3As (sodium arsenide), Zn_3Sb_2 (zinc antimonide) and Mg_3Bi_2 (magnesium bismuthide).

DINITROGEN**Preparation :****(a) Commercial preparation :**

Dinitrogen is produced **commercially** by the liquefaction and fractional distillation of air. Liquid dinitrogen (b.p. 77.2 K) distils out first leaving behind liquid oxygen (b.p. 90 K).


(b) Laboratory preparation :

(i) Dinitrogen is prepared by treating an aqueous solution of ammonium chloride with sodium nitrite.




Small amounts of NO and HNO_3 are also formed in this reaction; these impurities can be removed by passing the gas through aqueous sulphuric acid containing potassium dichromate.





(ii) Dinitrogen can also be obtained by the thermal decomposition of ammonium dichromate.



Note : Very pure nitrogen can be obtained by the thermal decomposition of sodium or barium azide.






(c) Other preparation




(red,overheated) (Black)

Cl_2 passed into liquid NH_3



In this method conc. of NH_3 should not be lowered down beyond a particular limit.



↓



(Tremendously explosive)

❑ **Physical properties :**


- (i) Dinitrogen is a colourless, odourless, tasteless and non-toxic gas.
- (ii) Nitrogen atom has two stable isotopes: ^{14}N and ^{15}N .
- (iii) It has a very low solubility in water (23.2 cm^3 per litre of water at 273 K and 1 bar pressure)
- (iv) Dinitrogen has low freezing and boiling points.

❑ **Chemical properties**


Reaction with metal : At higher temperatures, it directly combines with some metals to form predominantly ionic nitrides and with non-metals, covalent nitrides. A few typical reactions are:




Reaction with metal : It combines with hydrogen at about 773 K in the presence of a catalyst (Haber's Process) to form ammonia:



Dinitrogen combines with dioxygen only at very high temperature (at about 2000 K) to form nitric oxide, NO.



- * The optimum conditions for the production of ammonia are a pressure of 200×10^5 Pa (about 200 atm), a temperature of ~ 700 K.
- * Use of a catalyst such as iron oxide with small amounts of K_2O and Al_2O_3 to increase the rate of attainment of equilibrium.
- * The flow chart for the production of ammonia is shown in figure. Earlier, iron was used as a catalyst with molybdenum as a promoter.



Flow chart for the manufacture of ammonia


Other preparation :

- Nitrate or nitrite reduction : $NO_3^- / NO_2^- + Zn \text{ or } Al + NaOH \longrightarrow NH_3 + [Zn(OH)_4]^{2-} \text{ or } [Al(OH)_4]^-$
- Metal nitride hydrolysis : $N^{3-} + 3H_2O \longrightarrow NH_3 \uparrow + 3 OH^-$

Properties :

- Ammonia is a colourless gas with a pungent odour.
- Its freezing and boiling points are 198.4 and 239.7 K respectively.
- In the solid and liquid states, it is associated through hydrogen bonds as in the case of water and that accounts for its higher melting and boiling points than expected on the basis of its molecular mass.
- Ammonia gas is highly soluble in water.
- Basic character :**



Its aqueous solution is weakly basic due to the formation of OH^- ions.






It forms ammonium salts with acids, e.g., NH_4Cl , $(NH_4)_2 SO_4$, etc.

As a weak base, it precipitates the hydroxides (hydrated oxides in case of some metals) of many metals from their salt solutions.

For example,






In ammonia molecule the presence of a lone pair of electrons on the nitrogen atom of the makes it a Lewis base. It donates the electron pair and forms linkage with metal ions and the formation of such complex compounds finds applications in detection of metal ions such as Cu^{2+} , Ag^+ :



Other reactions



CH_3COOH is strong acid in liq. NH_3 while in water is weak acid.



Basicity order $\text{NH}_3 > \text{H}_2\text{O}$

more solvation of H^+ in NH_3 .

Hydrolysis and Ammonolysis occurs is a same way.



Rate of hydrolysis and Ammonolysis will be affected by the presence of HCl vapour & NH_4Cl vapour respectively.

Uses :

- (i) Ammonia is used to produce various nitrogenous fertilisers (ammonium nitrate, urea, ammonium phosphate and ammonium sulphate).
- (ii) In the manufacture of some inorganic nitrogen compounds, the most important one being nitric acid.
- (iii) Liquid ammonia is also used as a refrigerant.

BONDING IN AMMONIA :

The ammonia molecule is trigonal pyramidal with the nitrogen atom at the apex. It has three bond pairs and one lone pair of electrons.





OXIDES OF NITROGEN :

Nitrogen forms a number of oxides in different oxidation states. The names, formulas, preparation and physical appearance of these oxides are given in Table.


Oxides of Nitrogen				
Name	Formula	Oxidation state of nitrogen	Common methods of preparation	Physical appearance and chemical nature
Dinitrogen oxide [Nitrogen oxide]	N_2O	+1	$NH_4NO_3 \xrightarrow{\text{Heat}} N_2O + 2H_2O$	Colourless gas, neutral
Nitrogen monoxide [Nitrogen (II) oxide]	NO	+2	$2NaNO_2 + 2FeSO_4 + 3H_2SO_4 \rightarrow Fe_2(SO_4)_3 + 2NaHSO_4 + 2H_2O + 2NO$	Colourless gas, neutral
Dinitrogen trioxide [Nitrogen (III) oxide]	N_2O_3	+3	$2NO + N_2O_4 \xrightarrow{-30^\circ C} 2N_2O_3$	Pale Blue solid (MP = $-100.1^\circ C$), acidic, Intense blue liquid ($-30^\circ C$)
Nitrogen dioxide [Nitrogen (IV) oxide]	NO_2	+4	$2Pb(NO_3)_2 \xrightarrow{673K} 4NO_2 + 2PbO + O_2$	brown gas, acidic
Dinitrogen tetroxide [Nitrogen (IV) oxide]	N_2O_4	+4	$2NO_2 \xrightleftharpoons[\text{Heat}]{\text{Cool}} N_2O_4$	Colourless solid/liquid, acidic
Dinitrogen pentaoxide [Nitrogen(V) oxide]	N_2O_5	+5	$4HNO_3 + P_4O_{10} \rightarrow 4HPO_3 + 2N_2O_5$	colourless solid, acidic

Structure of Oxides of Nitrogen



Formula	Resonance structures	Bond Parameters
N_2O	$\ddot{N} = N = \ddot{O} \leftrightarrow :N \equiv N - \ddot{O}:$	$N-N-O$ 113 pm 119 pm Linear
NO	$:\dot{N} = \ddot{O} \leftrightarrow :\ddot{N} = \dot{O}:$	$N-O$ 115 pm
N_2O_3	$\begin{array}{c} \ddot{O} \\ \parallel \\ :\ddot{O}-N-N=\ddot{O} \\ \parallel \\ \ddot{O} \end{array} \leftrightarrow \begin{array}{c} \ddot{O} \\ \parallel \\ :\ddot{O}-N=N-\ddot{O} \\ \parallel \\ \ddot{O} \end{array}$	$N-O$ 115 pm 114 pm 105° 186 pm 117° 130° 121 pm
NO_2	$\begin{array}{c} \cdot \\ \cdot \\ :\ddot{O}-N=\ddot{O} \\ \cdot \\ \cdot \end{array} \leftrightarrow \begin{array}{c} \cdot \\ \cdot \\ :\ddot{O}=\ddot{N}-\ddot{O} \\ \cdot \\ \cdot \end{array}$	N 120 pm 134° Angular
N_2O_4	$\begin{array}{c} \ddot{O} \\ \parallel \\ :\ddot{O}-N-N=\ddot{O} \\ \parallel \\ \ddot{O} \end{array} \leftrightarrow \begin{array}{c} \ddot{O} \\ \parallel \\ :\ddot{O}-N=N-\ddot{O} \\ \parallel \\ \ddot{O} \end{array}$	N 175 pm 135° Planar
N_2O_5	$\begin{array}{c} \ddot{O} \\ \parallel \\ :\ddot{O}-N-\ddot{O}-N=\ddot{O} \\ \parallel \\ \ddot{O} \end{array} \leftrightarrow \begin{array}{c} \ddot{O} \\ \parallel \\ :\ddot{O}-N-\ddot{O}-N=\ddot{O} \\ \parallel \\ \ddot{O} \end{array}$	N 151 pm 119 pm 112° 134° Planar

Preparation:1. **N₂O**





(dil. & cold)

2. **NO**


hot





Industrial process.




3. **N₂O₃**4. **NO₂**

M = Pb, Cu, Ba, Ca






(hot & conc.)


5. **N₂O₅**

Properties:**(I) Decomposition Behaviour**







(Blue liq.) at (-30°C)

**(II) Reaction with H_2O & NaOH**

	H_2O	NaOH
(i) N_2O :	Fairly soluble in water and produces neutral solution	-----
(ii) NO :	Sparingly soluble in water and produces neutral sol ⁿ .	-----
(iii) N_2O_3 :	2HNO_2 Hence it is known as anhydride of HNO_2	NaNO_2
(iv) NO_2 :	$\text{HNO}_2 + \text{HNO}_3$ called as mixed anhydride	$\text{NaNO}_2 + \text{NaNO}_3$
(v) N_2O_5 :	2HNO_3 called as anhydride of HNO_3	NaNO_3

Other properties:

Hence, it is a better supporter for combustion.




- NO : (i) It burns : $\text{NO} + \frac{1}{2} \text{O}_2 \longrightarrow \text{NO}_2$
- (ii) It supports combustion also for molten sulphur and hot phosphorous.
 $\text{S} + 2\text{NO} \longrightarrow \text{SO}_2 + \text{N}_2$
 $2\text{P} + 5\text{NO} \longrightarrow \text{P}_2\text{O}_5 + \frac{5}{2} \text{N}_2$
- (iii) It is being absorbed by FeSO_4 solution.
- (iv) It is having reducing property.
 $\text{KMnO}_4 + \text{NO} + \text{H}_2\text{SO}_4 \longrightarrow \text{K}_2\text{SO}_4 + \text{MnSO}_4 + \text{HNO}_3 + \text{H}_2\text{O}$
 $\text{HOCl} + \text{NO} + \text{H}_2\text{O} \longrightarrow \text{HNO}_3 + \text{HCl}$
- (v) NO shows oxidising property also.
 $\text{SO}_2 + 2\text{NO} + \text{H}_2\text{O} \longrightarrow \text{H}_2\text{SO}_4 + \text{N}_2\text{O}$
 $\text{H}_2\text{S} + 2\text{NO} \longrightarrow \text{H}_2\text{O} + \text{S} \downarrow + \text{N}_2\text{O}$
 $3\text{SnCl}_2 + 2\text{NO} + 6\text{HCl} \longrightarrow 3\text{SnCl}_4 + 2\text{NH}_2\text{OH}$
 (Used for NH_2OH preparation)
- (vi) NO combines with X_2 ($\text{X}_2 = \text{Cl}_2, \text{Br}_2, \text{F}_2$) to produce NO X
 $2\text{NO} + \text{X}_2 \longrightarrow 2\text{NOX}$

N_2O_3 : No more properties.


- NO_2 : (1) It is having oxidising property.
 $\text{S} + \text{NO}_2 \longrightarrow \text{SO}_2 + \text{NO}$
 $\text{P} + \text{NO}_2 \longrightarrow \text{P}_2\text{O}_5 + \text{NO}$
 $\text{C} + \text{NO}_2 \longrightarrow \text{CO}_2 + \text{NO}$
 $\text{SO}_2 + \text{NO}_2 + \text{H}_2\text{O} \longrightarrow \text{H}_2\text{SO}_4 + \text{NO}$
 $\text{H}_2\text{S} + \text{NO}_2 \longrightarrow \text{H}_2\text{O} + \text{S} \downarrow + \text{NO}$
 $\text{CO} + \text{NO}_2 \longrightarrow \text{CO}_2 + \text{NO}$

NO not formed : $2\text{KI} + 2\text{NO}_2 \longrightarrow \text{I}_2 + 2\text{KNO}_2$

- (2) Reducing property of NO_2 .
 $\text{KMnO}_4 + \text{NO}_2 + \text{H}_2\text{SO}_4 \longrightarrow \text{K}_2\text{SO}_4 + \text{MnSO}_4 + \text{HNO}_3 + \text{H}_2\text{O}$



N_2O_5 :




This proves that N_2O_5 is consisting of ion pair of NO_2^+ & NO_3^-

OXOACIDS OF NITROGEN

$\text{H}_2\text{N}_2\text{O}_2$ (hyponitrous acid), HNO_2 (nitrous acid) and HNO_3 (nitric acid). Amongst them HNO_3 is the most important.







NITROUS ACID (HNO_2)

Preparation







Properties


- (a) **Oxidising property** : Because of its easy oxidation to liberate nascent oxygen, it acts as a strong oxidant $2\text{HNO}_2 \longrightarrow \text{H}_2\text{O} + 2\text{NO} + (\text{O})$





- (b) **Reducing property** : Nitrous acid also acts as a reducing agent as it can be oxidised into nitric acid.





- (c) **Reaction with NH_3 / $-\text{NH}_2$ compounds :**



Urea



Benzene diazonium chloride




NITRIC ACID

It was named aqua fortis (means strong water) by alchemists.


Preparation :

Laboratory Method : By heating KNO_3 or NaNO_3 and concentrated H_2SO_4 in a glass retort.




Large scale preparation (Ostwald's process) :


(i) This method is based upon catalytic oxidation of NH_3 by atmospheric oxygen.



(ii) Nitric oxide thus formed combines with oxygen giving NO_2 .







(iii) Nitrogen dioxide so formed, dissolves in water to give HNO_3 .



NO thus formed is recycled and the aqueous HNO_3 can be concentrated by distillation upto $\sim 68\%$ by mass. Further concentration to 98% can be achieved by dehydration with concentrated H_2SO_4 .

Birkeland Eyde Process or arc process



Properties


Physical properties

It has extremely corrosive action on the skin and causes painful sores.

(i) It is a colourless liquid (f.p. 231.4 K and b.p. 355.6 K).

(ii) Laboratory grade nitric acid contains $\sim 68\%$ of the HNO_3 by mass and has a specific gravity of 1.504.


(iii) Nitric acid usually acquires yellow or brown colour due to its decomposition by sunlight into NO_2 .



The yellow or brown colour of the acid can be removed by warming it to $60-80^\circ\text{C}$ and bubbling dry air through it.

Chemical properties

Acidic character in aqueous solution, nitric acid behaves as a strong acid giving hydronium and nitrate ions.




Oxidising nature: Nitric acid acts as a strong oxidising agent as it decomposes to give nascent oxygen easily.



(i) **Oxidation of non-metals :** The nascent oxygen oxidises various non-metals to their corresponding oxyacids of highest oxidation state.


- (1) Sulphur is oxidised to sulphuric acid




- (2) Carbon is oxidised to carbonic acid



- (3) Phosphorus is oxidised to orthophosphoric acid.




- (4) Iodine is oxidised to iodic acid




(ii) Oxidation of metalloids

Metalloids like non-metals also form oxyacids of highest oxidation state.

- (1) Arsenic is oxidised to arsenic acid



- (2) Antimony is oxidised to antimonic acid




- (3) Tin is oxidised to meta-stannic acid.




(iii) Oxidation of Compounds:


- (1) Sulphur dioxide is oxidised to sulphuric acid



- (2) Hydrogen sulphide is oxidised to sulphur




- (3) Ferrous sulphate is oxidised to ferric sulphate in presence of H_2SO_4




- (4) Iodine is liberated from KI.



- (5) HBr, HI are oxidised to Br_2 and I_2 , respectively.





- (6) Ferrous sulphide is oxidised to ferric sulphate



- (7) Stannous chloride is oxidised to stannic chloride in presence of HCl.



Hydroxylamine




(8) Oxidation of organic compounds .

Sawdust catches fire when nitric acid is poured on it.

Turpentine oil bursts into flames when treated with fuming nitric acid.




Toluene is oxidised to benzoic acid with dil. HNO_3 .

Cane sugar is oxidised to oxalic acid.








(ii) Reaction with metal concentrated nitric acid is a strong oxidising agent and attacks most metals except noble metals such as gold and platinum. Au & Pt dissolve in aqua regia a mixture of 25% conc. HNO_3 & 75% conc. HCl . The products of oxidation depend upon the concentration of the acid, temperature and the nature of the material undergoing oxidation.

Ex.



Some metals (e.g., Cr, Al) do not dissolve in concentrated nitric acid because of the formation of a passive film of oxide on the surface.

Armstrong postulated that primary action of nitric acid is to produce hydrogen in the nascent form. Before this hydrogen is allowed to escape, it reduces the nitric acid into number of products like NO_2 , NO , N_2O , N_2 or NH_3 according to the following reactions:



The progress of the reaction is controlled by a number of factors :

- (a) the nature of the metal,
- (b) the concentration of the acid,
- (c) the temperature of the reaction,
- (d) the presence of other impurities.

Concentration of nitric acid	Metal	Main Products
Very dilute HNO_3 (6%)	Mg, Mn	H_2 + Metal nitrate
	Fe, Zn, Sn	NH_4NO_3 + metal nitrate + H_2O
	Pb, Cu, Ag, Hg	NO + metal nitrate + H_2O

Dilute HNO_3 (20%) Fe, Zn $\text{N}_2\text{O} + \text{metal nitrate} + \text{H}_2\text{O}$

Sn $\text{NH}_4\text{NO}_3 + \text{Sn}(\text{NO}_3)_2$

Conc. HNO_3 (70%) Zn, Fe, Pb, Cu, Ag $\text{NO}_2 + \text{metal nitrate} + \text{H}_2\text{O}$

Sn $\text{NO}_2 + \text{H}_2\text{SnO}_3$
Metastannic acid

Action on Proteins :

Nitric acid attacks proteins forming a yellow nitro compound called xanthoprotein. It, therefore, stains skin and renders wool yellow. This property is utilized for the test of proteins.






Uses : The major use of nitric acid is in the manufacture of ammonium nitrate for fertilisers and other nitrates for use in explosives and pyrotechnics. It is also used for the preparation of nitroglycerin, trinitrotoluene and other organic nitro compounds. Other major uses are in the **pickling of stainless steel**, etching of metals and as an oxidiser in rocket fuels.

Pickling is a metal surface treatment used to remove impurities, such as stains, inorganic contaminants, rust or scale from ferrous metals, copper, precious metals and aluminium alloys. A solution called pickle liquor, which contains strong acids, is used to remove surface impurities.






ALLOTROPIC FORMS OF PHOSPHORUS

Comparison between White and Red Phosphorus



Property	White phosphorus	Red phosphorus
Physical state	Soft waxy solid.	Brittle powder.
Colour	White when pure. Attains yellow colour on standing.	Red.
Odour	Garlic	Odourless.
Solubility in water	Insoluble.	insoluble
Solubility in CS_2	Soluble.	Insoluble.
Physiological action	Poisonous.	Non-poisonous.
Chemical activity	Very active.	Less active.
Stability	Unstable.	Stable.
Phosphorescence	Glow in dark	Does not glow in dark.
Molecular formula	P_4	Complex polymer.

Preparation of white 'P'**Reactions of 'P'**


* Reaction with hot metal —

**PHOSPHINE****Preparation**





(i) Phosphine is prepared by the reaction of calcium phosphide with water or dilute HCl.



(ii) Laboratory preparation it is prepared by heating white phosphorus with concentrated NaOH solution in an inert atmosphere of CO_2 .



Pure PH_3 is non inflammable but becomes inflammable owing to the presence of P_2H_4 or P_4 vapours. To purify it from the impurities, it is absorbed in HI to form phosphonium iodide (PH_4I) which on treating with KOH gives off phosphine.

Other preparation



} Purest PH_3

Physical Properties :

- (i) It is a colourless gas with rotten fish smell and is highly poisonous.
- (ii) It explodes in contact with traces of oxidising agents like HNO_3 , Cl_2 and Br_2 vapours.
- (iii) It is slightly soluble in water but soluble in CS_2 . The solution of PH_3 in water decomposes in presence of light giving red phosphorus and H_2 .

Chemical Properties :

- (i) It absorbed in copper sulphate or mercuric chloride solution, the corresponding phosphides are obtained.



Phosphine is weakly basic and like ammonia, gives phosphonium compounds with acids e.g.,




- (ii) $\text{PH}_3 + \text{O}_2 \xrightarrow{150^\circ} \text{P}_2\text{O}_5 + \text{H}_2\text{O}$

- (iii) $\text{PH}_3 + 3\text{Cl}_2 \longrightarrow \text{PCl}_3 + 3\text{HCl}$

- (iv) $\text{PH}_3 + 4\text{N}_2\text{O} \xrightarrow[\text{sparking}]{\text{electrical}} \text{H}_3\text{PO}_4 + 4\text{N}_2$

- (v) $\text{PH}_3 + 6\text{AgNO}_3 \longrightarrow [\text{Ag}_3\text{P} \cdot 3\text{AgNO}_3\downarrow] + 3\text{HNO}_3$

yellow ppt.



Black ppt.

- (vi) $\text{PH}_3 + 4\text{HCHO} + \text{HCl} \longrightarrow [\text{P}(\text{CH}_2\text{OH})_4]^+\text{Cl}^-$



white/colourless solid

which is used for making



fire-proof cotton fabrics

Note :

Like NH_3 , PH_3 also can form addition product.



PH_3 can be absorbed by $\text{Ca}(\text{OCl})\text{Cl}$.

**Uses :**

(i) The spontaneous combustion of phosphine is technically used in Holme's signals. Containers containing calcium carbide and calcium phosphide are pierced and thrown in the sea when the gases evolved burn and serve as a signal.


(ii) It is also used in smoke screens.

PHOSPHORUS HALIDES

Phosphorus forms two types of halides, PX_3 ($X = F, Cl, Br, I$) and PX_5 ($X = F, Cl, Br$).

PHOSPHORUS TRICHLORIDE**Preparation**

- (i) By passing dry chlorine over heated white phosphorus.




- (ii) By the action of thionyl chloride with white phosphorus.



**Properties**

- (i) It is a colourless oily liquid


- (ii) Hydrolyses in the presence of moisture.




- (iii) It reacts with organic compounds containing $-OH$ group such as CH_3COOH , C_2H_5OH .

**PHOSPHORUS PENTACHLORIDE****Preparation**

- (i) By the reaction of white phosphorus with excess of dry chlorine.





- (ii) By the action of SO_2Cl_2 on phosphorus.


**Properties :**

- (i) PCl_5 is a yellowish white powder


- (ii) It hydrolysis in moist air to $POCl_3$ and finally gets converted to phosphoric acid.





- (iii) When heated, it sublimes but decomposes on stronger heating.



- (iv) It reacts with organic compounds containing $-OH$ group converting them to chloro derivatives.




- (v) Finely divided metals on heating with PCl_5 give corresponding chlorides.

**Uses :**

It is used in the synthesis of some organic compounds, e.g., C_2H_5Cl , CH_3COCl .

Note :

In gaseous and liquid phases, it has a trigonal bipyramidal structure as shown. The three equatorial $P-Cl$ bonds are equivalent, while the two axial bonds are longer than equatorial bonds. This is due to the fact that the axial bond pairs suffer more repulsion as compared to equatorial bond pairs.



In the solid state it exists as an ionic solid, $[PCl_4]^+[PCl_6]^-$ in which the cation, $[PCl_4]^+$ is tetrahedral and the anion, $[PCl_6]^-$ is octahedral.


OXIDES OF PHOSPHORUS

It forms two important oxides which exist in dimeric forms.

PHOSPHORUS TRIOXIDE (P_4O_6)


Preparation

Phosphorus trioxides is formed when phosphorus is burnt in a limited supply of air and inert atmosphere.




Properties

- (a) **Heating in air :** On heating in air, it forms phosphorus pentoxide.



- (b) **Action of water :** It dissolves in cold water to give phosphorus acid.



It is, therefore, considered as anhydride of phosphorus acid.


Note: With hot water, it gives phosphoric acid and inflammable phosphine.

Structure

- Each atom of phosphorus in P_4O_6 is present at the corner of a tetrahedron
- Each phosphorus atom is covalently bonded to three oxygen atoms and each oxygen atom is bonded to two phosphorus atoms.
- It is clear from the structure that the six oxygen atoms lie along the edges of the tetrahedron of P atoms.


PHOSPHORUS (V) OXIDE (P_4O_{10})

Preparation : It is prepared by heating white phosphorus in excess of air.



Properties

- It is snowy white solid.
- Action with water :** It readily dissolves in cold water forming metaphosphoric acid.






With hot water it gives phosphoric acid.



Phosphoric acid

- Dehydrating nature :** Phosphorus pentoxide has strong affinity for water and, therefore, acts as a powerful dehydrating agent. It extracts water from many inorganic and organic compounds.
- P_4O_{10} is a very strong dehydrating agent and extracts water from many compounds including sulphuric acid and nitric acid.



Structure

- (a) Its structure is similar to that of P_4O_6 .
- (b) In addition, each phosphorus atom forms a double bond with oxygen atom as shown in figure.

OXOACIDS OF PHOSPHORUS :

The important oxoacids of phosphorus with their formulae, methods of preparation and the presence of some characteristic bonds in their structures are given in a table.

Oxoacids of Phosphorus

Name	Formula	Oxidation state of Phosphorus	Characteristic bonds and their number	Preparation
Hypophosphorus (Phosphinic)	H_3PO_2	+ 1	One P — OH Two P — H One P = O	white P_4 + alkali
Orthophosphorous (Phosphonic)	H_3PO_3	+ 3	Two P — OH One P — H One P = O	$P_2O_3 + H_2O$
Pyrophosphorous	$H_4P_2O_5$	+ 3	Two P — OH Two P — H Two P = O	$PCl_3 + H_3PO_3$
Hypophosphoric	$H_4P_2O_6$	+ 4	Four P — OH Two P = O One P — P	red P_4 + alkali
Orthophosphoric	H_3PO_4	+ 5	Three P — OH One P = O	$P_4O_{10} + H_2O$
Pyrophosphoric	$H_4P_2O_7$	+ 5	Four P — OH Two P = O One P — O — P	heat phosphoric acid
Metaphosphoric*	$(HPO_3)_n$	+ 5	Three P — OH Three P = O Three P — O — P	phosphorous acid + Br_2 , heat in a sealed tube

STRUCTURE OF OXOACID :


In oxoacids phosphorus is tetrahedrally surrounded by other atoms. All these acids contain at least one P=O bond and one P—OH bond. The oxoacids in which phosphorus has lower oxidation state (less than +5) contain, in addition to P=O and P—OH bonds, either P—P (e.g., in $H_4P_2O_6$) or P—H (e.g., in H_3PO_3) bonds but not both. H_3PO_3 and H_3PO_2 are diacidic and monobasic respectively.

Note : -


- (i) These acids in +3 oxidation state of phosphorus tend to disproportionate to higher and lower oxidation states. For example, phosphorous acid on heating disproportionates to give orthophosphoric acid (or phosphoric acid) and phosphine.



(ii) The acids which contain P–H bond have strong reducing properties. Thus, hypophosphorous acid is a good reducing agent as it contains two P–H bonds and reduces, AgNO_3 to metallic silver.




Heating Effect :



Graham salt

Graham's salt is the best known of these long chain polyphosphates, and is formed by quenching molten NaPO_3 . Graham's salt is soluble in water. These solutions give precipitates with metal ions such as Pb^{2+} and Ag^+ but not with Ca^{2+} and Mg^{2+} . Graham's salt is sold commercially under the trade name Calgon. In industry it is incorrectly called sodium hexametaphosphate crystallizing. It is widely used for softening water.



OXYGEN FAMILY

GROUP 16 ELEMENTS (O, S, Se, Te, Po)

This is sometimes known as group of chalcogens.

❑ Occurrence

Oxygen is the most abundant of all the elements on earth crust. Oxygen forms about 46.6% by mass of earth's crust. Dry air contains 20.946% oxygen by volume. However, the abundance of sulphur in the earth's crust is only 0.03-0.1%. Combined sulphur exists primarily as sulphates such as gypsum $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$, epsom salt $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, baryte BaSO_4 and sulphides such as galena PbS , zinc blende ZnS , copper pyrites CuFeS_2 . Traces of sulphur occur as hydrogen sulphide in volcanoes. Organic materials such as eggs, proteins, garlic, onion, mustard, hair and wool contain sulphur.

Selenium and tellurium are also found as metal selenides and tellurides in sulphide ores. Polonium occurs in nature as a decay product of thorium and uranium minerals.

❑ Electronic Configuration

ns^2np^4 is the general valence shell electronic configuration.

❑ Atomic and Ionic Radii : Covalent radius : $\text{O} < \text{S} < \text{Se} < \text{Te}$

❑ Ionisation Enthalpy : $\text{O} > \text{S} > \text{Se} > \text{Te} > \text{Po}$ (IE_1 values)

❑ Most Negative Electron Gain Enthalpy: $\text{S} > \text{Se} > \text{Te} > \text{Po} > \text{O}$

❑ Electronegativity : $\text{O} > \text{S} > \text{Se} > \text{Te}$

❑ Metallic Character : $\text{O} < \text{S} < \text{Se} < \text{Te} < \text{Po}$

❑ Melting and Boiling points :

M.P. : $\text{Te} > \text{Po} > \text{Se} > \text{S} > \text{O}$

B.P. : $\text{Te} > \text{Po} > \text{Se} > \text{S} > \text{O}$

Elemental State

Oxygen exist as diatomic molecular gas in this case there is $p\pi - p\pi$ overlap thus two O atoms form double bond $\text{O} = \text{O}$. The intermolecular forces in O_2 are weak VB forces. $\therefore \text{O}_2$ exist as gas. On the other hand, other elements of family do not form stable $p\pi - p\pi$ bonds and do not exist as M_2 molecules. Other atoms are linked by single bonds and form poly atomic complex molecules for eg. $\text{S} - \text{S}_8$, $\text{Se} - \text{Se}_8$

Allotropy

All element exhibit allotropy for e.g.

Oxygen – O_2 and O_3

Liquid O_2 - pale blue

Solid O_2 - blue

Physical Properties

- (i) Oxygen and sulphur are non-metals, selenium and tellurium metalloids, whereas polonium is a metal.
- (ii) Polonium is radioactive and is short lived (Half-life 13.8 days).
- (iii) All these elements exhibit allotropy.
- (iv) The melting and boiling points increase with an increase in atomic number down the group. The large difference between the melting and boiling points of oxygen and sulphur may be explained on the basis of their atomicity; oxygen exists as diatomic molecule (O_2) whereas sulphur exists as polyatomic molecule (S_8).

Chemical Properties**Oxidation states and trends in chemical reactivity :**

- (i) The stability of -2 oxidation state decreases down the group. Polonium hardly shows -2 oxidation state.
- (ii) Electronegativity of oxygen is very high, it shows only negative oxidation state as -2 except Oxygen shows oxidation states of +2 and +1 in oxygen fluorides OF_2 and O_2F_2 respectively.
- (iii) Elements of the group exhibit +2, +4, +6 oxidation states but +4 and +6 are more common.
- (iv) Sulphur, selenium and tellurium usually show +4 oxidation state in their compounds with oxygen and +6 with fluorine.
- (v) The stability of +6 oxidation state decreases down the group and stability of +4 oxidation state increases (inert pair effect).
- (vi) Bonding in +4 and +6 oxidation states is primarily covalent.

Anomalous behaviour of oxygen

The anomalous behaviour of oxygen, like other members of p-block present in second period is due to its small size and high electronegativity. One typical example of effects of small size and high electronegativity is the presence of strong hydrogen bonding in H_2O which is not found in H_2S . The absence of d orbitals in oxygen limits its covalency to four and in practice, rarely exceeds two. On the other hand, in case of other elements of the group, the valence shells can be expanded and covalency exceeds four.

(I) Reactivity with hydrogen:

- (i) All the elements of Group 16 form hydrides of the type H_2E ($E = O, S, Se, Te, Po$).
- (ii) Their acidic character increases from H_2O to H_2Te . The increase in acidic character can be explained in terms of decrease in bond enthalpy for the dissociation of H-E bond down the group. Owing to the decrease in enthalpy for the dissociation of H-E bond down the group, the thermal stability of hydrides also decreases from H_2O to H_2Po .
- (iii) All the hydrides except water possess reducing property and this character increases from H_2S to H_2Te .

Properties of Hydrides of Group 16 Elements

Property	H ₂ O	H ₂ S	H ₂ Se	H ₂ Te
m.p./K	273	188	208	222
b.p./K	373	213	232	269
H—E distance /pm	96	134	146	169
HEH angle (°)	104	92	91	90
$\Delta_f H / \text{kJ mol}^{-1}$	− 286	− 20	73	100
$\Delta_{\text{diss}} H (\text{H} - \text{E}) / \text{kJ mol}^{-1}$	463	347	276	238
Dissociation constant ^a	1.8×10^{-16}	1.3×10^{-7}	1.3×10^{-4}	2.3×10^{-3}

(II) Reactivity with oxygen:

- All these elements form oxides of the EO₂ and EO₃ types where E = S, Se, Te or Po.
- Ozone (O₃) and sulphur dioxide (SO₂) are gases while selenium dioxide (SeO₂) is solid.
- Reducing property of dioxide decreases from SO₂ to TeO₂; SO₂ is reducing while TeO₂ is an oxidising agent.
- Besides EO₂ type, sulphur, selenium and tellurium also form EO₃ type oxides (SO₃, SeO₃, TeO₃). Both types of oxides are acidic in nature.

(III) Reactivity towards the halogens:


- Elements of Group 16 form a large number of halides of the type, EX₆, EX₄ and EX₂ where E is an element of the group and X is a halogen.
- The stability of the halides decreases in the order $\text{F}^- > \text{Cl}^- > \text{Br}^- > \text{I}^-$.
- Amongst hexahalides, hexafluorides are the only stable halides. All hexafluorides are gaseous in nature. They have octahedral structure. Sulphur hexafluoride, SF₆ is exceptionally stable for steric reasons.

Amongst tetrafluorides, SF₄ is a gas, SeF₄ a liquid and TeF₄ a solid. These fluorides have sp^3d hybridisation and thus, have trigonal bipyramidal structures in which one of the equatorial positions is occupied by a lone pair of electrons. This geometry is also regarded as see-saw geometry. All elements except oxygen form dichlorides and dibromides (because they form oxides). These dihalides are formed by sp^3 hybridisation and thus, have tetrahedral structure. The well known monohalides are dimeric in nature. Examples are S₂F₂, S₂Cl₂, S₂Br₂, Se₂Cl₂ and Se₂Br₂. These dimeric halides undergo disproportionation as given below : $2\text{Se}_2\text{Cl}_2 \rightarrow \text{SeCl}_4 + 3\text{Se}$



DIOXYGEN

(a) Laboratory method

(i) By heating oxygen containing salts such as chlorates, nitrates and permanganates.



(ii) By the thermal decomposition of the oxides of metals low in the electrochemical series and higher oxides of some metals.



(iii) Hydrogen peroxide is readily decomposed into water and dioxygen by catalysts such as finely divided metals and manganese dioxide.









(b) **Large scale preparation** : It can be prepared from water or air. Electrolysis of water leads to the release of hydrogen at the cathode and oxygen at the anode.



(c) **Industrially method** : Dioxygen is obtained from air by first removing carbon dioxide and water vapour and then, the remaining gases are liquefied and fractionally distilled to give dinitrogen and dioxygen.

Properties

- (i) Dioxygen is a colourless and odourless gas.
- (ii) Its solubility in water is to the extent of 3.08 cm^3 in 100 cm^3 water at 293 K which is just sufficient for the vital support of marine and aquatic life.
- (iii) It liquefies at 90 K and freezes at 55 K .
- (iv) Oxygen atom has three stable isotopes: ^{16}O , ^{17}O and ^{18}O . Molecular oxygen, O_2 is unique in being paramagnetic inspite of having even number of electrons.
- (v) Dioxygen directly reacts with nearly all metals and non-metals except some metals (e.g., Au, Pt) and some noble gases. Its combination with other elements is often strongly exothermic which helps in sustaining the reaction. However, to initiate the reaction, some external heating is required as bond dissociation enthalpy of oxygen-oxygen double bond is high ($493.4 \text{ kJ mol}^{-1}$). Some of the reactions of dioxygen with metals, non-metals and other compounds are as follows :



Some compounds are catalytically oxidised. For example,



Uses: (i) It's importance in normal respiration and combustion processes, oxygen is used in oxyacetylene welding, in the manufacture of many metals, particularly steel.

(ii) Oxygen cylinders are widely used in hospitals, high altitude flying and in mountaineering.


(iii) The combustion of fuels, e.g., hydrazines in liquid oxygen, provides tremendous thrust in rockets.

SIMPLE OXIDES

A binary compound of oxygen with another element is called oxide. In many cases one element forms two or more oxides. The oxides vary widely in their nature and properties. Oxides can be simple (e.g., MgO , Al_2O_3) or mixed (Pb_3O_4 , Fe_3O_4).


Types of simple oxide : Simple oxides can be classified on the basis of their acidic, basic or amphoteric character.

Acidic oxide : An oxide that combines with water to give an acid is termed acidic oxide (e.g., SO_2 , Cl_2O_7 , CO_2 , N_2O_5). For example, SO_2 combines with water to give H_2SO_3 , an acid.





As a general rule, only non-metal oxides are acidic but oxides of some metals in high oxidation state also have acidic character (e.g., Mn_2O_7 , CrO_3 , V_2O_5).

Basic oxide : The oxides which give a base with water are known as basic oxides (e.g., Na_2O , CaO , BaO). In general, metallic oxides are basic. For example, CaO combines with water to give Ca(OH)_2 , a base.



Amphoteric oxide :

Some metallic oxides exhibit a dual behaviour. They show characteristics of both acidic as well as basic oxides. Such oxides are known as amphoteric oxides. They react with acids as well as alkalies. For example, Al_2O_3 reacts with acids as well as alkalies.



Neutral oxide:

There are some oxides which are neither acidic nor basic. Such oxides are known as neutral oxides. Examples of neutral oxides are CO , NO and N_2O .


OZONE

(i) Ozone is an allotropic form of oxygen and is diamagnetic.

(ii) It is too reactive to remain for long in the atmosphere at sea level. At a height of about 20 kilometres, it is formed from atmospheric oxygen in the presence of sunlight. This ozone layer protects the earth's surface from an excessive concentration of ultraviolet (UV) radiations.

Threats to ozone layer


(i) Experiments have shown that nitrogen oxides (particularly nitric oxide) combine very rapidly with ozone and there is, thus, the possibility that nitrogen oxides emitted from the exhaust systems of supersonic jet aeroplanes might be slowly depleting the concentration of the ozone layer in the upper atmosphere.



(ii) Another threat to this ozone layer is probably posed by the use of freons which are used in aerosol sprays and as refrigerants.

Preparation

When a slow dry stream of oxygen is passed through a silent electrical discharge, conversion of oxygen to ozone (10%) occurs. The product is known as ozonised oxygen.




Since the formation of ozone from oxygen is an **endothermic** process, it is necessary to use a silent electrical discharge in its preparation to prevent its decomposition. If concentration of ozone greater than 10 percent is required, a battery of ozonisers can be used, and pure ozone (b.p. -112.4°C) can be condensed in a vessel surrounded by liquid oxygen.

Ques. Ozone is thermodynamically unstable with respect to oxygen. Explain ?

Sol. Because its decomposition into oxygen results in the liberation of heat (ΔH is negative) and an increase in entropy (ΔS is positive). These two effects reinforce each other, resulting in large negative Gibbs energy change (ΔG) for its conversion into oxygen.

Note :



Properties


- (i) Pure ozone is a pale blue gas, dark blue liquid and violet-black solid.
- (ii) Ozone has a characteristic fishy smell and in small concentrations it is harmless.

Toxic effect :

- (a) Toxic enough (more toxic than KCN). It's intense blue colour is due to the absorption of red light.
- (b) However, if the concentration rises above about 100 parts per million, breathing becomes uncomfortable resulting in headache and nausea.


Oxidizing properties

It is one of best oxidising agent, in acid solution, its standard, reduction potential value is 2.07 V.




It is next to F_2 . [above 2.07 V, only F_2 , F_2O are there]

It is not really surprising, therefore, high concentrations of ozone can be dangerously explosive. Due to the ease with which it liberates atoms of nascent oxygen ($O_3 \rightarrow O_2 + O$), it acts as a powerful oxidising agent. For example, it oxidises lead sulphide to lead sulphate and iodide ions to iodine.





- (i) Metal Sulphides to Sulphates.






- (ii) $2HX + O_3 \longrightarrow X_2 + H_2O + O_2 \quad [X = Cl, Br, I]$



- (iii) $NaNO_2 + O_3 \longrightarrow NaNO_3 + O_2$








- (iv) Moist S, P, As + $O_3 \Rightarrow$





- (v) Moist $I_2 \longrightarrow HIO_3$ whereas dry iodine $\longrightarrow I_4O_9$ (yellow)











(viii) Hg loses its fluidity (tailing of Hg)




Brown



In all above reaction O_3 gives up O_2 but some reactions are there which consumes all O-atom.



Bonding in ozone : In ozone the two oxygen-oxygen bond lengths in the ozone molecule are identical (128 pm) and the molecule is angular as expected with a bond angle of about 117° . It is a resonance hybrid of two main forms given below:



Absorbent : (i) Turpentine oil (ii) Oil of cinnamon

Quantitative method for the estimating on Ozone : Ozone reacts with an excess of potassium iodide solution buffered with a borate buffer (pH 9.2), iodine is liberated which can be titrated against a standard solution of sodium thiosulphate.

Uses : (i) Sterilising water

(ii) Detection of position of the double bond in the unsaturated compound.

(iii) It is used as a germicide, disinfectant and for sterilising water.

(iv) It is also used for bleaching oils, ivory, flour, starch, etc.


(v) It acts as an oxidising agent in the manufacture of potassium permanganate.

❑ HYDROGEN PEROXIDE (H₂O₂)


Hydrogen peroxide is an important chemical used in pollution control treatment of domestic and industrial effluents.

Preparation



It can be prepared by the following methods.






(ii) Acidifying barium peroxide and removing excess water by evaporation under reduced pressure gives hydrogen peroxide.




Instead of H₂SO₄, H₃PO₄ is added now-a-days because H₂SO₄ catalyses the decomposition of H₂O₂ whereas H₃PO₄ favours to restore it.




(iii) Peroxodisulphate, obtained by electrolytic oxidation of acidified sulphate solutions at high current density, on hydrolysis yields hydrogen peroxide.



This method is now used for the laboratory preparation of D₂O₂.



(iii) Industrially it is prepared by the autooxidation of 2-alkylanthraquinols.



In this case 1% H₂O₂ is formed. It is extracted with water and concentrated to ~30% (by mass) by distillation under reduced pressure. It can be further concentrated to ~85% by careful distillation under low pressure. The remaining water can be frozen out to obtain pure H₂O₂.

◆ Physical Properties

In the pure state H₂O₂ is an almost colourless (very pale blue) liquid. Its important physical properties are given in Table. H₂O₂ is miscible with water in all proportions and forms a hydrate H₂O₂·H₂O (mp 221K). A 30% solution of H₂O₂ is marketed as '100 volume' hydrogen peroxide. It means that one millilitre of 30% H₂O₂ solution will give 100 mL of oxygen at STP. Commercially marketed sample is 10 V, which means that the sample contains 3% H₂O₂.


30% (w / v) or "100 V" H₂O₂ solution is called **per hydrol**.

Problem

Calculate the strength of 10 volume solution of hydrogen peroxide.

Solution

10 volume solution of H_2O_2 means that 1L of this H_2O_2 solution will give 10 L of oxygen at STP



On the basis of above equation 22.7 L of O_2 is produced from 68 g H_2O_2 at STP 10 L of O_2 at STP is produced from

$$\frac{68 \times 10}{22.7} \text{ g} = 29.9 \text{ g} \quad 30 \text{ g } \text{H}_2\text{O}_2$$

Therefore, strength of H_2O_2 in 10 volume H_2O_2 solution = 30 g/L = 3% H_2O_2 solution

Physical Properties of Hydrogen Peroxide

Melting point/K	272.4	Density (liquid at 298K)/g cm ⁻³	1.44
Boiling point (extrapolated)/K	423	Viscosity (290 K)/centipoise	1.25
Vapour pressure (298K) mmHg	1.9	Dielectric constant (298K)/C ² /N m ²	70.7
Density (solid at 268.5K)/g cm ⁻³	1.64	Electrical conductivity (298K)/Ω ⁻¹ cm ⁻¹	5.1×10^{-8}

Structure

Hydrogen peroxide has a non-planar structure. The molecular dimensions in the gas phase and solid phase are shown in Fig.


































Fig. (a) H_2O_2 structure in gas phase, dihedral angle is 111.5° . (b) H_2O_2 structure in solid phase at 110K, dihedral angle is 90.2° .

Chemical Properties:



(ii) It is oxidant as well as reductant.

**Oxidising Properties:****Reducing properties:**







It acts as an oxidising as well as reducing agent in both acidic and alkaline media. Simple reactions are described below.



(i) Oxidising action in acidic medium




(ii) Reducing action in acidic medium



(iii) Oxidising action in basic medium



(iv) Reducing action in basic medium



◆ Storage

H_2O_2 decomposes slowly on exposure to light.



In the presence of metal surfaces or traces of alkali (present in glass containers), the above reaction is catalysed. It is, therefore, stored in wax-lined glass or plastic vessels in dark. Acetaldehyde or Glycerol or Urea can be added as a stabiliser. It is kept away from dust because dust can induce explosive decomposition of the compound.

Uses

Its wide scale use has led to tremendous increase in the industrial production of H_2O_2 .

Some of the uses are listed below :

- (i) In daily life it is used as a hair bleach and as a mild disinfectant. As an antiseptic it is sold in the market as perhydrol.

- (ii) It is used to manufacture chemicals like sodium perborate and per-carbonate, which are used in high quality detergents.
- (iii) It is used in the synthesis of hydroquinone, tartaric acid and certain food products and pharmaceuticals (cephalosporin) etc.
- (iv) It is employed in the industries as a bleaching agent for textiles, paper pulp, leather, oils, fats, etc.
- (v) As a rocket propellant:


$$\text{NH}_2\text{NH}_2 + 2\text{H}_2\text{O}_2 \longrightarrow \text{N}_2 + 4\text{H}_2\text{O} \text{ [highly exothermic and large increase in volume]}$$
- (vi) In detection of Cr^{+3} , Ti^{+4} etc.

$$\text{Ti}(\text{SO}_4)_2 + \text{H}_2\text{O}_2 + 2\text{H}_2\text{O} \longrightarrow \text{H}_2\text{TiO}_4 + 2\text{H}_2\text{SO}_4$$

Yellow or orange
Pertitanic acid
- (vii) Nowadays it is also used in Environmental (Green) Chemistry. For example, in pollution control treatment of domestic and industrial effluents, oxidation of cyanides, restoration of aerobic conditions to sewage wastes, etc.

ALLOTROPIC FORMS OF SULPHUR

Sulphur forms numerous allotropes of which the yellow rhombic (α -sulphur) and monoclinic (β -sulphur) forms are the most important. The stable form at room temperature is rhombic sulphur, which transforms to monoclinic sulphur when heated above 369 K.



At 369 K both the forms are stable. This temperature is called transition temperature.

Note: Viscosity of 'S' with temperature :


m.p. of 'S' $\longrightarrow$ 112.8°C.

$> 112.8^\circ\text{C}$ to $160^\circ\text{C} \Rightarrow$ slow decreases due to

S_8 rings slip and roll over one another easily.

$> 160^\circ\text{C}$, increases sharply due to breaking of

S_8 rings into chains and polymers into large size chain.




Amorphous forms are

- (i) Plastic sulphur (ii) Milk of sulphur (iii) Colloidal sulphur


SULPHUR DIOXIDE

Preparation






Sulphur dioxide is formed together with a little (6-8%) sulphur trioxide when sulphur is burnt in air or oxygen:




laboratory method by treating a sulphite with dilute sulphuric acid.



other preparation :




Industrial method, by-product of the roasting of sulphide ores.




The gas after drying is liquefied under pressure and stored in steel cylinders.

Properties



- (i) Sulphur dioxide is a colourless gas with pungent smell.
- (ii) It is highly soluble in water.
- (iii) It liquefies at room temperature under a pressure of two atmospheres and boils at 263 K.
- (iv) **Acidic character:** sulphur dioxide, when passed through water, forms a solution of sulphurous acid.



It reacts readily with sodium hydroxide solution, forming sodium sulphite, which then reacts with more sulphur dioxide to form sodium hydrogen sulphite.





In its reaction with water and alkalies, the behaviour of sulphur dioxide is very similar to that of carbon dioxide. Sulphur dioxide reacts with chlorine in the presence of charcoal (which acts as a catalyst) to give sulphuryl chloride, SO_2Cl_2 . It is oxidised to sulphur trioxide by oxygen in the presence of vanadium(V) oxide catalyst.




Reducing properties

When moist, sulphur dioxide behaves as a reducing agent. For example, it converts iron(III) ions to iron(II) ions and decolourises acidified potassium permanganate(VII) solution; the latter reaction is a convenient test for the gas.



Bonding in SO_2 : The molecule of SO_2 is angular. It is a resonance hybrid of the two canonical forms:




Uses:

- It is used refining petroleum and sugar
- It is used in bleaching wool and silk
- It is used as an anti-chlor, disinfectant and preservative. Sulphuric acid, sodium hydrogen sulphite and calcium hydrogen sulphite (industrial chemicals) are manufactured from sulphur dioxide. Liquid SO_2 is used as a solvent to dissolve a number of organic and inorganic chemicals.

HYDROGEN SULPHIDE (H_2S) SULPHURATED HYDROGEN

Preparation

By the action of dil. HCl or H_2SO_4 on iron pyrites.




Note: **Drying agent for this gas : fused CaCl_2 , Al_2O_3 (dehydrated) P_2O_5 etc. But not H_2SO_4 , because $\text{H}_2\text{SO}_4 + \text{H}_2\text{S} \longrightarrow 2\text{H}_2\text{O} + \text{SO}_2 + \text{S}$**


Properties

It is a colourless gas having an offensive smell of rotten eggs.

- (a) It burn in air with blue flame




If the air supply is in excess



- (b) It is a mild acid.




- (c) It act as a reducing agent. It reduces halogen into corresponding hydroacid.



Tests of H_2S

- (a) Unpleasant odour resembling that of rotten eggs.

- (b) It turns lead acetate into paper black



- (c) It gives a violet colouration with a alkaline solution of sodium nitroprusside.

Structure of H_2S


- (a) Similar to structure of water molecule i.e. V-shaped structure with bond length (H-S) $1.35\text{\AA}$ and bond angle (H-S-H) is 92.5°

Uses

- It is mainly employed in salt analysis for the detection of cation.
- Reducing agent for H_2SO_4 , KMnO_4 , $\text{K}_2\text{Cr}_2\text{O}_7$, O_3 , H_2O_2 , FeCl_3

OXOACIDS OF SULPHUR

Sulphur forms a number of oxoacids such as H_2SO_3 , $\text{H}_2\text{S}_2\text{O}_3$, $\text{H}_2\text{S}_2\text{O}_4$, $\text{H}_2\text{S}_2\text{O}_5$, $\text{H}_2\text{S}_x\text{O}_6$ ($x = 2$ to 5), H_2SO_4 , $\text{H}_2\text{S}_2\text{O}_7$, H_2SO_5 , $\text{H}_2\text{S}_2\text{O}_8$. Some of these acids are unstable and cannot be isolated. They are known in aqueous solution or in the form of their salts. Structures of some important oxoacids are shown in Fig.



Structures of some important oxoacids of sulphur

SULPHURIC ACID

Sulphuric acid is one of the most important industrial chemicals worldwide.


Industrial Manufacturing (Contact process)

Steps involved :

(i) **Burning of sulphur or sulphide ores in air to generate SO_2 .**

(ii) **Conversion of SO_2 to SO_3 by the reaction with oxygen in the presence of a catalyst (V_2O_5):**


The SO_2 produced is purified by removing dust and other impurities such as arsenic compounds. The key step in the manufacture of H_2SO_4 is the catalytic oxidation of SO_2 with O_2 to give SO_3 in the presence of V_2O_5 (catalyst).




The reaction is exothermic, reversible and the forward reaction leads to a decrease in volume. Therefore, low temperature and high pressure are the favourable conditions for maximum yield. But the temperature should not be very low otherwise rate of reaction will become slow. In practice, the plant is operated at a pressure of 2 bar and a temperature of 720 K.

(iii) **Absorption of SO_3 in H_2SO_4 to give Oleum ($\text{H}_2\text{S}_2\text{O}_7$) :**


The SO_3 gas from the catalytic converter is absorbed in concentrated H_2SO_4 to produce oleum. Dilution of oleum with water gives H_2SO_4 of the desired concentration. In the industry two steps are carried out simultaneously to make the process a continuous one and also to reduce the cost.





Flow diagram for the manufacture of sulphuric acid

The sulphuric acid obtained by Contact process is 96-98% pure.



Properties

- Sulphuric acid is a colourless, dense, oily liquid with a specific gravity of 1.84 at 298 K.
- The acid freezes at 283 K and boils at 611 K.
- It dissolves in water with the evolution of a large quantity of heat. Hence, care must be taken while preparing sulphuric acid solution from concentrated sulphuric acid. The concentrated acid must be added slowly into water with constant stirring.


Chemical properties

The chemical reactions of sulphuric acid are as a result of the following characteristics:

- | | |
|-----------------------------------|---|
| (a) low volatility | (b) strong acidic character |
| (c) strong affinity for water and | (d) ability to act as an oxidising agent. |


Acidic character :

In aqueous solution, sulphuric acid ionises in two steps.







The larger value of K_{a_1} ($K_{a_1} > 10$) means that H_2SO_4 is largely dissociated into H^+ and HSO_4^- . Greater the value of dissociation constant (K_a), the stronger is the acid.

The acid forms two series of salts: normal sulphates (such as sodium sulphate and copper sulphate) and acid sulphates (e.g., sodium hydrogen sulphate). Sulphuric acid, because of its low volatility can be used to manufacture more volatile acids from their corresponding salts.










Dehydrating Property :

Concentrated sulphuric acid is a strong dehydrating agent. Many wet gases can be dried by passing them through sulphuric acid, provided the gases do not react with the acid. Sulphuric acid removes water from organic compounds; it is evident by its charring action on carbohydrates.




**Oxidizing Nature :**

Hot concentrated sulphuric acid is a moderately strong oxidising agent. In this respect, it is intermediate between phosphoric and nitric acids. Both metals and non-metals are oxidised by concentrated sulphuric acid, which is reduced to SO_2 .







 **H_2SO_4 & SO_3 :**

Uses: Sulphuric acid is a very important industrial chemical. A nation's industrial strength can be judged by the quantity of sulphuric acid it produces and consumes. It is needed for the manufacture of hundreds of other compounds and also in many industrial processes. The bulk of sulphuric acid produced is used in the manufacture of fertilisers (e.g., ammonium sulphate, superphosphate). Other uses are in:








- (i) petroleum refining
- (ii) manufacture of pigments, paints and dyestuff intermediates
- (iii) detergent industry
- (iv) metallurgical applications (e.g., cleansing metals before enameling, electroplating and galvanising)
- (v) storage batteries
- (vi) in the manufacture of nitrocellulose products and
- (vii) as a laboratory reagent.

SODIUM THIOSULPHATE**Preparation:**

$\downarrow$ SO_2 passed into it



[Na_2S is readily oxidised in air giving rise to $\text{Na}_2\text{S}_2\text{O}_3$]



HALOGEN FAMILY**GROUP 17 ELEMENTS (F, Cl, Br, I, At)**

These are collectively known as the halogens (Greek halo means salt and genes means born i.e., salt producers). Astatine is a radioactive element.

□ Occurrence

- (i) Fluorine and chlorine are fairly abundant while bromine and iodine less so.
- (ii) Fluorine is present mainly as insoluble fluorides (fluorspar CaF_2 , cryolite Na_3AlF_6 and fluoroapatite $3\text{Ca}_3(\text{PO}_4)_2 \cdot \text{CaF}_2$) and small quantities are present in soil, river water plants and bones and teeth of animals.
- (iii) Sea water contains chlorides, bromides and iodides of sodium, potassium, magnesium and calcium, but is mainly sodium chloride solution (2.5% by mass).
- (iv) The deposits of dried up seas contain these compounds, e.g., sodium chloride and carnallite, $\text{KCl} \cdot \text{MgCl}_2 \cdot 6\text{H}_2\text{O}$. Certain forms of marine life contain iodine in their systems; various seaweeds, for example, contain upto 0.5% of iodine and Chile saltpetre contains upto 0.2% of sodium iodate.

Electronic Configuration

The electronic configuration of outermost shell 17th group element is (ns^2np^5) .

□ **Atomic and ionic radii :** $\text{F} < \text{Cl} < \text{Br} < \text{I}$

□ **Ionisation Enthalpy :** $\text{F} > \text{Cl} > \text{Br} > \text{I}$

□ **Most Negative Electron Gain Enthalpy:** $\text{Cl} > \text{F} > \text{Br} > \text{I}$

It is due to small size of fluorine atom. As a result, there are strong interelectronic repulsions in the relatively small 2p orbitals of fluorine and thus, the incoming electron does not experience much attraction.

□ **Electronegativity :** $\text{F} > \text{Cl} > \text{Br} > \text{I}$

□ Physical Properties

- (i) Their melting and boiling points steadily increase with atomic number.
- (ii) All halogens are coloured. For example, F_2 is a yellow gas, Cl_2 greenish yellow gas, Br_2 red liquid and I_2 violet coloured solid. Reason : Decrease in HOMO-LUMO gap.
- (iii) Fluorine and chlorine react with water. Bromine and iodine are only sparingly soluble in water but are soluble in various organic solvents such as chloroform, carbon tetrachloride, carbon disulphide and hydrocarbons to give coloured solutions.
- (iv) **Bond energy order ;** $\text{Cl}_2 > \text{Br}_2 > \text{F}_2 > \text{I}_2$

A reason for this anomaly is the relatively large electron-electron repulsion among the lone pairs in F_2 molecule where they are much closer to each other than in case of Cl_2 .

Chemical Properties

Oxidation states :




- (i) All the halogens exhibit -1 oxidation state. However, chlorine, bromine and iodine exhibit $+1$, $+3$, $+5$ and $+7$ oxidation states also as explained below:

Halogen atom in ground state (other than fluorine)	ns 	np 	nd 	1 unpaired electron accounts for -1 or $+1$ oxidation states
First excited state				3 unpaired electron accounts for $+3$ oxidation states
Second excited state				5 unpaired electron accounts for $+5$ oxidation states
Third excited state				7 unpaired electron accounts for $+7$ oxidation states


- (ii) The higher oxidation states of chlorine, bromine and iodine are realised mainly when the halogens are in combination with the small and highly electronegative fluorine and oxygen atoms. e.g., in interhalogens, oxides and oxoacids.
- (iii) The oxidation states of $+4$ and $+6$ occur in the oxides and oxoacids of chlorine and bromine.
- (iv) The fluorine atom has no d orbitals in its valence shell and therefore cannot expand its octet. Being the most electronegative, it exhibits only -1 oxidation state.

Chemical reactivity

- (i) All the halogens are highly reactive.
- (ii) They react with metals and non-metals to form halides and the reactivity of the halogens decreases down the group. i.e. the order is $F_2 > Cl_2 > Br_2 > I_2$
- (iii) The ready acceptance of an electron is the reason for the strong oxidising nature of halogens. F_2 is the strongest oxidising halogen and it oxidises other halide ions in solution or even in the solid phase. In general, a halogen oxidises halide ions of higher atomic number.





The decreasing oxidising ability of the halogens in aqueous solution down the group is evident from their standard electrode potentials, which are dependent on the parameters as follows :




(1) Reactivity towards water

The relative oxidising power of halogens can further be illustrated by their reactions with water. Fluorine oxidises water to oxygen whereas chlorine and bromine react with water to form corresponding hydrohalic and hypohalous acids. The reaction of iodine with water is nonspontaneous. In fact, I⁻ can be oxidised by oxygen in acidic medium; just the reverse of the reaction is observed with fluorine.



(where X = Cl or Br)

**Note : Anomalous behaviour of fluorine**

- (i) Fluorine is anomalous in many properties. For example, ionisation enthalpy, electronegativity, and electrode potentials are all higher for fluorine than expected from the trends set by other halogens. Also, ionic and covalent radii, m.p. and b.p., enthalpy of bond dissociation and electron gain enthalpy are quite lower than expected.
- (ii) Most of the reactions of fluorine are exothermic (due to the small and strong bond formed by it with other elements).
- (iii) It forms only one oxoacid while other halogens form a number of oxoacids.
- (iv) Hydrogen fluoride is a liquid (b.p. 293 K) due to strong hydrogen bonding. Other hydrogen halides are gases.

Reason for the anomalous behaviour of fluorine

The anomalous behaviour of fluorine is due to its small size, highest electronegativity, low F-F bond dissociation enthalpy, and non availability of d orbitals in valence shell.

Properties of Hydrogen Halides


Property	HF	HCl	HBr	HI
Melting point/K	190	159	185	222
Boiling point/K	293	189	206	238
Bond length (H – X)/pm	91.7	127.4	141.4	160.9
$\Delta_{\text{diss}}\text{H}^\ominus/\text{kJ mol}^{-1}$	57.4	432	363	295
pK _a	3.2	– 7.0	– 9.5	– 10.0

- (2) Reactivity towards hydrogen :** They all react with hydrogen to give hydrogen halides but affinity for hydrogen decreases from fluorine to iodine. Hydrogen halides dissolve in water to form hydrohalic acids. Some of the properties of hydrogen halides are :

- (i) The acidic strength order : $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$
- (ii) The stability order of these halides : $\text{H-F} > \text{H-Cl} > \text{H-Br} > \text{H-I}$.

(3) Reactivity towards oxygen :

- (i) Halogens form many oxides with oxygen but most of them are unstable. Fluorine forms two oxides OF_2 and O_2F_2 . However, only OF_2 is thermally stable at 298 K. These oxides are essentially oxygen fluorides because of the higher electronegativity of fluorine than oxygen. Both are strong fluorinating agents. O_2F_2 oxidises plutonium to PuF_6 and the reaction is used in removing plutonium as PuF_6 from spent nuclear fuel.
- (ii) Chlorine, bromine and iodine form oxides in which the oxidation states of these halogens range from +1 to +7.
- (iii) A combination of kinetic and thermodynamic factors lead to the generally decreasing order of stability of oxides formed by halogens, $\text{I} > \text{Cl} > \text{Br}$.
- (iv) The higher oxides of halogens tend to be more stable than the lower ones.
- (v) Chlorine oxides, Cl_2O , ClO_2 , Cl_2O_6 and Cl_2O_7 are highly reactive oxidising agents and tend to explode. ClO_2 is used as a bleaching agent for paper pulp and textiles and in water treatment.
- (vi) The bromine oxides, Br_2O , BrO_2 , BrO_3 are the least stable halogen oxides (middle row anomaly) and exist only at low temperatures. They are very powerful oxidising agents.
- (vii) The iodine oxides, I_2O_4 , I_2O_5 , I_2O_7 are insoluble solids and decompose on heating. I_2O_5 is a very good oxidising agent and is used in the estimation of carbon monoxide.




(4) Reactivity towards metals : Halogens react with metals to form metal halides. For example, bromine reacts with magnesium to give magnesium bromide.

The ionic character of the halides decreases in the order $\text{MF} > \text{MCl} > \text{MBr} > \text{MI}$ where M is monovalent metal. If a metal exhibits more than one oxidation state, the halides in higher oxidation state will be more covalent than the one in lower oxidation state. For example, SnCl_4 , PbCl_4 , SbCl_5 and UF_6 are more covalent than SnCl_2 , PbCl_2 , SbCl_3 and UF_4 respectively.

(5) Reactivity of halogens towards other halogens : Halogens combine amongst themselves to form a number of compounds known as interhalogens of the types XX' , XX_3' , XX_5' and XX_7' where X is a larger size halogen and X' is smaller size halogen.

FLUORINE**Method of Preparation :**

Moissan process : [By electrolysis of KHF_2 (which is obtained from CaF_2)]



**BROMINE****IODINE****CHLORINE**

Chlorine was discovered in 1774 by Scheele by the action of HCl on MnO_2 .

In 1810 Davy established its elementary nature and suggested the name chlorine on account of its colour (Greek, chloros = greenish yellow).

Preparation

(i) By electrolysis of aq. NaCl :



(ii) By heating manganese dioxide with concentrated hydrochloric acid.



However, a mixture of common salt and concentrated H_2SO_4 is used in place of HCl .




(iii) By the action of HCl on potassium permanganate.



❑ Manufacture of chlorine




- (i) **Deacon's process :** By oxidation of hydrogen chloride gas by atmospheric oxygen in the presence of CuCl_2 (catalyst) at 723 K.



- (ii) **Electrolytic process :** Chlorine is obtained by the electrolysis of brine (concentrated NaCl solution). Chlorine is liberated at anode. It is also obtained as a by-product in many chemical industries.




Properties

- (i) It is a greenish yellow gas with pungent and suffocating odour.
- (ii) It is about 2-5 times heavier than air.
- (iii) It can be liquefied easily into greenish yellow liquid which boils at 239 K.
- (iv) It is soluble in water. Chlorine reacts with a number of metals and non-metals to form chlorides.





Reaction with hydrogen

It has great affinity for hydrogen. It reacts with compounds containing hydrogen to form HCl .




Reaction with ammonia

With excess ammonia, chlorine gives nitrogen and ammonium chloride whereas with excess chlorine, nitrogen trichloride (explosive) is formed.




Reaction with alkalis

With cold and dilute alkalis chlorine produces a mixture of chloride and hypochlorite but with hot and concentrated alkalis it gives chloride and chlorate.





(cold and dilute)



(hot and conc.)


but on acidification the disproportionated product gives back the same element.



[X = Cl, Br, I]

Reaction with slaked lime



With dry slaked lime it gives bleaching powder.



The composition of bleaching powder is $\text{Ca(OCl)}_2 \cdot \text{CaCl}_2 \cdot \text{Ca(OH)}_2 \cdot 2\text{H}_2\text{O}$.

Reaction with hydrocarbon




Chlorine reacts with hydrocarbons and gives substitution products with saturated hydrocarbons and addition products with unsaturated hydrocarbons. For example,





Note :

Chlorine water on standing loses its yellow colour due to the formation of HCl and HOCl. Hypochlorous acid (HOCl) so formed, gives nascent oxygen which is responsible for oxidising and bleaching properties of chlorine.

- (i) It oxidises ferrous to ferric and sulphite to sulphate. Chlorine oxidises sulphur dioxide to sulphur trioxide and iodine to iodate. In the presence of water they form sulphuric acid and iodic acid respectively.















- (ii) It is a powerful bleaching agent; bleaching action is due to oxidation.



Uses: It is used (i) for bleaching woodpulp (required for the manufacture of paper and rayon), bleaching cotton and textiles, (ii) in the extraction of gold and platinum (iii) in the manufacture of dyes, drugs and organic compounds such as CCl_4 , CHCl_3 , DDT, refrigerants, etc. (iv) in sterilising drinking water and (v) preparation of poisonous gases such as phosgene (COCl_2), tear gas (CCl_3NO_2), mustard gas ($\text{ClCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{Cl}$). (vi) It bleaches vegetable or organic matter in the presence of moisture. Bleaching effect of chlorine is permanent.



- (a) On long standing it undergoes

**Oxidising Properties:****Reaction with acid:**

Note: ClO_2 does not dimerise because odd electron undergoes delocalisation (in its own vacant 3d-orbital)



Cl_2O_4 ($\text{Cl}.\text{ClO}_4$) is not the dimer of ClO_2 . Actually it is Cl-perchlorate.

HYDROGEN CHLORIDE

- (i) Glauber prepared this acid in 1648 by heating common salt with concentrated sulphuric acid.
- (ii) Davy in 1810 showed that it is a compound of hydrogen and chlorine.

Preparation


Laboratory method : it is prepared by heating sodium chloride with concentrated sulphuric acid.



HCl gas can be dried by passing through concentrated sulphuric acid.

Properties

- (i) It is a colourless and pungent smelling gas. Due to strong affinity for water, conc. HCl pulls moisture of air towards self. The moisture forms droplets of water and hence, cloudy white fumes appear.
- (ii) It is easily liquefied to a colourless liquid (b.p. 189 K) and freezes to a white crystalline solid (f.p. 159 K).
- (iii) It is extremely soluble in water
- (iv) **Acidic character** : It ionises as follows




Its aqueous solution is called hydrochloric acid. High value of dissociation constant (K_a) indicates that it is a strong acid in water. It reacts with NH_3 and gives white fumes of NH_4Cl .






Note : Aqua regia

When three parts of concentrated HCl and one part of concentrated HNO_3 are mixed, aqua regia is formed which is used for dissolving noble metals, e.g., gold, platinum.



Reaction with salts

Hydrochloric acid decomposes salts of weaker acids, e.g., carbonates, hydrogencarbonates, sulphites, etc.



Uses:





- (i) It is used in the manufacture of chlorine, NH_4Cl and glucose (from corn starch)
- (ii) It is used for extracting glue from bones and purifying bone black
- (iii) It is used in medicine and as a laboratory reagent.

OXOACIDS OF HALOGENS

- (i) Due to high electronegativity and small size, fluorine forms only one oxoacid, HOF known as fluoric (I) acid or hypofluorous acid.
- (ii) The other halogens form several oxoacids.
- (iii) Most of them cannot be isolated in pure state. They are stable only in aqueous solutions or in the form of their salts.

Oxoacids of Halogens


Halic (I) acid (Hypohalous acid)	HO₂F (Hypofluorous acid)	HOCl (Hypochlorous acid)	HOBr (Hypobromous acid)	HOI (Hypoiodous acid)
Halic (III) acid (Halous acid)	—	HOClO (Chlorous acid)	—	—
Halic (V) acid (Halic acid)	—	HOClO₂ (Chloric acid)	HOBrO₂ (Bromic acid)	HOIO₂ (Iodic acid)
Halic (VII) acid (Perhalic acid)	—	HOClO₃ (Perchloric acid)	HOBrO₃ (Perbromic acid)	HOIO₃ (Periodic acid)



The structures of oxoacids of chlorine

INTERHALOGEN COMPOUNDS

When two different halogens react with each other, interhalogen compounds are formed. They can be assigned general compositions as XX' , XX'_3 , XX'_5 and XX'_7 where X is halogen of larger size and X' of smaller size and X is more electropositive than X' . As the ratio between radii of X and X' increases, the number of atoms per molecule also increases. Thus, iodine (VII) fluoride should have maximum number of atoms as the ratio of radii between I and F should be maximum. That is why its formula is IF_7 (having maximum number of atoms).



There are never more than two halogens in a molecule.


Bonds are essentially covalent and b.p. increases as the E.N. difference increases.

AX_5 & AX_7 type formed by large atoms like Br & I to accommodate more atoms around it.

The interhalogens are generally more reactive than the halogens (except F_2) due to weaker A–X bonds compared to X–X bond.

Preparation

The interhalogen compounds can be prepared by the direct combination or by the action of halogen on lower interhalogen compounds. The product formed depends upon some specific conditions, For e.g.,



Properties


Some properties of interhalogen compounds are given in Table below

Some Properties of Interhalogen Compounds


Type	Formula	Physical state and colour	Structure
XX' ₁	ClF	colourless gas	—
	BrF	pale brown gas	—
	IF ^a	Detected spectroscopically	—
	BrCl ^b	gas	—
	ICl	ruby red solid (α-form)	—
		brown red solid (β-form)	—
XX' ₃	IBr	black solid	—
	ClF ₃	colourless gas	Bent T-shaped
	BrF ₃	yellow green liquid	Bent T-shaped
	IF ₃	yellow powder	Bent T-shaped
	ICl ₃ ^c	orange solid	Bent T-shaped
XX' ₅	IF ₅	colourless gas but solid below 77 K	square pyramidal
	BrF ₅	colourless liquid	square pyramidal
	ClF ₅	colourless liquid	square pyramidal
XX' ₇	IF ₇	colourless gas	pentagonal bipyramidal

^aVery unstable; ^bThe pure solid is known at room temperature; ^cDimerises as Cl-bridged dimer (I₂Cl₆)

One peculiarity with ICl :




- (i) These are all covalent molecules and are diamagnetic in nature.
- (ii) They are volatile solids or liquids at 298 K except ClF which is a gas.
- (iii) Their physical properties are intermediate between those of constituent halogens except that their m.p. and b.p. are a little higher than expected.
- (iv) Their chemical reactions can be compared with the individual halogens. In general, interhalogen compounds are more reactive than halogens (except fluorine). This is because X–X' bond in interhalogens is weaker than X–X bond in halogens except F–F bond.
- (v) All these undergo hydrolysis giving halide ion derived from the smaller halogen and a hypohalite (when XX'), halite (when XX'₃), halate (when XX'₅) and perhalate (when XX'₇) anion derived from the larger halogen.









Their molecular structures are very interesting which can be explained on the basis of VSEPR theory. The XX₃ compounds have the bent 'T' shape, XX₅ compounds square pyramidal and IF₇ has pentagonal bipyramidal structures.

Uses: (i) These compounds can be used as non aqueous solvents.

(ii) Interhalogen compounds are very useful fluorinating agents. ClF₃ and BrF₃ are used for the production of UF₆ in the enrichment of ²³⁵U.






POLYHALIDES



Here the products on heating depends on the lattice energy of the product halide. The lattice energy of alkali halide with smaller halogen is highest since the interatomic distance is least.

Structure of I₅⁻, I₇⁻, I₈²⁻



I₃⁻, Br₃⁻, Cl₃⁻, F₃⁻ are known Cl₃⁻ compounds are very less.

Stability order : I₃⁻ > Br₃⁻ > Cl₃⁻ > F₃⁻ depends upon the donating ability of X⁻.

PSEUDO HALOGENS

There are univalent ion consisting of two or more atoms of which at least one is N, that have properties similar to those of the halide ions. E.g.

(i) Na-salts are soluble in water but Ag-salts are insoluble in water.

(ii) H-compounds are acids like HX.

(iii) Some anions can be oxidised to give molecules X_2 .

Anions :

Acids

Dimer

CN^-

HCN

$(CN)_2$

SCN^-

HSCN(thiocyanic acid)

$(SCN)_2$

$SeCN^-$

$(SeCN)_2$

OCN^-

HOCN (cyanic acid)

NCN^{2-} (Bivalent)

H_2NCN (cyanamide)

ONC^-

HONC (Fulminic acid)

N_3^-

HN_3 (Hydrazoic acid)

$CN^\ominus$ shows maximum similarities with Cl^- , Br^- , I^-

(i) forms HCN

(ii) forms $(CN)_2$

(iii) AgCN, $Pb(CN)_2$, are insoluble

(iv) Interpseudo halogen compounds ClCN, BrCN, ICN can be formed

(v) AgCN is insoluble in H_2O but soluble in NH_3

(vi) forms large no. of complexes. e.g. $[Cu(CN)_4]^{3-}$ & $[CuCl_4]^{-3}$

NOBLE GASES FAMILY

GROUP 18 ELEMENTS (He, Ne, Ar, Kr, Xe, Rn)

Occurrence

(i) All the noble gases except radon occur in the atmosphere.

Relative abundance : Ar is highest (Ne, Kr, He, Rn)


(ii) Their atmospheric abundance in dry air is $\sim 1\%$ by volume of which argon is the major constituent.

(iii) Helium and sometimes neon are found in minerals of radioactive origin e.g., pitchblende, monazite, cleveite.

(iv) The main commercial source of helium is natural gas.

(v) Xenon and radon are the rarest elements of the group.

(vi) Radon is obtained as a decay product of ^{226}Ra .



(vii) He liquid can exist in two forms. I-form when changes to II-form at λ -point temperature many physical properties change abruptly.

e.g.

(i) Sp. heat changes by a factor of 10

(ii) Thermal conductivity increases by 10^6 and it becomes 800 times faster than Cu

(iii) It shows zero resistance

(iv) It can flow up the sides of the vessel

Qus. Why 18 group element are termed as noble gas ?

Ans. All these are gases and chemically unreactive. They form very few compounds. Because of this they are termed noble gases.


❑ **Electronic Configuration**

General electronic configuration of 18 group element is ns^2np^6 except helium which has $1s^2$.

Many of the properties of noble gases including their inactive nature are ascribed to their closed shell structures.

❑ **Ionisation Enthalpy**


Ionisation energy decreases down the group with increase in atomic size.



Due to stable electronic configuration these gases exhibit very high ionisation enthalpy.

❑ **Atomic Radii**

Atomic radii increase down the group with increase in atomic number.



❑ **Electron Gain Enthalpy**

They have large positive values of electron gain enthalpy due to stable electronic configurations, and therefore have no tendency to accept the electron

❑ **Melting point and boiling point**

$\text{He} < \text{Ne} < \text{Ar} < \text{Kr} < \text{Xe} < \text{Rn}$ (Melting point order)

↓

(-269°C)

B.P. order : $\text{He} < \text{Ne} < \text{Ar} < \text{Kr} < \text{Xe} < \text{Rn}$ (Boiling point order)

❑ **Density order :**



Physical properties :

(i) All the noble gases are monoatomic.

(ii) They are colourless, odourless and tasteless.

(iii) They are sparingly soluble in water.

(iv) They have very low melting and boiling points because the only type of interatomic interaction in these elements is weak dispersion forces.

(v) Helium has the lowest boiling point (4.2 K) of any known substance.

(vi) It has an unusual property of diffusing through most commonly used laboratory materials such as rubber, glass or plastics.

❑ Chemical Properties





In general, noble gases are least reactive. Their inertness to chemical reactivity is attributed to the following reasons:

- (i) The noble gases except helium ($1s^2$) have completely filled ns^2np^6 electronic configuration in their valence shell.
- (ii) They have high ionisation enthalpy and more positive electron gain enthalpy.

Note : The reactivity of noble gases has been investigated occasionally. In March 1962, Neil Bartlett, then at the University of British Columbia, observed the reaction of a noble gas. First, he prepared a red compound which is formulated as $O_2^+ PtF_6^-$. He, then realised that the first ionisation enthalpy of molecular oxygen (1175 kJ mol^{-1}) was almost identical with that of xenon (1170 kJ mol^{-1}). He made efforts to prepare same type of compound with Xe and was successful in preparing another red colour compound $Xe^+PtF_6^-$ by mixing PtF_6 and xenon. After this discovery, a number of xenon compounds mainly with most electronegative elements like fluorine and oxygen, have been synthesised. The compounds of krypton are fewer. Only the difluoride (KrF_2) has been studied in detail. Compounds of radon have not been isolated but only identified (e.g., RnF_2) by radiotracer technique. No true compounds of Ar, Ne or He are yet known.

FLUORIDES OF XENON

Preparation




Physical properties


XeF_2 , XeF_4 and XeF_6 are colourless crystalline solids and sublime readily at 298 K.


Chemical properties

(i) **Hydrolysis :** They are readily hydrolysed even by traces of water. For example, XeF_2 is hydrolysed to give Xe, HF and O_2 .





Hydrolysis of XeF_4 and XeF_6 with water gives XeO_3 .






Partial hydrolysis of XeF_6 gives oxyfluorides, XeOF_4 and XeO_2F_2 .






Note : **Hydrolysis in alkaline medium**





Xenate ion






(ii) **As fluorinating agents** : They are powerful fluorinating agents.



(iii) **As fluoride donor**





(iv) **As Fluoride acceptor**



(alkali metals fluoride)

(v) **Reaction with SiO_2**

SiO_2 also converts XeF_6 into XeOF_4





Similarly, $\text{XeO}_3 + \text{XeOF}_4 \longrightarrow 2\text{XeO}_2\text{F}_2$

(vi) Oxidizing properties

H_2 reduces Xe – fluorides to Xe



Xe - fluorides oxidise Cl^- to Cl_2 and I^- to I_2




Noble gas hydrate (clathrate compound) : Ar, Kr, Xe can form clathrate compounds but He, Ne cannot due to their smaller size.

eg. $\text{Xe} \cdot 6\text{H}_2\text{O}$	} formed only when
$\text{Ar} \cdot 6\text{H}_2\text{O}$	
$\text{Kr} \cdot 6\text{H}_2\text{O}$	


water freezes at high pressure together with noble gas

(a) Structure and bonding


XeF_2 and XeF_4 have linear and square planar structures respectively. XeF_6 has seven electron pairs (6 bonding pairs and one lone pair) and would, thus, have a distorted octahedral structure as found experimentally in the gas phase. Xenon fluorides react with fluoride ion acceptors to form cationic species and fluoride ion donors to form fluoroanions.



(a) Linear




(b) Square planar



(c) Distorted octahedral


(b) Xenon-oxygen compounds

XeO_3 is a colourless, white hygroscopic explosive solid and has a pyramidal molecular structure.



Pyramidal

XeOF_4 is a colourless volatile liquid and has a square pyramidal molecular structure.



Square pyramidal

Uses of helium :

- (i) He is a non-inflammable and light gas. Hence, it is used in filling balloons for meteorological observations.
- (ii) It is also used in gas-cooled nuclear reactors.
- (iii) It is used in cryoscopy to obtain the very low temperature required for superconductor and laser (b.p. 4.2 K) finds use as cryogenic agent for carrying out various experiments at low temperatures.
- (iv) It is used to produce and sustain powerful superconducting magnets which form an essential part of modern NMR spectrometers and Magnetic Resonance Imaging (MRI) systems for clinical diagnosis.
- (v) It is used as a diluent for oxygen in modern diving apparatus because of its very low solubility in blood. He is used in preference to N_2 to dil. O_2 in the gas cylinders used by divers. This is because N_2 is quite soluble in blood, so a sudden change in pressure causes degassing and gives bubbles of N_2 in the blood. This causes the painful condition called bends. He is slightly soluble so the risk of bends is reduced.

USES OF NEON :

- (i) Ne is used in discharge tubes and fluorescent bulbs for advertisement display purposes.
- (ii) Neon bulbs are used in botanical gardens and in green houses.

USES OF ARGON :

- (i) Argon is used mainly to provide an inert atmosphere in high temperature metallurgical processes (arc welding of metals or alloys) and for filling electric bulbs.
- (ii) It is also used in the laboratory for handling substances that are air-sensitive.

USES OF XENON AND KRYPTON :

There are no significant uses of Xenon and Krypton. They are used in light bulbs designed for special purposes.

SOLVED EXAMPLE

1. Though nitrogen exhibits +5 oxidation state, it does not form pentahalide. Give reason.


Sol. Nitrogen with $n = 2$, has s and p orbitals only. It does not have d orbitals to expand its covalency beyond four. That is why it does not form pentahalide.

2. PH_3 has lower boiling point than NH_3 . Why?

Sol. Unlike NH_3 , PH_3 molecules are not associated through hydrogen bonding in liquid state. That is why the boiling point of PH_3 is lower than NH_3 .

3. Write the reaction of thermal decomposition of sodium azide.

Sol. Thermal decomposition of sodium azide gives dinitrogen gas.





4. Why does NH_3 act as a Lewis base ?

Sol. Nitrogen atom in NH_3 has one lone pair of electrons which is available for donation. Therefore, it acts as a Lewis base.


5. NH_3 can't be dried by H_2SO_4 , P_2O_5 and anhydrous CaCl_2

Sol. because : $2\text{NH}_3 + \text{H}_2\text{SO}_4 \longrightarrow (\text{NH}_4)_2 \text{SO}_4$



forms adduct

Quick lime is used for this purpose



$\text{NH}_3 \text{ (base)}$ } *Hence no interaction*

6. Why does NO_2 dimerise ?

Sol. NO_2 contains odd number of valence electrons. It behaves as a typical odd molecule. On dimerisation, it is converted to stable N_2O_4 molecule with even number of electrons.

7. In what way can it be proved that PH_3 is basic in nature?


Sol. PH_3 reacts with acids like HI to form PH_4I which shows that it is basic in nature.



Due to lone pair on phosphorus atom, PH_3 is acting as a Lewis base in the above reaction.

8. Why does PCl_3 fume in moisture ?

Sol. PCl_3 hydrolyses in the presence of moisture giving fumes of HCl.



9. Are all the five bonds in PCl_5 molecule equivalent? Justify your answer.

Sol. PCl_5 has a trigonal bipyramidal structure and the three equatorial P-Cl bonds are equivalent, while the two axial bonds are different and longer than equatorial bonds.

10. How do you account for the reducing behaviour of H_3PO_2 on the basis of its structure?

Sol. In H_3PO_2 , two H atoms are bonded directly to P atom which imparts reducing character to the acid.

11. Elements of Group 16 generally show lower value of first ionisation enthalpy compared to the corresponding periods of group 15. Why?

Sol. Due to extra stable half-filled p orbitals electronic configurations of Group 15 elements, larger amount of energy is required to remove electrons compared to Group 16 elements.

12. H_2S is less acidic than H_2Te . Why?

Sol. Due to the decrease in bond (E–H) dissociation enthalpy down the group, acidic character increases.


13. Which form of sulphur shows paramagnetic behaviour ?

Sol. In vapour state sulphur partly exists as S_2 molecule which has two unpaired electrons in the antibonding π^* orbitals like O_2 and, hence, exhibits paramagnetism.


14. What happens when

- (i) Concentrated H_2SO_4 is added to calcium fluoride
- (ii) SO_3 is passed through water?

Sol. (i) It forms hydrogen fluoride



- (ii) It dissolves SO_3 to give H_2SO_4 .



15. Halogens have maximum negative electron gain enthalpy in the respective periods of the periodic table. Why?

Sol. Halogens have the smallest size in their respective periods and therefore high effective nuclear charge. As a consequence, they readily accept one electron to acquire noble gas electronic configuration.

16. Although electron gain enthalpy of fluorine is less negative as compared to chlorine, fluorine is a stronger oxidising agent than chlorine. Why?

Sol. It is due to

- (i) low enthalpy of dissociation of F–F bond
- (ii) high hydration enthalpy of F^-

17. Fluorine exhibits only –1 oxidation state whereas other halogens exhibit +1, +3, +5 and +7 oxidation states also. Explain.

Sol. Fluorine is the most electronegative element and cannot exhibit any positive oxidation state. Other halogens have d orbitals and therefore, can expand their octets and show +1, +3, +5 and +7 oxidation states also.

18. Write the balanced chemical equation for the reaction of Cl_2 with hot and concentrated NaOH . Is this reaction a disproportionation reaction? Justify.


Sol. $3\text{Cl}_2 + 6\text{NaOH} \rightarrow 5\text{NaCl} + \text{NaClO}_3 + 3\text{H}_2\text{O}$

Yes, chlorine from zero oxidation state is changed to –1 and +5 oxidation states.

19. CaF_2 used in HF prepⁿ. must be free from SiO_2 . Explain.

Ans. $\text{CaF}_2 + \text{H}_2\text{SO}_4 \longrightarrow \text{CaSO}_4 + \text{HF}$


If SiO_2 present as impurity



HF can not be stored in glass vessel due to same reason.

20. When HCl reacts with finely powdered iron, it forms ferrous chloride and not ferric chloride. Why?


Sol. Its reaction with iron produces H_2 .



Liberation of hydrogen prevents the formation of ferric chloride.

21. Discuss the molecular shape of BrF_3 on the basis of VSEPR theory.

Sol. The central atom Br has seven electrons in the valence shell. Three of these will form electronpair bonds with three fluorine atoms leaving behind four electrons. Thus, there are three bond pairs and two lone pairs. According to VSEPR theory, these will occupy the corners of a trigonal bipyramid. The two lone pairs will occupy the equatorial positions to minimise lone pair-lone pair and the bond pair-lone pair repulsions which are greater than the bond pair-bond pair repulsions. In addition, the axial fluorine atoms will be bent towards the equatorial fluorine in order to minimise the lone-pair-lone pair repulsions. The shape would be that of a slightly bent 'T'.



22. Why are the elements of Group 18 known as noble gases ?

Sol. The elements present in Group 18 have their valence shell orbitals completely filled and, therefore, react with a few elements only under certain conditions. Therefore, they are now known as noble gases.

23. Noble gases have very low boiling points. Why?

Sol. Noble gases being monoatomic have no interatomic forces except weak dispersion forces and therefore, they are liquefied at very low temperatures. Hence, they have low boiling points.

24. Does the hydrolysis of XeF_6 lead to a redox reaction?

Sol. No, the products of hydrolysis are XeOF_4 and XeO_2F_2 where the oxidation states of all the elements remain the same as it was in the reacting state.

25. Standard electrode potential values, $E^\ominus$ for Al^{3+}/Al is -1.66 V and that of Tl^{3+}/Tl is $+1.26\text{ V}$. Predict about the formation of M^{3+} ion in solution and compare the electropositive character of the two metals.

Sol. Standard electrode potential values for two half cell reactions suggest that aluminium has high tendency to make $\text{Al}^{3+}(\text{aq})$ ions, whereas Tl^{3+} is not only unstable in solution but is a powerful oxidising agent also. Thus Tl^+ is more stable in solution than Tl^{3+} . Aluminium being able to form +3 ions easily, is more electropositive than thallium.

26. White fumes appear around the bottle of anhydrous aluminium chloride. Give reason.

Sol. Anhydrous aluminium chloride is partially hydrolysed with atmospheric moisture to liberate HCl gas. Moist HCl appears white in colour.

27. Boron is unable to form BF_6^{3-} ion. Explain.

Sol. Due to non-availability of d orbitals, boron is unable to expand its octet. Therefore, the maximum covalence of boron cannot exceed 4.

28. Why is boric acid considered as a weak acid ?

Sol. Because it is not able to release H^+ ions on its own. It receives OH^- ions from water molecule to complete its octet and in turn releases H^+ ions.

29. Select the member(s) of group 14 that (i) forms the most acidic dioxide, (ii) is commonly found in +2 oxidation state, (iii) used as semiconductor.

Sol. (i) carbon (ii) lead (iii) silicon and germanium

30. $[\text{SiF}_6]^{2-}$ is known whereas $[\text{SiCl}_6]^{2-}$ not. Give possible reasons.

Sol. The main reasons are :

(i) six large chloride ions cannot be accommodated around Si^{4+} due to limitation of its size.

(ii) interaction between lone pair of chloride ion and Si^{4+} is not very strong.

31. Diamond is covalent, yet it has high melting point. Why ?

Sol. Diamond has a three-dimensional network involving strong $\text{C}-\text{C}$ bonds, which are very difficult to break and, in turn has high melting point.

32. SiH_4 is more reactive than CH_4 . Explain

Reasons

(i) $\text{Si}^{\delta+} - \text{H}^{\delta-}$ in $\text{C}^{\delta-} - \text{H}^{\delta+}$

C - more electronegative than H

Si less electronegative than H

So bond polarity is reversed when Nu^- attacks, it faces repulsion in C but not in Si

(ii) Silicon is having vacant d orbital which is not in case of carbon

(iii) Silicon is larger in size compared to C . By which the incoming Nu^- doesn't face any steric hindrance to attack at Si whereas CH_4 is tightly held from all sides.

QUESTION BANK ON p-BLOCK ELEMENTS

EXERCISE # I

Only one option is correct :

1. PH_3 (Phosphine) when passed in aqueous solution of CuSO_4 it produce -

(A) Blue precipitate of $\text{Cu}(\text{OH})_2$ (B) dark blue solution of $[\text{Cu}(\text{PH}_3)_4]\text{SO}_4$
 (C) Black precipitate of Cu_3P_2 (D) Colorless solution of $[\text{Cu}(\text{H}_2\text{O})_4]^+$
2. $\text{H}_3\text{PO}_2 \xrightarrow{\Delta} (\text{X}) + \text{PH}_3$; is

(A) Dehydration reaction (B) Oxidation reaction
 (C) Disproportionation reaction (D) Dephosphorelation reaction
3. Which of the following species is not a pseudohalide?

(A) CNO^- (B) RCOO^- (C) OCN^- (D) N_3^-
4. An orange solid (X) on heating, gives a colourless gas (Y) and a only green residue (Z). Gas (Y) on treatment with Mg, produces a white solid substance

(A) Mg_3N_2 (B) MgO (C) Mg_2O_3 (D) MgCl_2
5. Conc. HNO_3 is yellow coloured liquid due to

(A) dissolution of NO in conc. HNO_3 (B) dissolution of NO_2 in conc. HNO_3
 (C) dissolution of N_2O in conc. HNO_3 (D) dissolution of N_2O_3 in conc. HNO_3
6. A gas at low temperature does not react with the most of compounds. It is almost inert and is used to create inert atmosphere in bulbs. The combustion of this gas is exceptionally an endothermic reaction. Based on the given information, we can conclude that the gas is

(A) oxygen (B) nitrogen
 (C) carbon mono-oxide (D) hydrogen
7. When chlorine gas is passed through an aqueous solution of a potassium halide in the presence of chloroform, a violet colouration is obtained. On passing more of chlorine water, the violet colour is disappeared and solution becomes colourless. This test confirms the presence of in aqueous solution.

(A) chlorine (B) fluorine (C) bromine (D) iodine
8. $\text{H}_3\text{PO}_2 \xrightarrow{140^\circ\text{C}} \text{A} \xrightarrow{220^\circ\text{C}} \text{B} \xrightarrow{320^\circ\text{C}} \text{C}$
 Compound (C) is

(A) H_2PO_3 (B) H_3PO_3 (C) $(\text{HPO}_3)_n$ (D) $\text{H}_4\text{P}_2\text{O}_7$
9. An explosive compound (A) reacts with water to produce NH_4OH and HOCl . Then, the compound (A), is


(A) TNG (B) NCl_3 (C) PCl_3 (D) HNO_3
10. An inorganic compound (A) made of two most occurring elements into the earth crust, having a polymeric tetrahedral network structure. With carbon, compound (A) produces a poisonous gas (B) which is the most stable diatomic molecule. Compounds (A) and (B) will be

(A) SiO_2 , CO_2 (B) SiO_2 , CO (C) SiC , CO (D) SiO_2 , N_2


11. A sulphate of a metal (A) on heating evolves two gases (B) and (C) and an oxide (D). Gas (B) turns $K_2Cr_2O_7$ paper green while gas (C) forms a trimer in which there is no S–S bond. Compound (D) with HCl, forms a Lewis acid (E) which exists as a dimer. Compounds (A), (B), (C), (D) and (E) are respectively
 (A) $FeSO_4$, SO_2 , SO_3 , Fe_2O_3 , $FeCl_3$ (B) $Al_2(SO_4)_3$, SO_2 , SO_3 , Al_2O_3 , $FeCl_3$
 (C) FeS , SO_2 , SO_3 , $FeSO_4$, $FeCl_3$ (D) FeS , SO_2 , SO_3 , $Fe_2(PO_4)_3$, $FeCl_2$
12. A tetra-atomic molecule (A) on reaction with nitrogen(I)oxide, produces two substances (B) and (C). (B) is a dehydrating agent while substance (C) is a diatomic gas which shows almost inert behaviour. The substances (A) and (B) and (C) respectively will be
 (A) P_4 , P_4O_{10} , N_2 (B) P_4 , N_2O_5 , N_2 (C) P_4 , P_2O_3 , Ar (D) P_4 , P_2O_3 , H_2
13. First compound of inert gases was prepared by scientist Neil Bartlett in 1962. This compound is
 (A) $XePtF_6$ (B) XeO_3 (C) XeF_6 (D) $XeOF_4$
14. Carbongene has X% of CO_2 and is used as an antidote for poisoning of Y. Then, X and Y are
 (A) X = 95% and Y = lead poisoning (B) X = 5% and Y = CO poisoning
 (C) X = 30% and Y = CO_2 poisoning (D) X = 45% and Y = CO poisoning
15. The correct order of acidic strength of oxides of nitrogen is
 (A) $NO < NO_2 < N_2O < N_2O_3 < N_2O_5$ (B) $N_2O < NO < N_2O_3 < N_2O_4 < N_2O_5$
 (C) $NO < N_2O < N_2O_3 < N_2O_5 < N_2O_4$ (D) $NO < N_2O < N_2O_5 < N_2O_3 < N_2O_4$
16. $H_3BO_3 \xrightarrow{T_1} X \xrightarrow{T_2} Y \xrightarrow{\text{redhot}} B_2O_3$
 if $T_1 < T_2$ then X and Y respectively are
 (A) X = Metaboric acid and Y = Tetraboric acid
 (B) X = Tetraboric acid and Y = Metaboric acid
 (C) X = Borax and Y = Metaboric acid
 (D) X = Tetraboric acid and Y = Borax
17. When conc. H_2SO_4 was treated with $K_4[Fe(CN)_6]$, CO gas was evolved. By mistake, somebody used dilute H_2SO_4 instead of conc. H_2SO_4 then the gas evolved was
 (A) CO (B) HCN (C) N_2 (D) CO_2
18. An inorganic white crystalline compound (A) has a rock salt structure. (A) on reaction with conc. H_2SO_4 and MnO_2 , evolves a pungent smelling, greenish-yellow gas (B). Compound (A) gives white ppt. of (C) with $AgNO_3$ solution. Compounds (A), (B) and (C) will be respectively
 (A) $NaCl$, Cl_2 , $AgCl$ (B) $NaBr$, Br_2 , $NaBr$
 (C) $NaCl$, Cl_2 , Ag_2SO_4 (D) Na_2CO_3 , CO_2 , Ag_2CO_3
19. $RCl \xrightarrow[Si]{Cu\text{-powder}} R_2SiCl_2 \xrightarrow{H_2O} R_2Si(OH)_2 \xrightarrow{\text{condensation}} A$
 Compound (A) is
 (A) a linear silicone (B) a chlorosilane (C) a linear silane (D) a network silane

20. When oxalic acid reacts with conc. H_2SO_4 , two gases produced are of neutral and acidic in nature respectively. Potassium hydroxide absorbs one of the two gases. The product formed during this absorption and the gas which gets absorbed are respectively
(A) K_2CO_3 and CO_2 (B) KHCO_3 and CO_2
(C) K_2CO_3 and CO (D) KHCO_3 and CO
21. Conc. H_2SO_4 cannot be used to prepare HBr from NaBr because it
(A) reacts slowly with NaBr (B) oxidises HBr
(C) reduces HBr (D) disproportionates HBr
22. Ammonia can be dried by
(A) conc. H_2SO_4 (B) P_4O_{10} (C) CaO (D) anhydrous CaCl_2
23. When chlorine reacts with a gas X, an explosive inorganic compound Y is formed. Then X and Y will be
(A) $\text{X} = \text{O}_2$ and $\text{Y} = \text{NCl}_3$ (B) $\text{X} = \text{NH}_3$ and $\text{Y} = \text{NCl}_3$
(C) $\text{X} = \text{O}_2$ and $\text{Y} = \text{NH}_4\text{Cl}$ (D) $\text{X} = \text{NH}_3$ and $\text{Y} = \text{NH}_4\text{Cl}$
24. $\text{HNO}_3 + \text{P}_4\text{O}_{10} \longrightarrow \text{HPO}_3 + \text{A}$; the product A is
(A) N_2O (B) N_2O_3 (C) NO_2 (D) N_2O_5
25. Which of the following is the correct order of acidic strength?
(A) $\text{Cl}_2\text{O}_7 > \text{SO}_3 > \text{P}_4\text{O}_{10}$ (B) $\text{CO}_2 > \text{N}_2\text{O}_5 > \text{SO}_3$
(C) $\text{Na}_2\text{O} > \text{MgO} > \text{Al}_2\text{O}_3$ (D) $\text{K}_2\text{O} > \text{CaO} > \text{MgO}$
26. $\text{Ca} + \text{C}_2 \longrightarrow \text{CaC}_2 \xrightarrow{\text{N}_2} \text{A}$
Compound (A) is used as a/an
(A) fertilizer (B) dehydrating agent (C) oxidising agent (D) reducing agent
27. A gas which exists in three allotropic forms α , β and γ is
(A) SO_2 (B) SO_3 (C) CO_2 (D) NH_3
28. A red coloured mixed oxide (X) on treatment with conc. HNO_3 gives a compound (Y). (Y) with HCl produces a chloride compound (Z) which can also be produced by treating (X) with conc. HCl . Compounds (X), (Y), and (Z) will be
(A) Mn_3O_4 , MnO_2 , MnCl_2 (B) Pb_3O_4 , PbO_2 , PbCl_2
(C) Fe_3O_4 , Fe_2O_3 , FeCl_2 (D) Fe_3O_4 , Fe_2O_3 , FeCl_3
29. One mole of calcium phosphide on reaction with excess of water gives
(A) one mole of phosphine (B) two moles of phosphoric acid
(C) two moles of phosphine (D) one mole of phosphorus penta-oxide
30. $\text{NaH}_2\text{PO}_4 \xrightarrow{> 240^\circ\text{C}} (\text{NaPO}_3)_3 \xrightarrow{625^\circ\text{C}} \text{NaPO}_3(\text{liquid melt}) \xrightarrow[\text{cooling}]{\text{rapid}} \text{D}(\text{glass})$
Sodium trimetaphosphate
Compound (D) is known as
(A) Microcosmic salt (B) Graham's salt (C) Fischer's salt (D) Switzer's Salt

31. Three allotropes (A), (B) and (C) of phosphorous in the following change are respectively



- (A) white, β -black, red (B) β -black, white, red
 (C) red, β -black, white (D) red, violet, β -black
32. When an inorganic compound reacts with SO_2 in aqueous medium, produces (A). (A) on reaction with Na_2CO_3 , gives compound (B) which with sulphur, gives a substance (C) used in photography. Compound (C) is
- (A) Na_2S (B) $\text{Na}_2\text{S}_2\text{O}_7$ (C) Na_2SO_4 (D) $\text{Na}_2\text{S}_2\text{O}_3$
33. $\text{B(OH)}_3 + \text{NaOH} \rightleftharpoons \text{NaBO}_2 + \text{Na[B(OH)}_4] + \text{H}_2\text{O}$
 How can this reaction is made to proceed in forward direction?
- (A) addition of cis 1,2 diol (B) addititon of borax
 (C) addition of trans 1,2 diol (D) addition of Na_2HPO_4
34. Which is the compound responsible for the flickering light called **will-o-the-wisp**, some times seen in the Marsh.
- (A) PH_3 (B) P_2H_4 (C) H_2S (D) $\text{PH}_3 + \text{H}_2\text{S}$
35. The gun powder is consisting of '_____' + sulphur + Charcoal what is the missing substance for gun powder
- (A) LiNO_3 (B) NH_4NO_2 (C) KNO_3 (D) (A) and (B) mixture
36. An aqueous solution of borax is
- (A) Neutral (B) Amphoteric (C) Basic (D) Acidic
37. Boric acid is polymeric due to
- (A) Its acidic nature (B) The presence of hydrogen bonds
 (C) Its monobasic nature (D) Its geometry
38. The type of hybridisation of boron in diborane is
- (A) sp (B) sp^2 (C) sp^3 (D) dsp^2
39. Thermodynamically the most stable form of carbon is
- (A) Diamond (B) Graphite (C) Fullerenes (D) Coal
40. Elements of group 14
- (A) Exhibit oxidation state of +4 only (B) Exhibit oxidation state of +2 and +4 only
 (C) Form M^{2-} and M^{4+} ions (D) Form M^{2+} and M^{4+} ions



if A is a basic gas then identified (A) and (B)

- (A) NH_3 , NH_4Br (B) NH_3 , N_2O (C) NH_3 , N_2O_5 (D) None of these

Question No. 50 to 55 (6 questions)

Questions given below consist of two statements each printed as Assertion (A) and Reason (R); while answering these questions you are required to choose any one of the following four responses:

- (A) if both (A) and (R) are true and (R) is the correct explanation of (A)
 (B) if both (A) and (R) are true but (R) is not correct explanation of (A)
 (C) if (A) is true but (R) is false
 (D) if (A) is false and (R) is true

42. **Assertion :** Borax bead test is applicable only to coloured salt.

Reason : In borax bead test, coloured salts are decomposed to give coloured metal meta borates.

43. **Assertion :** Aluminium and zinc metal evolve H_2 gas from NaOH solution

Reason : Several non-metals such as P, S, Cl, etc. convert a hydride instead of producing H_2 gas from NaOH.

44. **Assertion :** Conc. H_2SO_4 can not be used to prepare pure HBr from NaBr

Reason : It reacts slowly with NaBr.

45. **Assertion :** Oxygen is more electronegative than sulphur, yet H_2S is acidic, while H_2O is neutral.

Reason : H–S bond is weaker than O–H bond.

46. **Assertion :** Chlorine gas disproportionates in hot & conc. NaOH solution.

Reason : NaCl and NaOCl are formed in the above reaction.

47. **Assertion :** Liquid IF_5 conducts electricity.


Reason : Liquid IF_5 self ionizes as, $2IF_5 \rightleftharpoons IF_4^+ + IF_6^-$

EXERCISE # II

One or more than one option may be correct :



1. When a compound X reacts with ozone in aqueous medium, a compound Y is produced. Ozone also reacts with Y and produces compound Z. Z acts as an oxidising agent, then X, Y and Z will be
 - (A) $X = \text{HI}$, $Y = \text{I}_2$ and $Z = \text{HIO}_3$
 - (B) $X = \text{KI}$, $Y = \text{I}_2$ and $Z = \text{HIO}_3$
 - (C) $X = \text{KI}$, $Y = \text{I}_2$ and $Z = \text{HIO}_4$
 - (D) $X = \text{HI}$, $Y = \text{I}_2$ and $Z = \text{HIO}_4$
2. Which of the following statements is/are correct regarding B_2H_6 ?
 - (A) banana bonds are longer but stronger than normal B–H bonds
 - (B) B_2H_6 is also known as 3c–2e compound
 - (C) the hybrid state of B in B_2H_6 is sp^3 while that of sp^2 in BH_3
 - (D) it cannot be prepared by reacting BF_3 with LiBH_4 in the presence of dry ether
3. Which of the following statements is/are correct regarding inter-halogen compounds of AB_x type?
 - (A) x may be 1,3,5 and 7
 - (B) A is a more electronegative halogen than B
 - (C) FBr_3 cannot exist
 - (D) The interhalogens are generally more reactive than the halogens (except F_2) due to weaker A–X bonds compared to X–X bond.
4. When an inorganic compound (X) having 3c-2e as well as 2c-2e bonds reacts with ammonia gas at a certain temperature, gives a compound (Y) iso-structural with benzene. Compound (X) with ammonia at a high temperature, produces a slippery substance (Z). Then
 - (A) (X) is B_2H_6
 - (B) (Z) is known as inorganic graphite
 - (C) (Z) having structure similar to graphite
 - (D) (Z) having structure similar to (X)
5. Boric acid
 - (A) exists in polymeric form due to inter-molecular hydrogen bonding.
 - (B) is used in manufacturing of optical glasses.
 - (C) is a tri-basic acid
 - (D) with borax, it is used in the preparation of a buffer solution.

6. The correct statement(s) related to allotropes of carbon is/are
- (A) graphite is the thermodynamically most stable allotrope of carbon and having a two dimensional sheet like structure of hexagonal rings of carbon (sp^2)
- (B) diamond is the hardest allotrope of carbon and having a three dimensional network structure of $C(sp^3)$
- (C) fullerene (C_{60}) is recently discovered non-crystalline allotrope of carbon having a football-like structure.
- (D) Vander Waal's force of attraction acts between the layers of graphite $6.14 \text{ \AA}$ away from each other
7. $Al_2(SO_4)_3 + NH_4OH \longrightarrow X$, then
- (A) X is a white coloured compound (B) X is insoluble in excess of NH_4OH
- (C) X is soluble in NaOH (D) X cannot be used as an antacid
8. The species that undergo(es) disproportionation in an alkaline medium is/are
- (A) Cl_2 (B) MnO_4^{2-} (C) P_4 (D) ClO_4^-
9. Select correct statement(s):
- (A) Borax is used as a buffer
- (B) 1 M borax solution reacts with equal volumes of 2 M HCl solution
- (C) Titration of borax can be made using methyl orange as the indicator
- (D) Coloured bead obtained in borax-bead test contains metaborate
10. Which of the following is / are correct for group 14 elements?
- (A) The stability of dihalides are in the order $CX_2 < SiX_2 < GeX_2 < SnX_2 < PbX_2$
- (B) The ability to form $p\pi-p\pi$ multiple bonds among themselves increases down the group
- (C) The tendency for catenation decreases down the group
- (D) They all form oxides with the formula MO_2 .
11. Zeolite is used in which of the following cases :
- (A) Conversion of alcohols into gasoline (B) Cracking of hydrocarbon
- (C) Isomerisation of hydrocarbons (D) Softening of hard water
12. Which of the following oxides are mixed oxide ?
- (A) PbO_2 (B) SnO_2 (C) Pb_2O_3 (D) Pb_3O_4
13. Which of the following oxide(s) gives brown ppt on reaction with conc. HNO_3 :
- (A) PbO (B) SnO (C) Pb_2O_3 (D) Pb_3O_4
14. Which of the following reaction produces PH_3 :
- (A) $Ca_3P_2 + H_2O \rightarrow$ (B) $P_4 + NaOH \rightarrow$ (C) $PH_4I + KOH \rightarrow$ (D) $H_3PO_2 \xrightarrow{\Delta}$
15. Which of the following halides is least stable and has doubtful existence ?
- (A) CCl_4 (B) GeI_4 (C) SnI_4 (D) PbI_4



Correct statement for [X]

- (A) Structure of anion of crystalline (X) has one boron atom sp^3 hybridised and other three boron atoms sp^2 hybridised
- (B) (X) with NaOH(aq.) gives a compound which on reaction with H_2O_2 in alkaline medium yields a compound used as brightner in soaps
- (C) Hydrolysis of (X) with HCl or H_2SO_4 yields a compound which on reaction with HF gives fluoroboric acid
- (D) [X] on heating with cobalt salt in oxidising flame gives blue coloured bead




If A is an example of 3-d silicate then select the correct statements about (C)

- (A) Central atom of C is sp^3 hybridised
- (B) (C) is non planar and all atoms are sp^3 hybridised
- (C) (C) has diamond like structure, and it is coloured when impurity is present but pale yellow to colourless solid at room temperature
- (D) (C) is silicon carbide (SiC) and it is not being affected by any acid except H_3PO_4



EXERCISE # III

Paragraph for Question No. 1 & 2



- When D react with $\text{C}_2\text{H}_5\text{OH}$ then product will be
 (A) $\text{C}_2\text{H}_5\text{Cl}$, H_3PO_4 (B) $\text{C}_2\text{H}_5\text{Cl}$, H_3PO_3
 (C) CH_3COCl , H_3PO_3 (D) Only H_3PO_3
- B can be absorbed by :
 (A) Ca(OCl)Cl (B) H_2S (C) Both (D) None

Paragraph for Question No. 3 to 6





- Select the correct statement about (X)
 (A) (X) is a colourless, odourless and almost water insoluble gas
 (B) X is highly poisonous and burns with blue flame
 (C) When (X) gas is passed through PdCl_2 solution giving rise to black ppt
 (D) All of these
- Mixture of (X) gas + H_2 is called
 (A) Water gas or synthesis gas (B) Producer gas
 (C) Methane gas (D) None of these
- In second reaction when air is used instead of steam a mixture of (X) gas and N_2 is produced which is called
 (A) Water gas (B) Synthesis gas (C) Producer gas (D) Carbon dioxide gas
- Select the correct statement about (X)
 (A) (X) gas is estimated by I_2O_5 (B) Cu_2Cl_2 is absorber of (X) gas
 (C) (X) gas is the purifying agent for Ni (D) All of these

Paragraph for Question No. 7 & 8


Read the following write-ups and answer the questions at the end of it.

Silicons are synthetic polymers containing repeated R_2SiO units. Since, the empirical formula is that of a ketone (R_2CO), the name silicone has been given to these materials. Silicones can be made into oils, rubbery elastomers and resins. They find a variety of applications because of their chemical inertness, water repelling nature, heat-resistance and good electrical insulating property.

Commercial silicon polymers are usually methyl derivatives and to a lesser extent phenyl derivatives and are synthesised by the hydrolysis of




7. If we mix Me_3SiCl with Me_2SiCl_2 , we get silicones of the type:



(C) both of the above

(D) none of the above

8. If we start with $MeSiCl_3$ as the starting material, silicones formed is:



(C) Both of the above

(D) None of the above

Paragraph for Question No. 9 & 10

CO gas is absorbed by aqueous suspension of cuprous chloride forming the complex like $[CuCl(CO)(H_2O)_2]$.

9. Comment on the shape of the above complex.

(A) Tetrahedral (B) TBP (C) Square planar (D) Cannot be predicted

10. Choose the correct statement regarding the above molecule


(A) Cl-atom is separated by equal angle from both of the water molecule

(B) Magnetic moment of the above complex is 1.73 B.M.

(C) There are two stereo isomer for the above complex.

(D) Both (A) and (C)

Paragraph for Question No. 11 to 12



11. Which of the following compound is not having +5 oxidation state on its central atom
 (A) B (B) D (C) C (D) E
12. "A" on reacting with Acetamide yields a compound "F". The compound "F" contains
 (A) 4 σ bonds, 2 π bonds
 (B) 5 σ bonds, 1 π bonds
 (C) 5 σ bonds, 2 π bonds
 (D) 2 σ bonds, 2 π bonds
13. Match List-I with List-II

List-I (Chemical reaction)

List-II (Name of process)

- (I) $4\text{NH}_3 + 5\text{O}_2 \xrightarrow{800^\circ\text{C}/\text{Pt}} 4\text{NO} + 6\text{H}_2\text{O}$ (a) Contact process
- (II) $4\text{HCl} + \text{O}_2 \xrightarrow{230^\circ\text{C}/\text{CuCl}_2} 2\text{Cl}_2 + 2\text{H}_2\text{O}$ (b) Ostwald's process
- (III) $2\text{SO}_2 + \text{O}_2 \xrightarrow{450-500^\circ/\text{V}_2\text{O}_5} 2\text{SO}_3$ (c) Deacon's process
- (IV) $2\text{N}_2 + 3\text{H}_2 \xrightarrow{\text{Fe}+\text{Mo}} 2\text{NH}_3$ (d) Haber's process
- (A) I-a, II-b, III-d, IV-c (B) I-b, II-c, III-a, IV-d
- (C) I-a, II-d, III-c, IV-b (D) I-a, II-c, III-b, IV-d

14. Column-I

Column-II

- (P) Dry ice (1) Used as antidote for CO-poisoning
- (Q) Carbongene (2) Used as nonstick coating
- (R) Carborundum (3) Used as refrigerant
- (S) Teflon (4) Used as abrasive

Code :

- | | | | | |
|-----|---|---|---|---|
| | P | Q | R | S |
| (A) | 4 | 1 | 3 | 2 |
| (C) | 3 | 1 | 4 | 2 |

- | | | | | |
|-----|---|---|---|---|
| | P | Q | R | S |
| (B) | 4 | 2 | 1 | 3 |
| (D) | 1 | 4 | 3 | 2 |

15. Column-I

Compound

- (P) SnCl_2
 (Q) Butter of tin
 (R) Mosaic gold
 (S) Pink's salt

Code :

	P	Q	R	S
(A)	4	3	2	1
(C)	2	1	3	4

Column-II

Correct statement for compounds given

- (1) Used in printing technology
 (2) Used for gilding purpose (in joining gold pieces)
 (3) Reducing agent
 (4) Mordant

	P	Q	R	S
(B)	3	4	2	1
(D)	1	3	4	2

16. Column-I (Metal)

- (P) Fe
 (Q) Cu
 (R) Pb
 (S) Sn

Code :

	P	Q	R	S
(A)	2, 1	1	3, 4	2, 3
(B)	2, 3	1, 3	1, 3	2, 3, 4
(C)	1, 3	1, 2	3, 4	2
(D)	1, 4	2, 3	1, 3	1, 4

Column-II (Correct statements)

- (1) Produces NO with 20% HNO_3
 (2) Produces NH_4NO_3 with 6% HNO_3
 (3) Produces NO_2 with 70% HNO_3
 (4) Produces NH_4NO_3 with 20% HNO_3

17. Column-I (Reactions)

- (P) $\text{XeF}_2 + \text{PF}_5 \rightarrow$
 (Q) $\text{XeF}_4 + \text{Pt} \rightarrow$
 (R) $\text{XeF}_4 + \text{H}_2\text{O} \rightarrow$
 (S) $\text{XeF}_6 + \text{CsF} \rightarrow$

Code :

	P	Q	R	S
(A)	4	2	3	1
(C)	4	3	2	1

Column-II (Correct statements)

- (1) Fluoride of Xe acts as fluoride acceptor
 (2) Fluoride of Xe undergoes disproportion
 (3) Fluoride of Xe acts as fluorinating agent
 (4) Fluoride of Xe act as fluoride donor

	P	Q	R	S
(B)	3	2	1	4
(D)	3	4	2	1

18. Column-I (Substances)

Column-II (Can be prepared by)

(P) O_3 (1) Acidification of BaO_2 with H_3PO_4

(Q) Bleaching powder

(2) Birkeland Eyde process

(R) H_2O_2 (3) Dry O_2 is passed through a silent electrical discharge(S) HNO_3 (4) Cl_2 gas is passed through slaked lime

Code :

	P	Q	R	S
(A)	3	4	1	2
(C)	2	1	3	4

	P	Q	R	S
(B)	1	3	4	2
(D)	4	1	2	3

MATCHING LIST TYPE 1 × 3 Q. (19 to 21)

For the following molecules in column-I, match the correct order of properties with column-II & column-III according to the questions asked.

Column - 1 Molecules	Column - 2 Properties	Column - 3 Correct order
(I) NH_3 , PH_3 , AsH_3 , SbH_3	(i) Bond angle	(p) Increasing order
(II) H_2O , H_2S , H_2Se , H_2Te	(ii) Reducing character	(q) Decreasing order
(III) HF , HCl , HBr , HI	(iii) Intermolecular force	(r) All equal
(IV) CH_4 , SiH_4 , GeH_4 , SnH_4	(iv) Boiling Point	(s) Irregular order

19. Which is the only **CORRECT** combination?

- (A) (I), (iv), (p) (B) (II), (i), (p) (C) (III), (iv), (q) (D) (IV), (iii), (p)

20. Which is the only **INCORRECT** combination?

- (A) (I), (ii), (p) (B) (IV), (i), (r) (C) (III), (iii), (q) (D) (II), (iv), (s)

21. In which combination, Drago's Rule plays an insignificant role in prediction of orders?

- (A) (I), (i), (q) (B) (II), (iv), (s) (C) (IV), (i), (r) (D) (III), (ii), (p)

MATCHING LIST TYPE 1 × 3 Q. (22 to 24)

Molecule	Hybridization of central atom(s)	Properties related to structure
(P) (HBNH) ₃	(I) sp ²	(i) Cyclic structure
(Q) C ₃ O ₂	(II) sp ³	(ii) Has (X- $\ddot{\text{O}}$ -X) linkage (X = central atom)
(R) P ₄ O ₁₂ ⁴⁻	(III) sp ³ d	(iii) Planar
(S) N ₂ O ₄	(IV)sp	(iv) Has (X – X) linkage

22. Identify **CORRECT** match

- (A) P, II, iii (B) P, I, i (C) P, IV, iii (D) Q, I, i

23. Identify **CORRECT** match

- (A) Q, IV, iii (B) Q, IV, i (C) Q, I, ii (D) R, I, ii

24. Identify **CORRECT** match

- (A) R, II, iv (B) R, II, ii (C) S, III, ii (D) S, IV, i

EXERCISE # JEE-MAINS

1. Which products are expected from the disproportionation of hypochlorous acid : [AIEEE-2002]
(1) HClO_3 and Cl_2O (2) HClO_2 and HClO (3) HCl and Cl_2O (4) HCl and HClO_3
2. Identify the **INCORRECT** statement among the following : [AIEEE-2002]
(1) Ozone reacts with SO_2 to give SO_3
(2) Silicon reacts with NaOH(aq.) in the presence of air to give Na_2SiO_3 and H_2O
(3) Cl_2 reacts with excess of NH_3 to give N_2 and HCl
(4) Br_2 reacts with hot and strong NaOH solution to give NaBr , NaBrO_4 and H_2O
3. Aluminium is industrially prepared by: [AIEEE-2002]
(1) Fused cryolite (2) Bauxite ore (3) Alunite (4) Borax
4. For making good quality mirrors, plates of float glass are used. These are obtained by floating molten glass over a liquid metal which does not solidify before glass. The metal used can be : [AIEEE-2003]
(1) Sodium (2) Magnesium (3) Mercury (4) Tin
5. What may be expected when phosphine gas is mixed with chlorine gas: [AIEEE-2003]
(1) PCl_5 and HCl are formed and mixture cools down
(2) $\text{PH}_3\cdot\text{Cl}_2$ is formed with warming up
(3) The mixture only cools down
(4) PCl_3 and HCl are formed and the mixture warms up
6. Graphite is a soft solid lubricant extremely difficult to melt. The reason for this anomalous behaviour is that graphite : [AIEEE-2003]
(1) Has molecules of variable molecular masses like polymers
(2) Has carbon atoms arranged in large plates of rings of strongly bonded carbon atoms with weak interplate bonds
(3) Is a non crystalline substance
(4) Is an allotropic form of diamond
7. Concentrated hydrochloric acid when kept in open air sometimes produces a cloud of white fumes. This is due to : [AIEEE-2003]
(1) Strong affinity of HCl gas for moisture in air results in forming of droplets of liquid solution which appears like a cloudy smoke
(2) Due to strong affinity for water, conc. HCl pulls moisture of air towards self. The moisture forms droplets of water and hence the cloud
(3) Conc. HCl emits strongly smelling HCl gas all the time
(4) Oxygen in air reacts with emitted HCl gas to form a cloud of Cl_2 gas

8. Aluminium chloride exists as dimer, Al_2Cl_6 in solid state as well as in solution of non-polar solvents such as benzene. When dissolved in water, it gives- [AIEEE-2004]
- (1) $\text{Al}^{3+} + 3\text{Cl}^-$ (2) $[\text{Al}(\text{H}_2\text{O})_6]^{3+} + 3\text{Cl}^-$
 (3) $[\text{Al}(\text{OH})_6]^{3-} + 3\text{HCl}$ (4) $\text{Al}_2\text{O}_3 + 6\text{HCl}$
9. The soldiers of Napoleon army while at Alps during freezing winter suffered a serious problem as regards to the tin buttons of their uniforms. White Metallic tin buttons get converted to grey powder. This transformation is related to:- [AIEEE-2004]
- (1) An interaction with water vapour contained in humid air
 (2) A change in crystalline structure of tin
 (3) A change in the partial pressure of O_2 in air
 (4) An interaction with N_2 of air at low temperature
10. Which one of the following statements regarding helium is incorrect [AIEEE-2004]
- (1) It is used to produce and sustain powerful superconducting magnets
 (2) It is used as a cryogenic agent for carrying out experiments at low temperatures
 (3) It is used to fill gas balloons instead of hydrogen because it is lighter than hydrogen and non-inflammable
 (4) It is used in gas-cooled nuclear reactors
11. The number of hydrogen atoms attached to phosphorus atom in hypophosphorous acid is : [AIEEE-2005]
- (1) Zero (2) Two (3) One (4) Three
12. Heating an aqueous solution of aluminium chloride to dryness will give :- [AIEEE-2005]
- (1) AlCl_3 (2) Al_2Cl_6 (3) Al_2O_3 (4) $\text{Al}(\text{OH})\text{Cl}_2$
13. Which one of the following is the correct statement [AIEEE-2005]
- (1) Boric acid is a protonic acid
 (2) Beryllium exhibits coordination number of six
 (3) Chlorides of both beryllium and aluminium have bridged chloride structures in solid phase
 (4) $\text{B}_2\text{H}_6 \cdot 2\text{NH}_3$ is known as "inorganic benzene"
14. In silicon dioxide : [AIEEE-2005]
- (1) Each silicon atom is surrounded by four oxygen atoms and each oxygen atom is bonded to two silicon atoms
 (2) Each silicon atom is surrounded by two oxygen atoms and each oxygen atom is bonded to two silicon atoms
 (3) Silicon atom is bonded to two oxygen atoms
 (4) There are double bonds between silicon and oxygen atoms

- 15.** Regular use of which of the following fertilizer increases the acidity of soil : [AIEEE-2007]
 (1) Potassium nitrate (2) Urea
 (3) Superphosphate of lime (4) Ammonium sulphate
- 16.** The stability of dihalides of Si, Ge, Sn and Pb increases steadily in the sequence: [AIEEE-2007]
 (1) $\text{GeX}_2 \ll \text{SiX}_2 \ll \text{SnX}_2 \ll \text{PbX}_2$ (2) $\text{SiX}_2 \ll \text{GeX}_2 \ll \text{PbX}_2 \ll \text{SnX}_2$
 (3) $\text{SiX}_2 \ll \text{GeX}_2 \ll \text{SnX}_2 \ll \text{PbX}_2$ (4) $\text{PbX}_2 \ll \text{SnX}_2 \ll \text{GeX}_2 \ll \text{SiX}_2$
- 17.** Among the following substituted silanes the one which will give rise to cross linked silicone polymer on hydrolysis is [AIEEE-2008]
 (1) R_4Si (2) RSiCl_3 (3) R_2SiCl_2 (4) R_3SiCl
- 18.** Which one of the following reactions of Xenon compounds is not feasible ? [AIEEE-2009]
 (1) $2\text{XeF}_2 + 2\text{H}_2\text{O} \rightarrow 2\text{Xe} + 4\text{HF} + \text{O}_2$ (2) $\text{XeF}_6 + \text{RbF} \rightarrow \text{Rb}[\text{XeF}_7]$
 (3) $\text{XeO}_3 + 6\text{HF} \rightarrow \text{XeF}_6 + 3\text{H}_2\text{O}$ (4) $3\text{XeF}_4 + 6\text{H}_2\text{O} \rightarrow 2\text{Xe} + \text{XeO}_3 + 12\text{HF} + 1.5\text{O}_2$
- 19.** Which of the following statement is wrong ? [AIEEE-2011]
 (1) Single N–N bond is weaker than the single P–P bond
 (2) N_2O_4 has two resonance structures
 (3) The stability of hydrides increases from NH_3 to BiH_3 in group 15 of the periodic table
 (4) Nitrogen cannot form $d\pi\text{-}p\pi$ bond
- 20.** Which of the following statements regarding sulphur is incorrect ? [AIEEE-2011]
 (1) At 600°C the gas mainly consists of S_2 molecules
 (2) The oxidation state of sulphur is never less than +4 in its compounds
 (3) S_2 molecule is paramagnetic
 (4) The vapour at 200°C consists mostly of S_8 rings
- 21.** Boron cannot form which one of the following anions ? [AIEEE-2011]
 (1) $\text{B}(\text{OH})_4^-$ (2) BO_2^- (3) BF_6^{3-} (4) BH_4^-
- 22.** In view of the signs of $\Delta_r G^\circ$ for the following reactions
 $\text{PbO}_2 + \text{Pb} \rightarrow 2 \text{PbO}, \Delta_r G^\circ < 0$
 $\text{SnO}_2 + \text{Sn} \rightarrow 2 \text{SnO}, \Delta_r G^\circ > 0,$
 Which oxidation states are more characteristic for lead and tin ? [AIEEE-2011]
 (1) For lead + 4, for tin + 2 (2) For lead + 2, for tin + 2
 (3) For lead + 4, for tin + 4 (4) For lead + 2, for tin + 4
- 23.** The number of S–S bonds in SO_3 , $\text{S}_2\text{O}_3^{2-}$, $\text{S}_2\text{O}_6^{2-}$ and $\text{S}_2\text{O}_8^{2-}$ respectively are :- [Jee Main(Online)-2012]
 (1) 1, 0, 1, 0 (2) 0, 1, 1, 0 (3) 1, 0, 0, 1 (4) 0, 1, 0, 1
- 24.** Which one of the following depletes ozone layer ? [Jee Main(Online)-2012]
 (1) NO and freons (2) SO_2 (3) CO (4) CO_2

25. In which of the following arrangements, the sequence is not strictly according to the property written against it ? [Jee Main(Online)-2012]
- (1) $\text{CO}_2 < \text{SiO}_2 < \text{SnO}_2 < \text{PbO}_2$: increasing oxidising power
 (2) $\text{B} < \text{C} < \text{O} < \text{N}$: increasing first ionisation enthalpy
 (3) $\text{NH}_3 < \text{PH}_3 < \text{AsH}_3 < \text{SbH}_3$: increasing basic strength
 (4) $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$: increasing acid strength
26. The formation of molecular complex $\text{BF}_3 - \text{NH}_3$ results in a change in hybridisation of boron :-
 (1) from sp^3 to sp^3d (2) from sp^2 to dsp^2 [JEE(Main) Online-2012]
 (3) from sp^3 to sp^2 (4) from sp^2 to sp^3
27. The catenation tendency of C, Si and Ge is in the order $\text{Ge} < \text{Si} < \text{C}$. The bond energies (in kJ mol^{-1}) of C—C, Si—Si and Ge—Ge bonds are respectively : [JEE(Main) Online-2013]
- (1) 348, 260, 297 (2) 348, 297, 260 (3) 297, 348, 260 (4) 260, 297, 348
28. The gas evolved on heating CaF_2 and SiO_2 with concentrated H_2SO_4 , on hydrolysis gives a white gelatinous precipitate. The precipitate is: [Jee Main(Online)-2014]
- (1) silica gel (2) silicic acid
 (3) hydrofluosilicic acid (4) calciumfluorosilicate
29. Which of the following series correctly represents relations between the elements from X to Y? [Jee Main(Online)-2014]
- $\text{X} \rightarrow \text{Y}$
- (1) ${}_{18}\text{Ar} \rightarrow {}_{54}\text{Xe}$ Noble character increases
 (2) ${}_3\text{Li} \rightarrow {}_{19}\text{K}$ Ionization enthalpy increases
 (3) ${}_6\text{C} \rightarrow {}_{32}\text{Ge}$ Atomic radii increases
 (4) ${}_9\text{F} \rightarrow {}_{35}\text{Br}$ Electron gain enthalpy with negative sign increases
30. Which of the following statements about the depletion of ozone layer is correct? [Jee Main(Online)-2014]
- (1) The problem of ozone depletion is more serious at poles because ice crystals in the clouds over poles act as catalyst for photochemical reactions involving the decomposition of ozone by $\text{Cl}^\bullet$ and $\text{ClO}^\bullet$ radicals
 (2) The problem of ozone depletion is less serious at poles because NO_2 solidifies and is not available for consuming $\text{ClO}^\bullet$ radicals
 (3) Oxides of nitrogen also do not react with ozone in stratosphere
 (4) Freons, chlorofluorocarbons, are inert chemically, they do not react with ozone in stratosphere
31. Which of the following xenon-OXO compounds may not be obtained by hydrolysis of xenon fluorides ? [Jee Main(Online)-2014]
- (1) XeO_2F_2 (2) XeO_3 (3) XeO_4 (4) XeOF_4

32. Hydrogen peroxide acts both as an oxidising and as a reducing agent depending upon the nature of the reacting species. In which of the following cases H_2O_2 acts as a reducing agent in acid medium ? **[Jee Main(Online)-2014]**
- (1) MnO_4^- (2) SO_3^{2-} (3) KI (4) $\text{Cr}_2\text{O}_7^{2-}$
33. Consider the reaction **[Jee Main(Online)-2014]**
- $$\text{H}_2\text{SO}_{3(\text{aq})} + \text{Sn}_{(\text{aq})}^{4+} + \text{H}_2\text{O}_{(\text{l})} \rightarrow \text{Sn}_{(\text{aq})}^{2+} + \text{HSO}_{4(\text{aq})}^- + 3\text{H}_{(\text{aq})}^+$$
- Which of the following statements is correct?
- (1) H_2SO_3 is the reducing agent because it undergoes oxidation
 (2) H_2SO_3 is the reducing agent because it undergoes reduction
 (3) Sn^{4+} is the reducing agent because it undergoes oxidation
 (4) Sn^{4+} is the oxidizing agent because it undergoes oxidation
34. In the following sets of reactants which two sets best exhibit the amphoteric character of $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}$? **[JEE(Main) Online-2014]**
- Set-1 :** $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}(\text{s})$ and $\text{OH}^- (\text{aq})$
Set-2 : $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}(\text{s})$ and $\text{H}_2\text{O} (\ell)$
Set-3 : $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}(\text{s})$ and $\text{H}^+ (\text{aq})$
Set-4 : $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}(\text{s})$ and $\text{NH}_3 (\text{aq})$
- (1) 1 and 2 (2) 2 and 4 (3) 1 and 3 (4) 3 and 4
35. Which of the following compounds has a P-P bond :- **[Jee Main(Online)-2015]**
- (1) $\text{H}_4\text{P}_2\text{O}_5$ (2) $(\text{HPO}_3)_3$ (3) $\text{H}_4\text{P}_2\text{O}_7$ (4) $\text{H}_4\text{P}_2\text{O}_6$
36. Chlorine water on standing loses its colour and forms :- **[Jee Main(Online)-2015]**
- (1) HCl and HClO_2 (2) HCl only
 (3) HOCl and HOCl_2 (4) HCl and HOCl
37. Which among the following is the most reactive ? **[Jee Main-2015]**
- (1) I_2 (2) ICl (3) Cl_2 (4) Br_2
38. Which one has the highest boiling point ? **[Jee Main-2015]**
- (1) Kr (2) Xe (3) He (4) Ne
39. From the following statements regarding H_2O_2 , choose the incorrect statement : **[Jee Main-2015]**
- (1) It has to be stored in plastic or wax lined glass bottles in dark
 (2) It has to be kept away from dust
 (3) It can act only as an oxidizing agent
 (4) It decomposes on exposure to light

40. The reaction of zinc with dilute and concentrated nitric acid, respectively produces :
[JEE (Main) 2016]
(1) NO_2 and N_2O (2) N_2O and NO_2 (3) NO_2 and NO (4) NO and N_2O
41. The non-metal that does not exhibit positive oxidation state is :
[JEE (Main) 2016]
(1) Oxygen (2) Fluorine (3) Iodine (4) Chlorine
42. Which intermolecular force is most responsible in allowing xenon gas to liquefy?
[JEE (Main) Online 2016]
(1) Ionic (2) Instantaneous dipole- induced dipole
(3) Dipole - dipole (4) Ion - dipole
43. The following statements concern elements in the periodic table. Which of the following is true ?
[JEE (Main) Online 2016]
(1) The group 13 elements are all metals.
(2) For group 15 elements, the stability of +5 oxidation state increases down the group.
(3) All the elements in Group 17 are gases.
(4) Elements of group 16 have lower ionization enthalpy values compared to those of group 15 in the corresponding periods.
44. **Assertion :** Among the carbon allotropes, diamond is an insulator, whereas, graphite is a good conductor of electricity.
[JEE (Main) Online 2016]
Reason : Hybridization of carbon in diamond and graphite are sp^3 and sp^2 , respectively.
(1) Assertion is incorrect statement, but the reason is correct.
(2) Both assertion and reason are correct, and the reason is the correct explanation for the assertion.
(3) Both assertion and reason are incorrect.
(4) Both assertion and reason are correct, but the reason is not the correct explanation for the assertion.
45. Identify the incorrect statement :
[JEE (Main) Online 2016]
(1) S_8 ring has a crown shape.
(2) The S-S-S bond angles in the S_8 and S_6 rings are the same
(3) S_2 is paramagnetic like oxygen
(4) Rhombic and monoclinic sulphur have S_8 molecules.
46. Which of the following reactions is an example of a redox reaction ?
[JEE (Main) 2017]
(1) $\text{XeF}_4 + \text{O}_2\text{F}_2 \rightarrow \text{XeF}_6 + \text{O}_2$
(2) $\text{XeF}_2 + \text{PF}_5 \rightarrow [\text{XeF}]^+\text{PF}_6^-$
(3) $\text{XeF}_6 + \text{H}_2\text{O} \rightarrow \text{XeOF}_4 + 2\text{HF}$
(4) $\text{XeF}_6 + 2\text{H}_2\text{O} \rightarrow \text{XeO}_2\text{F}_2 + 4\text{HF}$
47. The products obtained when chlorine gas reacts with cold and dilute aqueous NaOH are :-
[JEE (Main) 2017]
(1) ClO^- and ClO_3^- (2) ClO_2^- and ClO_3^-
(3) Cl^- and ClO^- (4) Cl^- and ClO_2^-

48. In which of the following reaction, hydrogen peroxide acts as an oxidizing agent ?
 (1) $I_2 + H_2O_2 + 2OH^- \rightarrow 2I^- + 2H_2O + O_2$ [JEE (Main) ONLINE 2017]
 (2) $HOCl + H_2O_2 \rightarrow H_3O^+ + Cl^- + O_2$
 (3) $PbS + 4H_2O_2 \rightarrow PbSO_4 + 4H_2O$
 (4) $2MnO_4^- + 3H_2O_2 \rightarrow 2MnO_2 + 3O_2 + 2H_2O + 2OH^-$
49. XeF_6 on partial hydrolysis with water produces a compound 'X'. The same compound 'X' is formed when XeF_6 reacts with silica. The compound 'X' is:- [JEE (Main) ONLINE 2017]
 (1) XeO_3 (2) XeF_4 (3) $XeOF_4$ (4) XeF_2
50. The compound that does not produce nitrogen gas by the thermal decomposition is [JEE (Main) OFFLINE 2018]
 (1) $(NH_4)_2Cr_2O_7$ (2) NH_4NO_2 (3) $(NH_4)_2SO_4$ (4) $Ba(N_3)_2$
51. For per gram of reactant, the maximum quantity of N_2 gas is produced in which of the following thermal decomposition reactions ? [JEE (Main) ONLINE 2018]
(Given : Atomic wt. – Cr = 52u, Ba = 137u)
 (1) $2NH_4NO_3(s) \rightarrow 2N_2(g) + 4H_2O(g) + O_2(g)$
 (2) $Ba(N_3)_2(s) \rightarrow Ba(s) + 3N_2(g)$
 (3) $(NH_4)_2Cr_2O_7(s) \rightarrow N_2(g) + 4H_2O(g)$
 (4) $2NH_3(g) \rightarrow N_2(g) + 3H_2(g)$
52. Lithium aluminium hydride reacts with silicon tetrachloride to form :- [JEE (Main) ONLINE 2018]
 (1) $LiCl, AlCl_3$ and SiH_4 (2) $LiCl, AlH_3$ and SiH_4
 (3) $LiH, AlCl_3$ and $SiCl_2$ (4) LiH, AlH_3 and SiH_4
53. Xenon hexafluoride on partial hydrolysis produces compounds 'X' and 'Y' Compounds 'X' and 'Y' and the oxidation state of Xe are respectively : [JEE (Main) ONLINE 2018]
 (1) $XeO_2F_2(+6)$ and $XeO_2(+4)$ (2) $XeOF_4(+6)$ and $XeO_2F_2(+6)$
 (3) $XeOF_4(+6)$ and $XeO_3(+6)$ (4) $XeO_2(+4)$ and $XeO_3(+6)$
54. Which of the following is a lewis acid ? [JEE (Main) ONLINE 2018]
 (1) NaH (2) NF_3 (3) PH_3 (4) $B(CH_3)_3$
55. Iodine reacts with concentrated HNO_3 to yield Y along with other products. The oxidation state of iodine in Y, is :- [JEE (Main) ONLINE 2019]
 (1) 5 (2) 3 (3) 1 (4) 7
56. Among the following reactions of hydrogen with halogens, the one that requires a catalyst is :
 (1) $H_2 + I_2 \rightarrow 2HI$ (2) $H_2 + F_2 \rightarrow 2HF$ [JEE (Main) ONLINE 2019]
 (3) $H_2 + Cl_2 \rightarrow 2HCl$ (4) $H_2 + Br_2 \rightarrow 2HBr$

57. Correct statements among a to d regarding silicones are : [JEE (Main) ONLINE 2019]
 (a) They are polymers with hydrophobic character
 (b) They are biocompatible.
 (c) In general, they have high thermal stability and low dielectric strength.
 (d) Usually, they are resistant to oxidation and used as greases.
 (1) (a), (b) and (c) only
 (2) (a), and (b) only
 (3) (a), (b), (c) and (d)
 (4) (a), (b) and (d) only
58. Diborane (B_2H_6) reacts independently with O_2 and H_2O to produce, respectively
 (1) HBO_2 and H_3BO_3 (2) H_3BO_3 and B_2O_3 [JEE (Main) ONLINE 2019]
 (3) B_2O_3 and H_3BO_3 (4) B_2O_3 and $[BH_4]^-$
59. The amorphous form of silica is : [JEE (Main) ONLINE 2019]
 (1) quartz (2) kieselguhr (3) cristobalite (4) tridymite
60. The correct statements among I to III regarding group 13 element oxides are, [JEE (Main) ONLINE 2019]
 (I) Boron trioxide is acidic.
 (II) Oxides of aluminium and gallium are amphoteric.
 (III) Oxides of indium and thallium are basic.
 (1) (I), (II) and (III) (2) (II) and (III) only (3) (I) and (III) only (4) (I) and (II) only
61. The redox reaction among the following is : [JEE (Main) ONLINE 2020]
 (1) Combination of dinitrogen with dioxygen at 2000 K
 (2) Formation of ozone from atmospheric oxygen in the presence of sunlight
 (3) Reaction of H_2SO_4 with NaOH
 (4) Reaction of $[Co(H_2O)_6]Cl_3$ with $AgNO_3$
62. In the following reactions products (A) and (B), respectively, are : [JEE (Main) ONLINE 2020]
 $NaOH + Cl_2 \rightarrow (A) + \text{side products}$
 (hot and conc.)
 $Ca(OH)_2 + Cl_2 \rightarrow (B) + \text{side products}$
 (dry)
 (1) $NaClO_3$ and $Ca(OCl)_2$ (2) $NaOCl$ and $Ca(ClO_3)_2$
 (3) $NaClO_3$ and $Ca(ClO_3)_2$ (4) $NaOCl$ and $Ca(OCl)_2$
63. Chlorine reacts with hot and concentrated NaOH and produces compounds (X) and (Y). Compound (X) gives white precipitate with silver nitrate solution. The average bond order between Cl and O atoms in (Y) is _____. [JEE (Main) ONLINE 2020]
64. White Phosphorus on reaction with concentrated NaOH solution in an inert atmosphere of CO_2 gives phosphine and compound (X). (X) on acidification with HCl gives compound (Y). The basicity of compound (Y) is : [JEE (Main) ONLINE 2020]
 (1) 4 (2) 1 (3) 2 (4) 3

65. When gypsum is heated to 393 K, it forms : [JEE (Main) ONLINE 2020]
(1) Dead burnt plaster
(2) Anhydrous CaSO_4
(3) $\text{CaSO}_4 \cdot 5\text{H}_2\text{O}$
(4) $\text{CaSO}_4 \cdot 0.5\text{H}_2\text{O}$
66. The reaction of $\text{H}_3\text{N}_3\text{B}_3\text{Cl}_3$ (A) with LiBH_4 in tetrahydrofuran gives inorganic benzene (B). Further, the reaction of (A) with (C) leads to $\text{H}_3\text{N}_3\text{B}_3(\text{Me})_3$. Compounds (B) and (C) respectively, are: [JEE (Main) ONLINE 2020]
(1) Boron nitride and MeBr
(2) Borazine and MeMgBr
(3) Borazine and MeBr
(4) Diborane and MeMgBr
67. 'X' melts at low temperature and is a bad conductor of electricity in both liquid and solid state. X is : [JEE (Main) ONLINE 2020]
(1) Carbon tetrachloride (2) Mercury
(3) Silicon carbide (4) Zinc sulphide
68. The compound that cannot act both as oxidising and reducing agent is : [JEE (Main) ONLINE 2020]
(1) H_2O_2 (2) H_2SO_3
(3) HNO_2 (4) H_3PO_4

EXERCISE # J-ADVANCED

(IIT JEE ASKED QUESTIONS)

Fill in the blanks

- Give reason : [IIT- 2000]
Why elemental nitrogen exists as a diatomic molecule whereas elemental phosphorus is a tetra atomic molecule.
- Give an example of oxidation of one halide by another halogen. Explain the feasibility of the reaction. [IIT- 2000]
- Compounds X on reduction with LiAlH_4 gives a hydride Y containing 21.72% hydrogen alongwith other products. The compound Y reacts with air explosively resulting in boron trioxide. Identify X and Y. Give balanced reactions involved in the formation of Y and its reaction with air Draw the structure of Y. [IIT- 2001]
- Starting from SiCl_4 , prepare the following in steps not exceeding the number given in parenthesis (reactions only) [IIT- 2001]
 - Silicon (1)
 - Linear silicon containing methyl group only (4)
 - Na_2SiO_3 (3)
- Write the balanced chemical equation for developing photographic films. [IIT- 2001]
- Identify (X) in the following synthetic scheme and write their structures. [IIT- 2001]
 $\text{BaCO}_3^* + \text{H}_2\text{SO}_4 \longrightarrow \text{X (gas)}$ (C denotes C^{14})
- Write the balanced equations for the reactions of the following compounds with water [2002]
 - Al_4Cl_3
 - CaNCN
 - BF_3
 - NCl_3
 - XeF_3
- Write the balanced equations for the reactions of the following compounds with water: [IIT- 2002]
 - Al_4C_3
 - CaNCN
 - BF_3
 - NCl_3
 - XeF_4
- Identify the following:

$$\text{Na}_2\text{CO}_3 \xrightarrow[\text{(aq)}]{\text{SO}_2} \text{A} \xrightarrow{\text{Na}_2\text{CO}_3} \text{B} \xrightarrow[\Delta]{\text{elemental S}} \text{C} \xrightarrow{\text{I}_2} \text{D}$$
 [IIT- 2003]
 Also mention the oxidation state of S in all the compounds.
- Arrange the following oxides in the increasing order of Bronsted basicity. [IIT- 2004]
 Cl_2O_7 , BaO , SO_3 , CO_2 , B_2O_3
- The number of P—O—P bonds in cyclic tetrametaphosphoric acid is – [IIT-2000]
 - Zero
 - Two
 - Three
 - Four
- The correct order of acidic strength is – [IIT- 2000]
 - $\text{Cl}_2\text{O}_7 > \text{SO}_2 > \text{P}_4\text{O}_{10}$
 - $\text{CO}_2 > \text{N}_2\text{O}_5 > \text{SO}_3$
 - $\text{Na}_2\text{O} > \text{MgO} > \text{Al}_2\text{O}_3$
 - $\text{K}_2\text{O} > \text{CaO} > \text{MgO}$

13. Amongst H_2O , H_2S , H_2Se and H_2Te , the one with the highest boiling point is – [IIT- 2000]
 (A) H_2O because of hydrogen bonding (B) H_2Te because of higher molecular weight
 (C) H_2S because of hydrogen bonding (D) H_2Se because of lower molecular weight.
14. Ammonia can be dried by – [IIT- 2000]
 (A) Conc. H_2SO_4 (B) P_4O_{10} (C) CaO (D) Anhydrous CaCl_2
15. Which of the following are hydrolysed – [REE 2000]
 (A) NCl_3 (B) BCl_3 (C) CCl_4 (D) SiCl_4
16. The set with correct order of acidity is – [IIT- 2001]
 (A) $\text{HClO} < \text{HClO}_2 < \text{HClO}_3 < \text{HClO}_4$ (B) $\text{HClO}_4 < \text{HClO}_3 < \text{HClO}_2 < \text{HClO}$
 (C) $\text{HClO} < \text{HClO}_4 < \text{HClO}_3 < \text{HClO}_2$ (D) $\text{HClO}_4 < \text{HClO}_2 < \text{HClO}_3 < \text{HClO}$
17. The reaction, $3\text{ClO}^- (\text{aq}) \longrightarrow \text{ClO}_3^- (\text{aq}) + 2\text{Cl}^- (\text{aq})$ is an example of – [IIT- 2001]
 (A) Oxidation reaction (B) reduction reaction
 (C) Disproportionation reaction (D) Decomposition reaction
18. The number of S–S bonds in sulphur trioxide trimer, (S_3O_9) is – [IIT- 2001]
 (A) Three (B) Two (C) One (D) Zero
19. **Statement-I :** Between SiCl_4 and CCl_4 , only SiCl_4 reacts with water [IIT- 2001]
Because :
Statement-II : SiCl_4 is ionic and CCl_4 is covalent
 (A) If both assertion and reason are correct and reason is the correct explanation of the assertion
 (B) If both assertion and reason are correct, but reason is not the correct explanation of the assertion
 (C) If assertion is correct, but reason is incorrect
 (D) If assertion is incorrect, but reason is correct.
20. Polyphosphates are used as water softening agents because they – [IIT- 2002]
 (A) Form soluble complexes with anionic species
 (B) Precipitate anionic species
 (C) Form soluble complexes with cationic species
 (D) Precipitate cationic species
21. Identify the correct order of solubility of Na_2S , CuS , and ZnS in aqueous medium – [IIT- 2002]
 (A) $\text{CuS} > \text{ZnS} > \text{Na}_2\text{S}$ (B) $\text{ZnS} > \text{Na}_2\text{S} > \text{CuS}$
 (C) $\text{Na}_2\text{S} > \text{CuS} > \text{ZnS}$ (D) $\text{Na}_2\text{S} > \text{ZnS} > \text{CuS}$
22. Identify, the correct order of acidic strength of CO_2 , CuO , CaO , H_2O – [IIT- 2002]
 (A) $\text{CaO} < \text{CuO} < \text{H}_2\text{O} < \text{CO}_2$ (B) $\text{H}_2\text{O} < \text{CuO} < \text{CaO} < \text{CO}_2$
 (C) $\text{CaO} < \text{H}_2\text{O} < \text{CuO} < \text{CO}_2$ (D) $\text{H}_2\text{O} < \text{CO}_2 < \text{CaO} < \text{CuO}$

23. H_3BO_3 is – [IIT- 2002, 3]
 (A) Monobasic acid and weak Lewis acid (B) Monobasic and weak Bronsted acid
 (C) Monobasic and strong Lewis acid (D) Tribasic and weak Bronsted acid
24. When Γ^- is oxidised by MnO_4^- in alkaline medium, Γ^- converts into – [IIT- 2003]
 (A) IO_3^- (B) I_2 (C) IO_4^- (D) IO^-
25. **Column-I (Change)** **Column-II (Given change is done by)**
 (A) $\text{Bi}^{3+} \longrightarrow (\text{BiO})^+$ (p) Heat [IIT- 2003]
 (B) $[\text{AlO}_2]^- \longrightarrow \text{Al}(\text{OH})_3$ (q) Hydrolysis
 (C) $\text{SiO}_4^{4-} \longrightarrow \text{Si}_2\text{O}_7^{6-}$ (r) Acidification
 (D) $(\text{B}_4\text{O}_7^{2-}) \longrightarrow [\text{B}(\text{OH})_3]$ (s) Dilution by water
26. $(\text{Me})_2\text{SiCl}_2$ on hydrolysis will produce – [IIT- 2003]
 (A) $(\text{Me})_2\text{Si}(\text{OH})_2$ (B) $(\text{Me})_2\text{Si}=\text{O}$
 (C) $[\text{—}(\text{Me})_2\text{Si—O—}]_n$ (D) $\text{Me}_2\text{SiCl}(\text{OH})$
27. Which is the most thermodynamically stable allotropic form of phosphorus? [IIT- 2004]
 (A) Red (B) White (C) Black (D) Yellow
28. When PbO_2 reacts with conc. HNO_3 the gas evolved may be : [2005]
 (A) NO_2 (B) O_2 (C) N_2 (D) N_2O
29. Which of the following is not oxidised by O_3 ? [IIT- 2005]
 (A) KI (B) FeSO_4 (C) KMnO_4 (D) K_2MnO_4
30. Which blue-liquid is obtained on reacting equimolar amounts of two gases at -30°C ? [IIT- 2005]
 (A) N_2O (B) N_2O_3 (C) N_2O_4 (D) N_2O_5
31. $\text{B}(\text{OH})_3 + \text{NaOH} \rightleftharpoons \text{NaBO}_2 + \text{Na}[\text{B}(\text{OH})_4] + \text{H}_2\text{O}$ how can this reaction is made to proceed in forward direction? [IIT- 2006]
 (A) Addition of cis 1, 2 diol (B) Addition of borax
 (C) Addition of trans 1, 2 diol (D) Addition of Na_2HPO_4
32. Among the following, the paramagnetic compound is – [IIT- 2007]
 (A) Na_2O_2 (B) O_3 (C) N_2O (D) KO_2
33. **Statement-I** : Boron always forms covalent bond [2007]
Because :

Statement-II : The small size of B^{3+} favours formation of covalent bond.

- (A) Statement-I is True, Statement-II is True, Statement-II is a correct explanation for Statement-I
 (B) Statement-I is True, Statement-II is True, Statement-II is not a correct explanation for Statement-II
 (C) Statement-I is True, Statement-II is False
 (D) Statement-I is False, Statement-II is True

34. **Statement-I** : In water, orthoboric acid behaves as a weak monobasic acid. [2007]
Statement-II : In water, orthoboric acid acts as a proton donor.
 (A) Statement-I is True, Statement-II is True, Statement-II is a correct explanation for Statement-I
 (B) Statement-I is True, Statement-II is True, Statement-II is not a correct explanation for Statement-II
 (C) Statement-I is True, Statement-II is False (D) Statement-I is False, Statement-II is True

Comprehension # 1 (Q. 35 to 37)

The noble gases have closed-shell electronic configuration and are monoatomic gases under normal conditions. The low boiling point of the lighter noble gases are due to weak dispersion forces between the atoms and the absence of other interatomic interactions. The direct reaction of xenon with fluorine leads to a series of compounds with oxidation number + 2, + 4 and + 6. XeF_4 reacts violently with water to give XeO_3 . The compounds of xenon exhibit rich stereochemistry and their geometries can be deduced considering the total number of electron pairs in the valence shell. [IIT- 2007]

35. Argon is used in arc welding because of its –
 (A) Low reactivity with metal (B) Ability to lower the melting point of metal
 (C) Flammability (D) High calorific value
36. The structure of XeO_3 is –
 (A) Linear (B) Planar (C) Pyramidal (D) T-shaped
37. XeF_4 and XeF_6 are expected to be –
 (A) Oxidising agent (B) Reducing agent (C) Unreactive (D) Strongly basic

Comprehension # 2 (Q.38 to 40)

There are some deposits of nitrates and phosphates in earth's crust. Nitrates are more soluble in water. Nitrates are difficult to reduce under the laboratory conditions but microbes do it easily. Ammonia forms large number of complexes with transition metal ions. Hybridization easily explains the ease of sigma donation capability of NH_3 and PH_3 . Phosphine is a flammable gas and is prepared from white phosphorous. [IIT- 2008]

38. Among the following, the correct statement is :-
 (A) Phosphates have no biological significance in humans
 (B) Between nitrates and phosphates, phosphates are less abundant in earth's crust
 (C) Between nitrates and phosphates, nitrates are less abundant in earth's crust
 (D) Oxidation of nitrates is possible in soil
39. Among the following, the correct statement is :-
 (A) Between NH_3 and PH_3 , NH_3 is a better electron donor because the lone pair of electrons occupies spherical 's' orbital and is less directional
 (B) Between NH_3 and PH_3 , PH_3 is a better electron donor because the lone pair of electrons occupies sp^3 orbital and is more directional
 (C) Between NH_3 and PH_3 , NH_3 is a better electron donor because the lone pair of electrons occupies sp^3 orbital and is more directional
 (D) Between NH_3 and PH_3 , PH_3 is a better electron donor because the lone pair of electrons occupies spherical 's' orbital and is less directional.

40. White phosphorus on reaction with NaOH gives PH_3 as one of the products. This is a :-
 (A) dimerization reaction (B) disproportionation reaction
 (C) condensation reaction (D) precipitation reaction
41. The reaction of P_4 with X leads selectively to P_4O_6 . The X is [JEE 2009]
 (A) Dry O_2 (B) A mixture of O_2 and N_2
 (C) Moist O_2 (D) O_2 in the presence of aqueous NaOH
42. The reaction of white phosphorus with aqueous NaOH gives phosphine along with another phosphorus containing compound. The reaction type ; the oxidation states of phosphorus in phosphine and the other product are respectively [JEE 2012]
 (A) redox reaction ; -3 and -5 (B) redox reaction ; +3 and +5
 (C) disproportionation reaction ; -3 and +1 (D) disproportionation reaction ; -3 and +3
43. Bleaching powder contains a salt of an oxoacid as one of its components. The anhydride of that oxoacid is :
 (A) Cl_2O (B) Cl_2O_7 (C) ClO_2 (D) Cl_2O_6 [JEE 2012]
44. With respect to graphite and diamond, which of the statement(s) given below is (are) correct ? [JEE 2012]
 (A) Graphite is harder than diamond.
 (B) Graphite has higher electrical conductivity than diamond.
 (C) Graphite has higher thermal conductivity than diamond.
 (D) Graphite has higher C-C bond order than diamond.
45. Concentrated nitric acid, upon long standing, turns yellow-brown due to the formation of -
 (A) NO (B) NO_2 (C) N_2O (D) N_2O_4 [JEE 2013]
46. The correct statement(s) about O_3 is(are) [JEE 2013]
 (A) O-O bond lengths are equal (B) Thermal decomposition of O_3 is endothermic
 (C) O_3 is diamagnetic in nature (D) O_3 has a bent structure

Comprehension # 3 (Q. 47 and 48)

The reaction of Cl_2 gas with cold dilute and hot concentrated NaOH in water give sodium salt of two (different) oxoacids of chlorine P and Q respectively. The Cl_2 gas reacts with SO_2 gas, in presence of charcoal to give a product R. R reacts with white phosphorous to give a compound S. On hydrolysis, S gives as oxoacid of phosphorous T.

47. R, S and T, respectively are - [JEE 2013]
 (A) SO_2Cl_2 , PCl_5 and H_3PO_4 (B) SO_2Cl_2 , PCl_3 and H_3PO_3
 (C) SOCl_2 , PCl_3 and H_3PO_2 (D) SOCl_2 , PCl_3 and H_3PO_3
48. P and Q, respectively, are the sodium salts of -
 (A) Hypochlorous and chloric acid (B) Hypochlorous and chlorus acid
 (C) Chloric and perchloric acids (D) Chloric and hypochlorous acids

49. The unbalanced chemical reactions given in List-I show missing reagent or condition (?) which are provided in List-II. Match List-I with List-II and select the correct answer using the code given below the lists
A [JEE 2013]

List-I

- (P) $\text{PbO}_2 + \text{H}_2\text{SO}_4 \xrightarrow{?} \text{PbSO}_4 + \text{O}_2 + \text{other product}$
 (Q) $\text{Na}_2\text{S}_2\text{O}_3 + \text{H}_2\text{O} \xrightarrow{?} \text{NaHSO}_4 + \text{other product}$
 (R) $\text{N}_2\text{H}_4 \xrightarrow{?} \text{N}_2 + \text{other product}$
 (S) $\text{XeF}_2 \xrightarrow{?} \text{Xe} + \text{other product}$

List-II


- (1) NO
 (2) I_2
 (3) Warm
 (4) Cl_2

Codes :

- | | P | Q | R | S |
|-----|---|---|---|---|
| (A) | 4 | 2 | 3 | 1 |
| (C) | 1 | 4 | 2 | 3 |

- | | P | Q | R | S |
|-----|---|---|---|---|
| (B) | 3 | 2 | 1 | 4 |
| (D) | 3 | 4 | 2 | 1 |

50. Under ambient conditions, the total number of gases released as products in the final step of the reaction scheme shown below is [JEE Adv. 2014]



- (A) 0 (B) 1 (C) 2 (D) 3

51. The product formed in the reaction of SOCl_2 with white phosphorous is [JEE Adv. 2014]

- (A) PCl_3 (B) SO_2Cl_2 (C) SCl_2 (D) POCl_3

52. The **CORRECT** statement(s) for orthoboric acid is / are - [JEE Adv. 2014]

- (A) It behaves as a weak acid in water due to self ionization
 (B) Acidity of its aqueous solution increases upon addition of ethylene glycol
 (C) It has a three dimensional structure due to hydrogen bonding.
 (D) It is a weak electrolyte in water

53. The correct statement(s) regarding, (i) HClO , (ii) HClO_2 , (iii) HClO_3 and (iv) HClO_4 , is(are)

- (A) The number of $\text{Cl}=\text{O}$ bonds in (ii) and (iii) together is two [JEE Adv. 2015]
 (B) The number of lone pairs of electrons on Cl in (ii) and (iii) together is three
 (C) The hybridization of Cl in (iv) is sp^3
 (D) Amongst (i) to (iv), the strongest acid is (i)

54. When O_2 is adsorbed on a metallic surface, electron transfer occurs from the metal to O_2 . The **TRUE**, statement (s) regarding this adsorption is (are) [JEE Adv. 2015]
 (A) O_2 is physisorbed (B) heat is released
 (C) occupancy of π_{2p}^* of O_2 is increased (D) bond length of O_2 is increased
55. Under hydrolytic conditions, the compounds used for preparation of linear polymer and for chain termination, respectively, are [JEE (Adv.) 2015]
 (A) CH_3SiCl_3 and $Si(CH_3)_4$ (B) $(CH_3)_2SiCl_2$ and $(CH_3)_3SiCl$
 (C) $(CH_3)_2SiCl_2$ and CH_3SiCl_3 (D) $SiCl_4$ and $(CH_3)_3SiCl$
56. Three moles of B_2H_6 are completely reacted with methanol. The number of moles of boron containing product formed is - [JEE (Adv.) 2015]
57. The increasing order of atomic radii of the following group 13 elements is : [JEE Adv. 2016]
 (A) $Al < Ga < In < Tl$ (B) $Ga < Al < In < Tl$
 (C) $Al < In < Ga < Tl$ (D) $Al < Ga < Tl < In$
58. The crystalline form of borax has [JEE Adv. 2016]
 (A) Tetranuclear $[B_4O_5(OH)_4]^{2-}$ unit
 (B) All boron atoms in the same plane
 (C) Equal number of sp^2 and sp^3 hybridized boron atoms
 (D) One terminal hydroxide per boron atom
59. The nitrogen containing compound produced in the reaction of HNO_3 with P_4O_{10} [JEE Adv. 2016]
 (A) can also be prepared by reaction of P_4 and HNO_3
 (B) is diamagnetic
 (C) contains one N-N bond
 (D) reacts with Na metal producing a brown gas
60. The correct statements(s) about the oxoacids, $HClO_4$ and $HClO$, is (are) - [JEE Adv. 2017]
 (A) $HClO_4$ is more acidic than $HClO$ because of the resonance stabilization of its anion
 (B) $HClO_4$ is formed in the reaction between Cl_2 and H_2O
 (C) The central atom in Both $HClO_4$ and $HClO$ is sp^3 hybridized
 (D) The conjugate base of $HClO_4$ is weaker base than H_2O
61. The colour of the X_2 molecules of group 17 elements changes gradually from yellow to violet down the group. This is due to - [JEE Adv. 2017]
 (A) the physical state of X_2 at room temperature changes from gas to solid down the group
 (B) decrease in HOMO-LUMO gap down the group
 (C) decrease in $\pi^*-\sigma^*$ gap down the group
 (D) decrease in ionization energy down the group

62. The order of the oxidation state of the phosphorus atom in H_3PO_2 , H_3PO_4 , H_3PO_3 and $\text{H}_4\text{P}_2\text{O}_6$ is [JEE Adv. 2017]
- (A) $\text{H}_3\text{PO}_4 > \text{H}_4\text{P}_2\text{O}_6 > \text{H}_3\text{PO}_3 > \text{H}_3\text{PO}_2$
 (B) $\text{H}_3\text{PO}_3 > \text{H}_3\text{PO}_2 > \text{H}_3\text{PO}_4 > \text{H}_4\text{P}_2\text{O}_6$
 (C) $\text{H}_3\text{PO}_2 > \text{H}_3\text{PO}_3 > \text{H}_4\text{P}_2\text{O}_6 > \text{H}_3\text{PO}_4$
 (D) $\text{H}_3\text{PO}_4 > \text{H}_3\text{PO}_2 > \text{H}_3\text{PO}_3 > \text{H}_4\text{P}_2\text{O}_6$
63. The option(s) with only amphoteric oxides is (are): [JEE Adv. 2017]
- (A) Cr_2O_3 , CrO , SnO , PbO
 (B) NO , B_2O_3 , PbO , SnO_2
 (C) Cr_2O_3 , BeO , SnO , SnO_2
 (D) ZnO , Al_2O_3 , PbO , PbO_2
64. Among the following, the correct statement(s) is are [JEE Adv. 2017]
- (A) $\text{Al}(\text{CH}_3)_3$ has the three-centre two-electron bonds in its dimeric structure
 (B) AlCl_3 has the three-centre two-electron bonds in its dimeric structure
 (C) BH_3 has the three-centre two-electron bonds in its dimeric structure
 (D) The Lewis acidity of BCl_3 is greater than that of AlCl_3

Paragraph for Q.65 & 66

Upon heating KClO_3 in the presence of catalytic amount of MnO_2 , a gas **W** is formed. Excess amount of **W** reacts with white phosphorus to give **X**. The reaction of **X** with pure HNO_3 gives **Y** and **Z**. [JEE Adv. 2017]

65. **W** and **X** are, respectively
- (A) O_3 and P_4O_6 (B) O_2 and P_4O_{10} (C) O_3 and P_4O_{10} (D) O_2 and P_4O_6
66. **Y** and **Z** are, respectively
- (A) N_2O_4 and H_3PO_3 (B) N_2O_4 and HPO_3
 (C) N_2O_5 and HPO_3 (D) N_2O_3 and H_3PO_4
67. The compound(s) which generate(s) N_2 gas upon thermal decomposition below 300°C is (are) (IIT JEE 2018)
- (A) NH_4NO_3 (B) $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$ (C) $\text{Ba}(\text{N}_3)_2$ (D) Mg_3N_2

INTEGER

68. The total number of compounds having at least one bridging oxo group among the molecules given below is _____. [JEE Adv. 2018]
- N_2O_3 , N_2O_5 , P_4O_6 , P_4O_7 , $\text{H}_4\text{P}_2\text{O}_5$, $\text{H}_5\text{P}_3\text{O}_{10}$, $\text{H}_2\text{S}_2\text{O}_3$, $\text{H}_2\text{S}_2\text{O}_5$

69. A tin chloride **Q** undergoes the following reactions (not balanced) [JEE Adv. 2019]
 $\text{Q} + \text{Cl}^- \rightarrow \text{X}$
 $\text{Q} + \text{Me}_3\text{N} \rightarrow \text{Y}$
 $\text{Q} + \text{CuCl}_2 \rightarrow \text{Z} + \text{CuCl}$
 X is a monoanion having pyramidal geometry. Both **Y** and **Z** are neutral compounds. Choose the correct option(s).
 (1) The central atoms in **X** is sp^3 hybridized
 (2) The oxidation state of the central atom in **Z** is +2
 (3) The central atom in **Z** has one lone pair of electrons
 (4) There is a coordinate bond in **Y**
70. At 143 K. the reaction of XeF_4 with O_2F_2 produces a xenon compound **Y**. The total number of lone pair(s) of electrons present on the whole molecule of **Y** is _____ [JEE Adv. 2019]
71. The amount of water produced (in g) in the oxidation of 1 mole of rhombic sulphur by conc. HNO_3 to a compound with the highest oxidation state of sulphur is ____ [JEE Adv. 2019]
 (Given data : Molar mass of water = 18 g mol^{-1})
72. Each of the following options contains a set of four molecules. Identify the option(s) where all four molecules possess permanent dipole moment at room temperature. [JEE Adv. 2019]
 (1) BeCl_2 , CO_2 , BCl_3 , CHCl_3 (2) SO_2 , $\text{C}_6\text{H}_5\text{Cl}$, H_2Se , BrF_5
 (3) BF_3 , O_3 , SF_6 , XeF_6 (4) NO_2 , NH_3 , POCl_3 , CH_3Cl
73. Among B_2H_6 , $\text{B}_3\text{N}_3\text{H}_6$, N_2O , N_2O_4 , $\text{H}_2\text{S}_2\text{O}_3$ and $\text{H}_2\text{S}_2\text{O}_8$, the total number of molecules containing covalent bond between two atoms of the same kind is _____ [JEE Adv. 2019]

55. Ans.(1)

56. Ans.(1)

57. Ans.(3)

58. Ans.(3)

59. Ans.(2)

60. Ans.(1)

61. Ans.(1)

62. Ans. (1)

63. Ans.(1.66 to 1.67)

64. Ans.(2)

65. Ans. (4)

66. Ans.(2)

67. Ans.(1)

68. Ans.(4)

EXERCISE # J- ADVANCED


Fill in the blanks

1. $P \equiv P$


$3p_{\pi}$ - $3p_{\pi}$ bond is not effective due to large size of P.

2. $2I^{-} \text{ (aqueous)} + Cl_2 \longrightarrow I_2 + 2Cl^{-} \text{ (aqueous)}$ (i) $2I^{-} \text{ (aqueous)} \longrightarrow I_2 \text{ (s)} + 2e^{-}$ (ii) $Cl_2 \text{ (g)} + 2e^{-} \longrightarrow 2Cl^{-} \text{ (aq)}$







Thus, I^{-} is oxidised into I_2 by Cl_2 due to higher oxidised potential of Cl_2 than I_2











3. X : BCl_3 Y : B_2H_6 

X





Y

4. (i) $3SiCl_4 + 4Al \longrightarrow 3Si + 4AlCl_3$ (in one step)(ii) $SiCl_4 + 2Mg \longrightarrow 2MgCl_2 + Si$ (iii) $SiCl_4 + 4H_2O \longrightarrow Si(OH)_4 + 4HCl$ 



9. Oxidation state




Objective



Boron always forms covalent bond because boron requires very high energy of form B^{3+} and again B^{3+} due to its very small size having high polarising power thus cause greater polarisation and eventually significant covalent characteristics-Fajans rule.

34. C



Comprehension # 1 (Q. 35 to 37)


35. A 36. C 37. A or B

Comprehension # 2 (Q.38 to 40)


38. C 39. C 40. B 41. B 42. C 43. A
 44. B,D 45. B 46. A,C,D 47. A 48. A 49. D
 50. C 51. A 52. B,D 53. B,C 54. B,C,D 55. B
 56. 6
 57. (B)

Sol. The order of radius of 13th group elements is $\text{Ga} < \text{Al} < \text{In} < \text{Tl}$.


Reason $\Rightarrow$ Due to poor shielding effect of d-orbital, radius of Ga is smaller than Al.

58. (A,C,D)

- (A) Having $[\text{B}_4\text{O}_5(\text{OH})_4]^{2-}$ tetranuclear (boron) unit
 (B) All boron atoms not in same plane
 (C) Two boron are sp^2 hybridised and two boron are sp^3 hybridised
 (D) One terminal hydroxide per boron atom is present.

59. (B,D)

- (A) $\text{P}_4 + 20\text{HNO}_3 \rightarrow 4\text{H}_3\text{PO}_4 + 20\text{NO}_2\uparrow + 4\text{H}_2\text{O}$
 (B) N_2O_5 is diamagnetic in nature



N_2O_5 contains one N-O-N bond not N-N bond.

- (D) $\text{Na} + \text{N}_2\text{O}_5 \rightarrow \text{NaNO}_3 + \text{NO}_2\uparrow$
 (Brown gas)

60. A,C,D 61. B,C 62. A 63. C,D
 64. A,C,D 65. B 66. C 67. B,C
 68. 5 or 6 69. (1,4) 70. (19.00) 71. (288.00)
 72. (2,4) 73. (4.00)