

## REDOX REACTION

### Question Stem for Question Nos. 1 and 2

#### Question Stem

A sample (5.6 g) containing iron is completely dissolved in cold dilute HCl to prepare a 250 mL of solution. Titration of 25.0 mL of this solution requires 12.5 mL of 0.03 M  $\text{KMnO}_4$  solution to reach the end point. Number of moles of  $\text{Fe}^{2+}$  present in 250 mL solution is  $x \times 10^{-2}$  (consider complete dissolution of  $\text{FeCl}_2$ ). The amount of iron present in the sample of  $y\%$  by weight.

(Assume :  $\text{KMnO}_4$  reacts only with  $\text{Fe}^{2+}$  in the solution)

Use : Molar mass of iron as  $56 \text{ g mol}^{-1}$

[JEE(Advanced) 2021]

- The value of  $x$  is \_\_\_\_\_.
- The value of  $y$  is \_\_\_\_\_.
- In the chemical reaction between stoichiometric quantities of  $\text{KMnO}_4$  and  $\text{KI}$  in weakly basic solution, what is the number of moles of  $\text{I}_2$  released for 4 moles of  $\text{KMnO}_4$  consumed? [JEE(Advanced) 2020]
- An acidified solution of potassium chromate was layered with an equal volume of amyl alcohol. When it was shaken after the addition of 1 mL of 3%  $\text{H}_2\text{O}_2$ , a blue alcohol layer was obtained. The blue color is due to the formation of a chromium (VI) compound 'X'. What is the number of oxygen atoms bonded to chromium through only single bonds in a molecule of X? [JEE(Advanced) 2020]
- The amount of water produced (in g) in the oxidation of 1 mole of rhombic sulphur by conc.  $\text{HNO}_3$  to a compound with the highest oxidation state of sulphur is \_\_\_\_\_.  
(Given data : Molar mass of water =  $18 \text{ g mol}^{-1}$ ) [JEE(Advanced) 2019]
- To measure the quantity of  $\text{MnCl}_2$  dissolved in an aqueous solution, it was completely converted to  $\text{KMnO}_4$  using the reaction,  
 $\text{MnCl}_2 + \text{K}_2\text{S}_2\text{O}_8 + \text{H}_2\text{O} \rightarrow \text{KMnO}_4 + \text{H}_2\text{SO}_4 + \text{HCl}$  (equation not balanced).  
Few drops of concentrated HCl were added to this solution and gently warmed. Further, oxalic acid (225 g) was added in portions till the colour of the permanganate ion disappeared. The quantity of  $\text{MnCl}_2$  (in mg) present in the initial solution is \_\_\_\_\_.  
(Atomic weights in  $\text{g mol}^{-1}$  : Mn = 55, Cl = 35.5) [JEE(Advanced) 2018]
- In neutral or faintly alkaline solution, 8 moles permanganate anion quantitatively oxidize thiosulphate anions to produce X moles of a sulphur containing product. the magnitude of X is [JEE(Advanced) 2016]
- For the reaction  
 $\text{I}^- + \text{ClO}_3^- + \text{H}_2\text{SO}_4 \rightarrow \text{Cl}^- + \text{HSO}_4^- + \text{I}_2$   
The correct statement(s) in the balanced equation is / are : [JEE(Advanced) 2014]  
(A) Stoichiometric coefficient of  $\text{HSO}_4^-$  is 6 (B) Iodide is oxidized  
(C) Sulphur is reduced (D)  $\text{H}_2\text{O}$  is one of the products
- Hydrogen peroxide in its reaction with  $\text{KIO}_4$  and  $\text{NH}_2\text{OH}$  respectively, is acting as a [JEE(Advanced) 2014]  
(A) reducing agent, oxidising agent (B) reducing agent, reducing agent  
(C) oxidising agent, oxidising agent (D) oxidising agent, reducing agent

## SOLUTIONS

1. Ans. (1.87 or 1.88)

2. Ans. (18.75)

Solution for Q.1 & Q.2



$$\frac{x}{10 \text{ mole}} \quad 12.5 \text{ ml}$$

$$0.03 \text{ M}$$

$$n_f = 1 \quad n_f = 5$$

$$\frac{x}{10} = \frac{12.5 \times 0.03 \times 5}{1000}$$

$$x = 0.01875 \quad (x = 1.88 \text{ or } 1.87)$$

$$\text{wt of Fe} = 1.05 \text{ g}$$

$$\% \text{ Fe} = \frac{1.05}{5.6} \times 100 = 18.75$$

3. Ans. (6)




$$\text{Eq of KMnO}_4 = \text{Eq of I}_2$$


$$4 \times 3 = n \times 2$$

$$n = 6$$

4. Ans. (4)




Here the structure of  $\text{CrO}_5$  is :-





Here, single bonded O-atoms with Cr is = 04

5. Ans. (288.00 to 288.30)



1 mole of rhombic sulphur produce 16 mole of  $\text{H}_2\text{O}$  i.e. 288 gm of  $\text{H}_2\text{O}$

6. Ans. (126)




$$m_{\text{eq}} \text{ of } \text{C}_2\text{O}_4^{--} = m_{\text{eq}} \text{ of } \text{MnO}_4^-$$

$$2 \times 0.225/90 = a \times 5$$

$$a = 1 \times [55 + 71] = 126 \text{ mg}$$

7. **Ans. (6)**



Equivalents of  $\text{MnO}_4^-$  = equivalents of  $\text{SO}_4^{2-}$

Moles of  $\text{MnO}_4^- \times \text{n-factor} = \text{moles of } \text{SO}_4^{2-} \times \text{n-factor}$

$$8 \times 3 = X \times 4$$


$$X = 6$$

8. **Ans. (A, B, D)**



**Sol.** Oxidation half reaction :



Reduction half reaction



Multiplying equation (1) by 3 and add in (2)



9. **Ans. (A)**


+7

**Sol.**  $\text{H}_2\text{O}_2 + \text{KIO}_4 \longrightarrow \text{O}_2 + \text{I}$  (with oxidation state lower than 7)

Reducing agent

-1

+3



Oxidising

agent