

Chapter 2

Structure of Atom





Solutions

SECTION - A

Objective Type Questions

(Discovery of Fundamental Particles, Nature of Electromagnetic Radiation)

1. Which is the correct graphical representation based on photoelectric effect?



(1) I & II

(2) II & III

(3) III & IV



(4) II & IV

Sol. Answer (4)

For photoelectric effect

$$KE = h(\nu - \nu_0)$$

$$KE = h\nu - h\nu_0$$



ν_0 = Threshold frequency

$\therefore$ [KE of e^- increases after crossing. Threshold frequency]

(Bohr's Model for Hydrogen atom)

2. What will be the longest wavelength line in Balmer series of spectrum of H-atom?

- (1) 546 nm (2) 656 nm (3) 566 nm (4) 556 nm

Sol. Answer (2)

All the wavelength are in visible region i.e. between 400 nm to 760 nm. Therefore maximum wavelength line will be 656 nm.

3. In hydrogen atom, energy of first excited state is -3.4 eV. Then find out KE of same orbit of hydrogen atom

- (1) $+3.4$ eV (2) $+6.8$ eV (3) -13.6 eV (4) $+13.6$ eV

Sol. Answer (1)

$$\frac{KE}{E_{\text{Total}}} = -1$$

Total energy = -3.4 eV (Given)

$$\therefore KE = -(-3.4 \text{ eV}) = +3.4 \text{ eV}$$

4. Total number of spectral lines in UV region, during transition from 5th excited state to 1st excited state

- (1) 10 (2) 3 (3) 4 (4) Zero

Sol. Answer (4)

As 1st excited state means $n_1 = 2$

For 5th excited state means $n_2 = 6$

$\therefore e^-$ will transit between 6th level to 2nd level

No transition will be upto 1st level. Because no line will appear in Lyman series i.e. UV region.

5. The first emission line in the atomic spectrum of hydrogen in the Balmer series appears at

- (1) $\frac{5R}{36} \text{ cm}^{-1}$ (2) $\frac{3R}{4} \text{ cm}^{-1}$ (3) $\frac{7R}{144} \text{ cm}^{-1}$ (4) $\frac{9R}{400} \text{ cm}^{-1}$

Sol. Answer (1)

1st line in the Balmer series means $n_1 = 2, n_2 = 3$

$$\bar{\nu} = \frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] z^2 \quad \text{for H } z = 1$$

$$\bar{\nu} = \frac{1}{\lambda} = R \left[\frac{1}{2^2} - \frac{1}{3^2} \right] \times 1^2 = R \left[\frac{1}{4} - \frac{1}{9} \right] = \frac{5R}{36} \text{ cm}^{-1}$$

6. In a hydrogen atom, if the energy of electron in the ground state is $-x$ eV., then that in the 2nd excited state of He^+ is

- (1) $-x$ eV (2) $-\frac{4}{9}x$ eV (3) $+2x$ eV (4) $-\frac{9}{4}x$ eV

Sol. Answer (2)

$$E_n = \frac{E_{\text{ground state}}}{n^2} \times Z^2 \quad \begin{cases} E_{\text{ground}} = x \text{ eV} & \text{given} \\ n = 3 & \text{because } 2^{\text{nd}} \text{ excited state} \\ z = 2 & \text{because} \end{cases}$$

$$= -\frac{x}{(3)^2} \times (2)^2 = -\frac{4}{9} x \text{ eV}$$

7. The wavelength of radiation emitted, when in He^+ electron falls from infinity to stationary state would be ($R = 1.097 \times 10^7 \text{ m}^{-1}$)

- (1) $2.2 \times 10^{-8} \text{ m}$ (2) $22 \times 10^{-9} \text{ m}$ (3) 120 m (4) $22 \times 10^7 \text{ m}$

Sol. Answer (1)

$$n_1 = 1 \quad \text{For } \text{He}^{\oplus} \quad z = 2$$

$$n_2 = \infty \text{ given}$$

$$\frac{1}{\lambda_{\text{He}^{\oplus}}} = R \left[\frac{1}{1^2} - \frac{1}{\infty^2} \right] \times (2)^2$$

$$\frac{1}{\lambda_{\text{He}^{\oplus}}} = 109678 \times 4 \text{ cm}^{-1}$$

$$\lambda_{\text{He}^{\oplus}} = \frac{1}{109678 \times 4} = \frac{1}{438712} = 2.2 \times 10^{-6} \text{ cm}$$

$$= 2.2 \times 10^{-8} \text{ m}$$

8. In Bohr series of lines of hydrogen spectrum, the third line from the red end corresponds to which one of the following inter-orbit jumps of the electron for Bohr orbits in an atom of hydrogen?

- (1) $3 \rightarrow 1$ (2) $5 \rightarrow 2$ (3) $2 \rightarrow 5$ (4) $3 \rightarrow 2$

Sol. Answer (2)

Third line means third excited state

i.e. $n_1 = 2$ Balmer series (visible region)

$n_2 = 5$ Third line

$\therefore$ Third line will appear when electron comes from 5^{th} energy level to 2^{nd} level.

9. The correct order of energy difference between adjacent energy levels in H atom

- (1) $E_2 - E_1 > E_3 - E_2 > E_4 - E_3$ (2) $E_2 - E_1 > E_4 - E_3 > E_3 - E_2$
 (3) $E_4 - E_3 > E_3 - E_2 > E_2 - E_1$ (4) $E_3 - E_2 > E_4 - E_3 > E_2 - E_1$

Sol. Answer (1)

In H atom

$$E_1 = \frac{-1312}{1^2} \quad E_2 = \frac{-1312}{4} \quad E_3 = \frac{-1312}{9} \quad E_4 = \frac{-1312}{16} \quad E_5 = \frac{-1312}{25}$$

$$\therefore (E_2 - E_1) > (E_3 - E_2) > (E_4 - E_3) \dots\dots$$

[Alternatively as the distance from the nucleus increases the value of ΔE (energy difference between two shell) decreases]

10. Which of the following electronic in a transition hydrogen atom will require the largest amount of energy?

- (1) $n = 1$ to $n = 2$ (2) $n = 2$ to $n = 3$ (3) $n = 1$ to $n = \infty$ (4) $n = 3$ to $n = 5$

Sol. Answer (3)

Largest amount of energy is required for the transition between $1 \rightarrow \infty$

$$\Delta E = hc \times \frac{1}{\lambda} \Rightarrow hcR \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \text{ [Large the difference between } n_1 \text{ and } n_2 \text{ large will be the value of DE]}$$

11. The time taken by the electron in one complete revolution in the n^{th} Bohr's orbit of the hydrogen atom is

- (1) Inversely proportional to n^2 (2) Directly proportional to n^3
 (3) Directly proportional to $\frac{h}{2\pi}$ (4) Inversely proportional to $\frac{n}{h}$

Sol. Answer (2)

$$\text{Time period} = \frac{\text{circumference}}{\text{velocity}} = \frac{2\pi r}{v_n} = \frac{n^3}{z^2} \times 1.5 \times 10^{-16} \text{ seconds}$$

$$\text{Time period} \propto n^3$$

12. What will be the ratio of the wavelength of the first line to that of the second line of Paschen series of H atom?

- (1) 256 : 175 (2) 175 : 256 (3) 15 : 16 (4) 24 : 27

Sol. Answer (1)

First time of paschen series $n_1 = 3$, $n_2 = 4$

$$\frac{1}{\lambda_1} = R \left[\frac{1}{9} - \frac{1}{16} \right]$$

$$\frac{1}{\lambda_1} = \frac{7R}{144} \quad \lambda_1 = \frac{144}{7R}$$

Second line of paschen series $n_1 = 3$, $n_2 = 5$

$$\frac{1}{\lambda_2} = R \left[\frac{1}{9} - \frac{1}{25} \right]$$

$$\frac{1}{\lambda_2} = \frac{16R}{225} \quad \lambda_2 = \frac{225}{16R}$$

$$\frac{\lambda_1}{\lambda_2} = \frac{144}{7R} \times \frac{16R}{225} = \frac{2304}{1575} = \frac{256}{175}$$

13. For the transition from $n = 2 \rightarrow n = 1$, which of the following will produce shortest wavelength?

- (1) H atom (2) D atom (3) He^+ ion (4) Li^{2+} ion

Sol. Answer (4)

$$\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] z^2 \text{ as } n_1 \propto n_2 \text{ are constant}$$

$$\therefore \lambda \propto \frac{1}{z^2} \text{ more the nuclear charge smaller will be the } \lambda$$

$$\text{H} = z = 1$$

$$\text{D} = z = 1 \quad \therefore \text{Li}^{2+} \text{ have shorter wavelength}$$

$$\text{He}^+ = z = 2$$

$$\text{Li}^{2+} = z = 3$$

(Towards Quantum Mechanical Model of the Atom (Dual behaviour of matter, Heisenberg's uncertainty Principle))

14. The uncertainty in momentum of an electron is $1 \times 10^{-5} \text{ kg-m/s}$. The uncertainty in its position will be ($h = 6.62 \times 10^{-34} \text{ kg-m}^2/\text{s}$)

(1) $5.27 \times 10^{-30} \text{ m}$ (2) $1.05 \times 10^{-26} \text{ m}$ (3) $1.05 \times 10^{-28} \text{ m}$ (4) $5.25 \times 10^{-28} \text{ m}$

Sol. Answer (1)

$$\Delta P = 10^{-5} \text{ kgms}^{-1}$$

$$\Delta x \times \Delta P = \frac{h}{4\pi}$$

$$\Delta x = \frac{6.6 \times 10^{-34} \text{ J}}{10^{-5} \times 4 \times 3.14} = 5.2 \times 10^{-30} \text{ m}$$

15. Two particles *A* and *B* are in motion. If the wavelength associated with particle *A* is $5 \times 10^{-8} \text{ m}$; calculate the wavelength associated with particle *B* if its momentum is half of *A*.

(1) $5 \times 10^{-8} \text{ m}$ (2) 10^{-5} cm (3) 10^{-7} cm (4) $5 \times 10^{-8} \text{ cm}$

Sol. Answer (2)

$$\lambda_A = \frac{h}{P_A} \quad \lambda_B = \frac{h}{P_B} \quad P_B = \frac{1}{2} P_A \quad (\text{Given})$$

$$\frac{\lambda_A}{\lambda_B} = \frac{\frac{h}{P_A}}{\frac{h}{P_B}} = \frac{P_B}{P_A} \quad \text{Putting } P_B = \frac{1}{2} P_A$$

$$\frac{\lambda_A}{\lambda_B} = \frac{1}{2} \frac{P_A}{P_A}$$

$$\lambda_B = 2\lambda_A \quad [\lambda_A = 5 \times 10^{-8} \text{ m}]$$

$$\lambda_B = 2 \times 5 \times 10^{-8}$$

$$= 10 \times 10^{-8} \text{ m} \quad \therefore 1 \text{ m} = 100 \text{ cm}$$

$$= 10^{-7} \text{ m} = 10^{-5} \text{ cm}$$

(Quantum Mechanical Model of the Atom)

16. Maximum number of electrons in a subshell with $l = 3$ and $n = 4$ is

(1) 10 (2) 12 (3) 14 (4) 16

Sol. Answer (3)

$$n = 4, l = 3 \text{ means } 4f$$

for $l = 3$, $m = -3, -2, -1, 0, 1, 2, 3 = 7$ orbital

Therefore, maximum 14 electrons are present.

17. The total number of subshells in fourth energy level of an atom is

- (1) 4 (2) 8 (3) 16 (4) 32

Sol. Answer (1)

18. For which of the following sets of four quantum numbers, an electron will have the highest energy?

n	l	m	s
(1) 3	2	1	+1/2
(2) 4	2	-1	+1/2
(3) 4	1	0	-1/2
(4) 5	0	0	-1/2

Sol. Answer (2)

Energy of an electron depends upon $(n + l)$ value

More the $(n + l)$ value more will be the energy

n	l	m	s	$(n + l)$
(1) 3	2	1	+1/2	5
(2) 4	2	-1	+1/2	6 Max. $(n + l)$. max. energy
(3) 4	1	0	-1/2	5
(4) 5	0	0	-1/2	5

19. A transition element X has a configuration $(\text{Ar})3d^4$ in its +3 oxidation state. Its atomic number is

- (1) 22 (2) 19 (3) 25 (4) 26

Sol. Answer (3)

$$\begin{aligned}\text{Total number of } e^- \text{ in } X^{+3} &= [\text{Ar}] 3d^4 \\ &= 18 + 4 = 22\end{aligned}$$

$$\therefore \boxed{\text{Number of electrons in X} = 22 + 3 = 25}$$

Atomic number = 25


20. Among the following which one is not paramagnetic? [Atomic numbers; Be = 4, Ne = 10, As = 33, Cl = 17]

- (1) Ne^{2+} (2) Be^+ (3) Cl^- (4) As^+

Sol. Answer (3)

Ions having all the electron paired will be non-paramagnetic or diamagnetic

$$\text{Ne}^{+2} = 8 = 1s^2, 2s^2, 2p^4$$






2 unpaired e^-

$$\text{Be}^{\oplus} = 3 = 1s^2, 2s^1$$



1 unpaired e^-


0 unpaired e^- 3 unpaired e^-

21. Isoelectronic species are

- (1) CO , CN^- , NO^+ , C_2^{2-}
- (2) CO^- , CN , NO , C_2^-
- (3) CO^+ , CN^+ , NO^- , C_2
- (4) CO , CN , NO , C_2

Sol. Answer (1)

Isoelectronic species have same number of electrons



22. Consider the following sets of quantum number

n	l	m	s
(i) 3	0	0	+1/2
(ii) 2	2	1	+1/2
(iii) 4	3	-2	-1/2
(iv) 1	0	-1	-1/2
(v) 3	2	3	+1/2

Which of the following sets of quantum number is not possible?

- (1) (i), (ii), (iii) and (iv) (2) (ii), (iv) and (v) (3) (i) and (iii) (4) (ii), (iii) and (iv)

Sol. Answer (2)

(ii), (iv), and (v) are not possible

- | | | | | |
|---------------|-------|--------|----------|---|
| (ii) $n = 2$ | l = 2 | m = 1 | s = +1/2 | l not equal to n not possible |
| (iii) $n = 1$ | l = 0 | m = -1 | s = -1/2 | Not possible because m = -1 where l = 0 |
| (iv) $n = 3$ | l = 2 | m = 3 | s = +1/2 | Not possible because m = 3 is not for l = 2 |

23. Any f -orbital can accommodate upto

- (1) 2 electrons with parallel spin
- (2) 6 electrons
- (3) 2 electrons with opposite spin
- (4) 14 electrons

Sol. Answer (3)

Any orbital have maximum of two electrons with opposite spin.

24. For principal quantum number $n = 5$, the total number of orbitals having $l = 3$ is

- (1) 7 (2) 14 (3) 9 (4) 18

Sol. Answer (1)For $l = 3$ $m = -3, -2, -1, 0, +1, +2, +3$ *i.e.*, 7 orbitals are present

25. The four quantum numbers of valence electron of potassium are

(1) $4, 0, 1, \frac{1}{2}$

(2) $4, 1, 0, \frac{1}{2}$

(3) $4, 0, 0, \frac{1}{2}$

(4) $4, 1, 1, \frac{1}{2}$

Sol. Answer (3) $K = 19 = 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^1$ last e^- Last electron $4s^1$

$$\therefore n = 4 \quad l = 0 \quad m = 0 \quad s = +\frac{1}{2}$$

26. In the ground state, an element has 13 electrons in its M-shell. The element is

(1) Manganese

(2) Cobalt

(3) Nickel

(4) Iron

Sol. Answer (1)M shell means 3rd orbit

$$Mn = 25 = 1s^2, 2s^2, 2p^6, \underbrace{3s^2, 3p^6}_8, \underbrace{4s^2, 3d^5}_5 \quad \text{total 13 } e^- \text{ in 3 orbit}$$

$$Co = 27 = 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^2, 3d^7 \quad \text{total 15 } e^- \text{ in 3 orbit}$$

$$Ni = 28 = 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^2, 3d^8 \quad \text{total 16 } e^- \text{ in 3 orbit}$$

$$Fe = 26 = 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^2, 3d^6 \quad \text{total 14 } e^- \text{ in 3 orbit}$$

27. Which combinations of quantum numbers n , l , m and s for the electron in an atom does not provide a permissible solutions of the wave equation?

(1) $3, 2, -2, \frac{1}{2}$

(2) $3, 3, 1, -\frac{1}{2}$

(3) $3, 2, 1, \frac{1}{2}$

(4) $3, 1, 1, -\frac{1}{2}$

Sol. Answer (2) $n = 3$ $l = 3$ [Not possible because value of l can never be equals to n]

28. The orbital angular momentum of electron in 4s orbital is

(1) $\frac{1}{2} \cdot \frac{h}{2\pi}$

(2) Zero

(3) $\frac{h}{2\pi}$

(4) $(2.5) \frac{h}{2\pi}$

Sol. Answer (2)

Orbital angular momentum = $\sqrt{l(l+1)} \frac{h}{4\pi}$

For 4s electron the value of $l = 0$ $\therefore$ [orbital angular momentum = zero]

29. Radial nodes present in 3s and 3p-orbitals are respectively

(1) 0, 2

(2) 2, 1

(3) 1, 1

(4) 2, 2

Sol. Answer (2)Radial nodes = $(n - l - 1)$ for 3s $(3 - 0 - 1) = 2$; For 3p $(3 - 1 - 1) = 1$

30. Quantum numbers for some electrons are given below

$$A : n = 4, l = 1$$

$$B : n = 4, l = 0$$

$$C : n = 3, l = 2$$

$$D : n = 3, l = 1$$

The correct increasing order of energy of electrons

$$(1) A < B < C < D$$

$$(2) D < C < B < A$$

$$(3) D < B < C < A$$

$$(4) C < B < A < D$$

Sol. Answer (3)

$$\text{Energy} = (n + l)$$

$$A = n = 4 \quad l = 1 \quad = 4 + 1 = 5$$

$$B = n = 4 \quad l = 0 \quad = 4 + 0 = 4$$

$$C = n = 3 \quad l = 2 \quad = 3 + 2 = 5$$

$$D = n = 3 \quad l = 1 \quad = 3 + 1 = 4$$

According to Pauli exclusion principle

(1) Larger the $(n + l)$; larger will be energy

(2) Same value of $(n + l)$; smaller n ; more will be energy

$$\therefore \boxed{D < B < C < A}$$

31. The number of lobes in most of the d -orbitals are

$$(1) 6$$

$$(2) 8$$

$$(3) 10$$

$$(4) 4$$

Sol. Answer (4)

32. For which of the following options $m = 0$ for all orbitals?

$$(1) 2s, 2p_x, 3d_{xy}$$

$$(2) 3s, 2p_z, 3d_{z^2}$$

$$(3) 2s, 2p_z, 3d_{x^2-y^2}$$

$$(4) 3s, 3p_x, 3d_{yz}$$

Sol. Answer (2)

$$\text{Value of } m = 0 \text{ for } 3s, 2p_z \text{ and } 3d_{z^2}$$

33. In any sub-shell, the maximum number of electrons having same value of spin quantum number is

$$(1) \sqrt{l(l+1)}$$

$$(2) l + 2$$

$$(3) 2l + 1$$

$$(4) 4l + 2$$

Sol. Answer (3)

$$\text{Total number of electron in subshell} = 2(2l + 1) \quad l = \text{angular quantum number}$$

$$\text{Number of electrons having same spin} = \frac{2(2l+1)}{2} = (2l+1)$$

[Because half e^- have clockwise and half e^- have anti clockwise spin]

34. If each orbital can hold a maximum of 3 electrons. The number of elements in 2nd period of periodic table (long form) is

$$(1) 27$$

$$(2) 9$$

$$(3) 18$$

$$(4) 12$$

Sol. Answer (4)

$$\text{For 2nd period electronic configuration} = 2s^2, 2p^6$$

$$\text{If each orbital have } 3e^- \text{ then electronic configuration} = 2s^3, 2p_x^3, 2p_y^3, 2p_z^3$$

Total 12 e^- will present

SECTION - B

Previous Years Questions


1. Which one is a **wrong** statement?

[NEET-2018]


- (1) Total orbital angular momentum of electron in 's' orbital is equal to zero
- (2) An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers
- (3) The value of m for d_{z^2} is zero
- (4) The electronic configuration of N atom is

**Sol.** Answer (4)

According to Hund's Rule of maximum multiplicity, the correct electronic configuration of N-atom is



OR



∴ Option (4) violates Hund's Rule.

2. Which one is the wrong statement?

[NEET-2017]

- (1) de-Broglie's wavelength is given by $\lambda = \frac{h}{mv}$, where m = mass of the particle, v = group velocity of the particle
- (2) The uncertainty principle is $\Delta E \times \Delta t \geq \frac{h}{4\pi}$
- (3) Half-filled and fully filled orbitals have greater stability due to greater exchange energy, greater symmetry and more balanced arrangement
- (4) The energy of 2s orbital is less than the energy of 2p orbital in case of Hydrogen like atoms

Sol. Answer (4)

Energy of 2s-orbital and 2p-orbital in case of hydrogen like atoms is equal.

3. How many electrons can fit in the orbital for which $n = 3$ and $l = 1$?

[NEET-Phase-2-2016]

- | | | | |
|-------|-------|--------|--------|
| (1) 2 | (2) 6 | (3) 10 | (4) 14 |
|-------|-------|--------|--------|

Sol. Answer (1)


An orbital can accommodate maximum of 2 electrons with anti-parallel spins.

4. Which of the following pairs of d-orbitals will have electron density along the axes?

[NEET-Phase-2-2016]

- | | | | |
|-----------------------|----------------------|----------------------------|---------------------------|
| (1) d_{z^2}, d_{xz} | (2) d_{xz}, d_{yz} | (3) $d_{z^2}, d_{x^2-y^2}$ | (4) $d_{xy}, d_{x^2-y^2}$ |
|-----------------------|----------------------|----------------------------|---------------------------|

Sol. Answer (3)



5. Two electrons occupying the same orbital are distinguished by

[NEET-2016]

- (1) Spin quantum number (2) Principal quantum number
(3) Magnetic quantum number (4) Azimuthal quantum number

Sol. Answer (1)

Fact.

6. The angular momentum of electron in 'd' orbital is equal to

[AIPMT-2015]

- (1) $0 \hbar$ (2) $\sqrt{6} \hbar$ (3) $\sqrt{2} \hbar$ (4) $2\sqrt{3} \hbar$

Sol. Answer (2)

$$\text{Angular momentum} = \sqrt{l(l+1)} \frac{\hbar}{2\pi} = \sqrt{l(l+1)} \hbar$$

7. What is the maximum number of orbitals that can be identified with the following quantum numbers?

$$n = 3, l = 1, m_l = 0$$

[AIPMT-2014]

- (1) 1 (2) 2 (3) 3 (4) 4

Sol. Answer (1)

It represents 3p orbital

8. Calculate the energy in joule corresponding to light of wavelength 45 nm: (Planck's constant $h = 6.63 \times 10^{-34}$ Js; speed of light $c = 3 \times 10^8$ ms⁻¹)

[AIPMT-2014]

- (1) 6.67×10^{15} (2) 6.67×10^{11} (3) 4.42×10^{-15} (4) 4.42×10^{-18}

Sol. Answer (4)

$$E = \frac{hc}{\lambda} = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{45 \times 10^{-9}} = \frac{6.63}{15} \times 10^{-17} = 4.42 \times 10^{-18} \text{ J}$$

9. Be²⁺ is isoelectronic with which of the following ions?

[AIPMT-2014]

- (1) H⁺ (2) Li⁺ (3) Na⁺ (4) Mg²⁺

Sol. Answer (2)

10. What is the maximum numbers of electrons that can be associated with the following set of quantum numbers?

$$n = 3, l = 1 \text{ and } m = -1$$

[NEET-2013]

- (1) 6 (2) 4 (3) 2 (4) 10

Sol. Answer (3)

Value of $m = -1$ represents one orbital. Therefore maximum number of electrons will be two

11. The value of Planck's constant is 6.63×10^{-34} Js. The speed of light is 3×10^{17} nm s⁻¹. Which value is closest to the wavelength in nanometer of a quantum of light with frequency of 6×10^{15} s⁻¹?

[NEET-2013]

- (1) 25 (2) 50 (3) 75 (4) 10

Sol. Answer (2)

$$\lambda = \frac{c}{\nu} = \frac{3.0 \times 10^{17} \text{ nms}^{-1}}{6 \times 10^{15} \text{ s}^{-1}} = \frac{1}{2} \times 10^2 \text{ nm}$$

$$= 0.5 \times 10^2 \text{ nm} = 50 \text{ nm}$$

12. Based on equation $E = -2.178 \times 10^{-18} \text{ J} \left(\frac{Z^2}{n^2} \right)$ certain conclusions are written. Which of them is **not** correct?

[NEET-2013]

- (1) Larger the value of n , the larger is the orbit radius
- (2) Equation can be used to calculate the change in energy when the electron changes orbit
- (3) For $n = 1$, the electron has a more negative energy than it does for $n = 6$ which means that the electron is more loosely bound in the smallest allowed orbit
- (4) The negative sign in equation simply means that the energy of electron bound to the nucleus is lower than it would be if the electrons were at the infinite distance from the nucleus

Sol. Answer (3)

In ($n = 1$) 1st shell e^- is tightly held compared to $n = 6$ (6th shell)

13. The correct set of four quantum numbers for the valence electron of rubidium atom ($Z = 37$) is

[AIPMT (Prelims)-2012]

- (1) 5, 0, 0, $+\frac{1}{2}$ (2) 5, 1, 0, $+\frac{1}{2}$ (3) 5, 1, 1, $+\frac{1}{2}$ (4) 6, 0, 0, $+\frac{1}{2}$

Sol. Answer (1)

Rb = 37 = [Ar] $4s^2, 3d^{10}, 4p^6$ last $5s^1 e^-$

$$\therefore n = 5, l = 0, m = 0, s = +\frac{1}{2}$$

14. Maximum number of electrons in a subshell with $l = 3$ and $n = 4$ is

[AIPMT (Prelims)-2012]

- (1) 10 (2) 12 (3) 14 (4) 16

Sol. Answer (3)

$n = 4$ $l = 3$ represents $4f$ subshell having 7 orbitals

$$\therefore \boxed{\text{Total number of electrons} = 14}$$

15. The orbital angular momentum of a p-electron is given as

[AIPMT (Mains)-2012]

- (1) $\frac{h}{\sqrt{2\pi}}$ (2) $\sqrt{3} \frac{h}{2\pi}$ (3) $\sqrt{\frac{3}{2}} \frac{h}{\pi}$ (4) $\sqrt{6} \frac{h}{2\pi}$

Sol. Answer (1)

$$\text{Angular momentum} = l(l+1) \frac{h}{2\pi}$$

16. The total number of atomic orbitals in fourth energy level of an atom is

[AIPMT (Prelims)-2011]

- (1) 4 (2) 8 (3) 16 (4) 32

Sol. Answer (3)

$$\boxed{\text{Number of orbitals} = x^2} \quad n = \text{number of orbit}$$

$$= 4^2 = 16$$

17. The energies E_1 and E_2 of two radiations are 25 eV and 50 eV respectively. The relation between their wavelengths i.e. λ_1 and λ_2 will be [AIPMT (Prelims)-2011]

(1) $\lambda_1 = \frac{1}{2}\lambda_2$ (2) $\lambda_1 = \lambda_2$ (3) $\lambda_1 = 2\lambda_2$ (4) $\lambda_1 = 4\lambda_2$

Sol. Answer (3)

$$\boxed{\lambda_1 = \frac{hc}{E_1}} \quad \lambda_2 = \frac{hc}{E_2} \quad E_1 = 25 \text{ eV} \quad E_2 = 50 \text{ eV}$$

$$\lambda_1 = \frac{hc}{25} \quad \dots\dots (1)$$

$$\lambda_2 = \frac{hc}{50} \quad \dots\dots (2)$$

$$\frac{\lambda_1}{\lambda_2} = \frac{\frac{hc}{25}}{\frac{hc}{50}} = 2 \quad \therefore \frac{\lambda_1}{\lambda_2} = 2 \quad \boxed{\lambda_1 = 2\lambda_2}$$

18. If $n = 6$, the correct sequence of filling of electrons will be

[AIPMT (Prelims)-2011]

- (1) $ns \rightarrow np \rightarrow (n-1)d \rightarrow (n-2)f$ (2) $ns \rightarrow (n-2)f \rightarrow (n-1)d \rightarrow np$
 (3) $ns \rightarrow (n-1)d \rightarrow (n-2)f \rightarrow np$ (4) $ns \rightarrow (n-2)f \rightarrow np \rightarrow (n-1)d$

Sol. Answer (2)

Putting the value of n and calculating the $(n + l)$ value

6s	<	4f	<	5d	<	6p
(6+0)		(4+3)		(5+2)		(6+1)
6		7		7		7
(lower energy)						(high energy)

→

19. According to the Bohr Theory, which of the following transitions in the hydrogen atom will give rise to the least energetic photon? [AIPMT (Mains)-2011]

- (1) $n = 6$ to $n = 5$ (2) $n = 5$ to $n = 3$ (3) $n = 6$ to $n = 1$ (4) $n = 5$ to $n = 4$

Sol. Answer (1)

$$\text{Because } (E_2 - E_1) > (E_3 - E_2) > (E_4 - E_3) > (E_5 - E_4) > (E_6 - E_5)$$

As the difference is of one energy levels

$$\therefore (E_6 - E_5) \text{ have less energy}$$

{Alternatively value of ΔE [difference between two successive energy level decreases] as the distance from the nucleus increases.}

20. A 0.66 kg ball is moving with a speed of 100 m/s. The associated wavelength will be

$$(h = 6.6 \times 10^{-34} \text{ Js})$$

[AIPMT (Mains)-2010]

- (1) $6.6 \times 10^{-32} \text{ m}$ (2) $6.6 \times 10^{-34} \text{ m}$ (3) $1.0 \times 10^{-35} \text{ m}$ (4) $1.0 \times 10^{-32} \text{ m}$

Sol. Answer (3)

$$\lambda = \frac{h}{mv} = \frac{6.6 \times 10^{-34} \text{ kgm}^2\text{s}^{-2} - \text{s}}{0.66 \text{ kg} \times 100 \text{ m/s}} = \boxed{1.0 \times 10^{-35} \text{ m}}$$

21. The energy absorbed by each molecule (A_2) of a substance is $4.4 \times 10^{-19} \text{ J}$ and bond energy per molecule is $4.0 \times 10^{-19} \text{ J}$. The kinetic energy of the molecule per atom will be [AIPMT (Prelims)-2009]
- (1) $2.2 \times 10^{-19} \text{ J}$ (2) $2.0 \times 10^{-19} \text{ J}$ (3) $4.0 \times 10^{-20} \text{ J}$ (4) $2.0 \times 10^{-20} \text{ J}$

Sol. Answer (4)

KE = Energy observed by molecule – Energy required to break one bond

$$KE = \frac{4.4 \times 10^{-19} \text{ J} - 4.0 \times 10^{-19} \text{ J}}{2}$$

$$KE \text{ per atom} = \left[\frac{0.4 \times 10^{-19}}{2} \right] = [0.2 \times 10^{-19} \text{ J}] = [2 \times 10^{-20} \text{ J}]$$

22. Which one of the elements with the following outer orbital configurations may exhibit the largest number of oxidation states? [AIPMT (Prelims)-2009]
- (1) $3d^5 4s^1$ (2) $3d^5 4s^2$ (3) $3d^2 4s^2$ (4) $3d^3 4s^2$

Sol. Answer (2)

23. Maximum number of electrons in a subshell of an atom is determined by the following [AIPMT (Prelims)-2009]
- (1) $2\ell + 1$ (2) $4\ell - 2$ (3) $2n^2$ (4) $4\ell + 2$

Sol. Answer (4)

24. Which of the following is not permissible arrangement of electrons in an atom? [AIPMT (Prelims)-2009]

- (1) $n = 5, \ell = 3, m = 0, s = +\frac{1}{2}$ (2) $n = 3, \ell = 2, m = -3, s = -\frac{1}{2}$
- (3) $n = 3, \ell = 2, m = -2, s = -\frac{1}{2}$ (4) $n = 4, \ell = 0, m = 0, s = +\frac{1}{2}$

Sol. Answer (2)

$$n = 3 \quad \ell = 2 \quad m = -3 \quad s = -\frac{1}{2}$$

Value of m (orbital) depends upon ℓ i.e., it cannot be more than ' ℓ '. Therefore is wrong.

25. A p-n photodiode is made of a material with a band gap of 2.0 eV. The minimum frequency of the radiation that can be absorbed by the material is nearly [AIPMT (Prelims)-2008]
- (1) $20 \times 10^{14} \text{ Hz}$ (2) $10 \times 10^{14} \text{ Hz}$ (3) $5 \times 10^{14} \text{ Hz}$ (4) $1 \times 10^{14} \text{ Hz}$

Sol. Answer (3)

26. If uncertainty in position and momentum are equal, then uncertainty in velocity is [AIPMT (Prelims)-2008]

- (1) $\sqrt{\frac{h}{\pi}}$ (2) $\frac{1}{2m} \sqrt{\frac{h}{\pi}}$ (3) $\sqrt{\frac{h}{2\pi}}$ (4) $\frac{1}{m} \sqrt{\frac{h}{\pi}}$

Sol. Answer (2)

$$\Delta x \times \Delta p = \frac{h}{2\pi}$$

27. The measurement of the electron position is associated with an uncertainty in momentum, which is equal to $1 \times 10^{-18} \text{ g cm s}^{-1}$. The uncertainty in electron velocity is, (Mass of an electron is $9 \times 10^{-28} \text{ g}$) [AIPMT (Prelims)-2008]

- (1) $1 \times 10^{11} \text{ cm s}^{-1}$ (2) $1 \times 10^9 \text{ cm s}^{-1}$ (3) $1 \times 10^6 \text{ cm s}^{-1}$ (4) $1 \times 10^5 \text{ cm s}^{-1}$

Sol. Answer (2)

28. Consider the following sets of quantum numbers

	n	ℓ	m	s
(a)	3	0	0	$+\frac{1}{2}$
(b)	2	2	1	$+\frac{1}{2}$
(c)	4	3	-2	$-\frac{1}{2}$
(d)	1	0	-1	$-\frac{1}{2}$
(e)	3	2	3	$+\frac{1}{2}$

Which of the following sets of quantum number is not possible?

[AIPMT (Prelims)-2007]

- (1) a and c (2) b, c and d (3) a, b, c and d (4) b, d and e

Sol. Answer (4)

29. With which of the following configuration an atom has the lowest ionization enthalpy?

[AIPMT (Prelims)-2007]

- (1) $1s^2 2s^2 2p^6$ (2) $1s^2 2s^2 2p^5$ (3) $1s^2 2s^2 2p^3$ (4) $1s^2 2s^2 2p^5 3s^1$

Sol. Answer (4)

30. Given : The mass of electron is 9.11×10^{-31} kg. Planck's constant is 6.626×10^{-34} Js, the uncertainty involved in the measurement of velocity within a distance of $0.1 \text{ \AA}$ is

[AIPMT (Prelims)-2006]

- (1) $5.79 \times 10^6 \text{ ms}^{-1}$ (2) $5.79 \times 10^7 \text{ ms}^{-1}$
 (3) $5.79 \times 10^8 \text{ ms}^{-1}$ (4) $5.79 \times 10^5 \text{ ms}^{-1}$

Sol. Answer (1)

$$\Delta x \times m \Delta v = \frac{h}{4\pi} \quad \Delta v = \frac{h}{\Delta x \times m \times 4 \times \pi}$$

$$\Delta v = \frac{6.6 \times 10^{-34} \text{ J-s}}{0.1 \times 10^{-10} \text{ m} \times 9.1 \times 10^{-31} \text{ kg} \times 4 \times 3.14} = 5.799 \times 10^6 \text{ m/s}$$

31. The orientation of an atomic orbital is governed by

[AIPMT (Prelims)-2006]

- (1) Azimuthal quantum number (2) Spin quantum number
 (3) Magnetic quantum number (4) Principal quantum number

Sol. Answer (3)

32. The energy of second Bohr orbit of the hydrogen atom is -328 kJ mol^{-1} ; hence the energy of fourth Bohr orbit would be

[AIPMT (Prelims)-2005]

- (1) -41 kJ mol^{-1} (2) $-1312 \text{ kJ mol}^{-1}$ (3) -164 kJ mol^{-1} (4) -82 kJ mol^{-1}

Sol. Answer (4)

$$E_n = -\frac{1312}{n^2} \text{ kJ mol}^{-1} \text{ for hydrogen}$$

33. Uncertainty in position of an electron (mass = 9.1×10^{-28} g) moving with a velocity of 3×10^4 cm/s accurate upto 0.001% will be (Use $h/(4\pi)$ in uncertainty expression where $h = 6.626 \times 10^{-27}$ erg-s)

- (1) 5.76 cm (2) 7.68 cm (3) 1.93 cm (4) 3.84 cm

Sol. Answer (3)

$$\Delta v = \frac{3 \times 10^4 \times 0.001}{10^3 \times 100} \text{ cm/second}$$

$$\Delta x = \frac{h}{4\pi m \Delta v} = \frac{6.6 \times 10^{-34} \text{ J-s}}{4 \times 3.14 \times 9.1 \times 10^{-28} \times \frac{0.001}{100} \times 3 \times 10^4} = \boxed{1.93 \text{ cm}}$$

34. The radius of hydrogen atom in the ground state is 0.53 Å. The radius of Li^{2+} ion (atomic number = 3) in a similar state is

- (1) 0.53 Å (2) 1.06 Å (3) 0.17 Å (4) 0.265 Å

Sol. Answer (3)

$$\boxed{r_n = \frac{r_0 \times n^2}{z}} = \frac{0.53 \times (3)^2}{3} = (0.53 \times 3) \text{ Å}$$

$$1.59 \text{ Å} \approx 1.7 \text{ Å}$$

$$n = 3 \text{ orbit}$$

$$z = 3 \text{ Li}^{2+}$$

35. In a Bohr's model of an atom, when an electron jumps from $n = 1$ to $n = 3$, how much energy will be emitted or absorbed?

- (1) 2.389×10^{-12} ergs (2) 0.239×10^{-10} ergs (3) 2.15×10^{-11} ergs (4) 0.1936×10^{-10} ergs

Sol. Answer (4)

$$\text{Energy of electron when } n = 1 \quad E_1 = -\frac{1312}{(1)^2} \text{ kJ/mol}$$

$$\text{Energy of electron when } n = 3 \quad E_3 = -\frac{1312}{(3)^2} = -\frac{1312}{9} \text{ kJ/mol}$$

$$\boxed{\Delta E = E_3 - E_1}$$

$$= \frac{-1312}{9} - \left(\frac{-1312}{1} \right) = +1166 \text{ kJ}$$

$$= 1166 \times 10^3 \text{ J}$$

$$= 1166 \times 10^{+10} \text{ erg}$$

Alternatively

$$\Delta E = \frac{hc}{\lambda} = hc \times R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$

$$= 3.0 \times 10^8 \times 6.6 \times 10^{-34} \times 8.314 \left[\frac{1}{1^2} - \frac{1}{3^2} \right] \text{ J}$$


$$= 3.0 \times 10^8 \times 6.6 \times 10^{-34} \times 8.314 \left[\frac{8}{9} \right] \text{ J}$$

36. The electronic configuration of gadolinium (Atomic No. 64) is

- (1) $[\text{Xe}]4f^3 5d^5 6s^2$ (2) $[\text{Xe}]4f^6 5d^2 6d^2$ (3) $[\text{Xe}]4f^8 5d^9 6s^2$ (4) $[\text{Xe}] 4f^7 5d^1 6s^2$

Sol. Answer (4)

Gd have exceptional configuration e^- will enter in $5d$ because $4f$ have 7 electrons and have half filled stability



37. The ion that is isoelectronic with CO is

- (1) CN^- (2) N_2^+ (3) O_2^- (4) N_2^-

Sol. Answer (1)

Isoelectronic means same number of electrons

CO = Number of electrons = 14

$\text{CN}^- = 6 + 7 + 1 = 14$

38. The Bohr orbit radius for the hydrogen atom ($n = 1$) is approximately 0.530 Å. The radius for the first excited state ($n = 2$) orbit is (in Å)

- (1) 4.77 (2) 1.06 (3) 0.13 (4) 2.12

Sol. Answer (4)

$$\boxed{r_n = \frac{r_0 \times n^2}{z}}$$

n = Number of orbit, z = charge on nucleus

$$\begin{aligned} \text{1st excited state for H} &= \frac{n=2}{z=1} \\ &= \frac{0.53 \times (2)^2}{1} \\ &= 0.53 \times 4 = 2.12 \text{ Å} \end{aligned}$$

39. The position of both, an electron and a helium atom is known within 1.0 nm. Further the momentum of the electron is known within $5.0 \times 10^{-26} \text{ kg ms}^{-1}$. The minimum uncertainty in the measurement of the momentum of the helium atom is

- (1) $8.0 \times 10^{-26} \text{ kg ms}^{-1}$ (2) 80 kg ms^{-1}
(3) 50 kg ms^{-1} (4) $5.0 \times 10^{-26} \text{ kg ms}^{-1}$

Sol. Answer (4)

$$\boxed{\Delta x_{\text{electron}} \times \Delta P_{\text{electron}} = \frac{h}{4\pi}}$$

$$\Delta x_{\text{electron}} = \frac{h}{4\pi \Delta P_{\text{electron}}}$$

$$\Delta x_{\text{electron}} = \Delta x_{\text{He}} = 1.0 \text{ nm}$$

$$\Delta x_{\text{He}} \times \Delta P_{\text{He}} = \frac{h}{4\pi}$$

$$\Delta x_{\text{He}} = \frac{h}{4\pi \Delta P_{\text{He}}}$$

$$\therefore \frac{h}{4\pi \Delta P_{\text{electron}}} = \frac{h}{4\pi \Delta P_{\text{He}}}$$

$$\Delta P_{\text{He}} = 5.0 \times 10^{-26} \text{ kg ms}^{-1}$$

40. Which of the following configuration is correct for iron?

(1) $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 d^7$

(2) $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 d^5$

(3) $1s^2 2s^2 2p^6 3s^2 3p^6 d^5$

(4) $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 d^6$

Sol. Answer (4)

$$\text{Fe} = 26 = 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^2, 3d^6$$

41. Which of the following has maximum number of unpaired *d*-electrons?

(1) N^{3+}

(2) Fe^{2+}

(3) Zn^+


(4) Cu^+

Sol. Answer (2)

$$\text{N}^{3+} = 4 = 1s^2, 2s^2$$

Zero unpaired

$$\text{Fe}^{2+} = 24 = 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 3d^6$$



$$\text{Zn}^0 = 29 = 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^1, 3d^{10}$$

One unpaired

$$\text{Cu}^0 = 28 = 1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^0, 3d^{10}$$

Zero unpaired

42. Who modified Bohr's theory by introducing elliptical orbits for electron path?

(1) Rutherford


(2) Thomson

(3) Hund

(4) Sommerfeld

Sol. Answer (4)

$$\frac{n}{k} = \frac{\text{length of major axis}}{\text{length of minor axis}}$$



$$\text{Number of elliptical orbit in shell} = (n - 1)$$

43. The de Broglie wavelength of a particle with mass 1 g and velocity 100 m/s is

(1) $6.63 \times 10^{-35} \text{ m}$

(2) $6.63 \times 10^{-34} \text{ m}$

(3) $6.63 \times 10^{-33} \text{ m}$

(4) $6.65 \times 10^{-35} \text{ m}$

Sol. Answer (3)

$$\lambda = \frac{h}{mv}$$

$$m = 1 \text{ g} = 0.001 \text{ kg}$$

$$v = 100 \text{ m/s}$$

$$\lambda = \frac{6.6 \times 10^{-34} \text{ kg m}^2 \text{ s}^{-2}}{0.001 \text{ kg} \times 100 \text{ m/s}}$$

$$\lambda = 6.63 \times 10^{-33} \text{ m}$$

44. The following quantum numbers are possible for how many orbitals : $n = 3, l = 2, m = +2$?

(1) 1

(2) 2

(3) 3

(4) 4

Sol. Answer (1)

As the value of $m = +2$

i.e. one value

Therefore one orbital is represented.

45. The frequency of radiation emitted when the electron falls from $n = 4$ to $n = 1$ in a hydrogen atom will be (Given ionization energy of H = 2.18×10^{-18} J atom $^{-1}$ and $h = 6.625 \times 10^{-34}$ Js)

(1) $1.54 \times 10^{15} \text{ s}^{-1}$ (2) $1.03 \times 10^{15} \text{ s}^{-1}$ (3) $3.08 \times 10^{15} \text{ s}^{-1}$ (4) $2.00 \times 10^{15} \text{ s}^{-1}$

Sol. Answer (3)

$$v = \frac{c}{\lambda} = c \times R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] Z^2$$

$$v = 3.0 \times 10^{10} \times 109678 \left[\frac{1}{1^2} - \frac{1}{4^2} \right] \times (1)^2$$

$$v = 3 \times \frac{10}{10^{10}} \times 109678 \text{ cm}^{-1} \left[\frac{15}{16} \right] = 3.09 \times 10^{15} \text{ s}^{-1}$$

Alternatively

$$I.E = E_{\infty} - E_1$$

$$2.18 \times 10^{-18} = E_{\infty} - E_1$$

$$\boxed{E_1 = -2.18 \times 10^{-18} \text{ J}}$$

$$E_4 = -\frac{2.18 \times 10^{-18}}{(4)^2} = -\frac{2.18 \times 10^{-18}}{16} = -0.136 \times 10^{-18} \text{ J}$$

$$\Delta E = E_4 - E_1 [-0.136 - (-2.18)] \times 10^{-18}$$

$$= 2.04 \times 10^{-18} \quad \lambda = \frac{hc}{\Delta E} \quad \dots (1)$$

$$v = \frac{c}{\lambda} \quad \dots (2)$$

Put (1) in (2)

$$v = \frac{c}{hc} \times \Delta E = \frac{\Delta E}{h}$$

$$v = \frac{2.04 \times 10^{-18}}{6.6 \times 10^{-34}} = 0.309 \times 10^{+16} = 3.09 \times 10^{15} \text{ s}^{-1}$$

46. Which one of the following ions has electronic configuration $[\text{Ar}]3d^6$?

- (1) Co^{3+} (2) Ni^{3+}
 (3) Mn^{3+} (4) Fe^{3+}

(At. nos. Mn = 25, Fe = 26, Co = 27, Ni = 28)

Sol. Answer (1)

Electronic configuration $[\text{Ar}] 3d^6$ represents 24 electrons

i.e. $\boxed{\text{Co}^{3+} = 24 e^-}$

$$\text{Ni}^{3+} = 28 - 3 = 25 e^-$$

$$\text{Mn}^{3+} = 25 - 3 = 22 e^-$$

$$\text{Fe}^{3+} = 26 - 3 = 23 e^-$$

47. Which of the following is not among shortcomings of Bohr's model?

- (1) Bohr theory could not account for the fine lines in the atomic spectrum
 (2) Bohr theory was unable to account for the splitting of the spectral lines in the presence of magnetic field
 (3) Bohr theory failed for He atom
 (4) It did not give information about energy level

Sol. Answer (4)

Bohr's model explain the energy level i.e. Energy of electron in each orbital is quantized.

$$\boxed{E_n = \frac{-1312}{x^2} z^2 \text{ kJ/mol}}$$

48. Number of spectral lines falling in Balmer series when electrons are de-excited from n^{th} shell will be given as

- (1) $(n - 2)$ in UV (2) $(n - 2)$ in visible region
 (3) $(n - 3)$ in near IR (4) $(n - 3)$ in far IR

Sol. Answer (2)

49. The ratio of the energy required to remove an electron from the first three Bohr's orbits of hydrogen is

- (1) 3 : 2 : 1 (2) 9 : 4 : 1
 (3) 36 : 9 : 4 (4) 1 : 4 : 9

Sol. Answer (3)

$$E_n = \frac{-1312}{x^2} = \frac{x}{x^2}$$

$$E_1 = +\frac{x}{1^2} \quad E_2 = \frac{x}{2^2} \quad E_3 = \frac{x}{3^2}$$

$$\frac{1}{1} : \frac{1}{4} : \frac{1}{9} = \frac{36 : 9 : 4}{36} = \boxed{36 : 9 : 4}$$

SECTION - C

Assertion - Reason Type Questions

1. A : Orbital angular momentum of (1s, 2s, 3s etc.) all s electrons is same.

R : Orbital angular momentum depends on orientation of orbitals.

Sol. Answer (3)

2. A : Energy of electron is taken negative.

R : Energy of electron at infinity is zero.

Sol. Answer (1)

3. A : Bohr's orbits are also called stationary states.

R : Electrons are stationary in an orbit.

Sol. Answer (3)

4. A : K.E. of two subatomic particles, having same de-Broglie's wavelength is same.

R : de-Broglie's wavelength is directly related to mass of subatomic particles.

Sol. Answer (4)

5. A : Electronic energy for hydrogen atom of different orbitals follow the sequence :
 $1s < 2s = 2p < 3s = 3p = 3d$.

R : Electronic energy for hydrogen atom depends only on n and is independent of ' l ' & ' m ' values.

Sol. Answer (1)

6. A : Wavelength for first line of any series in hydrogen spectrum is biggest among all other lines of the same series.

R : Wavelength of spectral line for an electronic transition is inversely related to difference in the energy levels involved in the transition.

Sol. Answer (1)

7. A : Zn(II) salts are diamagnetic.

R : Zn^{2+} ion has one unpaired electron.

Sol. Answer (3)

8. A : In third energy level there is no f -subshell.

R : For $n = 3$, the possible values of l are 0, 1, 2 and for f -subshell $l = 3$.

Sol. Answer (1)

9. A : The charge to mass ratio of the particles in anode rays depends on nature of gas taken in the discharge tube.

R : The particles of anode rays carry positive charge.

Sol. Answer (2)

10. A : Angular momentum of an electron in an atom is quantized.

R : In an atom only those orbits are permitted in which angular momentum of the electron is a natural number multiple of $\frac{h}{2\pi}$.

Sol. Answer (1)

11. A : The radius of second orbit of He^+ is equal to that of first orbit of hydrogen.
R : The radius of an orbit in hydrogen like species is directly proportional to n and inversely proportional to Z .

Sol. Answer (4)

12. A : The orbitals having equal energy are known as degenerate orbitals.
R : The three $2p$ orbitals are degenerate in the presence of external magnetic field.

Sol. Answer (3)

13. A : In a multielectron atom, the electrons in different sub-shells have different energies.
R : Energy of an orbital depends upon $n + l$ value.

Sol. Answer (1)

14. A : Isotopes of an element have almost similar chemical properties.
R : Isotopes have same electronic configuration.

Sol. Answer (1)

15. A : The number of angular nodes in $3d_{z^2}$ is zero.
R : Number of angular nodes of atomic orbitals is equal to value of l .

Sol. Answer (2)

