

SOME BASIC CONCEPTS OF CHEMISTRY

1

MCQs with One Correct Answer

- Which of the following statements is correct about the reaction given below ?
$$4\text{Fe(s)} + 3\text{O}_2\text{(g)} \longrightarrow 2\text{Fe}_2\text{O}_3\text{(g)}$$
 - Total mass of iron and oxygen in reactants = total mass of iron and oxygen in product therefore, it follows law of conservation of mass.
 - Total mass of reactants = total mass of product; therefore, law of multiple proportions is followed.
 - Amount of Fe_2O_3 can be increased by reducing the amount of any one of the reactants (iron or oxygen).
 - Amount of Fe_2O_3 produced will decrease if the amount of any one of the reactants (iron or oxygen) is taken in excess.
- Which of the following statements indicates that law of multiple proportion is being followed.
 - Sample of carbon dioxide taken from any source will always have carbon and oxygen in the ratio 1 : 2.
 - Carbon forms two oxides namely CO_2 and CO , where masses of oxygen which combine with fixed mass of carbon are in the simple ratio 2 : 1.
 - When magnesium burns in oxygen, the amount of magnesium taken for the reaction is equal to the amount of magnesium in magnesium oxide formed.
 - At constant temperature and pressure, 200 mL of hydrogen will combine with 100 mL oxygen to produce 200 mL of water vapour.
- The number of water molecules in 250 mL of water is closest to [Given, density of water is 1.0 g mL^{-1} ; Avogadro's number = 6.023×10^{23}]
 - 83.6×10^{23}
 - 13.9×10^{23}
 - 1.5×10^{23}
 - 33.6×10^{23}
- A metal oxide has the formula Z_2O_3 . It can be reduced by hydrogen to give free metal and water. 0.1596 g of the metal oxide requires 6 mg of hydrogen for complete reduction. The atomic weight of the metal is
 - 27.9
 - 159.6
 - 79.8
 - 55.8
- 4 g of a hydrated crystal of formula $\text{A} \cdot x\text{H}_2\text{O}$ has 0.8 g of water. If the molar mass of the anhydrous crystal (A) is 144 g mol^{-1} , The value of x is
 - 4
 - 1
 - 2
 - 3
- A solution of 20.2 g of 1,2-dibromopropane in MeOH upon heating with excess Zn produces 3.58 g of an unsaturated compound X. The yield (%) of X is closest to [Atomic weight of Br is 80]
 - 18
 - 85
 - 89
 - 30

7. 5.0 g of a certain element X forms 10.0 g of its oxide having the formula X_4O_6 . The atomic mass of X is
 (a) 12.0 amu (b) 24.0 amu
 (c) 30.0 amu (d) 32.0 amu
8. The maximum number of molecules are present in
 (a) 15 L of H_2 gas at STP
 (b) 5 L of N_2 gas at STP
 (c) 0.5 g of H_2 gas
 (d) 10 g of O_2 gas
9. Number of moles of MnO_4^- required to oxidize one mole of ferrous oxalate completely in acidic medium will be
 (a) 0.6 moles (b) 0.4 moles
 (c) 7.5 moles (d) 0.2 moles
10. A metallic chloride contain 47.22% metal. Calculate the equivalent weight of metal.
 (a) 39.68 (b) 31.76 (c) 36.35 (d) 33.46
11. The number of moles of water present in a spherical water droplet of radius 1.0 cm is [Given: density of water in the droplet = 1.0 g cm^{-3}]
 (a) $\frac{\pi}{18}$ (b) $\frac{2\pi}{27}$ (c) 24π (d) $\frac{2\pi}{9}$
12. If potassium chlorate is 80% pure, then 48 g of oxygen would be produced from (atomic mass of K = 39)
 (a) 153.12 g of $KClO_3$ (b) 122.5 g of $KClO_3$
 (c) 245 g of $KClO_3$ (d) 98 g of $KClO_3$
13. 10 g $CaCO_3$ were dissolved in 250 mL of 100 M HCl or the solution was boiled. What volume of 2M KOH would be required to equivalence point after boiling? Assume no change in volume during boiling.
 (a) 50 mL (b) 25 mL (c) 75 mL (d) 60 mL
14. The density of 3M solution of sodium chloride is 1.252 g mL^{-1} . The molality of the solution will be :
 (molar mass, $NaCl = 58.5 \text{ g mol}^{-1}$)
 (a) 260m (b) 2.18m (c) 2.79m (d) 3.00m
15. 2 g of a mixture of CO and CO_2 on reaction with excess I_2O_5 produced 2.54 g of I_2 . What would be the mass % of CO_2 in the original mixture?
 (a) 60 (b) 30 (c) 70 (d) 35
16. 3.0 g of oxalic acid $[(CO_2H)_2 \cdot 2H_2O]$ is dissolved in a solvent to prepare a 250 mL solution. The density of the solution is 1.9 g/mL . The molality and normality of the solution, respectively, are closest to
 (a) 0.10 and 0.38
 (b) 0.10 and 0.19
 (c) 0.05 and 0.19
 (d) 0.05 and 0.09
17. How many moles of P_4O_6 and P_4O_{10} will be produced by the combustion of 12.4 g of phosphorous (atomic mass 31) in 12.8 g of oxygen, leaving no P_4 or O_2 ?
 (a) 0.1 and 0.3 mol
 (b) 0.15 mol and 0.25 mol
 (c) 0.05 mol each
 (d) 0.1 mol each
18. 5 g sample contain only Na_2CO_3 and Na_2SO_4 . This sample is dissolved and the volume made up to 250 mL. 25 mL of this solution neutralizes 20 mL of 0.1 M H_2SO_4 . Calculate the % of Na_2SO_4 in the sample.
 (a) 42.4 (b) 57.6 (c) 36.2 (d) 0.576
19. 1 mole of equimolar mixture of ferric oxalate and ferrous oxalate requires x mole of $KMnO_4$ in acidic medium for complete oxidation. x is:
 (a) 0.5 mole (b) 0.9 mole
 (c) 1.2 mole (d) 4.5 mole
20. Two solutions of a substance (non electrolyte) are mixed in the following manner. 480 mL of 1.5 M first solution + 520 mL of 1.2 M second solution. What is the molarity of the final mixture?
 (a) 2.70 M (b) 1.344 M
 (c) 1.50 M (d) 1.20 M

Numeric Value Answer

21. A mixture of $HCOOH$ and $H_2C_2O_4$ is heated with concentrated H_2SO_4 . The gas produced is collected and on treating with KOH solution, the volume of gas decreases by one-sixth. Calculate the molar ratio of the two acids ($HCOOH : H_2C_2O_4$) in the original mixture.
22. One gram of a metallic chloride was found to contain 0.835 g of chlorine. Its vapour density is 85.5. If its molecular formula is M_xCl_y , then what is value of $(x + y)$?

23. 0.7875 g of crystalline barium hydroxide is dissolved in water. For the neutralization of this solution 20 mL of $N/4$ HNO_3 is required. How many moles of water of crystallization are present in one mole of this base? (Given : Atomic mass Ba = 137, O = 16, N = 14, H = 1)
24. A mixture contains 1.0 mole each of NaOH , Na_2CO_3 and NaHCO_3 . When half of mixture is titrated with HCl , it required x mole of HCl in presence of phenolphthalein. In another experiment, half of mixture required y mole of same HCl in presence of methyl orange. Find the value of $(x + y)$.
25. A 0.276 g impure sample of copper ore is dissolved and Cu^{2+} is titrated with KI solution. I_2 liberated required 40 mL of 0.1M $\text{Na}_2\text{S}_2\text{O}_3$ solution for titration. What is the % of impurities in the ore?
26. Density of a sulphuric acid solution is 1.225g/mL. and it is 40% H_2SO_4 by weight. Determine molarity of this solution.
27. A 16.24 mL sample of vinegar of density 1.06 g/mL required 48.24 mL of 0.36 N solution of a standard alkali. Determine percentage weight of acetic acid in vinegar.
28. How much volume of sulphur dioxide at STP will be obtained by completely burning 10 g of pure sulphur?
29. How much volume of 4.0 M HNO_3 is required to prepare 60 mL of 0.2 M HNO_3 from a stock solution of 4.0 M HNO_3 ?
30. The minimum number of moles of O_2 required for complete combustion of 1 mole of propane and 2 moles of butane is _____.

ANSWER KEY

| | | | | | | | | | | | | | | | | | | | |
|---|-----|---|-----|---|-----|----|-----|----|-----|----|-----|----|-----|----|-----|----|-----|----|------|
| 1 | (a) | 4 | (d) | 7 | (b) | 10 | (b) | 13 | (b) | 16 | (c) | 19 | (b) | 22 | (5) | 25 | (8) | 28 | (7) |
| 2 | (b) | 5 | (c) | 8 | (a) | 11 | (b) | 14 | (c) | 17 | (c) | 20 | (b) | 23 | (8) | 26 | (5) | 29 | (3) |
| 3 | (a) | 6 | (b) | 9 | (b) | 12 | (a) | 15 | (b) | 18 | (b) | 21 | (4) | 24 | (3) | 27 | (6) | 30 | (18) |