

Chapter 16. Solutions

- If molality of the dilute solution is doubled, the value of molal depression constant (K_f) will be
 (a) halved (b) tripled
 (c) unchanged (d) doubled.
 (NEET 2017)
- Which of the following is dependent on temperature?
 (a) Molarity (b) Mole fraction
 (c) Weight percentage (d) Molality
 (NEET 2017)
- The van't Hoff factor (i) for a dilute aqueous solution of the strong electrolyte barium hydroxide is
 (a) 0 (b) 1
 (c) 2 (d) 3
 (NEET-II 2016)
- Which one of the following is incorrect for ideal solution?
 (a) $\Delta H_{\text{mix}} = 0$ (b) $\Delta U_{\text{mix}} = 0$
 (c) $\Delta P = P_{\text{obs}} - P_{\text{calculated by Raoult's law}} = 0$
 (d) $\Delta G_{\text{mix}} = 0$ (NEET-II 2016)
- Which of the following statements about the composition of the vapour over an ideal 1 : 1 molar mixture of benzene and toluene is correct? Assume that the temperature is constant at 25°C. (Given, vapour pressure data at 25°C, benzene = 12.8 kPa, toluene = 3.85 kPa)
 (a) The vapour will contain equal amounts of benzene and toluene.
 (b) Not enough information is given to make a prediction.
 (c) The vapour will contain a higher percentage of benzene.
 (d) The vapour will contain a higher percentage of toluene. (NEET-I 2016)
- At 100°C the vapour pressure of a solution of 6.5 g of a solute in 100 g water is 732 mm. If $K_b = 0.52$, the boiling point of this solution will be
 (a) 102°C (b) 103°C
 (c) 101°C (d) 100°C
 (NEET-I 2016)
- What is the mole fraction of the solute in a 1.00 m aqueous solution?
 (a) 1.770 (b) 0.0354
 (c) 0.0177 (d) 0.177
 (2015, 2011)
- The boiling point of 0.2 mol kg⁻¹ solution of X in water is greater than equimolar solution of Y in water. Which one of the following statements is true in this case?
 (a) Molecular mass of X is less than the molecular mass of Y .
 (b) Y is undergoing dissociation in water while X undergoes no change.
 (c) X is undergoing dissociation in water.
 (d) Molecular mass of X is greater than the molecular mass of Y . (2015, Cancelled)
- Which one of the following electrolytes has the same value of van't Hoff factor (i) as that of $\text{Al}_2(\text{SO}_4)_3$ (if all are 100% ionised)?
 (a) $\text{Al}(\text{NO}_3)_3$ (b) $\text{K}_4[\text{Fe}(\text{CN})_6]$
 (c) K_2SO_4 (d) $\text{K}_3[\text{Fe}(\text{CN})_6]$
 (2015, Cancelled)
- Which of them is not equal to zero for an ideal solution?
 (a) ΔV_{mix}
 (b) $\Delta P = P_{\text{observed}} - P_{\text{Raoult}}$
 (c) ΔH_{mix} (d) ΔS_{mix}
 (2015, Cancelled)
- Of the following 0.10 m aqueous solutions, which one will exhibit the largest freezing point depression?
 (a) KCl (b) $\text{C}_6\text{H}_{12}\text{O}_6$
 (c) $\text{Al}_2(\text{SO}_4)_3$ (d) K_2SO_4
 (2014)

12. How many grams of concentrated nitric acid solution should be used to prepare 250 mL of 2.0 M HNO_3 ? The concentrated acid is 70% HNO_3 .
 (a) 70.0 g conc. HNO_3
 (b) 54.0 g conc. HNO_3
 (c) 45.0 g conc. HNO_3
 (d) 90.0 g conc. HNO_3 (NEET 2013)
13. Which condition is not satisfied by an ideal solution?
 (a) $\Delta_{\text{mix}} V = 0$ (b) $\Delta_{\text{mix}} S = 0$
 (c) Obeys Raoult's Law
 (d) $\Delta_{\text{mix}} H = 0$ (Karnataka NEET 2013)
14. p_A and p_B are the vapour pressure of pure liquid components, A and B, respectively of an ideal binary solution. If x_A represents the mole fraction of component A, the total pressure of the solution will be
 (a) $p_A + x_A(p_B - p_A)$ (b) $p_A + x_A(p_A - p_B)$
 (c) $p_B + x_A(p_B - p_A)$ (d) $p_B + x_A(p_A - p_B)$ (2012)
15. Vapour pressure of chloroform (CHCl_3) and dichloromethane (CH_2Cl_2) at 25°C are 200 mm Hg and 41.5 mm Hg respectively. Vapour pressure of the solution obtained by mixing 25.5 g of CHCl_3 and 40 g of CH_2Cl_2 at the same temperature will be
 (Molecular mass of $\text{CHCl}_3 = 119.5$ u and molecular mass of $\text{CH}_2\text{Cl}_2 = 85$ u)
 (a) 173.9 mm Hg (b) 615.0 mm Hg
 (c) 347.9 mm Hg (d) 285.5 mm Hg (Mains 2012)
16. The freezing point depression constant for water is $-1.86^\circ\text{C m}^{-1}$. If 5.00 g Na_2SO_4 is dissolved in 45.0 g H_2O , the freezing point is changed by -3.82°C . Calculate the van't Hoff factor for Na_2SO_4 .
 (a) 2.05 (b) 2.63
 (c) 3.11 (d) 0.381 (2011)
17. The van't Hoff factor i for a compound which undergoes dissociation in one solvent and association in other solvent is respectively
 (a) less than one and greater than one
 (b) less than one and less than one
 (c) greater than one and less than one
 (d) greater than one and greater than one. (2011)
18. A 0.1 molal aqueous solution of a weak acid is 30% ionized. If K_f for water is 1.86°C/m , the freezing point of the solution will be
 (a) -0.18°C (b) -0.54°C
 (c) -0.36°C (d) -0.24°C (Mains 2011)
19. 200 mL of an aqueous solution of a protein contains its 1.26 g. The osmotic pressure of this solution at 300 K is found to be 2.57×10^{-3} bar. The molar mass of protein will be
 ($R = 0.083 \text{ L bar mol}^{-1} \text{ K}^{-1}$)
 (a) 51022 g mol^{-1} (b) 122044 g mol^{-1}
 (c) 31011 g mol^{-1} (d) 61038 g mol^{-1} (Mains 2011)
20. An aqueous solution is 1.00 molal in KI. Which change will cause the vapour pressure of the solution to increase?
 (a) Addition of NaCl
 (b) Addition of Na_2SO_4
 (c) Addition of 1.00 molal KI
 (d) Addition of water (2010)
21. A solution of sucrose (molar mass = 342 g mol^{-1}) has been prepared by dissolving 68.5 g of sucrose in 1000 g of water. The freezing point of the solution obtained will be (K_f for water = $1.86 \text{ K kg mol}^{-1}$)
 (a) -0.372°C (b) -0.520°C
 (c) $+0.372^\circ\text{C}$ (d) -0.570°C (2010)
22. A 0.0020 m aqueous solution of an ionic compound $[\text{Co}(\text{NH}_3)_5(\text{NO}_2)]\text{Cl}$ freezes at -0.00732°C . Number of moles of ions which 1 mol of ionic compound produces on being dissolved in water will be
 ($K_f = -1.86^\circ\text{C/m}$)
 (a) 3 (b) 4
 (c) 1 (d) 2 (2009)
23. 0.5 molal aqueous solution of a weak acid (HX) is 20% ionised. If K_f for water is $1.86 \text{ K kg mol}^{-1}$, the lowering in freezing point of the solution is
 (a) 0.56 K (b) 1.12 K
 (c) -0.56 K (d) -1.12 K (2007)
24. Concentrated aqueous sulphuric acid is 98% H_2SO_4 by mass and has a density of 1.80 g mL^{-1} . Volume of acid required to make one litre of 0.1 M H_2SO_4 solution is
 (a) 16.65 mL (b) 22.20 mL
 (c) 5.55 mL (d) 11.10 mL (2007)

25. During osmosis, flow of water through a semipermeable membrane is
 (a) from solution having lower concentration only
 (b) from solution having higher concentration only
 (c) from both sides of semipermeable membrane with equal flow rates
 (d) from both sides of semipermeable membrane with unequal flow rates. (2006)
26. A solution of acetone in ethanol
 (a) obeys Raoult's law
 (b) shows a negative deviation from Raoult's law
 (c) shows a positive deviation from Raoult's law
 (d) behaves like a near ideal solution. (2006)
27. 1.00 g of a non-electrolyte solute (molar mass 250 g mol^{-1}) was dissolved in 51.2 g of benzene. If the freezing point depression constant, K_f of benzene is $5.12 \text{ K kg mol}^{-1}$, the freezing point of benzene will be lowered by
 (a) 0.2 K (b) 0.4 K
 (c) 0.3 K (d) 0.5 K (2006)
28. A solution containing 10 g per dm^3 of urea (molecular mass = 60 g mol^{-1}) is isotonic with a 5% solution of a nonvolatile solute is
 (a) 200 g mol^{-1} (b) 250 g mol^{-1}
 (c) 300 g mol^{-1} (d) 350 g mol^{-1} (2006)
29. The mole fraction of the solute in one molal aqueous solution is
 (a) 0.009 (b) 0.018
 (c) 0.027 (d) 0.036 (2005)
30. The vapour pressure of two liquids P and Q are 80 and 60 torr, respectively. The total vapour pressure of solution obtained by mixing 3 mole of P and 2 mol of Q would be
 (a) 72 torr (b) 140 torr
 (c) 68 torr (d) 20 torr (2005)
31. A solution has a 1 : 4 mole ratio of pentane to hexane. The vapour pressures of the pure hydrocarbons at 20°C are 440 mm Hg for pentane and 120 mm Hg for hexane. The mole fraction of pentane in the vapour phase would be
 (a) 0.200 (b) 0.549
 (c) 0.786 (d) 0.478 (2005)
32. A solution of urea (mol. mass 56 g mol^{-1}) boils at 100.18°C at the atmospheric pressure. If K_f and K_b for water are 1.86 and $0.512 \text{ K kg mol}^{-1}$ respectively, the above solution will freeze at
 (a) 0.654°C (b) -0.654°C
 (c) 6.54°C (d) -6.54°C (2005)
33. A solution containing components A and B follows Raoult's law
 (a) $A - B$ attraction force is greater than $A - A$ and $B - B$
 (b) $A - B$ attraction force is less than $A - A$ and $B - B$
 (c) $A - B$ attraction force remains same as $A - A$ and $B - B$
 (d) volume of solution is different from sum of volume of solute and solvent. (2002)
34. A solution contains non volatile solute of molecular mass M_2 . Which of the following can be used to calculate the molecular mass of solute in terms of osmotic pressure?
 (a) $M_2 = \left(\frac{m_2}{\pi}\right)VRT$
 (b) $M_2 = \left(\frac{m_2}{V}\right)\frac{RT}{\pi}$
 (c) $M_2 = \left(\frac{m_2}{V}\right)\pi RT$
 (d) $M_2 = \left(\frac{m_2}{V}\right)\frac{\pi}{RT}$
 (m_2 = mass of solute, V = volume of solution, π = osmotic pressure) (2002)
35. 2.5 litre of 1 M NaOH solution is mixed with another 3 litre of 0.5 M NaOH solution. Then find out molarity of resultant solution.
 (a) 0.80 M (b) 1.0 M
 (c) 0.73 M (d) 0.50 M (2002)
36. The beans are cooked earlier in pressure cooker because
 (a) boiling point increases with increasing pressure
 (b) boiling point decreases with increasing pressure
 (c) extra pressure of pressure cooker softens the beans
 (d) internal energy is not lost while cooking in pressure cooker. (2001)
37. Pure water can be obtained from sea water by
 (a) centrifugation (b) plasmolysis
 (c) reverse osmosis (d) sedimentation. (2001)

38. From the colligative properties of solution, which one is the best method for the determination of molecular weight of proteins and polymers?
(a) Osmotic pressure
(b) Lowering in vapour pressure
(c) Lowering in freezing point
(d) Elevation in boiling point (2000)
39. The vapour pressure of benzene at a certain temperature is 640 mm of Hg. A non-volatile and non-electrolyte solid, weighing 2.175 g is added to 39.08 of benzene. The vapour pressure of the solution is 600 mm of Hg. What is the molecular weight of solid substance?
(a) 69.5 (b) 59.6
(c) 49.50 (d) 79.8 (1999)
40. How many g of dibasic acid (mol. weight 200) should be present in 100 mL of the aqueous solution to give strength of 0.1 N?
(a) 10 g (b) 2 g
(c) 1 g (d) 20 g (1999)
41. If 0.15 g of a solute, dissolved in 15 g of solvent, is boiled at a temperature higher by 0.216°C , than that of the pure solvent. The molecular weight of the substance (Molal elevation constant for the solvent is 2.16°C) is
(a) 10.1 (b) 100
(c) 1.01 (d) 1000 (1999)
42. The vapour pressure of a solvent decreased by 10 mm of mercury when a non-volatile solute was added to the solvent. The mole fraction of the solute in the solution is 0.2. What should be the mole fraction of the solvent if the decrease in the vapour pressure is to be 20 mm of mercury?
(a) 0.4 (b) 0.6
(c) 0.8 (d) 0.2 (1998)
43. A 5% solution of cane sugar (mol. wt. = 342) is isotonic with 1% solution of a substance X. The molecular weight of X is
(a) 68.4 (b) 171.2
(c) 34.2 (d) 136.8 (1998)
44. Which of the following 0.10 m aqueous solution will have the lowest freezing point?
(a) KI (b) $\text{C}_{12}\text{H}_{22}\text{O}_{11}$
(c) $\text{Al}_2(\text{SO}_4)_3$ (d) $\text{C}_5\text{H}_{10}\text{O}_5$ (1997)
45. What is the molarity of H_2SO_4 solution, that has a density 1.84 g/cc at 35°C and contains 98% by weight?
(a) 18.4 M (b) 18 M
(c) 4.18 M (d) 8.14 M (1996)
46. The vapour pressure of CCl_4 at 25°C is 143 mm Hg. If 0.5 g of a non-volatile solute (mol. weight = 65) is dissolved in 100 mL CCl_4 , the vapour pressure of the solution will be
(a) 199.34 mm Hg (b) 143.99 mm Hg
(c) 141.43 mm Hg (d) 94.39 mm Hg. (1996)
47. The relationship between osmotic pressure at 273 K when 10 g glucose (p_1), 10 g urea (p_2), and 10 g sucrose (p_3) are dissolved in 250 mL of water is
(a) $p_2 > p_1 > p_3$
(b) $p_2 > p_3 > p_1$
(c) $p_1 > p_2 > p_3$
(d) $p_3 > p_1 > p_2$ (1996)
48. The concentration unit, independent of temperature, would be
(a) normality
(b) weight volume percent
(c) molality
(d) molarity. (1995, 1992)
49. According to Raoult's law, the relative lowering of vapour pressure for a solution is equal to
(a) mole fraction of solute
(b) mole fraction of solvent
(c) moles of solute
(d) moles of solvent. (1995)
50. How many grams of CH_3OH should be added to water to prepare 150 mL solution of 2 M CH_3OH ?
(a) 9.6×10^3 (b) 2.4×10^3
(c) 9.6 (d) 2.4 (1994)
51. At 25°C , the highest osmotic pressure is exhibited by 0.1 M solution of
(a) glucose (b) urea
(c) CaCl_2 (d) KCl. (1994)
52. Which of the following salt has the same value of vant's Hoff factor (i) as that of $\text{K}_3[\text{Fe}(\text{CN})_6]$?
(a) Na_2SO_4 (b) $\text{Al}(\text{NO}_3)_3$
(c) $\text{Al}_2(\text{SO}_4)_3$ (d) NaCl (1994)

53. If 0.1 M solution of glucose and 0.1 M solution of urea are placed on two sides of the semipermeable membrane to equal heights, then it will be correct to say that
 (a) there will be no net movement across the membrane
 (b) glucose will flow towards glucose solution
 (c) urea will flow towards glucose solution
 (d) water will flow from urea solution to glucose. (1992)
54. Which one is a colligative property?
 (a) Boiling point (b) Vapour pressure
 (c) Osmotic pressure (d) Freezing point (1992)
55. Blood cells retain their normal shape in solution which are
 (a) hypotonic to blood
 (b) isotonic to blood
 (c) hypertonic to blood
 (d) equinormal to blood. (1991)
56. Which of the following aqueous solution has minimum freezing point?
 (a) 0.01 m NaCl
 (b) 0.005 m C_2H_5OH
 (c) 0.005 m MgI_2
 (d) 0.005 m $MgSO_4$ (1991)
57. The relative lowering of the vapour pressure is equal to the ratio between the number of
 (a) solute molecules to the solvent molecules
 (b) solute molecules to the total molecules in the solution
 (c) solvent molecules to the total molecules in the solution
 (d) solvent molecules to the total number of ions of the solute. (1991)
58. All form ideal solution except
 (a) C_6H_6 and $C_6H_5CH_3$
 (b) C_2H_6 and C_2H_5I
 (c) C_6H_5Cl and C_6H_5Br
 (d) C_2H_5I and C_2H_5OH (1988)
59. An ideal solution is formed when its components
 (a) have no volume change on mixing
 (b) have no enthalpy change on mixing
 (c) have both the above characteristics
 (d) have high solubility. (1988)

Answer Key


1. (c) 2. (a) 3. (d) 4. (d) 5. (c) 6. (c) 7. (c) 8. (c) 9. (b) 10. (d)
 11. (c) 12. (c) 13. (b) 14. (d) 15. (None) 16. (b) 17. (c) 18. (d) 19. (d)
 20. (d) 21. (a) 22. (d) 23. (b) 24. (c) 25. (d) 26. (c) 27. (b) 28. (c) 29. (b)
 30. (a) 31. (d) 32. (b) 33. (c) 34. (b) 35. (c) 36. (a) 37. (c) 38. (a) 39. (a)
 40. (c) 41. (b) 42. (b) 43. (a) 44. (c) 45. (a) 46. (c) 47. (a) 48. (c) 49. (a)
 50. (c) 51. (c) 52. (b) 53. (a) 54. (c) 55. (b) 56. (a) 57. (a) 58. (d) 59. (c)
-

EXPLANATIONS

1. (c) : The value of molal depression constant, K_f is constant for a particular solvent, thus, it will be unchanged when molality of the dilute solution is doubled.

2. (a)

3. (d) : Being a strong electrolyte, Ba(OH)_2 undergoes 100% dissociation in a dilute aqueous solution,



Thus, van't Hoff factor $i = 3$.

4. (d) : For an ideal solution, $\Delta H_{\text{mix}} = 0$, $\Delta V_{\text{mix}} = 0$,
Now, $\Delta U_{\text{mix}} = \Delta H_{\text{mix}} - P\Delta V_{\text{mix}}$

$$\therefore \Delta U_{\text{mix}} = 0$$

Also, for an ideal solution,

$$p_A = x_A p_A^\circ, p_B = x_B p_B^\circ$$

$$\therefore \Delta P = P_{\text{obs}} - P_{\text{calculated by Raoult's law}} = 0$$

$$\Delta G_{\text{mix}} = \Delta H_{\text{mix}} - T\Delta S_{\text{mix}}$$

For an ideal solution, $\Delta S_{\text{mix}} \neq 0$

$$\therefore \Delta G_{\text{mix}} \neq 0$$

5. (c) : $p_{\text{Benzene}} = x_{\text{Benzene}} p_{\text{Benzene}}^\circ$

$$p_{\text{Toluene}} = x_{\text{Toluene}} p_{\text{Toluene}}^\circ$$

For an ideal 1 : 1 molar mixture of benzene and

$$\text{toluene, } x_{\text{Benzene}} = \frac{1}{2} \text{ and } x_{\text{Toluene}} = \frac{1}{2}$$

$$p_{\text{Benzene}} = \frac{1}{2} p_{\text{Benzene}}^\circ = \frac{1}{2} \times 12.8 \text{ kPa} = 6.4 \text{ kPa}$$

$$p_{\text{Toluene}} = \frac{1}{2} p_{\text{Toluene}}^\circ = \frac{1}{2} \times 3.85 \text{ kPa} = 1.925 \text{ kPa}$$

Thus, the vapour will contain a high percentage of benzene as the partial vapour pressure of benzene is higher as compared to that of toluene.

6. (c) : Given : $W_B = 6.5 \text{ g}$, $W_A = 100 \text{ g}$,

$p_s = 732 \text{ mm}$, $K_b = 0.52$, $T_b^\circ = 100^\circ\text{C}$, $p^\circ = 760 \text{ mm}$

$$\frac{p^\circ - p_s}{p^\circ} = \frac{n_2}{n_1} \Rightarrow \frac{760 - 732}{760} = \frac{n_2}{100/18}$$

$$\Rightarrow n_2 = \frac{28 \times 100}{760 \times 18} = 0.2046 \text{ mol}$$

$$\Delta T_b = K_b \times m$$

$$T_b - T_b^\circ = K_b \times \frac{n_2 \times 1000}{W_A (\text{g})}$$

$$T_b - 100^\circ\text{C} = \frac{0.52 \times 0.2046 \times 1000}{100} = 1.06$$

$$T_b = 100 + 1.06 = 101.06^\circ\text{C}$$




7. (c) : 1 molal aqueous solution means 1 mole of solute is present in 1000 g of water.

$$\therefore x_{\text{solute}} = \frac{1}{1 + \frac{1000}{18}} = \frac{1}{56.5} = 0.0177$$

8. (c) : $\Delta T_b = i K_b m$

For equimolal solutions, elevation in boiling point will be higher if solution undergoes dissociation i.e., $i > 1$.

9. (b) : $\text{Al}_2(\text{SO}_4)_3 \longrightarrow 2\text{Al}^{3+} + 3\text{SO}_4^{2-}$, $i = 5$



10. (d) : For an ideal solution, $\Delta S_{\text{mix}} > 0$ while ΔH_{mix}

ΔV_{mix} and $\Delta P = 0$.

11. (c) : $\Delta T_f = i \times K_f \times m$

So, $\Delta T_f \propto i$ (van't Hoff factor)

Salt	i
KCl	2
$\text{C}_6\text{H}_{12}\text{O}_6$	1
$\text{Al}_2(\text{SO}_4)_3$	5
K_2SO_4	3

Hence, i is maximum i.e., 5 for $\text{Al}_2(\text{SO}_4)_3$.

$$12. (c) : \text{Molarity} = \frac{W \times 1000}{M_w \times V_{\text{sol}}(\text{mL})} \Rightarrow 2 = \frac{W}{63} \times \frac{1000}{250}$$

$$W = \frac{63}{2} \text{ g; Mass of acid} \times \frac{70}{100} = \frac{63}{2}$$

Mass of acid = 45 g

13. (b) : An ideal solution is follow:

- Volume change (ΔV) of mixing should be zero
- Heat change (ΔH) on mixing should be zero.
- Obey Raoult's law at every range of concentration.

14. (d) : According to Raoult's law,

$$P = x_A p_A + x_B p_B \quad \dots (i)$$

For binary solutions,

$$x_A + x_B = 1, \quad x_B = 1 - x_A \quad \dots (ii)$$

Putting value of x_B from eqn. (ii) to eqn. (i)

$$P = x_A p_A + (1 - x_A) p_B = x_A p_A + p_B - x_A p_B$$

$$P = p_B + x_A (p_A - p_B)$$

15. (None) :

$$P^\circ_{\text{CHCl}_3} = 200 \text{ mm Hg, } P^\circ_{\text{CH}_2\text{Cl}_2} = 41.5 \text{ mm Hg}$$

Moles of CHCl_3

$$= \frac{\text{Weight}}{\text{Molecular weight}} = \frac{25.5}{119.5} = 0.213$$

$$\text{Moles of CH}_2\text{Cl}_2 = \frac{40}{85} = 0.470$$

$$X_{\text{CHCl}_3} = \frac{0.213}{0.213 + 0.470} = 0.31$$

$$X_{\text{CH}_2\text{Cl}_2} = \frac{0.470}{0.213 + 0.470} = 0.69$$

$$P_T = P_{\text{CHCl}_3}^{\circ} X_{\text{CHCl}_3} + P_{\text{CH}_2\text{Cl}_2}^{\circ} X_{\text{CH}_2\text{Cl}_2}$$

$$= 200 \times 0.31 + 41.5 \times 0.69 = 62 + 28.63 = 90.63$$

16. (b) : We know that $\Delta T_f = i \times K_f \times \frac{w_B \times 1000}{m_B \times w_A}$

Given : $\Delta T_f = 3.82$, $K_f = 1.86$,

$w_B = 5$, $m_B = 142$, $w_A = 45$

$$i = \frac{\Delta T_f \times m_B \times w_A}{K_f \times w_B \times 1000} = \frac{3.82 \times 142 \times 45}{1.86 \times 5 \times 1000} = 2.63$$

17. (c) : From the value of van't Hoff factor i it is possible to determine the degree of dissociation or association. In case of dissociation, i is greater than 1 and in case of association i is less than 1.

18. (d) : We know that $\Delta T_f = i \times K_f \times m$

Here i is van't Hoff's factor.

i for weak acid is $1 + \alpha$.

Here α is degree of dissociation *i.e.*, $30/100 = 0.3$

$$\therefore i = 1 + \alpha = 1 + 0.3 = 1.3$$

$$\Delta T_f = i \times K_f \times m = 1.3 \times 1.86 \times 0.1 = 0.24$$

$$\therefore \text{Freezing point} = -0.24$$

19. (d) : We know that $\pi V = nRT$, where $n = \frac{w}{M}$

$$\pi V = \frac{w}{M} RT$$

$$M = \frac{wRT}{\pi V} = \frac{1.26 \times 0.083 \times 300}{2.57 \times 10^{-3} \times \frac{200}{1000}} = \frac{1.26 \times 0.083 \times 300}{2.57 \times 10^{-3} \times 0.2}$$

$$= 61038 \text{ g mol}^{-1}$$

20. (d) : Addition of water to an aqueous solution of KI causes the concentration of the solution to decrease thereby increasing the vapour pressure. In the other three options the electrolytes undergo ionization, which leads to lowering of vapour pressure.

21. (a) : We know, $\Delta T_f = K_f m$

$$m = \frac{w_B}{M_B} \times \frac{1000}{W_A} = \frac{68.5 \times 1000}{342 \times 1000} = \frac{68.5}{342}$$

$$\Delta T_f = 1.86 \times \frac{68.5}{342} = 0.372^\circ\text{C}$$


$$\therefore T_f = 0 - 0.372^\circ\text{C} = -0.372^\circ\text{C}$$

22. (d) : The number of moles of ions produced by 1 mol of ionic compound = i

Applying, $\Delta T_f = i \times K_f \times m$

$$0.00732 = i \times 1.86 \times 0.002$$

$$\Rightarrow i = \frac{0.00732}{1.86 \times 0.002} = 1.96 \approx 2$$



$$\begin{matrix} 1 - \alpha & \alpha & \alpha \end{matrix}$$

$$\text{Total} = 1 + \alpha$$

$$\therefore i = 1 + \alpha = 1 + 0.2 = 1.2$$

$$\Delta T_f = i \times K_f \times m = 1.2 \times 1.86 \times 0.5$$

$$= 1.116 \text{ K} \approx 1.12 \text{ K}$$

24. (c) : H_2SO_4 is 98% by weight.

Weight of $\text{H}_2\text{SO}_4 = 98 \text{ g}$

Weight of solution = 100 g

$$\therefore \text{Volume of solution} = \frac{\text{mass}}{\text{density}} = \frac{100}{1.80} \text{ mL}$$

$$= 55.55 \text{ mL} = 0.0555 \text{ L}$$

$$\text{Molarity of solution} = \frac{98}{98 \times 0.0555} \text{ M} = 18.02 \text{ M}$$

Let V mL of this H_2SO_4 are used to prepare 1 litre of 0.1 M H_2SO_4 .

$$\therefore \text{mM of concentrated } \text{H}_2\text{SO}_4 = \text{mM of dilute } \text{H}_2\text{SO}_4$$

$$\text{or, } V \times 18.02 = 1000 \times 0.1 \Rightarrow V = \frac{1000 \times 0.1}{18.02} = 5.55 \text{ mL}$$

25. (d) : Osmosis is the phenomenon of flow of pure solvent from the solvent to the solution or from a less concentrated solution to a more concentrated solution through a semipermeable membrane. Common semipermeable membranes are permeable to certain solute particles also. Infact, there is no perfect semipermeable membrane. Therefore we can say that flow of water through a semipermeable membrane takes place both sides with unequal rates.

26. (c) : Both the components escape easily showing higher vapour pressure than the expected value.

27. (b) : $m = \frac{1000 \times K_f \times w}{W \times \Delta T}$ or, $250 = \frac{1000 \times 5.12 \times 1}{51.2 \times \Delta T}$

$$\therefore \Delta T = \frac{1000 \times 5.12 \times 1}{51.2 \times 250} = 0.4 \text{ K}$$

28. (c) : For isotonic solution, osmotic pressure of urea = osmotic pressure of non-volatile solute

$$\frac{10}{60 \times 1000} = \frac{5}{m \times 100} \Rightarrow m = 300 \text{ g mol}^{-1}$$

29. (b) : 1 mole of solute present in 1 kg of H_2O

1 mole of solute present in $\frac{1000}{18}$ g mole of H_2O

$$X_{\text{solute}} = \frac{1}{\frac{1000}{18} + 1} = \frac{18}{1018} = 0.01768 \approx 0.018$$

30. (a) : By Raoult's Law

$$P_T = P_P^{\circ} X_P^{\circ} + P_Q^{\circ} X_Q^{\circ}$$

where $P_P^\circ = 80$ torr, $P_Q^\circ = 60$ torr, $X_P = \frac{3}{5}$, $X_Q = \frac{2}{5}$

$$P_T = 80 \times \frac{3}{5} + 60 \times \frac{2}{5} = 48 + 24 = 72 \text{ torr}$$

$$31. (d) : \frac{n_{C_5H_{12}}}{n_{C_6H_{14}}} = \frac{1}{4}$$

$$\Rightarrow X_{C_5H_{12}} = \frac{1}{5} \text{ and } X_{C_6H_{14}} = \frac{4}{5}$$

$$P_{C_5H_{12}}^\circ = 440 \text{ mm Hg}; P_{C_6H_{14}}^\circ = 120 \text{ mm Hg}$$

$$P_T = P_{C_5H_{12}}^\circ X_{C_5H_{12}} + P_{C_6H_{14}}^\circ X_{C_6H_{14}}$$

$$= 440 \times \frac{1}{5} + 120 \times \frac{4}{5} = 88 + 96 = 184 \text{ mm of Hg}$$

By Raoult's Law, $P_{C_5H_{12}} = P_{C_5H_{12}}^\circ X_{C_5H_{12}} \dots (1)$

$X_{C_5H_{12}} \rightarrow$ mole fraction of pentane in solution

By Dalton's Law, $P_{C_5H_{12}} = X'_{C_5H_{12}} P \dots (2)$

$X'_{C_5H_{12}} \rightarrow$ mole fraction of pentane above the solution.

From (1) and (2),

$$P_{C_5H_{12}} = 440 \times \frac{1}{5} = 88 \text{ mm of Hg} \Rightarrow 88 = X'_{C_5H_{12}} \times 184$$

$$X' = \frac{88}{184}; X' = 0.478$$

$$32. (b) : \left. \begin{array}{l} \Delta T_f = K_f m \dots (1) \\ \Delta T_b = K_b m \dots (2) \end{array} \right\} \Rightarrow \frac{\Delta T_f}{\Delta T_b} = \frac{K_f}{K_b} \dots (3)$$

$\Delta T_f \rightarrow$ depression in freezing point

$\Delta T_b \rightarrow$ elevation in b.p.

b.p. of water = 100°C ; $K_f = 1.86 \text{ kg mol}^{-1}$

b.p. of urea in water = 100.18°C

$K_b = 0.512 \text{ kg mol}^{-1} \Rightarrow \Delta T_b = 0.18$

f.p. of water = 0°C

f.p. of urea in water = $-T^\circ\text{C}$

$$\Rightarrow \Delta T_f = T$$

$$\Rightarrow \text{from eq. (3), } \frac{T}{0.18} = \frac{1.86}{0.512} \Rightarrow T = 0.6539$$

$$\Rightarrow \text{f.p. of urea in water} = -0.654^\circ\text{C}$$

33. (c) : Raoult's law is valid for ideal solution only. The element of non-ideality enters into the picture when the molecules of the solute and solvent affect each others intermolecular forces. A solution containing components of A and B behaves as ideal solution when A-B attraction force remains same as A-A and B-B.

34. (b) : For dilute solution, the van't Hoff equation

$$\text{is } \pi = \frac{n}{V} RT$$

$$\Rightarrow \pi V = \frac{m_2}{M} RT \Rightarrow M = \frac{m_2 RT}{\pi V}$$

35. (c) : Molecular weight of NaOH = 40

2.5 litre of 1 M NaOH solution contain

$$40 \times 2.5 \text{ g of NaOH}$$

3 litre of 0.5 M NaOH solution contain

$$40 \times 0.5 \times 3 \text{ g of NaOH}$$

If these two solutions are mixed, the volume of the resultant solution = $(2.5 + 3) = 5.5$ litre.

5.5 litre of the resultant solution contain

$$40(2.5 + 1.5) \text{ g of NaOH}$$

1 litre of the resultant solution contain

$$\frac{40 \times 4}{5.5} \text{ g of NaOH} = \frac{40 \times 4}{5.5 \times 40} \text{ mole of NaOH}$$

The molarity of the resultant solution = $0.727 \approx 0.73 \text{ M}$

36. (a) : More is the pressure, greater will be the boiling point.

37. (c)

38. (a) : Polymers and proteins are covalent compounds, therefore by measuring osmotic pressure of their solution, the molecular weight can be determined.

$$PV = nRT \text{ or, } \pi V = \frac{W}{M} RT$$

$$\Rightarrow M = \frac{WRT}{\pi V} = \frac{CRT}{\pi}$$

39. (a) : $P^\circ = 640$ mm, $P_s = 600$ mm, $w = 2.175$ g. $W = 39.08$ g

From Raoult's law

$$\frac{P^\circ - P_s}{P^\circ} = \frac{w \times M}{W \times m} \Rightarrow \frac{640 - 600}{640} = \frac{2.175 \times 78}{39.08 \times m}$$

$$\Rightarrow m = 69.5$$

40. (c) : The strength of the solution is 0.1 N

$$\frac{w}{E} = \frac{V \times N}{1000} \text{ (Equivalent weight} = \frac{200}{2} = 100)$$

$$\Rightarrow w = \frac{100 \times 0.1 \times 100}{1000} = 1 \text{ g}$$

41. (b) : $w = 0.15$ g, $W = 15$ g, $\Delta T_b = 0.216^\circ\text{C}$

$K_b = 2.16$, $m = ?$

$$\text{As } \Delta T_b = \frac{1000 \times K_b \times w}{m \times W}$$

$$\Rightarrow m = \frac{1000 \times 2.16 \times 0.15}{0.216 \times 15} = 100$$

42. (b) : x_1 (mole fraction of solute) = 0.2

From Raoult's law,

$$\frac{P^\circ - P_s}{P^\circ} = x_1 \Rightarrow \frac{10}{P^\circ} = 0.2 \Rightarrow P^\circ = 50 \text{ mm}$$

Again, when $P^\circ - P_s = 20$ mm, then

$$\frac{P^\circ - P_s}{P^\circ} = \text{mole fraction of solute} = \frac{20}{50} = 0.4$$

$$\Rightarrow \text{mole fraction of solvent} = 1 - 0.4 = 0.6$$

43. (a) : 'Cane Sugar'

$$W_1 = 5 \text{ g}$$

$$V_1 = 100 \text{ mL}$$

$$= 0.1 \text{ L}$$

$$M_1 = 342$$

'X'

$$W_2 = 1 \text{ g}$$

$$V_2 = 100 \text{ mL}$$

$$= 0.1 \text{ L}$$

$$M_2 = ?$$

For isotonic solutions, $C_1 = C_2$

$$\frac{W_1}{M_1 V_1} = \frac{W_2}{M_2 V_2} \Rightarrow \frac{5}{342 \times 0.1} = \frac{1}{M_2 \times 0.1}$$

$$\Rightarrow M_2 = \frac{342}{5} = 68.4$$

44. (c) : Since $\text{Al}_2(\text{SO}_4)_3$ gives maximum number of ions on dissociation, therefore it will have the lowest freezing point.

45. (a) : We know that 98% H_2SO_4 by weight means 98 g of H_2SO_4 is present in 100 g of solution in water.

Therefore its weight is 98 and moles of H_2SO_4

$$= \frac{\text{Weight of } \text{H}_2\text{SO}_4}{\text{Molecular Weight}} = \frac{98}{98} = 1$$

and volume of solution = $\frac{\text{Mass}}{\text{Density}}$

$$= \frac{100}{1.84} = 54.35 \text{ mL} = \frac{54.35}{1000} \text{ L}$$

Therefore molarity of H_2SO_4

$$= \frac{\text{Moles of } \text{H}_2\text{SO}_4}{\text{Volume (in litres)}} = \frac{1 \times 1000}{54.35} = 18.4 \text{ M}$$

46. (c) : Vapour pressure of pure solvent (P_A°) = 143 mm Hg, weight of solute (w) = 0.5 gram, weight of solvent (W) = 100 grams, molecular weight of solute (m) = 65 and molecular weight of solvent (M) = 154.

$$\frac{P_A^\circ - P}{P_A^\circ} = \frac{wM}{mW} \quad \text{or} \quad \frac{143 - P}{143} = \frac{0.5 \times 154}{65 \times 100} = 0.01$$

or $P = 141.31 \text{ mm Hg}$

47. (a) : Weight of glucose = 10 g,
Weight of urea = 10 g and weight of sucrose = 10 g.
We know that the number of moles of glucose

$$(n_1) = \frac{\text{Weight}}{\text{Molecular weight}} = \frac{10}{180} = 0.05$$

Similarly number of moles of urea (n_2) = $\frac{10}{60} = 0.16$
and the number of moles of sucrose

$$(n_3) = \frac{10}{342} = 0.03$$

The osmotic pressure is a colligative property and it depends upon the number of moles of a solute. Since $n_2 > n_1 > n_3$, therefore $p_2 > p_1 > p_3$.

48. (c) : The molality involves weights of the solute and the solvent. Since the weight does not change with the temperatures, therefore molality does not depend upon the temperature.

49. (a) : Since relative lowering of vapour pressure is a colligative property, therefore it depends upon the number of solute particles or mole fraction of solute.

50. (c) : Since the molecular mass of CH_3OH is 32, therefore quantity of CH_3OH to prepare 150 mL

$$\text{solution of } 2 \text{ M } \text{CH}_3\text{OH} = \left(\frac{2}{1000} \right) \times 150 \times 32 = 9.6 \text{ g}$$

51. (c) : In solution, CaCl_2 gives three ions, KCl gives two ions while glucose and urea are covalent molecules so they do not undergo ionisation. Since osmotic pressure is a colligative property and it depends upon the number of solute particles (ions), therefore, 0.1 M solution of CaCl_2 exhibits the highest osmotic pressure.

52. (b) : $\text{K}_3[\text{Fe}(\text{CN})_6] \rightleftharpoons 3\text{K}^+ + [\text{Fe}(\text{CN})_6]^{3-}$ and
 $\text{Al}(\text{NO}_3)_3 \rightleftharpoons \text{Al}^{3+} + 3\text{NO}_3^-$

Since both $\text{Al}(\text{NO}_3)_3$ and $\text{K}_3[\text{Fe}(\text{CN})_6]$ give the same number of ions, therefore they have the same van't Hoff factor.

53. (a) : There is no net movement of the solvent through the semipermeable membrane between two solutions of equal concentration.

54. (c) : The properties which depend only upon the number of solute particles present in the solution irrespective of their nature are called as colligative properties. Lowering in vapour pressure, elevation in boiling point, depression in freezing point and osmotic pressure are colligative properties.

55. (b) : Blood cells neither swell nor shrink in isotonic solution.

The solutions having same osmotic pressure are called isotonic solutions.

56. (a) : Here, $\Delta T_f = i \times K_f \times m$

Van't Hoff factor, $i = 2$ for NaCl, so conc. = 0.02, which is maximum in the present case.

Hence, ΔT_f is maximum or freezing point is minimum in 0.01 m NaCl.

$$\text{57. (a) : } \frac{P^\circ_{\text{solvent}} - P_{\text{solution}}}{P^\circ_{\text{solvent}}} = x_{\text{solute}}$$

Here, $P_A^\circ - P_A$ (difference in vapour pressure of pure solvent and solution) = lowering in vapour pressure

... (i)

Now, P_A° = vapour pressure of pure solvent ... (ii)

by dividing (i) and (ii), we get $\frac{P_A^\circ - P_A}{P_A^\circ}$

which is relative lowering in vapour pressure

58. (d) : Because $\text{C}_2\text{H}_5\text{I}$ and $\text{C}_2\text{H}_5\text{OH}$ are dissimilar liquids.

59. (c) : For ideal solution,

$$\Delta V_{\text{mixing}} = 0 \text{ and } \Delta H_{\text{mixing}} = 0$$

