

DAY FOURTEEN

Maxima and Minima

Learning & Revision for the Day

• Maxima and Minima of a Function

• Concept of Global Maximum/Minimum

Maxima and Minima of a Function

A function $f(x)$ is said to attain a **maximum** at $x = a$, if there exists a neighbourhood $(a - \delta, a + \delta)$, $x \neq a$ i.e. $f(x) < f(a)$, $\forall x \in (a - \delta, a + \delta)$,


$x \neq a \cdot h > 0$ (very small quantity)

In such a case $f(a)$ is said to be the maximum value of $f(x)$ at $x = a$.

A function $f(x)$ is said to attain a **minimum** at $x = a$, if there exists a neighbourhood $(a - \delta, a + \delta)$, $x \neq a$ such that $f(x) > f(a)$, $\forall x \in (a - \delta, a + \delta)$, $x \neq a$.

Graph of a continuous function explained local maxima (minima) and absolute maxima (minima). In such a case $f(a)$ is said to be the minimum value of $f(x)$ at $x = a$.

The points at which a function attains either the maximum or the minimum values are known as the **extreme points** or **turning points** and both minimum and maximum values of $f(x)$ are called extreme values. The turning points A and C are called **local maximum** and points B and D are called **local minimum**.



Critical Point

- A point c in the domain of a function f at which either $f'(c) = 0$ or f is not differentiable is called a **critical point** of f . Note that, if f is continuous at point c and $f'(c) = 0$, then there exists $h > 0$ such that f is differentiable in the interval $(c - h, c + h)$.
- The converse of above theorem need not be true, that is a point at which the derivative vanishes need not be a point of local maxima or local minima.


Method to Find Local Maxima or Local Minima

First Derivative Test

Let f be a function defined on an open interval I and f be continuous at a critical point c in I . Then,

- (i) If $f'(x)$ changes sign from positive to negative as x increases through c , i.e. if $f'(x) > 0$ at every point sufficiently close to and to the left of c and $f'(x) < 0$ at every point sufficiently close to and to the right of c , then c is a point of **local maxima**.

- (ii) If $f'(x)$ changes sign from negative to positive as x increases through point c , i.e. if $f'(x) < 0$ at every point sufficiently close to and to the left of c and $f'(x) > 0$ at every point sufficiently close to and to the right of c , then c is a point of **local minima**.
- (iii) If $f'(x)$ does not change sign as x increases through c , then c is neither a point of local maxima nor a point of local minima. Infact, such a point is called **point of inflection**.



Graph of f around c explained following points.

- (iv) If c is a point of local maxima of f , then $f(c)$ is a local maximum value of f . Similarly, if c is a point of local minima of f , then $f(c)$ is a local minimum value of f .

Second or Higher Order Derivative Test

- (i) Find $f'(x)$ and equate it to zero. Solve $f'(x) = 0$ let its roots be $x = a_1, a_2, \dots$
- (ii) Find $f''(x)$ and at $x = a_1$,
 - (a) if $f''(a_1)$ is positive, then $f(x)$ is minimum at $x = a_1$.
 - (b) if $f''(a_1)$ is negative, then $f(x)$ is maximum at $x = a_1$.
- (iii) (a) If at $x = a_1, f''(a_1) = 0$, then find $f'''(x)$. If $f'''(a_1) \neq 0$, then $f(x)$ is neither maximum nor minimum at $x = a_1$.
 - (b) If $f'''(a_1) = 0$, then find $f^{iv}(x)$.
 - (c) If $f^{iv}(x)$ is positive (minimum value) and $f^{iv}(x)$ is negative (maximum value).
- (iv) If at $x = a_1, f^{iv}(a_1) = 0$, then find $f^v(x)$ and proceed similarly.

Point of Inflection

At point of inflection

- (i) It is not necessary that 1st derivative is zero.
- (ii) 2nd derivative must be zero or 2nd derivative changes sign in the neighbourhood of point of inflection.

n th Derivative Test

Let f be a differentiable function on an interval I and a be an interior point of I such that

- (i) $f'(a) = f''(a) = f'''(a) = \dots = f^{n-1}(a) = 0$ and
- (ii) $f^n(a)$ exists and is non-zero.

Important Results

- If n is even and $f^n(a) < 0 \Rightarrow x = a$ is a point of local maximum.
- If n is even and $f^n(a) > 0 \Rightarrow x = a$ is a point of local minimum.
- If n is odd $\Rightarrow x = a$ is a point of neither local maximum nor a point of local minimum.
- The function $f(x) = \frac{ax + b}{cx + d}$ has no local maximum or minimum regardless of values of a, b, c and d .
- The function $f(\theta) = \sin^m \theta \cdot \cos^n \theta$ attains maximum values at $\theta = \tan^{-1} \left(\sqrt{\frac{m}{n}} \right)$.
- If AB is diameter of circle and C is any point on the circumference, then area of the ΔABC will be maximum, if triangle is isosceles.

Concept of Global Maximum/Minimum

- Let $y = f(x)$ be a given function with domain D and $[a, b] \subseteq D$, then global maximum/minimum of $f(x)$ in $[a, b]$ is basically the greatest / least value of $f(x)$ in $[a, b]$.
- Global maxima/minima in $[a, b]$ would always occur at critical points of $f(x)$ within $[a, b]$ or at end points of the interval.

Global Maximum/Minimum in $[a, b]$

In order to find the global maximum and minimum of $f(x)$ in $[a, b]$.

- Step I** Find out all critical points of $f(x)$ in $[a, b]$ [i.e. all points at which $f'(x) = 0$] and let these points are c_1, c_2, \dots, c_n .
- Step II** Find the value of $f(c_1), f(c_2), \dots, f(c_n)$ and also at the end points of domain i.e. $f(a)$ and $f(b)$.
- Step III** Find $M_1 \rightarrow$ Global maxima or greatest value and $M_2 \rightarrow$ Global minima or least value. where, $M_1 = \max \{f(a), f(c_1), f(c_2), \dots, f(c_n), f(b)\}$ and $M_2 = \min \{f(a), f(c_1), f(c_2), \dots, f(c_n), f(b)\}$

Some Important Results on Maxima and Minima

- (i) Maxima and minima occur alternatively i.e. between two maxima there is one minimum and *vice-versa*.
- (ii) If $f(x) \rightarrow \infty$ as $x \rightarrow a$ or b and $f'(x) = 0$ only for one value of x (say c) between a and b , then $f(c)$ is necessarily the minimum and the least value.
- (iii) If $f(x) \rightarrow -\infty$ as $x \rightarrow a$ or b , then $f(c)$ is necessarily the maximum and greatest value.
- (iv) The **stationary points** are the points of the domain, where $f'(x) = 0$.

DAY PRACTICE SESSION 1

FOUNDATION QUESTIONS EXERCISE

- 1 If f is defined as $f(x) = x + \frac{1}{x}$, then which of the following is true?
→ NCERT Exemplar

(a) Local maximum value of $f(x)$ is -2
(b) Local minimum value of $f(x)$ is 2
(c) Local maximum value of $f(x)$ is less than local minimum value of $f(x)$
(d) All the above are true

- 2 If the sum of two numbers is 3 , then the maximum value of the product of the first and the square of second is
→ NCERT Exemplar

(a) 4 (b) 1 (c) 3 (d) 0

- 3 If $y = a \log x + bx^2 + x$ has its extremum value at $x = 1$ and $x = 2$, then (a, b) is equal to

(a) $\left(1, \frac{1}{2}\right)$ (b) $\left(\frac{1}{2}, 2\right)$ (c) $\left(2, \frac{-1}{2}\right)$ (d) $\left(\frac{-2}{3}, \frac{-1}{6}\right)$

- 4 The function $f(x) = a \cos x + b \tan x + x$ has extreme values at $x = 0$ and $x = \frac{\pi}{6}$, then

(a) $a = -\frac{2}{3}, b = -1$ (b) $a = \frac{2}{3}, b = -1$
(c) $a = -\frac{2}{3}, b = 1$ (d) $a = \frac{2}{3}, b = 1$

- 5 The minimum radius vector of the curve

$$\frac{4}{x^2} + \frac{9}{y^2} = 1 \text{ is of length}$$

(a) 1 (b) 5 (c) 7 (d) None of these

- 6 The function $f(x) = 4x^3 - 18x^2 + 27x - 7$ has
→ NCERT Exemplar

(a) one local maxima
(b) one local minima
(c) one local maxima and two local minima
(d) neither maxima nor minima

- 7 The function $f(x) = \frac{x^2 - 2}{x^2 - 4}$ has

(a) no point of local minima
(b) no point of local maxima
(c) exactly one point of local minima
(d) exactly one point of local maxima

- 8 Let $f: R \rightarrow R$ be defined by $f(x) = \begin{cases} k - 2x, & \text{if } x \leq -1 \\ 2x + 3, & \text{if } x > -1 \end{cases}$

If f has a local minimum at $x = -1$, then a possible value of k is
→ AIEEE 2010

(a) 1 (b) 0 (c) $-\frac{1}{2}$ (d) -1

- 9 The minimum value of $9x + 4y$, where $xy = 16$ is

(a) 48 (b) 28 (c) 38 (d) 18

- 10 If the function $f(x) = 2x^3 - 9ax^2 + 12a^2x + 1$, where $a > 0$ attains its maximum and minimum at p and q respectively such that $p^2 = q$, then a is equal to

(a) 3 (b) 1 (c) 2 (d) $\frac{1}{2}$

- 11 If $f(x) = x^2 + 2bx + 2c^2$ and $g(x) = -x^2 - 2cx + b^2$ such that minimum $f(x) >$ maximum $g(x)$, then the relation between b and c is

(a) $0 < c < b\sqrt{2}$ (b) $|c| < |b|\sqrt{2}$
(c) $|c| > |b|\sqrt{2}$ (d) No real values of b and c

- 12 Let $f(x)$ be a polynomial of degree four having extreme values at $x = 1$ and $x = 2$. If $\lim_{x \rightarrow 0} \left[1 + \frac{f(x)}{x^2}\right] = 3$, then $f(2)$ is equal to
→ JEE Mains 2015

(a) -8 (b) -4
(c) 0 (d) 4

- 13 If a differential function $f(x)$ has a relative minimum at $x = 0$, then the function $\phi(x) = f(x) + ax + b$ has a relative minimum at $x = 0$ for

(a) all a and all b (b) all b , if $a = 0$
(c) all $b > 0$ (d) all $a > 0$

- 14 The denominator of a fraction is greater than 16 of the square of numerator, then least value of fraction is

(a) $-1/4$ (b) $-1/8$
(c) $1/12$ (d) $1/16$

- 15 The function $f(x) = ax + \frac{b}{x}$, $b, x > 0$ takes the least value at x equal to

(a) b (b) \sqrt{a} (c) \sqrt{b} (d) $\sqrt{\frac{b}{a}}$

- 16 Let f be a function defined by $f(x) = \begin{cases} \frac{\tan x}{x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$

Statement I $x = 0$ is point of minima of f .

Statement II $f'(0) = 0$.

→ AIEEE 2011

(a) Statement I is true, Statement II is true; Statement II is a correct explanation for Statement I
(b) Statement I is true, Statement II is true; Statement II is not a correct explanation for Statement I
(c) Statement I is true; Statement II is false
(d) Statement I is false; Statement II is true

- 17 The absolute maximum and minimum values of the function f given by $f(x) = \cos^2 x + \sin x$, $x \in [0, \pi]$

→ NCERT Exemplar

(a) 2.25 and 2 (b) 1.25 and 1
(c) 1.75 and 1.5 (d) None of these

18 The maximum value of $f(x) = \frac{x}{4+x+x^2}$ on $[-1, 1]$ is

- (a) $-\frac{1}{4}$ (b) $-\frac{1}{3}$ (c) $\frac{1}{6}$ (d) $\frac{1}{5}$

19 In interval $[1, e]$, the greatest value of $x^2 \log x$ is

- (a) e^2 (b) $\frac{1}{e} \log \frac{1}{\sqrt{e}}$ (c) $e^2 \log \sqrt{e}$ (d) None of these

20 If x is real, the maximum value of $\frac{3x^2 + 9x + 17}{3x^2 + 9x + 7}$ is

- (a) 41 (b) 1 (c) $\frac{17}{7}$ (d) $\frac{1}{4}$ **→ AIEEE 2007**

21 The maximum and minimum values of $f(x) = \sec x + \log \cos^2 x$, $0 < x < 2\pi$ are respectively

- (a) $(1, -1)$ and $\{2(1 - \log 2), 2(1 + \log 2)\}$
(b) $(1, -1)$ and $\{2(1 - \log 2), 2(1 - \log 2)\}$
(c) $(1, -1)$ and $(2, -3)$
(d) None of the above **→ NCERT Exemplar**

22 The difference between greatest and least values of the function $f(x) = \sin 2x - x$, on $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ is

- (a) π (b) 2π (c) 3π (d) $\frac{\pi}{2}$ **→ NCERT Exemplar**

23 The point of inflection for the curve $y = x^{5/2}$ is

- (a) $(1, 1)$ (b) $(0, 0)$ (c) $(1, 0)$ (d) $(0, 1)$

24 The maximum area of a right angled triangle with hypotenuse h is

- (a) $\frac{h^3}{2\sqrt{2}}$ (b) $\frac{h^2}{2}$ (c) $\frac{h^2}{\sqrt{2}}$ (d) $\frac{h^2}{4}$ **→ JEE Main 2013**

25 A straight line is drawn through the point $P(3, 4)$ meeting the positive direction of coordinate axes at the points A and B . If O is the origin, then minimum area of $\triangle OAB$ is equal to

- (a) 12 sq units (b) 6 sq units
(c) 24 sq units (d) 48 sq units

26 Suppose the cubic $x^3 - px + q$ has three distinct real roots, where $p > 0$ and $q > 0$. Then, which one of the following holds?

- (a) The cubic has maxima at both $\sqrt{\frac{p}{3}}$ and $-\sqrt{\frac{p}{3}}$
(b) The cubic has minima at $\sqrt{\frac{p}{3}}$ and maxima at $-\sqrt{\frac{p}{3}}$
(c) The cubic has minima at $-\sqrt{\frac{p}{3}}$ and maxima at $\sqrt{\frac{p}{3}}$
(d) The cubic has minima at both $\sqrt{\frac{p}{3}}$ and $-\sqrt{\frac{p}{3}}$

27 If $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$ are the vertices of a $\triangle ABC$. A parallelogram $AFDE$ is drawn with D , E and F on the line segment BC , CA and AB , respectively. Then, maximum area of such parallelogram is

- (a) $\frac{1}{2}$ (area of $\triangle ABC$) (b) $\frac{1}{4}$ (area of $\triangle ABC$)
(c) $\frac{1}{6}$ (area of $\triangle ABC$) (d) $\frac{1}{8}$ (area of $\triangle ABC$)

28 If $y = f(x)$ is a parametrically defined expression such that $x = 3t^2 - 18t + 7$ and $y = 2t^3 - 15t^2 + 24t + 10$, $\forall x \in [0, 6]$.

Then, the maximum and minimum values of $y = f(x)$ are

- (a) 36, 3 (b) 46, 6 (c) 40, -6 (d) 46, -6

29 The value of a , so that the sum of the squares of the roots of the equation $x^2 - (a - 2)x - a + 1 = 0$ assume the least value is

- (a) 2 (b) 1 (c) 3 (d) 0

30 The minimum intercepts made by the axes on the tangent to the ellipse $\frac{x^2}{16} + \frac{y^2}{9} = 1$ is

- (a) 25 (b) 7 (c) 1 (d) None of these

31 The curved surface of the cone inscribed in a given sphere is maximum, if

- (a) $h = \frac{4R}{3}$ (b) $h = \frac{R}{3}$ (c) $h = \frac{2R}{3}$ (d) None of these

32 The volume of the largest cone that can be inscribed in a sphere of radius R is **→ NCERT**

- (a) $\frac{3}{8}$ of the volume of the sphere
(b) $\frac{8}{27}$ of the volume of the sphere
(c) $\frac{2}{7}$ of the volume of the sphere
(d) None of the above

33 Area of the greatest rectangle that can be inscribed in the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is

- (a) \sqrt{ab} (b) $\frac{a}{b}$ (c) $2ab$ (d) ab

34 The real number x when added to its inverse gives the minimum value of the sum at x equal to **→ AIEEE 2003**

- (a) 2 (b) 1 (c) -1 (d) -2

35 The greatest value of

$f(x) = (x + 1)^{1/3} - (x - 1)^{1/3}$ on $[0, 1]$ is **→ AIEEE 2002**

- (a) 1 (b) 2 (c) 3 (d) $\frac{1}{3}$

36 The coordinate of a point on the parabola $y^2 = 8x$ whose distance from the circle $x^2 + (y + 6)^2 = 1$ is minimum, is

- (a) $(2, -4)$ (b) $(2, 4)$ (c) $(18, -12)$ (d) $(8, 8)$

37 The volume of the largest cylinder that can be inscribed in a sphere of radius r cm is

- (a) $\frac{4\pi r^3}{\sqrt{3}}$ (b) $\frac{4\pi r^3}{3\sqrt{3}}$ (c) $\frac{4\pi r^3}{2\sqrt{3}}$ (d) $\frac{4\pi r^3}{5\sqrt{2}}$

38 Maximum slope of the curve $y = -x^3 + 3x^2 + 9x - 27$ is

- (a) 0 (b) 12 (c) 16 (d) 32

39 If $ab = 2a + 3b$, $a > 0$, $b > 0$, then the minimum value of ab is

- (a) 12 (b) 24
(c) $\frac{1}{4}$ (d) None of these

40 The perimeter of a sector is p . The area of the sector is maximum, when its radius is

- (a) \sqrt{p} (b) $\frac{1}{\sqrt{p}}$ (c) $\frac{p}{2}$ (d) $\frac{p}{4}$

DAY PRACTICE SESSION 2

PROGRESSIVE QUESTIONS EXERCISE

1 The minimum radius vector of the curve $\frac{a^2}{x^2} + \frac{b^2}{y^2} = 1$ is of

- length
(a) $a - b$ (b) $a + b$
(c) $2a + b$ (d) None of these

2 $f(x) = x^2 - 4|x|$ and

$g(x) = \begin{cases} \min \{f(t) : -6 \leq t \leq x\}, & x \in [-6, 0] \\ \max \{f(t) : 0 < t \leq x\}, & x \in (0, 6] \end{cases}$, then $g(x)$ has

- (a) exactly one point of local minima
(b) exactly one point of local maxima
(c) no point to local maxima but exactly one point of local minima
(d) neither a point of local maxima nor minima

3 $f(x) = \begin{cases} 4x - x^3 + \log(a^2 - 3a + 3), & 0 \leq x < 3 \\ x - 18, & x \geq 3 \end{cases}$

Complete the set of values of a such that $f(x)$ has a local maxima at $x = 3$, is

- (a) $[-1, 2]$ (b) $(-\infty, 1) \cup (2, \infty)$
(c) $[1, 2]$ (d) $(-\infty, -1) \cup (2, \infty)$

4 The point in the interval $[0, 2\pi]$, where $f(x) = e^x \sin x$ has maximum slope is

- (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{2}$
(c) π (d) None of these

5 The total number of local maxima and local minima of the function $f(x) = \begin{cases} (2+x)^3, & -3 < x \leq -1 \\ x^{2/3}, & -1 < x < 2 \end{cases}$ is

- (a) 0 (b) 1 (c) 2 (d) 3

6 If 20 m of wire is available for fencing off a flower-bed in the form of a circular sector, then the maximum area (in sq m) of the flower-bed is → JEE Mains 2017

- (a) 12.5 (b) 10 (c) 25 (d) 30

7 The cost of running a bus from A to B, is ₹ $\left(av + \frac{b}{v}\right)$,

where v km/h is the average speed of the bus. When the bus travels at 30 km/h, the cost comes out to be ₹ 75

₹ 75 while at 40 km/h, it is ₹ 65. Then, the most economical speed (in km/h) of the bus is → JEE Mains 2013

- (a) 45 (b) 50 (c) 60 (d) 40

8 If $f(x) = \begin{cases} |x^2 - 2|, & -1 \leq x < \sqrt{3} \\ \frac{x}{\sqrt{3}}, & \sqrt{3} \leq x < 2\sqrt{3} \\ 3 - x, & 2\sqrt{3} \leq x \leq 4 \end{cases}$, then the points,

where $f(x)$ takes maximum and minimum values, are

- (a) 1, 4 (b) 0, 4
(c) 2, 4 (d) None of these

9 Let $f(x) = \begin{cases} |x^3 + x^2 + 3x + \sin x| \left(3 + \sin \frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$, then

number of points [where, $f(x)$ attains its minimum value] is

- (a) 1 (b) 2
(c) 3 (d) infinite many

10 A wire of length 2 units is cut into two parts which are bent respectively to form a square of side = x units and a circle of radius = r units. If the sum of the areas of the square and the circle so formed is minimum, then

- (a) $2x = (\pi + 4)r$ (b) $(4 - \pi)x = \pi r$
(c) $x = 2r$ (d) $2x = r$

11 Let $f(x) = x^2 + \frac{1}{x^2}$ and $g(x) = x - \frac{1}{x}$, $x \in \mathbb{R} - \{-1, 0, 1\}$. If

$h(x) = \frac{f(x)}{g(x)}$, then the local minimum value of $h(x)$ is

- (a) 3 (b) -3 (c) $-2\sqrt{2}$ (d) $2\sqrt{2}$ → JEE Mains 2018

12 The largest term in the sequence $a_n = \frac{n^2}{n^3 + 200}$ is given by


- (a) $\frac{529}{49}$ (b) $\frac{8}{89}$
(c) $\frac{49}{543}$ (d) None of these

13 All possible values of the parameter a so that the function $f(x) = x^3 - 3(7-a)x^2 - 3(9-a^2)x + 2$ has a negative point of local minimum are

- (a) all real values (b) no real values
(c) $(0, \infty)$ (d) $(-\infty, 0)$

- 14 The circle $x^2 + y^2 = 1$ cuts the X-axis at P and Q. Another circle with centre at Q and variable radius intersects the first circle at R above the X-axis and the line segment PQ at S. Then, the maximum area of the ΔQSR is

- (a) $4\sqrt{3}$ sq units
(b) $14\sqrt{3}$ sq units
(c) $\frac{4\sqrt{3}}{9}$ sq units
(d) $15\sqrt{3}$ sq units



- 15 Given, $P(x) = x^4 + ax^3 + bx^2 + cx + d$ such that $x = 0$ is the only real root of $P'(x) = 0$. If $P(-1) < P(1)$, then in the interval $[-1, 1]$.

→ AIEEE 2009

- (a) $P(-1)$ is the minimum and $P(1)$ is the maximum of P
(b) $P(-1)$ is not minimum but $P(1)$ is the maximum of P
(c) $P(-1)$ is the minimum and $P(1)$ is not the maximum of P
(d) Neither $P(-1)$ is the minimum nor $P(1)$ is the maximum of P

ANSWERS

SESSION 1

1 (d)	2 (a)	3 (d)	4 (a)	5 (b)	6 (d)	7 (d)	8 (d)	9 (a)	10 (c)
11 (c)	12 (c)	13 (b)	14 (b)	15 (d)	16 (b)	17 (b)	18 (c)	19 (a)	20 (a)
21 (b)	22 (a)	23 (b)	24 (d)	25 (c)	26 (b)	27 (a)	28 (d)	29 (b)	30 (b)
31 (a)	32 (b)	33 (c)	34 (b)	35 (b)	36 (a)	37 (b)	38 (b)	39 (b)	40 (d)

SESSION 2

1 (b)	2 (d)	3 (c)	4 (b)	5 (c)	6 (c)	7 (c)	8 (b)	9 (a)	10 (c)
11 (d)	12 (c)	13 (b)	14 (c)	15 (b)					

Hints and Explanations

SESSION 1

- 1 Let $y = x + \frac{1}{x} \Rightarrow \frac{dy}{dx} = 1 - \frac{1}{x^2}$

Now, $\frac{dy}{dx} = 0 \Rightarrow x^2 = 1$

$\Rightarrow x = \pm 1$

$\Rightarrow \frac{d^2y}{dx^2} = \frac{2}{x^3}$, therefore

$\frac{d^2y}{dx^2}$ (at $x = 1$) > 0

and $\frac{d^2y}{dx^2}$ (at $x = -1$) < 0

Hence, local maximum value of y is at $x = -1$ and the local maximum value $= -2$.

Local minimum value of y is at $x = 1$ and local minimum value $= 2$.

Therefore, local maximum value -2 is less than local minimum value 2 .

- 2 Let two numbers be x and $(3-x)$.

Then, product $P = x(3-x)^2$

$\frac{dP}{dx} = -2x(3-x) + (3-x)^2$

$\frac{dP}{dx} = (3-x)(3-3x)$ and $\frac{d^2P}{dx^2} = 6x - 12$

For maxima or minima, put $\frac{dP}{dx} = 0$

$\Rightarrow (3-x)(3-3x) = 0 \Rightarrow x = 3, 1$

At $x = 3$,
 $\frac{d^2P}{dx^2} = 18 - 12 = 6 > 0$ [minima]

At $x = 1$,
 $\frac{d^2P}{dx^2} = -6 < 0$

So, P is maximum at $x = 1$.

\therefore Maximum value of $P = 1(3-1)^2 = 4$

3 $\therefore \frac{dy}{dx} = \frac{a}{x} + 2bx + 1$

$\Rightarrow \left(\frac{dy}{dx}\right)_{x=1} = a + 2b + 1 = 0$

$\Rightarrow a = -2b - 1$

and $\left(\frac{dy}{dx}\right)_{x=2} = \frac{a}{2} + 4b + 1 = 0$

$\Rightarrow \frac{-2b-1}{2} + 4b + 1 = 0$

$\Rightarrow -b + 4b + \frac{1}{2} = 0 \Rightarrow 3b = -\frac{1}{2}$

$\Rightarrow b = -\frac{1}{6}$ and $a = \frac{1}{3} - 1 = -\frac{2}{3}$

4 $f'(x) = -a \sin x + b \sec^2 x + 1$

Now, $f'(0) = 0$ and $f'\left(\frac{\pi}{6}\right) = 0$

$\Rightarrow b + 1 = 0$ and $-\frac{a}{2} + \frac{4b}{3} + 1 = 0$

$\Rightarrow b = -1, a = -\frac{2}{3}$

- 5 The given curve is $\frac{4}{x^2} + \frac{9}{y^2} = 1$

Put $x = r \cos \theta$, $y = r \sin \theta$, we get

$r^2 = (2 \sec \theta)^2 + (3 \csc \theta)^2$

So, r^2 will have minimum value $(2 + 3)^2$.

or r have minimum value equal to 5.

6 $f(x) = 4x^3 - 18x^2 + 27x - 7$

$f'(x) = 12x^2 - 36x + 27$

$= 3(4x^2 - 12x + 9) = 3(2x - 3)^2$

$f'(x) = 0 \Rightarrow x = \frac{3}{2}$ (critical point)

Since, $f'(x) > 0$ for all $x < \frac{3}{2}$ and for all

$x > \frac{3}{2}$

Hence, $x = \frac{3}{2}$ is a point of inflection i.e.,

neither a point of maxima nor a point of minima.

$x = \frac{3}{2}$ is the only critical point and f

has neither maxima nor minima.

7 For $y = \frac{x^2 - 2}{x^2 - 4} \Rightarrow \frac{dy}{dx} = \frac{-4x}{(x^2 - 4)^2}$

$$\Rightarrow \frac{dy}{dx} > 0, \text{ for } x < 0$$

$$\text{and } \frac{dy}{dx} < 0, \text{ for } x > 0$$


Thus, $x = 0$ is the point of local maxima for y . Now, $(y')_{x=0} = \frac{1}{2}$ (positive). Thus,

$x = 0$ is also the point of local

$$\text{maximum for } y = \left| \frac{x^2 - 2}{x^2 - 4} \right|$$

- 8** If $f(x)$ has a local minimum at $x = -1$, then

$$\begin{aligned} \lim_{x \rightarrow -1^+} f(x) &= \lim_{x \rightarrow -1^-} f(x) \\ \Rightarrow \lim_{x \rightarrow -1^+} 2x + 3 &= \lim_{x \rightarrow -1^-} 1 < -2x \\ \Rightarrow -2 + 3 &= k + 2 \Rightarrow k = -1 \end{aligned}$$



- 9** Let $S = 9x + 4y$

Since, $xy = 16$ is given.

$$\therefore y = \frac{16}{x} \text{ or } S = 9x + \frac{64}{x}$$

On differentiating both sides, we get

$$\frac{dS}{dx} = 9 - \frac{64}{x^2} \quad \dots(i)$$

$$\therefore \frac{dS}{dx} = 0 \Rightarrow \frac{64}{x^2} = 9 \Rightarrow x = \pm \frac{8}{3}$$

Again, on differentiating Eq. (i)

$$\text{w.r.t. } x, \text{ we get } \frac{d^2S}{dx^2} = \frac{128}{x^3}$$

Hence, it is minimum at $x = \frac{8}{3}$ and minimum value of S is

$$S_{\min} = 9\left(\frac{8}{3}\right) + 4(6) = 48$$

- 10** We have,

$$f(x) = 2x^3 - 9ax^2 + 12a^2x + 1$$

$$f'(x) = 6x^2 - 18ax + 12a^2$$

$$f''(x) = 12x - 18a$$

For maximum and minimum,

$$6x^2 - 18ax + 12a^2 = 0$$

$$\Rightarrow x^2 - 3ax + 2a^2 = 0$$

$$\Rightarrow x = a \text{ or } x = 2a$$

At $x = a$ maximum and at $x = 2a$ minimum.

$$\therefore p^2 = q$$

$$\therefore a^2 = 2a \Rightarrow a = 2 \text{ or } a = 0$$

But $a > 0$, therefore $a = 2$

- 11** Minimum of $f(x) = -\frac{D}{4a}$
- $$= \frac{-(4b^2 - 8c^2)}{4}$$
- $$= 2c^2 - b^2$$

$$\text{and maximum of } g(x) = -\frac{(4c^2 + 4b^2)}{4(-1)}$$

$$= b^2 + c^2$$

Since, $\min f(x) > \max g(x)$

$$\Rightarrow 2c^2 - b^2 > b^2 + c^2$$

$$\Rightarrow c^2 > 2b^2$$

$$\Rightarrow |c| > \sqrt{2}|b|$$

- 12 Central Idea** Any function have extreme values (maximum or minimum) at its critical points, where $f'(x) = 0$. Since, the function have extreme values at $x = 1$ and $x = 2$.

$$\therefore f'(x) = 0 \text{ at } x = 1 \text{ and } x = 2$$

$$\Rightarrow f'(1) = 0 \text{ and } f'(2) = 0$$

Also it is given that

$$\lim_{x \rightarrow 0} \left[1 + \frac{f(x)}{x^2} \right] = 3 \Rightarrow 1 + \lim_{x \rightarrow 0} \frac{f(x)}{x^2} = 3$$

$$\Rightarrow \lim_{x \rightarrow 0} \frac{f(x)}{x^2} = 2$$

$\Rightarrow f(x)$ will be of the form

$$ax^4 + bx^3 + 2x^2$$

[$\because f(x)$ is of four degree polynomial]

$$\text{Let } f(x) = ax^4 + bx^3 + 2x^2 \Rightarrow f'(x)$$

$$= 4ax^3 + 3bx^2 + 4x$$

$$\Rightarrow f'(1) = 4a + 3b + 4 = 0 \quad \dots(i)$$

$$\text{and } f'(2) = 32a + 12b + 8 = 0$$

$$\Rightarrow 8a + 3b + 2 = 0 \quad \dots(ii)$$

On solving Eqs. (i) and (ii), we get

$$a = \frac{1}{2}, b = -2$$

$$\therefore f(x) = \frac{x^4}{2} - 2x^3 + 2x^2$$

$$\Rightarrow f(2) = 8 - 16 + 8 = 0$$

- 13** $\phi'(x) = f'(x) + a$

$$\therefore \phi'(0) = 0 \Rightarrow f'(0) + a = 0$$

$$\Rightarrow a = 0 \quad [\because f'(0) = 0]$$

Also, $\phi'(0) > 0$ [$\because f''(0) > 0$]

$\Rightarrow \phi(x)$ has relative minimum at $x = 0$ for all b , if $a = 0$

- 14** Let the number be x , then

$$f(x) = \frac{x}{x^2 + 16}$$

On differentiating w.r.t. x , we get

$$\begin{aligned} f'(x) &= \frac{(x^2 + 16) \cdot 1 - x(2x)}{(x^2 + 16)^2} \\ &= \frac{x^2 + 16 - 2x^2}{(x^2 + 16)^2} = \frac{16 - x^2}{(x^2 + 16)^2} \quad \dots(i) \end{aligned}$$

Put $f'(x) = 0$ for maxima or minima

$$f'(x) = 0 \Rightarrow 16 - x^2 = 0$$

$$\Rightarrow x = 4, -4$$

Again, on differentiating w.r.t. x , we get

$$\begin{aligned} (x^2 + 16)^2 (-2x) - (16 - x^2) \\ f''(x) = \frac{2(x^2 + 16)2x}{(x^2 + 16)^4} \text{ At} \end{aligned}$$

$$x = 4, f''(x) < 0$$

$\therefore f(x)$ is maximum at $x = 4$.

and at $x = -4$, $f''(x) > 0$, $f(x)$ is minimum.

\therefore Least value of

$$f(x) = \frac{-4}{16 + 16} = -\frac{1}{8}$$

- 15** Given, $f(x) = ax + \frac{b}{x}$

On differentiating w.r.t. x , we get

$$f'(x) = a - \frac{b}{x^2}$$

For maxima or minima, put $f'(x) = 0$

$$\Rightarrow x = \sqrt{\frac{b}{a}}$$

Again, differentiating w.r.t. x , we get

$$f''(x) = \frac{2b}{x^3}$$

At $x = \sqrt{\frac{b}{a}}$, $f''(x)$ is positive

$\Rightarrow f(x)$ is minimum at $x = \sqrt{\frac{b}{a}}$.

$\therefore f(x)$ has the least value at $x = \sqrt{\frac{b}{a}}$.

- 16** $f(x) = \begin{cases} \frac{\tan x}{x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$

$$\text{As } \frac{\tan x}{x} > 1, \forall x \neq 0$$

$\therefore f(0 + h) > f(0)$ and $f(0 - h) > f(0)$

At $x = 0$, $f(x)$ attains minima.

$$\text{Now, } f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{\tan h}{h} - 1}{h} = \lim_{h \rightarrow 0} \frac{\tan h - h}{h^2}$$

[using L' Hospital's rule]

$$= \lim_{h \rightarrow 0} \frac{\sec^2 h - 1}{2h} \quad [\because \tan^2 \theta = \sec^2 \theta - 1]$$

$$= \lim_{h \rightarrow 0} \frac{\tan^2 h}{2h^2} \cdot h = \frac{1}{2} \cdot 0 = 0$$

Therefore, Statement II is true.

Hence, both statements are true but Statement II is not the correct explanation of Statement I.

- 17** Given, $f(x) = \cos^2 x + \sin x$, $x \in [0, \pi]$

Now,

$$\begin{aligned} f'(x) &= 2 \cos x (-\sin x) + \cos x \\ &= -2 \sin x \cos x + \cos x \end{aligned}$$

For maximum or minimum put $f'(x) = 0$

$$\Rightarrow -2 \sin x \cos x + \cos x = 0$$

$$\Rightarrow \cos x (-2 \sin x + 1) = 0$$

$$\Rightarrow \cos x = 0 \text{ or } \sin x = \frac{1}{2}$$

$$\Rightarrow x = \frac{\pi}{6}, \frac{\pi}{2}$$

For absolute maximum and absolute minimum, we have to evaluate

$$f(0), f\left(\frac{\pi}{6}\right), f\left(\frac{\pi}{2}\right), f(\pi)$$

At $x = 0$,

$$f(0) = \cos^2 0 + \sin 0 = 1^2 + 0 = 1$$

$$\begin{aligned} \text{At } x = \frac{\pi}{6}, f\left(\frac{\pi}{6}\right) &= \cos^2\left(\frac{\pi}{6}\right) + \sin \frac{\pi}{6} \\ &= \left(\frac{\sqrt{3}}{2}\right)^2 + \frac{1}{2} = \frac{5}{4} = 1.25 \end{aligned}$$

At $x = \frac{\pi}{2}$,

$$f\left(\frac{\pi}{2}\right) = \cos^2\left(\frac{\pi}{2}\right) + \sin \frac{\pi}{2} = 0^2 + 1 = 1$$

At $x = \pi$,

$$f(\pi) = \cos^2 \pi + \sin \pi = (-1)^2 + 0 = 1$$

Hence, the absolute maximum value of f is 1.25 occurring at $x = \frac{\pi}{6}$ and the absolute minimum value of f is 1 occurring at $x = 0, \frac{\pi}{2}$ and π .

Note If close interval is given, to determine global maximum (minimum), check the value at all critical points as well as end points of a given interval.

$$18 \because f(x) = \frac{x}{4 + x + x^2}$$

On differentiating w.r.t. x , we get

$$f'(x) = \frac{4 + x + x^2 - x(1 + 2x)}{(4 + x + x^2)^2}$$

For maximum, put $f'(x) = 0$

$$\Rightarrow \frac{4 - x^2}{(4 + x + x^2)^2} = 0 \Rightarrow x = 2, -2$$

Both the values of x are not in the interval $[-1, 1]$.

$$\therefore f(-1) = \frac{-1}{4 - 1 + 1} = \frac{-1}{4}$$

$$f(1) = \frac{1}{4 + 1 + 1} = \frac{1}{6} \text{ (maximum)}$$

$$19 \text{ Given, } f(x) = x^2 \log x$$

On differentiating w.r.t. x , we get

$$f'(x) = (2 \log x + 1)x$$

For a maximum, put $f'(x) = 0$

$$\Rightarrow (2 \log x + 1)x = 0$$

$$\Rightarrow x = e^{-1/2}, 0$$

$$\because 0 < e^{-1/2} < 1$$

None of these critical points lies in the interval $[1, e]$.

So, we only compute the value of $f(x)$ at the end points 1 and e .

$$\text{We have, } f(1) = 0, f(e) = e^2$$

Hence, greatest value of $f(x) = e^2$

$$20 \text{ Let } f(x) = 1 + \frac{10}{3\left(x^2 + 3x + \frac{7}{3}\right)}$$

$$= 1 + \frac{10}{3\left[\left(x + \frac{3}{2}\right)^2 + \frac{1}{12}\right]}$$

So, the maximum value of $f(x)$ at

$$x = -\frac{3}{2} \text{ is}$$

$$f\left(-\frac{3}{2}\right) = 1 + \frac{10}{3\left(\frac{1}{12}\right)} = 1 + 40 = 41$$

$$21 \text{ Given, } f(x) = \sec x + \log \cos^2 x$$

$$\Rightarrow f(x) = \sec x + 2 \log(\cos x)$$

Therefore,

$$\begin{aligned} f'(x) &= \sec x \tan x - 2 \tan x \\ &= \tan x (\sec x - 2) \end{aligned}$$

$$f'(x) = 0$$

$$\Rightarrow \tan x = 0 \text{ or } \sec x = 2 \Rightarrow \cos x = \frac{1}{2}$$

Therefore, possible values of x are

$$x = 0, x = \pi \text{ and } x = \frac{\pi}{3} \text{ or } x = \frac{5\pi}{3}.$$

$$\text{Again, } f''(x) = \sec^2 x (\sec x - 2)$$

$$+ \tan x (\sec x \tan x)$$

$$= \sec^3 x + \sec x \tan^2 x - 2 \sec^2 x$$

$$= \sec x (\sec^2 x + \tan^2 x - 2 \sec x)$$

$$\Rightarrow f''(0) = 1(1 + 0 - 2) = -1 < 0$$

Therefore, $x = 0$ is a point of maxima.

$$f''(\pi) = -1(1 + 0 + 2) = -3 < 0$$

Therefore, $x = \pi$ is a point of maxima.

$$f''\left(\frac{\pi}{3}\right) = 2(4 + 3 - 4) = 6 > 0$$

Therefore, $x = \frac{\pi}{3}$ is a point of minima.

$$f''\left(\frac{5\pi}{3}\right) = 2(4 + 3 - 4) = 6 > 0$$

Therefore, $x = \frac{5\pi}{3}$ is a point of minima.

Maximum value of y at $x = 0$ is

$$1 + 0 = 1.$$

Maximum value of y at $x = \pi$ is

$$-1 + 0 = -1.$$

Minimum value of y at $x = \frac{\pi}{3}$ is

$$2 + 2 \log \frac{1}{2} = 2(1 - \log 2).$$

Minimum value of y at $x = \frac{5\pi}{3}$ is

$$2 + 2 \log \frac{1}{2} = 2(1 - \log 2).$$

$$22 \text{ Given, } f(x) = \sin 2x - x$$

$$\Rightarrow f'(x) = 2 \cos 2x - 1$$

$$\text{Put } f'(x) = 0 \Rightarrow \cos 2x = \frac{1}{2}$$

$$\Rightarrow 2x = -\frac{\pi}{3} \text{ or } \frac{\pi}{3} \Rightarrow x = -\frac{\pi}{6} \text{ or } \frac{\pi}{6}$$

$$\text{Now, } f\left(-\frac{\pi}{6}\right) = \sin\left(-\frac{\pi}{3}\right) + \frac{\pi}{6} = -\frac{\sqrt{3}}{2} + \frac{\pi}{6}$$

$$f\left(-\frac{\pi}{6}\right) = \sin\left(-\frac{2\pi}{6}\right) + \frac{\pi}{6} = -\frac{\sqrt{3}}{2} + \frac{\pi}{6}$$

$$f\left(\frac{\pi}{6}\right) = \sin\left(\frac{2\pi}{6}\right) - \frac{\pi}{6} = \frac{\sqrt{3}}{2} - \frac{\pi}{6}$$

$$\text{and } f\left(\frac{\pi}{2}\right) = \sin(\pi) - \frac{\pi}{2} = -\frac{\pi}{2}$$

Clearly, $\frac{\pi}{2}$ is the greatest value and $-\frac{\pi}{2}$ is the least.

$$\text{Therefore, difference} = \frac{\pi}{2} + \frac{\pi}{2} = \pi$$

$$23 \text{ Given, } y = x^{3/2}$$

$$\therefore \frac{dy}{dx} = \frac{3}{2} x^{1/2}, \frac{d^2y}{dx^2} = \frac{15}{4} x^{-1/2}$$

$$\text{At } x = 0, \frac{dy}{dx} = 0, \frac{d^2y}{dx^2} = 0$$

and $\frac{d^3y}{dx^3}$ is not defined,


when $x = 0, y = 0$

$\therefore (0, 0)$ is a point of inflection.

$$24 \text{ Area of triangle, } \Delta = \frac{1}{2} x \sqrt{h^2 - x^2}$$

$$\frac{d\Delta}{dx} = \frac{1}{2} \left[\sqrt{h^2 - x^2} + \frac{x(-2x)}{2\sqrt{h^2 - x^2}} \right] = 0$$

$$\Rightarrow x = \frac{h}{\sqrt{2}}$$



$$\Rightarrow \frac{d^2\Delta}{dx^2} < 0 \text{ at } x = \frac{h}{\sqrt{2}}$$

$$\therefore \Delta = \frac{1}{2} \times \frac{h}{\sqrt{2}} \sqrt{h^2 - \frac{h^2}{2}} = \frac{h^2}{4}$$

$$25 \text{ Let the equation of drawn line be}$$

$$\frac{x}{a} + \frac{y}{b} = 1, \text{ where } a > 3,$$

$b > 4$, as the line passes through $(3, 4)$ and meets the positive direction of coordinate axes.

$$\text{We have, } \frac{3}{a} + \frac{4}{b} = 1 \Rightarrow b = \frac{4a}{a-3}$$

Now, area of ΔAOB ,


$$\Delta = \frac{1}{2} ab = \frac{2a^2}{a-3}$$

$$\frac{d\Delta}{da} = \frac{2a(a-6)}{(a-3)^2}$$

Clearly, $a = 6$ is the point of minima for Δ .


$$\text{Thus, } \Delta_{\min} = \frac{2 \times 36}{3} = 24 \text{ sq units}$$

$$26 \text{ Let } f(x) = x^3 - px + q$$



Then, $f'(x) = 3x^2 - p$
 Put $f'(x) = 0$
 $\Rightarrow x = \sqrt{\frac{p}{3}}, -\sqrt{\frac{p}{3}}$
 Now, $f''(x) = 6x$
 At $x = \sqrt{\frac{p}{3}}$, $f''(x) = 6\sqrt{\frac{p}{3}} > 0$ [minima]
 and at $x = -\sqrt{\frac{p}{3}}$, $f''(x) < 0$ [maxima]

27 We have, $AF \parallel DE$ and $AE \parallel FD$



Now, in $\triangle ABC$ and $\triangle DEC$,
 $\angle DEC = \angle BAC$, $\angle ACB$ is common.

$\Rightarrow \triangle ABC \cong \triangle DEC$
 Now, $\frac{b-y}{b} = \frac{x}{c} \Rightarrow x = \frac{c}{b}(b-y)$


Now, $S = \text{Area of parallelogram } AFDE = 2 (\text{area of } \triangle AEF)$

$$\Rightarrow S = 2 \left(\frac{1}{2} xy \sin A \right)$$

$$= \frac{c}{b} (b-y)y \sin A$$

$$\frac{dS}{dy} = \left(\frac{c}{b} \sin A \right) (b - 2y)$$

Sign scheme of $\frac{dS}{dy}$,



Hence, S is maximum when $y = \frac{b}{2}$.

$$\therefore S_{\max} = \frac{c}{b} \left(\frac{b}{2} \right) \times \frac{b}{2} \sin A$$

$$= \frac{1}{2} \left(\frac{1}{2} bc \sin A \right) = \frac{1}{2} (\text{area of } \triangle ABC)$$

28 We have,

$$\frac{dy}{dt} = 6t^2 - 30t + 24 = 6(t-1)(t-4)$$

$$\text{and } \frac{dx}{dt} = 6t - 18 = 6(t-3)$$

$$\text{Thus, } \frac{dy}{dx} = \frac{(t-1)(t-4)}{(t-3)}$$

which indicates that $t = 1, 3$ and 4 are the critical points of $y = f(x)$.

$$\text{Now, } \frac{d^2y}{dx^2} = \frac{d}{dt} \left(\frac{dy}{dx} \right) \cdot \frac{dt}{dx}$$

$$= \frac{t^2 - 6t + 11}{(t-3)^2} \times \frac{1}{6(t-3)}$$

At $(t=1)$, $\frac{d^2y}{dx^2} < 0$
 $\Rightarrow t = 1$ is a point of local maxima.

At $(t=4)$, $\frac{d^2y}{dx^2} > 0$
 $\Rightarrow t = 4$ is a point of local minima.

At $(t=3)$, $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ are not defined and change its sign.

$\frac{d^2y}{dx^2}$ is unknown in the vicinity of $t = 3$, thus $t = 3$ is a point of neither maxima nor minima.

Finally, maximum and minimum values of expression $y = f(x)$ are 46 and -6 , respectively.

29 Let α and β be the roots of the equation

$$x^2 - (a-2)x - a + 1 = 0$$

Then, $\alpha + \beta = a - 2$, $\alpha\beta = -a + 1$

$$\text{Let } z = \alpha^2 + \beta^2$$

$$= (\alpha + \beta)^2 - 2\alpha\beta$$

$$= (a-2)^2 + 2(a-1)$$

$$= a^2 - 2a + 2$$

$$\Rightarrow \frac{dz}{da} = 2a - 2$$

Put $\frac{dz}{da} = 0$, then

$$\Rightarrow a = 1$$

$$\therefore \frac{d^2z}{da^2} = 2 > 0$$

So, z has minima at $a = 1$.

So, $\alpha^2 + \beta^2$ has least value for $a = 1$.

This is because we have only one stationary value at which we have minima.

Hence, $a = 1$.

30 Any tangent to the ellipse is

$\frac{x}{4} \cos t + \frac{y}{3} \sin t = 1$, where the point of contact is $(4 \cos t, 3 \sin t)$

$$\text{or } \frac{x}{4 \sec t} + \frac{y}{3 \csc t} = 1,$$


It means the axes $Q(4 \sec t, 0)$ and $R(0, 3 \csc t)$.

\therefore The distance of the line segment QR is

$$QR^2 = D = 16 \sec^2 t + 9 \csc^2 t$$

So, the minimum value of D is $(4 + 3)^2$ or $QR = 7$.

31 Let S be the curved surface area of a cone.



$$OA = AC - OC = h - R$$

$$\text{In } \triangle OAB, R^2 = r^2 + (h - R)^2$$

$$\Rightarrow r = \sqrt{2Rh - h^2}$$

$$\therefore S = \pi r l = \pi (\sqrt{2Rh - h^2}) (\sqrt{h^2 + r^2})$$

$$= (\pi \sqrt{2Rh - h^2}) (\sqrt{2Rh})$$

Let $S^2 = P$

$$\therefore P = \pi^2 2R(2Rh^2 - h^3)$$

Since, S is maximum, if P is maximum, then

$$\frac{dP}{dh} = 2\pi^2 R(4Rh - 3h^2) = 0$$

$$\therefore h = 0, \frac{4R}{3}$$

Again, on differentiating $\frac{dP}{dh}$, we get

$$\frac{d^2P}{dh^2} = 2\pi^2 R(4R - 6h)$$

$$\frac{d^2P}{dh^2} < 0 \text{ at } h = \frac{4R}{3}$$


32 Let $OC = x$, $CQ = r$

Now, $OA = R$ [given]

Height of the cone = $h = x + R$

\therefore Volume of the cone

$$= V = \frac{1}{3} \pi r^2 h \quad \dots(i)$$



Also, in right angled $\triangle OCQ$,

$$OC^2 + CQ^2 = OQ^2$$

$$\Rightarrow x^2 + r^2 = R^2$$

$$\Rightarrow r^2 = R^2 - x^2 \quad \dots(ii)$$

From Eqs. (i) and (ii),

$$V = \frac{1}{3} \pi (R^2 - x^2)(x + R) \quad \dots(iii)$$

$$[\because h = x + R]$$

On differentiating Eq. (iii) w.r.t. x , we get

$$\frac{dV}{dx} = \frac{1}{3} \pi [(R^2 - x^2) - 2x(x + R)]$$

$$\Rightarrow \frac{dV}{dx} = \frac{\pi}{3} (R^2 - x^2 - 2x^2 - 2xR)$$

$$\Rightarrow \frac{dV}{dx} = \frac{\pi}{3} (R^2 - 2xR - 3x^2)$$

$$\Rightarrow \frac{dV}{dx} = \frac{\pi}{3} (R - 3x)(R + x) \quad \dots(iv)$$

For maxima, put $\frac{dV}{dx} = 0$

$$\Rightarrow \frac{\pi}{3} (R - 3x)(R + x) = 0$$

$$\Rightarrow x = \frac{R}{3} \text{ or } x = -R \Rightarrow x = \frac{R}{3}$$

[since, x cannot be negative]

On differentiating Eq. (iv) w.r.t. x , we get

$$\frac{d^2V}{dx^2} = \frac{\pi}{3} [(-3)(R+x) + (R-3x)]$$

$$= \frac{\pi}{3} (-2R-6x) = -\frac{\pi}{3} (2R+6x)$$

$$\text{At } x = \frac{R}{3}, \frac{d^2V}{dx^2} = -\frac{\pi}{3} \left(2R + \frac{6R}{3} \right)$$

$$= -\frac{4\pi}{3} R < 0$$

So, V has a local maxima at $x = R/3$.


Now, on substituting the value of x in Eq. (iii), we get

$$V = \frac{\pi}{3} \left(R^2 - \frac{R^2}{9} \right) \left(R + \frac{R}{3} \right)$$

$$= \frac{\pi}{3} \cdot \frac{8R^2}{9} \cdot \frac{4R}{3} = \frac{8}{27} \left(\frac{4}{3} \pi R^3 \right)$$

$$\Rightarrow V = \frac{8}{27} \times \text{Volume of sphere}$$

33



Area of rectangle $ABCD$

$$= (2a \cos \theta)(2b \sin \theta) = 2ab \sin 2\theta$$

Hence, area of greatest rectangle is equal to $2ab$ when $\sin 2\theta = 1$.

34 Let

$$f(x) = x + \frac{1}{x}$$

$$f'(x) = 1 - \frac{1}{x^2}$$

For maxima and minima, put $f'(x) = 0$

$$\Rightarrow 1 - \frac{1}{x^2} = 0 \Rightarrow x = \pm 1$$

$$\text{Now, } f''(x) = \frac{2}{x^3}$$

At $x = 1$, $f''(x) = +ve$ [minima]
and at $x = -1$, $f''(x) = -ve$ [maxima]

Thus, $f(x)$ attains minimum value at $x = 1$.

35 Given that, $f(x) = (x+1)^{1/3} - (x-1)^{1/3}$

On differentiating w.r.t. x , we get

$$f'(x) = \frac{1}{3} \left[\frac{1}{(x+1)^{2/3}} - \frac{1}{(x-1)^{2/3}} \right]$$

$$= \frac{(x-1)^{2/3} - (x+1)^{2/3}}{3(x^2-1)^{2/3}}$$

Clearly, $f'(x)$ does not exist at $x = \pm 1$.

Now, put $f'(x) = 0$, then

$$(x-1)^{2/3} = (x+1)^{2/3} \Rightarrow x = 0$$

At $x = 0$

$$f(x) = (0+1)^{1/3} - (0-1)^{1/3} = 2$$

Hence, the greatest value of $f(x)$ is 2.

36 $\because y^2 = 8x$. But $y^2 = 4ax$

$$\Rightarrow 4a = 8 \Rightarrow a = 2$$

Any point on parabola is $(at^2, 2at)$, i.e., $(2t^2, 4t)$.

For its minimum distance from the circle means its distance from the centre $(0, -6)$ of the circle.

$$\text{Let } z = (2t^2)^2 + (4t+6)^2$$

$$= 4(t^4 + 4t^2 + 12t + 9)$$

$$\therefore \frac{dz}{dt} = 4(4t^3 + 8t + 12)$$

$$\Rightarrow 16(t^3 + 2t + 3) = 0$$

$$\Rightarrow (t+1)(t^2-t+3) = 0$$


$$\Rightarrow t = -1$$

$$\Rightarrow \frac{d^2z}{dt^2} = 16(3t^2 + 2) > 0, \text{ hence minimum.}$$

So, point is $(2, -4)$.

37 We know that, volume of cylinder,

$$V = \pi R^2 h$$



$$\text{In } \triangle OCA, r^2 = \left(\frac{h}{2} \right)^2 + R^2$$

$$\Rightarrow R^2 = r^2 - \frac{h^2}{4}$$

$$\therefore V = \pi \left(r^2 - \frac{h^2}{4} \right) h$$

$$\Rightarrow V = \pi r^2 h - \frac{\pi}{4} h^3 \dots (i)$$

On differentiating Eq. (i) both sides w.r.t. h , we get

$$\frac{dV}{dh} = \pi r^2 - \frac{3\pi h^2}{4}$$

$$\Rightarrow \frac{d^2V}{dh^2} = -\frac{3\pi h}{2}$$

For maximum or minimum value of V ,

$$\frac{dV}{dh} = 0 \Rightarrow \pi r^2 - \frac{3\pi h^2}{4} = 0$$

$$\Rightarrow h^2 = \frac{4r^2}{3} \Rightarrow h = \frac{2}{\sqrt{3}} r$$

$$\text{Now, } \left(\frac{d^2V}{dh^2} \right)_{h=\frac{2r}{\sqrt{3}}} = -\sqrt{3}\pi r < 0$$

Thus, V is maximum when $h = \frac{2r}{\sqrt{3}}$, then

$$R^2 = r^2 - \frac{h^2}{4} = r^2 - \frac{1}{4} \left(\frac{2r}{\sqrt{3}} \right)^2 = \frac{2}{3} r^2$$

$$\text{Max } V = \pi R^2 h = \frac{4\pi r^3}{3\sqrt{3}}$$

38 Let $f(x) = -x^3 + 3x^2 + 9x - 27$

The slope of this curve

$$f'(x) = -3x^2 + 6x + 9$$

$$\text{Let } g(x) = f'(x) = -3x^2 + 6x + 9$$

On differentiating w.r.t. x , we get

$$g'(x) = -6x + 6$$

For maxima or minima put $g'(x) = 0$
 $\Rightarrow x = 1$

Now, $g''(x) = -6 < 0$ and hence, at $x = 1$, $g(x)$ (slope) will have maximum value.

$$\therefore [g(1)]_{\max} = -3 \times 1 + 6(1) + 9 = 12$$

39 Given,

$$ab = 2a + 3b \Rightarrow (a-3)b = 2a$$

$$\Rightarrow b = \frac{2a}{a-3}$$

$$\text{Now, let } z = ab = \frac{2a^2}{a-3}$$

On differentiating w.r.t. x , we get

$$\frac{dz}{da} = \frac{2[(a-3)2a - a^2]}{(a-3)^2} = \frac{2[a^2 - 6a]}{(a-3)^2}$$

$$\text{For a minimum, put } \frac{dz}{da} = 0$$

$$\Rightarrow a^2 - 6a = 0$$

$$\Rightarrow a = 0, 6$$

$$\text{At } a = 6, \frac{d^2z}{da^2} = \text{positive}$$

$$\text{When } a = 6, b = 4$$

$$\therefore (ab)_{\min} = 6 \times 4 = 24$$

40 \therefore Perimeter of a sector = p

Let AOB be the sector with radius r .

If angle of the sector be θ radians, then area of sector,

$$A = \frac{1}{2} r^2 \theta \dots (i)$$

$$\text{and length of arc, } s = r\theta \Rightarrow \theta = \frac{s}{r}$$

\therefore Perimeter of the sector

$$p = r + s + r = 2r + s \dots (ii)$$

On substituting $\theta = \frac{s}{r}$ in Eq. (i), we get

$$A = \left(\frac{1}{2} r^2 \right) \left(\frac{s}{r} \right) = \frac{1}{2} rs \Rightarrow s = \frac{2A}{r}$$

Now, on substituting the value of s in Eq. (ii), we get

$$p = 2r + \left(\frac{2A}{r} \right) \Rightarrow 2A = pr - 2r^2$$

On differentiating w.r.t. r , we get


$$2 \frac{dA}{dr} = p - 4r$$

For the maximum area, put

$$\frac{dA}{dr} = 0$$

$$\Rightarrow p - 4r = 0$$

$$\Rightarrow r = \frac{p}{4}$$



SESSION 2

- 1 Let radius vector is r .

$$\therefore r^2 = x^2 + y^2$$

$$\Rightarrow r^2 = \frac{a^2 y^2}{y^2 - b^2} + y^2 \quad \left(\because \frac{a^2}{x^2} + \frac{b^2}{y^2} = 1 \right)$$

For minimum value of r ,


$$\frac{d(r^2)}{dy} = 0 \Rightarrow \frac{-2yb^2a^2}{(y^2 - b^2)^2} + 2y = 0$$

$$\Rightarrow y^2 = b(a + b)$$

$$\therefore x^2 = a(a + b)$$

$$\Rightarrow r^2 = (a + b)^2 \Rightarrow r = a + b$$

- 2 Bold line represents the graph of $y = g(x)$, clearly $g(x)$ has neither a point of local maxima nor a point of local minima.



- 3 Clearly, $f(x)$ in increasing just before $x = 3$ and decreasing after $x = 3$. For $x = 3$ to be the point of local maxima, $f(3) \geq f(3 - 0)$
- $$\Rightarrow -15 \geq 12 - 27 + \log(a^2 - 3a + 3)$$
- $$\Rightarrow 0 < a^2 - 3a + 3 \leq 1 \Rightarrow 1 \leq a \leq 2$$

- 4 (Slope) $f'(x) = e^x \cos x + \sin x e^x$
- $$= e^x \sqrt{2} \sin(x + \pi/4)$$
- $$f''(x) = \sqrt{2}e^x \{\sin(x + \pi/4) + \cos(x + \pi/4)\}$$
- $$= 2e^x \cdot \sin(x + \pi/2)$$

For maximum slope, put $f''(x) = 0$

$$\Rightarrow \sin(x + \pi/2) = 0$$

$$\Rightarrow \cos x = 0$$


$$\therefore x = \pi/2, 3\pi/2$$

$$f'''(x) = 2e^x \cos(x + \pi/2)$$

$$f'''(\pi/2) = 2e^x \cdot \cos \pi = -ve$$

Maximum slope is at $x = \pi/2$.


- 5 $f'(x) = \begin{cases} 3(2 + x)^2, & -3 < x \leq -1 \\ \frac{2}{3}x^{-1/3}, & -1 < x < 2 \end{cases}$



Clearly, $f'(x)$ changes its sign at $x = -1$ from positive to negative and so $f(x)$ has local maxima at $x = -1$.

Also, $f'(0)$ does not exist but $f'(0^-) < 0$ and $f'(0^+) > 0$. It can only be inferred that $f(x)$ has a possibility of a minimum at $x = 0$. Hence, it has one local maxima at $x = -1$ and one local minima at $x = 0$. So, total number of local maxima and local minima is 2.

- 6 Total length = $2r + r\theta = 20$



$$\Rightarrow \theta = \frac{20 - 2r}{r}$$

Now, area of flower-bed,

$$A = \frac{1}{2} r^2 \theta$$

$$\Rightarrow A = \frac{1}{2} r^2 \left(\frac{20 - 2r}{r} \right)$$

$$\Rightarrow A = 10r - r^2$$

$$\therefore \frac{dA}{dr} = 10 - 2r$$

For maxima or minima, put $\frac{dA}{dr} = 0$.

$$\Rightarrow 10 - 2r = 0 \Rightarrow r = 5$$

$$\therefore A_{\max} = \frac{1}{2} (5)^2 \left[\frac{20 - 2(5)}{5} \right]$$

$$= \frac{1}{2} \times 25 \times 2 = 25 \text{ sq m}$$

- 7 Let $c = av + \frac{b}{v}$... (i)

When $v = 30$ km/h, then $c = ₹ 75$

$$\therefore 75 = 30a + \frac{b}{30} \quad \dots (ii)$$

When $v = 40$ km/h, then $c = ₹ 65$

$$\therefore 65 = 40a + \frac{b}{40} \quad \dots (iii)$$

On solving Eqs. (ii) and (iii), we get

$$a = \frac{1}{2} \quad \text{and} \quad b = 1800$$

On differentiating w.r.t. v in Eq. (i),

$$\frac{dc}{dv} = a - \frac{b}{v^2}$$

For maximum or minimum c ,

$$\frac{dc}{dv} = 0 \Rightarrow v = \pm \sqrt{\frac{b}{a}}$$

$$\Rightarrow \frac{d^2c}{dv^2} = \frac{2b}{v^3} \quad \text{at } v = \sqrt{\frac{b}{a}}, \frac{d^2c}{dv^2} > 0$$

So, at $v = \sqrt{\frac{b}{a}}$ the speed is most economical.


\therefore Most economical speed is

$$c = a\sqrt{\frac{b}{a}} + b\sqrt{\frac{a}{b}} = 2\sqrt{ab}$$

$$c = 2\sqrt{\frac{1}{2} \times 1800} = 2 \times 30$$

$$\Rightarrow c = 60$$

$$8 \quad f(x) = \begin{cases} |x^2 - 2|, & -1 \leq x < \sqrt{3} \\ \frac{x}{\sqrt{3}}, & \sqrt{3} \leq x < 2\sqrt{3} \\ 3 - x, & 2\sqrt{3} \leq x \leq 4 \end{cases}$$



From the above graph,

Maximum occurs at $x = 0$ and minimum at $x = 4$.

$$9 \quad f(x) = \begin{cases} |x^3 + x^2 + 3x + \sin x|, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

$$\text{Let } g(x) = x^3 + x^2 + 3x + \sin x$$

$$g'(x) = 3x^2 + 2x + 3 + \cos x$$

$$= 3 \left(x^2 + \frac{2x}{3} + 1 \right) + \cos x$$

$$= 3 \left\{ \left(x + \frac{1}{3} \right)^2 + \frac{8}{9} \right\} + \cos x > 0$$

$$\text{and } 2 < 3 + \sin \left(\frac{1}{x} \right) < 4$$

Hence, minimum value of $f(x)$ is 0 at $x = 0$.

Hence, number of points = 1

- 10 According to given information, we have

$$\text{Perimeter of square} + \text{Perimeter of circle} = 2 \text{ units}$$

$$\Rightarrow 4x + 2\pi r = 2$$

$$\Rightarrow r = \frac{1 - 2x}{\pi} \quad \dots (i)$$

Now, let A be the sum of the areas of the square and the circle. Then,

$$A = x^2 + \pi r^2$$

$$= x^2 + \pi \frac{(1 - 2x)^2}{\pi^2}$$

$$\Rightarrow A(x) = x^2 + \frac{(1 - 2x)^2}{\pi}$$

Now, for minimum value of $A(x)$,

$$\begin{aligned}\frac{dA}{dx} &= 0 \\ \Rightarrow 2x + \frac{2(1-2x)}{\pi} \cdot (-2) &= 0 \\ \Rightarrow x &= \frac{2-4x}{\pi} \\ \Rightarrow \pi x + 4x &= 2 \\ \Rightarrow x &= \frac{2}{\pi+4} \quad \dots(ii)\end{aligned}$$

Now, from Eq. (i), we get

$$\begin{aligned}r &= \frac{1-2 \cdot \frac{2}{\pi+4}}{\pi} \\ &= \frac{\pi+4-4}{\pi(\pi+4)} = \frac{1}{\pi+4} \quad \dots(iii)\end{aligned}$$

From Eqs. (ii) and (iii), we get
 $x = 2r$

11 We have,

$$\begin{aligned}f(x) &= x^2 + \frac{1}{x^2} \text{ and } g(x) = x - \frac{1}{x} \\ \Rightarrow h(x) &= \frac{f(x)}{g(x)} \\ \therefore h(x) &= \frac{x^2 + \frac{1}{x^2}}{x - \frac{1}{x}} = \frac{\left(x - \frac{1}{x}\right)^2 + 2}{x - \frac{1}{x}} \\ \Rightarrow h(x) &= \left(x - \frac{1}{x}\right) + \frac{2}{x - \frac{1}{x}} \\ x - \frac{1}{x} &> 0, \left(x - \frac{1}{x}\right) + \frac{2}{x - \frac{1}{x}} \in [2\sqrt{2}, \infty) \\ x - \frac{1}{x} &< 0, \left(x - \frac{1}{x}\right) + \frac{2}{x - \frac{1}{x}} \in (-\infty, 2\sqrt{2}]\end{aligned}$$

\therefore Local minimum value is $2\sqrt{2}$.

12 Consider the function

$$\begin{aligned}f(x) &= \frac{x^2}{(x^3 + 200)} \\ f'(x) &= x \frac{(400 - x^3)}{(x^3 + 200)^2} = 0\end{aligned}$$

when $x = (400)^{1/3}$, ($\because x \neq 0$)

$$x = (400)^{1/3} - h \Rightarrow f'(x) > 0$$

$$x = (400)^{1/3} + h \Rightarrow f'(x) < 0$$

$\therefore f(x)$ has maxima at $x = (400)^{1/3}$

Since, $7 < (400)^{1/3} < 8$, either a_7 or a_8 is the greatest term of the sequence.

$$\therefore a_7 = \frac{49}{543}$$

$$\text{and } a_8 = \frac{8}{89}$$

$$\text{and } \frac{49}{543} > \frac{8}{89}$$

$$\Rightarrow a_7 = \frac{49}{543} \text{ is the greatest term.}$$

13 $f(x) = x^3 - 3(7-a)x^2 - 3(9-a^2)x + 2$

$$f'(x) = 3x^2 - 6(7-a)x - 3(9-a^2)$$

For real root $D \geq 0$,

$$\Rightarrow 49 + a^2 - 14a + 9 - a^2 \geq 0$$

$$\Rightarrow a \leq \frac{58}{14}$$

For local minimum

$$f''(x) = 6x - 6(7-a) > 0$$

$$\Rightarrow x > 7 - a$$

has x must be negative

$$\Rightarrow 7 - a < 0$$

$$\Rightarrow a > 7$$

Thus contradictory, i.e., for real roots


$a \leq \frac{58}{14}$ and for negative point of local

minimum $a > 7$.

No possible values of a .

14 From the given figure coordinate of Q is $(-1, 0)$.

The equation of circle centre at Q with variable radius r is



$$(x+1)^2 + y^2 = r^2 \quad \dots(i)$$

This circle meets the line segment QP at S , where $QS = r$

It meets the circle $x^2 + y^2 = 1$ at $\dots(ii)$

$$R\left(\frac{r^2-2}{2}, \frac{r}{2}\sqrt{4-r^2}\right)$$

[on solving Eqs. (i) and (ii)]

$A = \text{Area of } \triangle QSR$

$$= \frac{1}{2} \times QS \times RT$$

$$= \frac{1}{2} r \left(\frac{r}{2} \cdot \sqrt{4-r^2} \right)$$

[since, RT is the y -coordinate of R]

$$= \frac{1}{4} [r^2 \sqrt{4-r^2}]$$

$$\begin{aligned}\therefore \frac{dA}{dr} &= \frac{1}{4} \left\{ 2r \sqrt{4-r^2} + \frac{r^2(-r)}{\sqrt{4-r^2}} \right\} \\ &= \frac{\{2r(4-r^2) - r^3\}}{4\sqrt{4-r^2}}\end{aligned}$$

$$= \frac{8r - 3r^3}{4\sqrt{4-r^2}}$$

$$\frac{dA}{dr} = 0, \text{ when } r(8-3r^2) = 0 \text{ giving}$$

$$r = \sqrt{\frac{8}{3}}$$

$$4\sqrt{4-r^2}(8-9r^2)$$

$$\Rightarrow \frac{d^2A}{dr^2} = \frac{-(8r-3r^3) \frac{(-4r)}{\sqrt{4-r^2}}}{16(4-r^2)}$$

$$\text{When } r = \sqrt{\frac{8}{3}}, \text{ then } \frac{d^2A}{dr^2} < 0$$

Hence, A is maximum when $r = \sqrt{\frac{8}{3}}$.

$$\begin{aligned}\text{Then, maximum area} &= \frac{8}{4 \times 3} \sqrt{4 - \frac{8}{3}} = \frac{4\sqrt{3}}{9} \text{ sq unit}\end{aligned}$$

15 Given,

$$P(x) = x^4 + ax^3 + bx^2 + cx + d$$

$$\Rightarrow P'(x) = 4x^3 + 3ax^2 + 2bx + c$$

Since, $x = 0$ is a solution for

$$P'(x) = 0, \text{ then}$$

$$c = 0$$

$$\therefore P(x) = x^4 + ax^3 + bx^2 + d \quad \dots(i)$$

Also, we have $P(-1) < P(1)$

$$\Rightarrow 1 - a + b + d < 1 + a + b + d$$

$$\Rightarrow a > 0$$

Since, $P'(x) = 0$, only when $x = 0$ and

$P(x)$ is differentiable in $(-1, 1)$, we

should have the maximum and minimum at the points $x = -1, 0$ and 1 .

Also, we have $P(-1) < P(1)$

\therefore Maximum of $P(x) = \text{Max}\{P(0), P(1)\}$

and minimum of $P(x) = \text{Min}$

$\{P(-1), P(0)\}$

In the interval $[0, 1]$,

$$P'(x) = 4x^3 + 3ax^2 + 2bx$$

$$= x(4x^2 + 3ax + 2b)$$

Since, $P'(x)$ has only one root $x = 0$,

then $4x^2 + 3ax + 2b = 0$ has no real roots.

$$\therefore (3a)^2 - 32b < 0$$

$$\Rightarrow \frac{9a^2}{32} < b$$

$$\therefore b > 0$$

Thus, we have $a > 0$ and $b > 0$.

$$\therefore P'(x) = 4x^3 + 3ax^2 + 2bx > 0,$$

$$\forall x \in (0, 1)$$

Hence, $P(x)$ is increasing in $[0, 1]$.

\therefore Maximum of $P(x) = P(1)$

Similarly, $P(x)$ is decreasing in $[-1, 0]$.

Therefore, minimum $P(x)$ does not occur at $x = -1$.