

DPP No. 53

Total Marks : 27

Max. Time : 28 min.

Topic : Straight Lines

Type of Questions		M.M	., Min.
Single choice Objective (no negative marking) Q.1,2,3	(3 marks, 3 min.)	[6,	6]
Multiple choice objective (no negative marking) Q.4	(5 marks, 4 min.)	[5,	4]
Subjective Questions (no negative marking) Q.5,6	(4 marks, 5 min.)	[8,	10]
Match the Following (no negative marking) Q.7	(8 marks, 8 min.)	[8,	8]

1. A is a point on either of two rays $y + \sqrt{3} |x| = 2$ at a distance of $\frac{4}{\sqrt{3}}$ units from their point of intersection. The co-ordinates of the foot of perpendicular from A on the bisector of the angle between them are

(A)
$$\left(-\frac{2}{\sqrt{3}},2\right)$$
 (B) (0, 0) (C) $\left(\frac{2}{\sqrt{3}},2\right)$ (D) (0, 4)

- 2. The base BC of a \triangle ABC is bisected at the point (p, q) & the equation to the side AB & AC are px + qy = 1 & qx + py = 1. The equation of the median through A is : (A) (p - 2q)x + (q - 2p)y + 1 = 0(B) (p + q)(x + y) - 2 = 0(C) $(2pq - 1)(px + qy - 1) = (p^2 + q^2 - 1)(qx + py - 1)$ (D) none of these
- 3. If the line y = x cuts the curve $x^3 + 3y^3 30xy + 72x 55 = 0$ in points A, B and C, then the value of $\frac{4\sqrt{2}}{55}$ OA.OB.OC (where O is the origin), is
 - (A) 55 (B) $\frac{1}{4\sqrt{2}}$ (C) 2 (D) 4
- 4. The equation of lines passing through point of intersection of lines 3x y 20 = 0 and x 2y 5 = 0, which are at a distance of 5 units from origin, is/are : (A) 4x + 3y = 25 (B) 3x - 4y = 25 (C) 4x - 3y = 25 (D) 3x + 4y = 25
- 5. A circle with centre in the first quadrant is tangent to y = x + 10, y = x 6, and the y-axis. Let (h, k) be the centre of the circle. If the value of $(h + k) = a + b\sqrt{a}$ where $(a, b \in Q)$, find the value of a + b.
- 6. If the variable line 3x 4y + k = 0 lies between the circles $x^2 + y^2 2x 2y + 1 = 0$ and $x^2 + y^2 16x 2y + 61 = 0$ without intersecting or touching either circle, then the range of k is (a, b) where a, b \in I. Find the value of (b a)

7. Match the column

Column – I Column – II Minimum possible number of positive roots of (A) (p) 2 $x^{2} - (1 + b)x + b - 2 = 0$ is $(b \in R)$ (B) In a \triangle ABC, co-ordinates of orthocentre, centroid and vertex A are (q) 0 (3, 2), (3, 1) and (1, 2) respectively. Then x-coordinate of vertex B is If $\log_1 \log_2 \log_2 (2x^2) = 0$, then x = (C) (r) 1 If there are three non concurrent and non parallel lines, (D) (s) 4 then number of points which are equidistant from all the three lines are

Answers Key

- **1.** (B)
- **2.** (C)
- **3.** (D)
- **4.** (C)(D)
- **5.** 10
- **6.** 6
- 7. (A) \rightarrow (r), (B) \rightarrow (s), (C) \rightarrow (p), (D) \rightarrow (s)