

# Periodic Table

---

## Modern Periodic Law:

- Properties of elements are the periodic function to their atomic numbers.
- The periodicity in properties is due to repetition of similar outer shell electronic configuration at a certain regular intervals.
- In modern periodic table is based on modern periodic law in which elements are arranged in increasing order of their atomic numbers.
- In the modern periodic table, the elements are arranged in rows and columns. These rows and columns are known as periods and groups respectively.
- The table consists of 7 periods and 18 groups
- Period indicates the value of 'n' (principal quantum number) for the outermost or valence shell.
- Same number of electrons is present in the outer orbitals (that is, similar valence shell electronic configuration)

## IUPAC Nomenclature for Elements with Atomic Number $\geq 100$

Digit	Name	Abbreviation
0	nil	n
1	un	u
2	bi	b
3	tri	t
4	quad	q
5	pent	p
6	hex	h
7	sept	s
8	oct	o
9	enn	e

### Classification of Elements:

Elements:	Valence Shell Electronic Configuration	Nature	Position in Modern Periodic Table
<b>s-block elements</b>	$ns^{1-2}$ (n = 1 to 7).	Metals	1 and 2 group elements
<b>p – block elements</b>	$ns^2 np^{1-6}$ (n = 2 to 7).	Metalloids & non metals but some of them are metals also.	groups 13 to 18
<b>d-Block Elements</b>	$(n-1)d^{1-10} ns^{1-2}$ (n = 4 to 7).	Metals	3 to 12 groups 3d series – Sc(21) to Zn (30) 4d series – Y (39) to Cd (48) 5d series – La (57), Hf (72) to Hg (80)
<b>f-Block Elements</b>	$(n-2)f^{1-14} (n-1)s^2 (n-1)p^6 (n-1)d^{0-1} ns^2$ (n = 6 and 7).	Radioactive	group 3 4f series – Lanthanides – 14 Elements Ce (58) to Lu (71)  5f series – Actinides – 14 Elements Th (90) to Lw (103)

### Periodicity in Atomic Properties:

#### 1. Atomic Radius:

- Within a given period atomic radius **decreases from left to right**. This is due to the effect of increase in nuclear charge while the electrons are being added to the same shell.
- Within a given group atomic radius increases down the group. This is due to the increase in number of shells.

Period	Group							
	IA	IIA	IIIA	IVA	VA	VIA	VIIA	Zero
<b>1.</b>	H 0.37							He 0.93
<b>2.</b>	Li 1.34	Be 0.90	B 0.82	C 0.77	N 0.73	O 0.74	F 0.72	Ne 1.31
<b>3.</b>	Na 1.54	Mg 1.30	Al 1.18	Si 1.11	P 1.06	S 1.02	Cl 0.99	Ar 1.74

- In the first transition series the atomic size slightly decreases from Sc to Mn because effect of effective nuclear charge is stronger than the shielding effect. The atomic size from Fe to Ni remains almost the same because both the effects balance each other.
- The atomic size from Cu to Zn slightly increases because shielding effect is more than effective nuclear charge due to  $d^{10}$  structure of Cu and Zn.
- Inner transition elements – As we move along the lanthanide series, there is a decrease in atomic as well as ionic radius. The decrease in size is regular in ions but not so regular in atoms. This is called lanthanide contraction.

<div>Increases</div>	1																	18																												
	H	2											13	14	15	16	17	He																												
	Li	Be											B	C	N	O	F	Ne																												
	Na	Mg	3	4	5	6	7	8	9	10	11	12	Al	Si	P	S	Cl	Ar																												
	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr																												
	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe																												
	Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn																												
	Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Uut	Uuq	Uup	Uuh	Uus	Uuo																												
<div>Increases</div>																																														
<div>Atomic Size</div>																																														
<table><tr><td>Ce</td><td>Pr</td><td>Nd</td><td>Pm</td><td>Sm</td><td>Eu</td><td>Gd</td><td>Tb</td><td>Dy</td><td>Ho</td><td>Er</td><td>Tm</td><td>Yb</td><td>Lu</td></tr><tr><td>Th</td><td>Pa</td><td>U</td><td>Np</td><td>Pu</td><td>Am</td><td>Cm</td><td>Bk</td><td>Cf</td><td>Es</td><td>Fm</td><td>Md</td><td>No</td><td>Lr</td></tr></table>																			Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu																																	
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr																																	

## 2) Ionisation potential or Ionisation Energy:

Ionization energy increases along the period while decreases down the group.

	1																	18																												
	H	2																	He																											
	Li	Be																	B	C	N	O	F	Ne																						
	Na	Mg	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	Ar																												
	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr																												
	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe																												
	Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn																												
	Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Uut	Uuq	Uup	Uuh	Uus	Uuo																												
	<table><tr><td>Ce</td><td>Pr</td><td>Nd</td><td>Pm</td><td>Sm</td><td>Eu</td><td>Gd</td><td>Tb</td><td>Dy</td><td>Ho</td><td>Er</td><td>Tm</td><td>Yb</td><td>Lu</td></tr><tr><td>Th</td><td>Pa</td><td>U</td><td>Np</td><td>Pu</td><td>Am</td><td>Cm</td><td>Bk</td><td>Cf</td><td>Es</td><td>Fm</td><td>Md</td><td>No</td><td>Lr</td></tr></table>																		Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu																																	
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr																																	

Increases

Increases

Increases

Ionization Energy

### Factors which influence I.E.

- Atomic size: the larger the size of the atom, the smaller the I.E. i.e., I.E.  $\mu$
- Effective nuclear charge: The greater the effective charge on the nucleus of an atom, the more difficult it would be to remove an electron from the atom because electrostatic force of attraction between the nucleus and the outermost electron increases. So greater energy will be required to remove the electron.
- Penetration effect of orbitals: The order of energy required to remove electron from s,p,d-and f-orbitals of a shell is  $s > p > d > f$ .
- Shielding or screening effect: Screening effect results in decrease of force of attraction between the nucleus and the outermost electron and lesser energy is required to separate the electron. Thus the value of I.P. decreases.
- Stability of half-filled and fully-filled orbitals: According to Hund's rule the stability of half filled or completely filled degenerate orbitals is comparatively high. So comparatively more energy is required to separate the electron from such atoms.

### Successive Ionisation Energies

Element	IE <sub>1</sub>	IE <sub>2</sub>	IE <sub>3</sub>	IE <sub>4</sub>
Li	520.1	7297	11813	—
C	1086.2	2352	4620	6221
N	1402.1	2856	4577	7474

### 3) Electron Affinity:

Electron affinity increases along the period while decreases down the group.

#### Factors affecting the magnitude of electron affinity

- Atomic size – In general electron affinity value decreases with the increasing atomic radius because electrostatic force of attraction decreases between the electron being added and the atomic nucleus due to increase of distance between them.
- Effective nuclear charge – Electron affinity value of the element increase as the effective nuclear charge on the atomic nucleus increases because electrostatic force of attraction between the electron being added and the nucleus increases. As the electrostatic force of attraction increases, amount of energy released is more.
- Screening or Shielding effect – Electron affinity value of the elements decreases with the increasing shielding or screening effect. The shielding effect between the outer electrons and the nucleus increases as the number of electrons increases in the inner shells.

Stability of half filled and completely filled orbitals – The stability of half filled and completely filled degenerate orbitals of a sub shell is comparatively more, so it is difficult to add electron in such orbitals and lesser energy is released on addition of electron hence the electron affinity value will decrease.

**Electron Affinity**

1	2											13	14	15	16	17	18
H												B	C	N	O	F	He
Li	Be											Al	Si	P	S	Cl	Ar
Na	Mg	3	4	5	6	7	8	9	10	11	12	Ga	Ge	As	Se	Br	Kr
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	In	Sn	Sb	Te	I	Xe
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	Tl	Pb	Bi	Po	At	Rn
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Uut	Uuq	Uup	Uuh	Uus	Uuo

  

Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr

? Electron Affinities ( $\text{kJ mol}^{-1}$ ),  $M_{(g)} + e^{-} \rightarrow M_{(g)}^{-} + \text{energy}$

Group	1	2	13	14	15	16	17	18
<b>H</b>								He
<b>72</b>								< 0
<b>Li</b>		Be	B	C	N	O	F	
<b>57</b>		< 0	27	122	< 0	142	333	
<b>Na</b>		Mg	Al	Si	P	S	Cl	Ne
<b>53</b>		< 0	44	134	72	200	349	< 0
<b>K</b>		Ca	Ga	Ge	As	Se	Br	Ar
<b>48</b>		< 0	29	116	77	195	324	< 0
<b>Cs</b>					Bi	Te	1	Xe
<b>43</b>								