

INVERSE TRIGONOMETRIC FUNCTIONS

DOMAIN, RANGE AND GRAPH OF INVERSE TRIGONOMETRIC FUNCTIONS



What you already know

- Function and classification of functions
- Basic graph of functions



What you will learn

- Condition for existence of inverse of a function
- Domain, range, and graph of inverse trigonometric functions

Inverse function recap

If $f : A \rightarrow B$ is a one-one and an onto function, then $f^{-1} : B \rightarrow A$ is its inverse.
If f and f^{-1} are the inverse of each other, then we get the following:

- Domain of f = Range of f^{-1}
- Range of f = Domain of f^{-1}
- $(f^{-1} \text{ of } (x)) = f^{-1}(f(x)) = x$


Condition for Existence of Inverse of a Function

An inverse of a function exists only if the function is one-one and onto, i.e., bijective.
If a function is not bijective in the given domain, then its domain is modified to get the inverse.
The modified domain is known as a restricted domain.

Example

Let us consider the following function: $f : \mathbb{R} \rightarrow \mathbb{R} : f(x) = \sin x$

From the given graph, we can see that the given function is neither one-one nor onto as a horizontal line will cut the graph at more than one point and codomain (\mathbb{R}) is not equal to the range of the function.




To make the function bijective domain and codomain of the given function is modified to

$$f : \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow [-1, 1] \text{ as}$$

shown in the figure.

Now, we can see that in the modified domain and codomain, the given function $f(x) = \sin x$ is both one-one and onto function, i.e., bijective.



Note

- Inverse trigonometric function gives angle in radians. For example, $\sin^{-1}x$ is the measure of the angle in radian.
- There is a difference between $\sin^{-1}x$ (or) $(\arcsin x)$ and $(\sin x)^{-1}$
- $(\sin x)^{-1} = \frac{1}{\sin x}$ (wherever it exists)

Domain, Range, and Graph of Inverse Trigonometric Functions

Domain, range, and graph of $f(x) = \sin^{-1}x$


For the function $f(x) = \sin x$,

$$\text{Restricted domain} = \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

$$\text{Range} = [-1, 1]$$

Here, we can see that the nature of the graph of $f(x)$ is strictly increasing.



By taking the mirror image of $f(x)$ about the line $y = x$, we get the graph of inverse of $f(x)$, i.e., $\sin^{-1}x$.



For the inverse trigonometric function $g(x) = \sin^{-1} x$, we get,

$$\text{Domain} = [-1, 1]$$

$$\text{Range} = \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$



Note

For the inverse trigonometric function $g(x) = \sin^{-1} x$, if x is positive, then angle $g(x)$ lies in the first quadrant, and if x is negative, then angle $g(x)$ lies in the fourth quadrant.

Example

$$(i) \sin^{-1} 1 = \theta \Rightarrow \sin \theta = 1 \Rightarrow \theta = \frac{\pi}{2}$$

$$(ii) \sin^{-1} \left(-\frac{1}{2}\right) = \alpha \Rightarrow \sin \alpha = -\frac{1}{2} \Rightarrow \alpha = -\frac{\pi}{6}$$

$$(iii) \sin^{-1}(\pi) = \text{Not defined as } \pi \notin [-1, 1]$$


Domain, range, and graph of $f(x) = \cos^{-1} x$

For the cosine function $f(x) = \cos x$,


Restricted domain = $[0, \pi]$

Range = $[-1, 1]$

Here, we can see that the nature of the graph of $f(x)$ is strictly decreasing.




By taking the mirror image of $f(x)$ about the line $y = x$, we get the graph of inverse of $f(x)$, i.e., $\cos^{-1} x$.



For the inverse trigonometric function, $g(x) = \cos^{-1} x$, we get,

Domain = $[-1, 1]$

Range = $[0, \pi]$



Note

For the inverse trigonometric function $g(x) = \cos^{-1} x$, if x is positive, then angle $g(x)$ lies in the first quadrant, and if x is negative, then angle $g(x)$ lies in the second quadrant.

Example

$$\cos^{-1}\left(\frac{\sqrt{3}}{2}\right) = \theta \Rightarrow \cos \theta = \frac{\sqrt{3}}{2} \Rightarrow \theta = \frac{\pi}{6}$$



If $\cos^{-1}x + \cos^{-1}y + \cos^{-1}z = 3\pi$, then find the value of $x^{50} + y^{50} + z^{50}$.

Solution

We know, $0 \leq \cos^{-1}x \leq \pi$

$$\text{Given, } \cos^{-1}x + \cos^{-1}y + \cos^{-1}z = 3\pi$$

It is only possible if

$$\cos^{-1}x = \cos^{-1}y = \cos^{-1}z = \pi$$

$$\Rightarrow x = y = z = \cos \pi = -1$$

$$\begin{aligned}\therefore x^{50} + y^{50} + z^{50} &= (-1)^{50} + (-1)^{50} + (-1)^{50} \\ &= 1 + 1 + 1 = 3\end{aligned}$$


Domain, range, and graph of $f(x) = \tan^{-1}x$

For the tangent function $f(x) = \tan x$,


$$\text{Restricted domain} = \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$


$$\text{Range} = \mathbb{R}$$

Here, we can see that the nature of the graph of $f(x)$ is strictly increasing.



By taking the mirror image of $f(x)$ about the line $y = x$, we get the graph of the inverse of $f(x)$, i.e., $\tan^{-1}x$.






For the inverse trigonometric function $g(x) = \tan^{-1} x$, we get,

Domain = \mathbb{R}

Range = $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$



Note

For the inverse trigonometric function $g(x) = \tan^{-1} x$, if x is positive, then $g(x)$ lies in the first quadrant, and if x is negative, then $g(x)$ lies in the fourth quadrant.



What is the value of $\tan^{-1}(1) + \cos^{-1}\left(-\frac{1}{2}\right) + \sin^{-1}\left(-\frac{1}{2}\right)$?

Solution

$$\tan^{-1}(1) = \frac{\pi}{4}$$

$$\cos^{-1}\left(-\frac{1}{2}\right) = \frac{2\pi}{3}$$

$$\sin^{-1}\left(-\frac{1}{2}\right) = -\frac{\pi}{6}$$

$$\therefore \tan^{-1}(1) + \cos^{-1}\left(-\frac{1}{2}\right) + \sin^{-1}\left(-\frac{1}{2}\right) = \frac{\pi}{4} + \frac{2\pi}{3} + \left(-\frac{\pi}{6}\right) = \frac{3\pi}{4}$$


Domain, range, and graph of $f(x) = \cot^{-1} x$

For the cotangent function $f(x) = \cot x$,


Restricted domain = $(0, \pi)$

Range = \mathbb{R}

Here, we can see that the nature of the graph of $f(x)$ is strictly decreasing.




By taking the mirror image of $f(x)$ about the line $y = x$, we get the graph of the inverse of $f(x)$, i.e., $\cot^{-1} x$.



For the inverse trigonometric function $g(x) = \cot^{-1} x$, we get,

Domain = \mathbb{R}

Range = $(0, \pi)$






Note

For the inverse trigonometric function $g(x) = \cot^{-1} x$, if x is positive, then $g(x)$ lies in the first quadrant, and if x is negative, then $g(x)$ lies in the second quadrant.

Domain, range, and graph of $f(x) = \sec^{-1} x$


For the secant function $f(x) = \sec x$,

Restricted domain = $[0, \pi] - \left\{\frac{\pi}{2}\right\}$
 Range = $(-\infty, -1] \cup [1, \infty)$



By taking the mirror image of $f(x)$ about the line $y = x$, we get the graph of the inverse of $f(x)$, i.e., $\sec^{-1} x$.


For the function ()




For the inverse trigonometric function $g(x) = \sec^{-1} x$, we get,

Domain = $(-\infty, -1] \cup [1, \infty)$

Range = $[0, \pi] - \left\{\frac{\pi}{2}\right\}$





Note

For the inverse trigonometric function $g(x) = \sec^{-1}x$, if x is positive, then $g(x)$ lies in the first quadrant, and if x is negative, then $g(x)$ lies in the second quadrant.



Domain, range, and graph of $f(x) = \operatorname{cosec}^{-1}x$

For the cosecant function $f(x) = \operatorname{cosec} x$,

Restricted domain = $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right) - \{0\}$

Range = $(-\infty, -1] \cup [1, \infty)$


By taking the mirror image of $f(x)$ about the line $y = x$, we get the graph of the inverse of $f(x)$, i.e., $\operatorname{cosec}^{-1}x$.



For the inverse trigonometric function
 $g(x) = \operatorname{cosec}^{-1} x$,

Domain = $(-\infty, -1] \cup [1, \infty)$

Range = $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$



Note

For the inverse trigonometric function $g(x) = \operatorname{cosec}^{-1} x$, if x is positive, then $g(x)$ lies in the first quadrant, and if x is negative, then $g(x)$ lies in the fourth quadrant.

Example

(i) $\operatorname{cosec}^{-1}(1) = \frac{\pi}{2}$

(ii) $\operatorname{cosec}^{-1}\left(\frac{1}{2}\right)$ = Not defined as $\frac{1}{2} \notin (-\infty, -1] \cup [1, \infty)$



Find domain and range of $\cos^{-1}[x]$, where $[.]$ represents the greatest integer function.

Solution

Step 1

Given function, $y = \cos^{-1}[x]$
 for the function $y = \cos^{-1}x$, $-1 \leq x \leq 1$

$$\Rightarrow -1 \leq [x] \leq 1$$

$$\Rightarrow -1 \leq x < 2$$

Step 2

$$-1 \leq [x] \leq 1 \Rightarrow [x] = -1, 0, 1$$

$$[x] = -1 \Rightarrow \cos^{-1}(-1) = \pi$$

$$[x] = 0 \Rightarrow \cos^{-1}(0) = \frac{\pi}{2}$$

$$[x] = 1 \Rightarrow \cos^{-1}(1) = 0$$

The range of the function is $\{0, \frac{\pi}{2}, \pi\}$.



Concept Check

1. Find the domain and range of $y = \sin^{-1}(e^x)$

2. Find the domain of $f(x) = \sqrt{\sin^{-1}(2x) + \frac{\pi}{6}}$



JEE ADVANCED 2003



Summary Sheet



Key Takeaways

- An inverse of a function exists only if the function is one-one and onto, i.e., bijective.
- If a function is not bijective in the given domain, then its domain is modified to get the inverse. Thus, a modified domain is known as a restricted domain.
- For $g(x) = \sin^{-1} x$,
Domain = $[-1, 1]$
Range = $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
- For $g(x) = \cos^{-1} x$,
Domain = $[-1, 1]$
Range = $[0, \pi]$
- For $g(x) = \tan^{-1} x$,
Domain = \mathbb{R}
Range = $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
- For $g(x) = \cot^{-1} x$,
Domain = \mathbb{R}
Range = $(0, \pi)$
- For $g(x) = \sec^{-1} x$,
Domain = $(-\infty, -1] \cup [1, \infty)$
Range = $[0, \pi] - \left\{\frac{\pi}{2}\right\}$
- For $g(x) = \operatorname{cosec}^{-1} x$,
Domain = $(-\infty, -1] \cup [1, \infty)$
Range = $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$



Mind Map

Condition for
existence of
inverse

Domain, range and graph
of inverse trigonometric
functions

$$f(x) = \sin^{-1} x$$

$$f(x) = \cos^{-1} x$$

$$f(x) = \tan^{-1} x$$

$$f(x) = \cot^{-1} x$$

$$f(x) = \sec^{-1} x$$

$$f(x) = \operatorname{cosec}^{-1} x$$



Self-Assessment

Find the value of $\tan^{-1} \sqrt{3} - \sec^{-1}(-2)$.



Answers

Concept Check

1.

Step 1:

The domain of $\sin^{-1} x$ is $-1 \leq x \leq 1$

\Rightarrow For $\sin^{-1} e^x$, we have $-1 \leq e^x \leq 1$

$\Rightarrow 0 < e^x \leq 1$ (Since exponential function cannot be negative)

$\Rightarrow x \leq 0 \rightarrow$ Domain of the function


Step 2:

$$0 < e^x \leq 1$$

$$\Rightarrow \sin^{-1}(0) < \sin^{-1}(e^x) \leq \sin^{-1} 1$$

$$\Rightarrow 0 < \sin^{-1}(e^x) \leq \frac{\pi}{2}$$

$$\Rightarrow \text{Range} = \left(0, \frac{\pi}{2}\right]$$



2.

$$\text{Given, } y = f(x) = \sqrt{\sin^{-1}(2x) + \frac{\pi}{6}}$$

For the real value of y , we get,

$$\sin^{-1}(2x) + \frac{\pi}{6} \geq 0$$

$$\Rightarrow \sin^{-1}(2x) \geq -\frac{\pi}{6}$$

Step 2:

We know that the maximum value of

$$\sin^{-1}x \text{ is } \frac{\pi}{2}.$$

So,


$$\Rightarrow \frac{\pi}{2} \geq \sin^{-1}(2x) \geq -\frac{\pi}{6}$$

$$\Rightarrow \sin \frac{\pi}{2} \geq 2x \geq \sin\left(-\frac{\pi}{6}\right)$$

$$\Rightarrow 1 \geq 2x \geq -\frac{1}{2}$$

$$\Rightarrow \frac{1}{2} \geq x \geq -\frac{1}{4}$$

$$\Rightarrow x \in \left[-\frac{1}{4}, \frac{1}{2}\right]$$



Self-Assessment

$$\tan^{-1} \sqrt{3} - \sec^{-1}(-2)$$

$$= \frac{\pi}{3} - \frac{2\pi}{3} = -\frac{\pi}{3}$$

INVERSE TRIGONOMETRIC FUNCTIONS

PROPERTIES OF INVERSE TRIGONOMETRIC FUNCTIONS



What you already know

- Conditions for the existence of inverse functions
- Domain, range, and graph of ITF



What you will learn

- Properties of inverse function $f^{-1}(-x)$
- Graphs of inverse function $f(f^{-1}(x))$
- Properties of inverse function $f(f^{-1}(x))$
- Properties and graphs of inverse function $f^{-1}(f(x))$

Property 1: Properties of Inverse Trigonometric Function $f^{-1}(-x)$

1. $\sin^{-1}(-x) = -\sin^{-1}x; |x| \leq 1$
2. $\tan^{-1}(-x) = -\tan^{-1}x; x \in \mathbb{R}$
3. $\operatorname{cosec}^{-1}(-x) = -\operatorname{cosec}^{-1}x; |x| \geq 1$
4. $\cos^{-1}(-x) = \pi - \cos^{-1}x; |x| \leq 1$
5. $\cot^{-1}(-x) = \pi - \cot^{-1}x; x \in \mathbb{R}$
6. $\sec^{-1}(-x) = \pi - \sec^{-1}x; |x| \geq 1$

To prove $\sin^{-1}(-x) = -\sin^{-1}x; |x| \leq 1$

Proof

Let $\sin^{-1}(-x) = \theta$, where $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
 $\Rightarrow -x = \sin \theta$ or $x = -\sin \theta$

We know that $\sin(-\theta) = -\sin \theta$

Hence, $x = -\sin \theta$ can be written as $x = \sin(-\theta)$

$\Rightarrow \sin^{-1}x = -\theta$

$\Rightarrow \sin^{-1}x = -\sin^{-1}(-x)$ or $\sin^{-1}(-x) = -\sin^{-1}x$

Hence proved.

To prove $\cot^{-1}(-x) = \pi - \cot^{-1}x; x \in \mathbb{R}$

Proof

Let $\cot^{-1}(-x) = \theta$, $\theta \in (0, \pi)$

$\Rightarrow -x = \cot \theta$ or $x = -\cot \theta$

We know that $\cot(\pi - \theta) = -\cot \theta$

Hence, $x = -\cot \theta$ can be written as $x = \cot(\pi - \theta)$

$\Rightarrow \cot^{-1}x = (\pi - \theta)$ or $\theta = \pi - \cot^{-1}x$

$\Rightarrow \cot^{-1}(-x) = \pi - \cot^{-1}x$

Hence proved.



Note

1. $\sin^{-1}(x)$, $\tan^{-1}(x)$, $\operatorname{cosec}^{-1}(x)$ are **odd functions** ($f(-x) = -f(x)$)
2. $\cos^{-1}(x)$, $\cot^{-1}(x)$, $\sec^{-1}(x)$ are **neither even nor odd functions**.



Evaluate: $\sin^{-1}\left(\frac{-1}{2}\right)$

Solution

$$\sin^{-1}\left(\frac{-1}{2}\right) = -\sin^{-1}\left(\frac{1}{2}\right) = -\frac{\pi}{6}$$

Property 2: Properties of Inverse Function $f(f^{-1}(x))$

1. $\sin(\sin^{-1}(x)) = x; x \in [-1, 1]$
2. $\cos(\cos^{-1}(x)) = x; x \in [-1, 1]$
3. $\tan(\tan^{-1}(x)) = x; x \in \mathbb{R}$
4. $\cot(\cot^{-1}(x)) = x; x \in \mathbb{R}$
5. $\operatorname{cosec}(\operatorname{cosec}^{-1}(x)) = x; |x| \geq 1$
6. $\sec(\sec^{-1}(x)) = x; |x| \geq 1$

Proof

To prove $\sin(\sin^{-1}(x)) = x; x \in [-1, 1]$

Let $y = \sin(\sin^{-1}(x))$ and $\sin^{-1}(x) = \theta$, $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

$$\Rightarrow x = \sin \theta \dots\dots(1)$$

Substituting $\sin^{-1}(x) = \theta$ in y , we get,

$$y = \sin \theta \dots(2)$$

From (1) and (2), we get,

$$y = x$$


$$\therefore \sin(\sin^{-1}(x)) = x$$

Graphs of Inverse Function $f(f^{-1}(x))$

1. $y = \sin(\sin^{-1}(x))$ and $y = \cos(\cos^{-1}(x))$ for $x \in [-1, 1]$

We know that $y = \sin(\sin^{-1}(x)) = x; x \in [-1, 1]$ and $y = \cos(\cos^{-1}(x)) = x; x \in [-1, 1]$


So, the graph will be the line $y = x; x \in [-1, 1]$



2. $y = \tan(\tan^{-1}(x))$ and $y = \cot(\cot^{-1}(x))$ for $x \in \mathbb{R}$

We know that $y = \tan(\tan^{-1}(x)) = x$; $x \in \mathbb{R}$ and $y = \cot(\cot^{-1}(x)) = x$; $x \in \mathbb{R}$


So, the graph will be the line $y = x$; $x \in \mathbb{R}$



3. $y = \operatorname{cosec}(\operatorname{cosec}^{-1}(x))$ and $y = \sec(\sec^{-1}(x))$ for $|x| \geq 1$

We know that $y = \operatorname{cosec}(\operatorname{cosec}^{-1}(x)) = x$; $|x| \geq 1$ and $y = \sec(\sec^{-1}(x)) = x$; $|x| \geq 1$

So, the graph will be the line $y = x$; $|x| \geq 1$



Evaluate: $\cos\left(\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)\right)$

Solution


$$\cos(\cos^{-1}(x)) = x; x \in [-1, 1]$$

$$\Rightarrow \cos\left(\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)\right) = -\frac{\sqrt{3}}{2}$$

Property 3: Properties of Inverse Function $f^{-1}(f(x))$

1. $\sin^{-1}(\sin(x))$

Graph of $y = \sin^{-1}(\sin(x))$



Proof

Consider $y = \sin^{-1}(\sin(x))$

We know that $\sin^{-1}(\sin(x)) \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

$$\Rightarrow y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

Also, $y = \sin^{-1}(\sin(x)) \Rightarrow \sin y = \sin x$

That is, $y = n\pi + (-1)^n x$, $n \in \mathbb{Z} \dots (i)$

$$\sin^{-1}(\sin(x)) = n\pi + (-1)^n x, n \in \mathbb{Z}$$

The graph of $\sin^{-1}(\sin(x))$ will be a straight line.

We have to ensure that $(n\pi + (-1)^n x) \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ as $\sin^{-1}(\sin(x)) \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

Case 1: $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

If $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$, then $n = 0$ in (i)


$$\Rightarrow y = x$$

Case 2: $x \in \left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$

If $x \in \left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$, then $n = 1$ in (i)


$$\Rightarrow y = \pi - x$$

Let us plot the graph of $y = \sin^{-1}(\sin(x))$ for $x \in \left[-\frac{\pi}{2}, \frac{3\pi}{2}\right]$



We also know that if $g(x)$ is periodic with period T , then $f(g(x))$ is also periodic with period T . That means the period of $\sin^{-1}(\sin(x))$ is 2π , and we have already plotted the graph for 2π length. So, the graph will simply repeat itself after every 2π interval.

The graph of $y = \sin^{-1}(\sin(x))$ is as follows:




Evaluate: $\sin^{-1}(\sin 165^\circ)$.

Solution

Step 1: 165° lies between 90° and 180° .

From the graph, it is clear that $y = \sin^{-1}(\sin(x))$ will follow $y = \pi - x$




Step 2:

$$\sin^{-1}(\sin 165^\circ) = 180^\circ - 165^\circ = 15^\circ$$

2. $\cos^{-1}(\cos(x))$

Graph of $y = \cos^{-1}(\cos(x))$



Proof

Consider $y = \cos^{-1}(\cos(x))$

We know that $\cos^{-1}(\cos(x)) \in [0, \pi]$

$\Rightarrow y \in [0, \pi]$

Also, $y = \cos^{-1}(\cos(x)) \Rightarrow \cos y = \cos x$

That is, $y = 2n\pi \pm x, n \in \mathbb{Z} \dots (i)$

$\cos^{-1}(\cos(x)) = 2n\pi \pm x, n \in \mathbb{Z}$

\Rightarrow The graph of $\cos^{-1}(\cos(x))$ will be a straight line.

We have to ensure that $(2n\pi \pm x) \in [0, \pi]$ as $\cos^{-1}(\cos(x)) \in [0, \pi]$

Case 1: $x \in [0, \pi]$

If $x \in [0, \pi]$, then $n = 0$ in (i)


$\Rightarrow y = x$

Case 2: $x \in [\pi, 2\pi]$


If $x \in [\pi, 2\pi]$, then $n = 1$ in (i)

$\Rightarrow y = 2\pi - x$

Let us plot the graph of $y = \cos^{-1}(\cos(x))$ for $x \in [0, 2\pi]$



We also know that if $g(x)$ is periodic with period T , then $f(g(x))$ is also periodic with period T . That means the period of $\cos^{-1}(\cos(x))$ is 2π , and we have already plotted the graph for 2π length. So, the graph will simply repeat itself after every 2π interval. The graph of $y = \cos^{-1}(\cos(x))$ is as follows:




Evaluate: $\cos^{-1}(\cos 225^\circ)$.

Solution

Step 1: 225° lies between 180° and 360° .

From the graph, it is clear that $y = \cos^{-1}(\cos(x))$ will follow $y = 2\pi - x$



Step 2:

$$\cos^{-1}(\cos 225^\circ) = 360^\circ - 225^\circ = 135^\circ$$




Let $f : [0, 4\pi] \rightarrow [0, \pi]$ be defined by $f(x) = \cos^{-1}(\cos x)$. Find the number of points for $x \in [0, 4\pi]$ satisfying the equation $f(x) = \frac{(10-x)}{10}$

Solution

Step 1: Let us plot $y = \cos^{-1}(\cos x)$ and $y = \frac{(10-x)}{10}$ for $x \in [0, 4\pi]$. The number of solutions will be equal to the number of intersection points.

Step 2:



Number of intersection points = 3

Hence, $f(x) = \frac{(10-x)}{10}$ has 3 solutions in $x \in [0, 4\pi]$.



Concept Check

1. Evaluate the following:

(a) $2 \cot^{-1}(-\sqrt{3})$

(b) $\sin\left(\frac{\pi}{2} - \sin^{-1}\left(-\frac{\sqrt{3}}{2}\right)\right)$

2. Evaluate the following:

(a) $\operatorname{cosec}\left(\operatorname{cosec}^{-1}\left(\frac{1}{2}\right)\right)$

(b) $\sin(\sin^{-1}(2))$

3. Evaluate $\sin^{-1}(\sin 7)$.

4. Evaluate $\cos^{-1}(\cos(7))$.



Summary Sheet



Key formulae

- **Properties of inverse function $f^{-1}(-x)$**


1. $\sin^{-1}(-x) = -\sin^{-1} x; |x| \leq 1$
2. $\tan^{-1}(-x) = -\tan^{-1} x; x \in \mathbb{R}$
3. $\operatorname{cosec}^{-1}(-x) = -\operatorname{cosec}^{-1} x; |x| \geq 1$
4. $\cos^{-1}(-x) = \pi - \cos^{-1} x; |x| \leq 1$
5. $\cot^{-1}(-x) = \pi - \cot^{-1} x; x \in \mathbb{R}$
6. $\sec^{-1}(-x) = \pi - \sec^{-1} x; |x| \geq 1$

- **Properties of inverse function $f(f^{-1}(x))$**


1. $\sin(\sin^{-1}(x)) = x; x \in [-1, 1]$
2. $\cos(\cos^{-1}(x)) = x; x \in [-1, 1]$
3. $\tan(\tan^{-1}(x)) = x; x \in \mathbb{R}$
4. $\cot(\cot^{-1}(x)) = x; x \in \mathbb{R}$
5. $\operatorname{cosec}(\operatorname{cosec}^{-1}(x)) = x; |x| \geq 1$
6. $\sec(\sec^{-1}(x)) = x; |x| \geq 1$

Graphs


1. $y = \sin(\sin^{-1}(x))$ and $y = \cos(\cos^{-1}(x))$ for $x \in [-1, 1]$




2. $y = \tan(\tan^{-1}(x))$ and $y = \cot(\cot^{-1}(x))$ for $x \in \mathbb{R}$




3. $y = \operatorname{cosec}(\operatorname{cosec}^{-1}(x))$ and $y = \sec(\sec^{-1}(x))$ for $|x| \geq 1$



4. $y = \sin^{-1}(\sin(x))$



5. $y = \cos^{-1}(\cos(x))$





Mind Map

Properties of ITF

Properties of $f^{-1}(-x)$

Properties of $f(f^{-1}(x))$

Properties of $f^{-1}(f(x))$



Self-Assessment

Evaluate $\cos^{-1} \cos\left(\frac{7\pi}{6}\right)$.



Answers

Concept Check

1.

(a) Step 1:

$$2 \cot^{-1}(-\sqrt{3}) = 2(\pi - \cot^{-1}\sqrt{3})$$

Step 2:

$$= 2\left(\pi - \frac{\pi}{6}\right) = \frac{5\pi}{3}$$

(b) Step 1:

$$\sin\left(\frac{\pi}{2} - \sin^{-1}\left(-\frac{\sqrt{3}}{2}\right)\right) = \sin\left(\frac{\pi}{2} + \sin^{-1}\left(\frac{\sqrt{3}}{2}\right)\right)$$

Step 2:

$$= \sin\left(\frac{\pi}{2} + \frac{\pi}{3}\right) = \cos\frac{\pi}{3} = \frac{1}{2}$$

2.

(a)

Domain of $\operatorname{cosec}^{-1} x$ is $|x| \geq 1$

$\therefore \operatorname{cosec}\left(\operatorname{cosec}^{-1}\left(\frac{1}{2}\right)\right)$ is not defined.

(b)

Domain of $\sin^{-1} x$ is $x \in [-1, 1]$

$\therefore \sin(\sin^{-1}(2))$ is not defined.


3.

Step 1: 7 lies between 2π and $\frac{5\pi}{2}$.

From the graph, it is clear that

$y = \sin^{-1}(\sin(x))$ will follow

$y = x - 2\pi$



Step 2: $\sin^{-1}(\sin 7) = 7 - 2\pi$


4.

Step 1: 7 lies between 2π and 3π .

From the graph, it is clear that

$y = \cos^{-1}(\cos(x))$ will follow

$y = x - 2\pi$




Step 2: $\cos^{-1}(\cos 7) = 7 - 2\pi$

Self-Assessment

Step 1: $\frac{7\pi}{6}$ lies between π and 2π .

From the graph, it is clear that $y = \cos^{-1}(\cos(x))$ will follow

$y = 2\pi - x$



Step 2: $\cos^{-1} \cos\left(\frac{7\pi}{6}\right) = 2\pi - \frac{7\pi}{6} = \frac{5\pi}{6}$

INVERSE TRIGONOMETRIC FUNCTIONS

MORE ON PROPERTIES OF INVERSE TRIGONOMETRIC FUNCTIONS



What you already know

- Function and classification of functions
- Basic graphs of functions
- Domain, range, and graph of inverse trigonometric functions



What you will learn

- Function of the form $f^{-1}(f(x))$
- Properties of inverse trigonometric function $f^{-1}\left(\frac{1}{x}\right)$

Properties of Inverse Function $f^{-1}(f(x))$ (cont.)

Let us consider function $y = \tan^{-1}(\tan x)$.

Given function is an inverse tangent function

\therefore Range of the function is $y \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Also, x is an argument of the tangent function

\therefore Domain of the function is $x \in \mathbb{R} - \left\{(2n+1)\frac{\pi}{2}\right\}$

As the value of the function oscillates in the range $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, the function is periodic and

depends on the period of $\tan x$.

We know that the period of $\tan x$ is π .

\therefore Period of $y = \tan^{-1}(\tan x)$ is also π

So, let's plot the graph of $y = \tan^{-1}(\tan x)$ for the length π .

$$y = \tan^{-1}(\tan x)$$

$$\Rightarrow \tan y = \tan x$$

$$\Rightarrow y = n\pi + x \dots (i)$$


\Rightarrow Graph of $y = \tan^{-1}(\tan x)$ will be a straight line with slope 1.

Now, by substituting $n = 0$, we get the relation between x and y for the first interval,

$$y = x$$


So, for the first interval, the function

$y = \tan^{-1}(\tan x)$ can be plotted as shown in the figure above.



As the function is periodic with the period π , the slope of the graph is always 1.

Hence, for all the intervals graph of the function $y = \tan^{-1}(\tan x)$ can be plotted as shown in the figure below.



Let us consider the function $y = \cot^{-1}(\cot x)$.

Given function is an inverse cotangent function

\therefore Range of the function is $y \in (0, \pi)$

Also, x is the argument of the tangent function


\therefore Domain of the function is $x \in \mathbb{R} - \{n\pi\}; n \in \mathbb{Z}$

As the value of the function oscillates in the range $(0, \pi)$, the function is periodic and depends on the period of $\cot x$.

We know that the period of $\cot x$ is π .

\therefore Period of $y = \cot^{-1}(\cot x)$ is also π

So, let's plot the graph of $y = \cot^{-1}(\cot x)$ for the length π .



$$y = \cot^{-1}(\cot x)$$

$$\Rightarrow \cot y = \cot x$$

$$\Rightarrow \tan x = \tan y$$

$$\Rightarrow y = n\pi + x \dots (i)$$

\Rightarrow Graph of $y = \cot^{-1}(\cot x)$ will be a straight line with slope 1.


Now, by substituting $n = 0$, we get the relation between x and y for the first interval,

$$y = x$$

So, the first interval the function $y = \cot^{-1}(\cot x)$ can be plotted as shown in the figure above.

As the function is periodic with the period π , the slope of the graph is always 1.

Hence, all the interval graphs of the function $y = \cot^{-1}(\cot x)$ can be plotted as shown in the adjacent figure.



Find $\sin^{-1}(\sin 5) + \cos^{-1}(\cos 10) + \tan^{-1}(\tan (-6)) + \cot^{-1}(\cot (-10))$.

Solution


Step 1:

From the graph of $y = \sin^{-1}(\sin x)$, we get,

$$\text{For } x = 5, y = x - 2\pi$$

$$\Rightarrow y = 5 - 2\pi$$

$$\Rightarrow \sin^{-1}(\sin 5) = 5 - 2\pi \dots (i)$$




Step 2:

From the graph of $y = \cos^{-1}(\cos x)$,
we get,

$$\text{For } x = 10, y = 4\pi - x$$

$$\Rightarrow y = 4\pi - 10$$

$$\Rightarrow \cos^{-1}(\cos 10) = 4\pi - 10 \dots (ii)$$


**Step 3:**

From the graph of $y = \tan^{-1}(\tan(x))$,

For $x = -6$, $y = x + 2\pi$, we get,

$$\Rightarrow y = 2\pi - 6$$

$$\Rightarrow \tan^{-1}(\tan(-6)) = 2\pi - 6 \dots (iii)$$


**Step 4:**

From the graph of $y = \cot^{-1}(\cot(x))$,

For $x = -10$, $y = x + 4\pi$, we get,

$$\Rightarrow y = 4\pi - 10$$

$$\Rightarrow \cot^{-1}(\cot(-10)) = 4\pi - 10 \dots (iv)$$

**Step 5:**

$$\sin^{-1}(\sin 5) + \cos^{-1}(\cos 10) + \tan^{-1}(\tan(-6)) + \cot^{-1}(\cot(-10))$$

$$= 5 - 2\pi + 4\pi - 10 + 2\pi - 6 + 4\pi - 10 \text{ (from (i), (ii), (iii), (iv))}$$

$$= 8\pi - 21$$

Let us consider the function $y = \sec^{-1}(\sec x)$.

Given function is an inverse secant function

$$\therefore \text{Range of the function is } y \in [0, \pi] - \left\{ \frac{\pi}{2} \right\}$$

Also, x is the argument of the secant function

$$\therefore \text{Domain of the function is } x \in \mathbb{R} - \left\{ (2n+1)\frac{\pi}{2} \right\}$$

We know that the period of $\sec x$ is 2π .

\therefore Period of $y = \sec^{-1}(\sec x)$ is also 2π

So, let's plot the graph of $y = \sec^{-1}(\sec x)$ for the length 2π .

$$\Rightarrow y = \sec^{-1}(\sec x)$$

$$\Rightarrow \sec y = \sec x$$

$$\Rightarrow \cos x = \cos y$$

$$\Rightarrow y = 2n\pi \pm x \dots (i)$$

Hence, the graph of the function

$y = \sec^{-1}(\sec x)$ will be a replica of the graph of the function $y = \cos^{-1}(\cos x)$, except that

$y = \sec^{-1}(\sec x)$ is not defined for

$$x = \left\{ (2n+1)\frac{\pi}{2} \right\}; n \in \mathbb{Z}$$

Now, let us consider the function

$$y = \operatorname{cosec}^{-1}(\operatorname{cosec} x).$$

Given function is an inverse cosecant function

$$\therefore \text{Range of the function is } y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$$

Also, x is the argument of the cosecant function

$$\therefore \text{Domain of the function is } x \in \mathbb{R} - \{n\pi\}$$

We know that the period of $\operatorname{cosec} x$ is 2π

$$\therefore \text{Period of } y = \operatorname{cosec}^{-1}(\operatorname{cosec} x) \text{ is also } 2\pi$$

So, let's plot the graph of

$$y = \operatorname{cosec}^{-1}(\operatorname{cosec} x) \text{ for the length } 2\pi.$$

$$\Rightarrow y = \operatorname{cosec}^{-1}(\operatorname{cosec} x)$$

$$\Rightarrow \operatorname{cosec} y = \operatorname{cosec} x$$

$$\Rightarrow \sin y = \sin x$$


$$\Rightarrow y = n\pi + (-1)^n x \dots (i)$$

Hence, the graph of the function

$y = \operatorname{cosec}^{-1}(\operatorname{cosec} x)$ will be a replica of the graph of the function $y = \sin^{-1}(\sin x)$, except

$y = \operatorname{cosec}^{-1}(\operatorname{cosec} x)$ is not defined for

$$x = n\pi; n \in \mathbb{Z}$$



Property 3: Function of the Form $f^{-1}(f(x))$ (For the Principal Values of x Only)

- $\sin^{-1}(\sin(x)) = x; \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
- $\cos^{-1}(\cos(x)) = x; \forall x \in [0, \pi]$
- $\tan^{-1}(\tan(x)) = x; \forall x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
- $\cot^{-1}(\cot(x)) = x; \forall x \in (0, \pi)$
- $\operatorname{cosec}^{-1}(\operatorname{cosec} x) = x; \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
- $\sec^{-1}(\sec(x)) = x; \forall x \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$



If $x^2 + 2x + n > 10 + \sin^{-1}(\sin 9) + \tan^{-1}(\tan 9)$ for all real numbers x , then the possible value of n can be:

- (a) 11 (b) 12 (c) 13 (d) 14

Solution


Step 1:

From the graph of $y = \sin^{-1}(\sin x)$, we get,

$$\text{For } x = 9, y = 3\pi - x$$

$$\Rightarrow y = 3\pi - 9$$

$$\Rightarrow \sin^{-1}(\sin 9) = 3\pi - 9 \dots (i)$$




Step 2:

From the graph of $y = \tan^{-1}(\tan x)$

$$\text{For } x = 9, y = x - 3\pi, \text{ we get,}$$

$$\Rightarrow y = 9 - 3\pi$$

$$\Rightarrow \tan^{-1}(\tan 9) = 9 - 3\pi \dots (ii)$$



Step 3:

$$x^2 + 2x + n > 10 + \sin^{-1}(\sin 9) + \tan^{-1}(\tan 9)$$

$$x^2 + 2x + n > 10 + 3\pi - 9 + 9 - 3\pi \text{ (from (i) and (ii))}$$

$$x^2 + 2x + n > 10$$

$$x^2 + 2x + (n - 10) > 0$$

$$\text{Here, } a > 0$$

We know that for $a > 0, y = ax^2 + bx + c > 0 \forall x \in \mathbb{R}$ only if $D < 0$

$$2^2 - 4(n - 10) < 0$$

$$4 - 4n + 40 < 0$$

$$n > 11$$

\therefore Options (b), (c), and (d) are the correct answers.

Property 4: Properties of Inverse Function $f^{-1}\left(\frac{1}{x}\right)$

$$\bullet \operatorname{cosec}^{-1}(x) = \sin^{-1}\left(\frac{1}{x}\right); |x| \geq 1$$

Proof

Let $\operatorname{cosec}^{-1}(x) = \theta$; $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
 $\operatorname{cosec} \theta = x$

$$\Rightarrow \frac{1}{\sin \theta} = x$$

$$\Rightarrow \sin \theta = \left(\frac{1}{x}\right)$$

$$\Rightarrow \theta = \sin^{-1}\left(\frac{1}{x}\right)$$

$$\Rightarrow \operatorname{cosec}^{-1}(x) = \sin^{-1}\left(\frac{1}{x}\right)$$

As $\frac{1}{x} \neq 0$, so $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$

Similarly,

$$\bullet \sec^{-1}(x) = \cos^{-1}\left(\frac{1}{x}\right); |x| \geq 1$$

$$\bullet \cot^{-1}(x) = \begin{cases} \tan^{-1}\left(\frac{1}{x}\right); x > 0 \\ \pi + \tan^{-1}\left(\frac{1}{x}\right); x < 0 \end{cases}$$

Justification:

Range of $\cot^{-1}x$ is $(0, \pi)$, but the range of $\tan^{-1}x$ is $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$. So, we can not equate L.H.S. and R.H.S. directly.

For $x > 0$, both L.H.S. and R.H.S. lie in the interval $\left(0, \frac{\pi}{2}\right)$.

For $x < 0$, L.H.S. lies in the interval $\left(\frac{\pi}{2}, \pi\right)$, and R.H.S. lies in the interval $\left(-\frac{\pi}{2}, 0\right)$. So, π is added to the R.H.S. to make the range of L.H.S. and R.H.S. equal



Note

$\sin^{-1}x = \operatorname{cosec}^{-1}\left(\frac{1}{x}\right) \rightarrow$ Not identical, because L.H.S. $\sin^{-1}x$ is defined for $x = 0$, but R.H.S. $\operatorname{cosec}^{-1}\left(\frac{1}{x}\right)$ is not defined for $x = 0$.

Due to the same reason, $\sec^{-1}(x) = \cos^{-1}\left(\frac{1}{x}\right) \rightarrow$ Not identical



Find the value of $\sec^{-1}(\sqrt{2}) + \cot^{-1}(-\sqrt{2}) + \tan^{-1}\left(\frac{1}{\sqrt{2}}\right)$.

Solution

Step 1:

Given,

$$\begin{aligned} & \sec^{-1}(\sqrt{2}) + \cot^{-1}(-\sqrt{2}) + \tan^{-1}\left(\frac{1}{\sqrt{2}}\right) \\ &= \sec^{-1}(\sqrt{2}) + \pi + \tan^{-1}\left(-\frac{1}{\sqrt{2}}\right) + \tan^{-1}\left(\frac{1}{\sqrt{2}}\right) \quad (\cot^{-1}(x) = \pi + \tan^{-1}\left(\frac{1}{x}\right); x < 0) \\ &= \sec^{-1}(\sqrt{2}) + \pi - \tan^{-1}\left(\frac{1}{\sqrt{2}}\right) + \tan^{-1}\left(\frac{1}{\sqrt{2}}\right) \quad (\tan^{-1}(-x) = -\tan^{-1}(x)) \\ &= \sec^{-1}(\sqrt{2}) + \pi \end{aligned}$$

Step 2:

$$\text{Let } \sec^{-1}(\sqrt{2}) = \alpha$$

$$\Rightarrow \sec \alpha = \sqrt{2}$$

$$\Rightarrow \cos \alpha = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \alpha = \frac{\pi}{4}$$

$$\therefore \sec^{-1}(\sqrt{2}) + \pi = \frac{\pi}{4} + \pi = \frac{5\pi}{4}$$

Property 5: Properties of some particular Inverse Functions

- $\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}; x \in [-1, 1]$
- $\sec^{-1}x + \operatorname{cosec}^{-1}x = \frac{\pi}{2}; |x| \geq 1$
- $\tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}; x \in \mathbb{R}$

Proof

$$\text{Let } \sin^{-1}x = \theta ; \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

$$\Rightarrow \sin \theta = x$$

$$\Rightarrow \cos\left(\frac{\pi}{2} - \theta\right) = x$$

$$\because -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

$$\Rightarrow \frac{\pi}{2} \geq -\theta \geq -\frac{\pi}{2}$$

$$\Rightarrow \frac{\pi}{2} + \frac{\pi}{2} \geq -\theta + \frac{\pi}{2} \geq -\frac{\pi}{2} + \frac{\pi}{2}$$

$$\Rightarrow \pi \geq \frac{\pi}{2} - \theta \geq 0$$

Which is the range of the inverse cosine function

$$\Rightarrow \frac{\pi}{2} - \sin^{-1}x = \cos^{-1}x$$

$$\Rightarrow \sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}$$

Similarly, the other two results can also be proved.



Find the range of $f(x) = \sin^{-1}x + \cos^{-1}x + \tan^{-1}x$.

Solution

Step 1:

Domain of $\sin^{-1}x$ is $x \in [-1, 1]$

Domain of $\cos^{-1}x$ is $x \in [-1, 1]$

Domain of $\tan^{-1}x$ is $x \in \mathbb{R}$

\therefore Domain of $f(x) = \sin^{-1}x + \cos^{-1}x + \tan^{-1}x$ is $x \in [-1, 1]$

Step 2:


We know that for $x \in [-1, 1]$,

$$\sin^{-1}(x) + \cos^{-1}(x) = \frac{\pi}{2}$$

From the graph of the inverse tangent function,

$$\text{for } x \in [-1, 1], -\frac{\pi}{4} \leq \tan^{-1}(x) \leq \frac{\pi}{4}$$

$$\Rightarrow -\frac{\pi}{4} + \frac{\pi}{2} \leq \sin^{-1}(x) + \cos^{-1}(x) + \tan^{-1}(x) \leq \frac{\pi}{4} + \frac{\pi}{2}$$



$$\Rightarrow \frac{\pi}{4} \leq f(x) \leq \frac{3\pi}{4}$$

$$\therefore \text{Range of the function } f(x) \text{ is } \left[\frac{\pi}{4}, \frac{3\pi}{4}\right]$$



The greatest and least value of $(\sin^{-1}(x))^2 + (\cos^{-1}(x))^2$ are ____ and ____, respectively.

(a) $\frac{5\pi^2}{4}, \frac{\pi^2}{8}$

(b) $\frac{\pi}{2}, -\frac{\pi}{2}$

(c) $\frac{\pi^2}{4}, -\frac{\pi^2}{4}$

(d) $\frac{\pi^2}{4}, 0$

Solution

Step 1:

Given,

$$y = (\sin^{-1}x)^2 + (\cos^{-1}x)^2$$

$$\Rightarrow y = (\sin^{-1}x)^2 + (\cos^{-1}x)^2 + 2 \sin^{-1}x \cos^{-1}x - 2 \sin^{-1}x \cos^{-1}x$$

$$\Rightarrow y = (\sin^{-1}x + \cos^{-1}x)^2 - 2 \sin^{-1}x \cos^{-1}x$$

$$\Rightarrow y = \left(\frac{\pi}{2}\right)^2 - 2 \sin^{-1}x \left(\frac{\pi}{2} - \sin^{-1}x\right)$$

$$\Rightarrow y = \frac{\pi^2}{4} - \pi \sin^{-1}x + 2(\sin^{-1}x)^2$$

$$\Rightarrow y = 2 \left\{ (\sin^{-1}x)^2 - \frac{\pi}{2} \sin^{-1}x + \frac{\pi^2}{8} \right\}$$

$$\Rightarrow y = 2 \left[(\sin^{-1}x)^2 - 2(\sin^{-1}x) \frac{\pi}{4} + \left(\frac{\pi}{4}\right)^2 - \left(\frac{\pi}{4}\right)^2 + \frac{\pi^2}{8} \right]$$

$$\Rightarrow y = 2 \left[\left(\sin^{-1}x - \frac{\pi}{4} \right)^2 + \frac{\pi^2}{16} \right]$$

Step 2:

$$\therefore y_{\min} = 2 \left[0 + \frac{\pi^2}{16} \right] = \frac{\pi^2}{8}$$

$$\text{and } y_{\max} = 2 \left[\left(-\frac{\pi}{2} - \frac{\pi}{4} \right)^2 + \frac{\pi^2}{16} \right]$$

$$= 2 \left(\frac{9\pi^2}{16} + \frac{\pi^2}{16} \right) = \frac{5\pi^2}{4}$$

\therefore Option (a) is the correct option.



Concept Check



AIEEE 2007

1. If $\sin^{-1}\left(\frac{x}{5}\right) + \operatorname{cosec}^{-1}\left(\frac{5}{4}\right) = \frac{\pi}{2}$, then the value of x is:

(a) 1

(b) 3

(c) 4

(d) 5

2. Solve $\sin^{-1}(x^2 - 2x + 1) + \cos^{-1}(x^2 - x) = \frac{\pi}{2}$



Summary Sheet



Key Takeaways

- $\sin^{-1}(\sin(x)) = x; \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
- $\cos^{-1}(\cos(x)) = x; \forall x \in [0, \pi]$
- $\tan^{-1}(\tan(x)) = x; \forall x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
- $\cot^{-1}(\cot(x)) = x; \forall x \in (0, \pi)$
- $\operatorname{cosec}^{-1}(\operatorname{cosec} x) = x; \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
- $\sec^{-1}(\sec(x)) = x; \forall x \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$
- $\operatorname{cosec}^{-1}(x) = \sin^{-1}\left(\frac{1}{x}\right); |x| \geq 1$
- $\sec^{-1}(x) = \cos^{-1}\left(\frac{1}{x}\right); |x| \geq 1$
- $\cot^{-1}(x) = \begin{cases} \tan^{-1}\left(\frac{1}{x}\right); x > 0 \\ \pi + \tan^{-1}\left(\frac{1}{x}\right); x < 0 \end{cases}$
- $\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}; x \in [-1, 1]$
- $\sec^{-1}x + \operatorname{cosec}^{-1}x = \frac{\pi}{2}; |x| \geq 1$
- $\tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}; x \in \mathbb{R}$



Mind Map

Properties of inverse function

Properties of inverse function

$$f^{-1}(x) + g^{-1}(x) = \frac{\pi}{2}$$

Properties of inverse function $f^{-1}\left(\frac{1}{x}\right)$

Function of the form $f^{-1}(f(x))$



Self-Assessment

A value of x satisfying the equation $\sin[\cot^{-1}(1+x)] = \cos[\tan^{-1}x]$ is:

- (a) $\frac{1}{2}$ (b) -1 (c) 0 (d) $-\frac{1}{2}$



Answers

Concept Check

1.

Step 1:

Given,

$$\sin^{-1}\left(\frac{x}{5}\right) + \operatorname{cosec}^{-1}\left(\frac{5}{4}\right) = \frac{\pi}{2}$$

$$\sin^{-1}\left(\frac{x}{5}\right) = \frac{\pi}{2} - \operatorname{cosec}^{-1}\left(\frac{5}{4}\right)$$

$$\sin^{-1}\left(\frac{x}{5}\right) = \sec^{-1}\left(\frac{5}{4}\right) \quad (\because \sec^{-1}x + \operatorname{cosec}^{-1}x = \frac{\pi}{2}, |x| \geq 1)$$


$$\sin^{-1}\left(\frac{x}{5}\right) = \cos^{-1}\left(\frac{4}{5}\right) \quad (\because \sec^{-1}x = \cos^{-1}\left(\frac{1}{x}\right); |x| \geq 1)$$

Step 2:

$$\text{Let } \cos^{-1}\left(\frac{4}{5}\right) = \theta, \theta \in \left(0, \frac{\pi}{2}\right)$$

$$\Rightarrow \cos \theta = \frac{4}{5}$$

Here, θ is an acute angle in the right angle triangle, as shown in the figure.



$$\sin^{-1}\left(\frac{x}{5}\right) = \theta$$

$$\Rightarrow \frac{x}{5} = \sin \theta$$

$$\Rightarrow \frac{x}{5} = \frac{3}{5} \quad (\text{From the triangle})$$

$$\Rightarrow x = 3$$

\therefore Option (b) is the correct answer.

2.

$$-1 \leq x^2 - 2x + 1 \leq 1 \dots (i)$$

$$-1 \leq x^2 - x \leq 1 \dots (ii)$$

$$\text{Given, } \sin^{-1}(x^2 - 2x + 1) + \cos^{-1}(x^2 - x) = \frac{\pi}{2}$$

$$\Rightarrow x^2 - 2x + 1 = x^2 - x \quad [\sin^{-1}(A) + \cos^{-1}(A) = \frac{\pi}{2}]$$

$$\Rightarrow 2x - x = 1$$

$$\Rightarrow x = 1$$

$x = 1$ satisfies both (i) and (ii)

$$\text{Hence, } x = 1 \text{ is the solution of } \sin^{-1}(x^2 - 2x + 1) + \cos^{-1}(x^2 - x) = \frac{\pi}{2}$$

Self Assessment

Step 1:

Given equation is $\sin[\cot^{-1}(1+x)] = \cos[\tan^{-1}x]$

Let $\cot^{-1}(1+x) = a$

$$\Rightarrow \cot a = 1+x$$

We know,

$$\operatorname{cosec} a = \sqrt{1 + \cot^2 a} = \sqrt{1 + (1+x)^2} = \sqrt{x^2 + 2x + 2}$$

$$\text{Also, } \sin a = \frac{1}{\operatorname{cosec} a}$$

$$\Rightarrow \sin a = \frac{1}{\sqrt{x^2 + 2x + 2}}$$

$$\Rightarrow a = \sin^{-1}\left(\frac{1}{\sqrt{x^2 + 2x + 2}}\right)$$

Let $\tan^{-1}x = b$

$$\Rightarrow x = \tan b$$

We know that $\sec b = \sqrt{1 + \tan^2 b} = \sqrt{1 + x^2}$

$$\text{Also, } \cos b = \frac{1}{\sec b}$$

$$\Rightarrow \cos b = \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow b = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)$$

Step 2:

Given equation is $\sin[\cot^{-1}(1+x)] = \cos[\tan^{-1}x]$

$$\Rightarrow \sin\left(\sin^{-1}\left(\frac{1}{\sqrt{x^2 + 2x + 2}}\right)\right) = \cos\left(\cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right)\right)$$

$$\Rightarrow \frac{1}{\sqrt{x^2 + 2x + 2}} = \frac{1}{\sqrt{1+x^2}}$$

$$\Rightarrow \sqrt{1+x^2} = \sqrt{x^2 + 2x + 2}$$

$$\Rightarrow 1 + x^2 = x^2 + 2x + 2 \quad (\text{Squaring on both sides})$$

$$\Rightarrow 2x = -1$$

$$\Rightarrow x = -\frac{1}{2}$$