

CBSE Board Class 10 Maths Chapter 8- Introduction to Trigonometry
Objective Questions

Introduction

1. In a right triangle ABC, the right angle is at B. Which of the following is true about the other two angles A and C?

- (A) There is no restriction on the measure of the angles
- (B) Both the angles should be obtuse
- (C) Both the angles should be acute
- (D) One of the angles is acute and the other is obtuse


Answer: (C) Both the angles should be acute

Solution: In triangle ABC, $\angle A + \angle B + \angle C = 180^\circ$

$$\angle A + \angle C = 180^\circ - 90^\circ = 90^\circ \Rightarrow \text{None of the angles can be } \geq 90^\circ$$

\therefore The other 2 angles must be acute angles.

2. In a right triangle ABC, the right angle is at B. What is the length of missing side in the figure?




- (A) 25 cm
- (B) 12cm
- (C) 7cm
- (D) 5cm

Answer: (D) 5cm

Solution: Pythagoras theorem: In a right angled triangle,

Hypotenuse² = Sum of squares of other 2 sides

That is,



$$c^2 = a^2 + b^2$$

Here $a = 4$ cm and $b = 3$ cm,

$$\text{So the missing side} = c = \sqrt{3^2 + 4^2} = 5 \text{ cm}$$

3. Which of the following numbers can form sides of a right angled triangle?

- (A) 13 cm , 27 cm , 15 cm
- (B) 4 cm , 5 cm , 9 cm
- (C) 2 cm , 17 cm , 9 cm
- (D) 10 cm , 6 cm , 8 cm

Answer: (D) 10 cm, 6 cm, 8 cm

Solution: The basic condition for any type of triangle is:


- (i) The sum of 2 sides of a triangle should be greater than the third side
- (ii) The difference of any 2 sides should be less than the third side.

For a triangle to be a right angled triangle, there is an additional condition.

Pythagoras theorem: In a right angled triangle,
Hypotenuse²= Sum of squares of other 2 sides

That is, $c^2=a^2+b^2$; Also note that the hypotenuse is the largest side in a right triangle.

Considering each of the given options,



$$10^2=6^2+8^2$$

$$17^2\neq 2^2+9^2$$

$$9^2\neq 5^2+4^2$$

$$27^2\neq 13^2+15^2$$

So, A is the correct option.

4. Which of the following are Pythagorean triplets?

(A) 4 cm , 6 cm , 8 cm

(B) 24 cm , 10 cm , 26 cm

(C) 13 cm , 27 cm , 30 cm

(D) 2 cm , 17 cm , 9 cm

Answer: (B) 24 cm, 10 cm, 26 cm

Solution: Pythagorean triplets are those set of numbers which satisfy the Pythagoras theorem.

Considering the options given to us –

$$8^2\neq 4^2+6^2$$

$$17^2\neq 2^2+9^2$$

$$26^2 = 24^2 + 10^2$$

$$30^2 \neq 27^2 + 13^2$$

Therefore, 24, 10 and 26 are Pythagorean triplets.

Trigonometric Identities

5. If $\sec\theta + \tan\theta = x$, then $\tan\theta$ is:

(A) $(x^2 - 1) / 2x$

(B) $(x^2 + 1) / 2x$

(C) $(x^2 - 1) / x$

(D) $(x^2 + 1) / x$

Answer: (A) $(x^2 - 1) / 2x$

Solution: We know that, $\sec^2\theta - \tan^2\theta = 1$

$$\text{Therefore, } (\sec\theta + \tan\theta)(\sec\theta - \tan\theta) = 1$$

$$\text{Since, } (\sec\theta + \tan\theta) = x$$

$$\text{Thus, } (\sec\theta - \tan\theta) = 1/x$$

Solving both equations

$$\text{We get } \tan\theta = (x^2 - 1) / 2x$$

6. If $p \cot\theta = \sqrt{q^2 - p^2}$ then the value of $\sin\theta$ is _____. (θ being an acute angle)

(A) $q/3p$

(B) $q/2p$

(C) p/q

(D) 0

Answer: (C) p/q

$$\text{Given, } p \cot\theta = \sqrt{q^2 - p^2}$$

$$\therefore \cot \theta = (\sqrt{q^2 - p^2}) / 2$$

Using the identity, $\operatorname{cosec}^2 \theta = 1 + \cot^2 \theta$

$$= 1 + \frac{\sqrt{q^2 - p^2}}{p^2}$$

$$= q^2 / p^2$$

Hence, $\operatorname{cosec} \theta = q/p$

$$\therefore \sin \theta = p/q$$

7. If $\sin A = 8/17$, find the value of $\sec A \cos A + \operatorname{cosec} A \cos A$.

(A) 23/8

(B) 15/8

(C) 8/15

(D) 6/23

Answer: (A) 23/8

Solution: $\sin A = 8/17$

$$\operatorname{cosec} A = 17/8$$

$$\cos A = \sqrt{1 - \sin^2 A}$$

$$= \sqrt{1 - \frac{64}{289}} = \sqrt{\frac{225}{289}} = 15/17$$

$$\sec A = 17/15$$

$$\sec A \cos A + \operatorname{cosec} A \cos A = (17/15) * (15/17) + (17/15) * (15/17)$$

$$= 1 + (15/8)$$

$$= 23/8$$

8. $(\sin A - 2 \sin^3 A) / (2 \cos^3 A - \cos A) =$

- (A) $\tan A$
- (B) $\cot A$
- (C) $\sec A$
- (D) 1

Answer: (A) $\tan A$

Solutions: $(\sin A - 2 \sin^3 A) / (2 \cos^3 A - \cos A) = (\sin A (1 - 2 \sin^2 A)) / (\cos A (2 \cos^2 A - 1))$

$$= (\sin A (\sin^2 A + \cos^2 A - 2 \sin^2 A)) / (\cos A (2 \cos^2 A - (\sin^2 A + \cos^2 A)))$$

$$= (\sin A (\cos^2 A - \sin^2 A)) / (\cos A (\cos^2 A - \sin^2 A))$$

$$= \tan A$$

Trigonometric Ratios

9. $(\cos A / \cot A) + \sin A =$ _____

- (A) $\cot A$
- (B) $2 \sin A$
- (C) $2 \cos A$
- (D) $\sec A$

Answer: (B) $2 \sin A$

Solution: $(\cos A / \cot A) + \sin A$

$$= \cos A / (\cos A / \sin A) + \sin A$$

$$= \sin A + \sin A$$

$$= 2 \sin A$$

10. If $5 \tan \theta = 4$, then value of $(5 \sin \theta - 4 \cos \theta) / (5 \sin \theta + 4 \cos \theta)$ is:

- (A) $1/6$
- (B) $5/6$
- (C) 0
- (D) $5/3$

Answer: (C) 0

Solution: Divide both numerator and denominator by $\cos \theta$ and solve

$$(5 \sin \theta - 4 \cos \theta) / (5 \sin \theta + 4 \cos \theta)$$

$$= \frac{\frac{(5 \sin \theta - 4 \cos \theta)}{\cos \theta}}{\frac{(5 \sin \theta + 4 \cos \theta)}{\cos \theta}}$$

$$= \frac{5 \tan \theta - 4}{5 \tan \theta + 4}$$

$$= \frac{4 - 4}{4 + 4}$$

$$= 0$$

(Since, given that $5 \tan \theta = 4$)

11. In $\triangle PQR$, $PQ = 12$ cm and $PR = 13$ cm. $\angle Q = 90^\circ$ Find $\tan P - \cot R$

(A) $-(119/60)$


(B) $119/60$

(C) 0

(D) 1

Answer: (C) 0

Solution:



Given that in $\triangle PQR$, $PQ = 12$ cm and $PR = 13$ cm.

Now, from Pythagoras theorem,

$$PQ^2 + QR^2 = PR^2$$

$$\Rightarrow QR^2 = PR^2 - PQ^2$$

$$\Rightarrow QR^2 = 13^2 - 12^2$$

$$\Rightarrow QR^2 = 169 - 144 = 25$$

$$\Rightarrow QR = \sqrt{25} = 5 \text{ cm}$$

Now, $\tan P = \text{opposite side} / \text{adjacent side} = QR/PQ = 5/12$

$\cot R = \text{adjacent side} / \text{opposite side} = QR/PQ = 5/12$

$$\therefore \tan P - \cot R = (5/12) - (5/12) = 0$$

12. If $\tan \theta = (x \sin \phi) / (1 - x \cos \phi)$ and, $\tan \phi = (y \sin \theta) / (1 - y \cos \theta)$ then $x/y =$

(A) $\sin \theta / (1 - \cos \phi)$

(B) $\sin \theta / (1 - \cos \theta)$

(C) $\sin \theta / \sin \phi$

(D) $\sin \phi / \sin \theta$

Answer: (C) $\sin \theta / \sin \phi$

Solution: We have, $\tan \theta = (x \sin \phi) / (1 - x \cos \phi)$

$$\Rightarrow (1 - x \cos \phi) / (x \sin \phi) = 1 / \tan \theta \Rightarrow (1 / x \sin \phi) - \cot \phi = \cot \theta$$

$$\Rightarrow 1 / x \sin \phi = \cot \theta + \cot \phi$$

$$\text{and } \tan \phi = y \sin \theta / (1 - y \cos \theta) \Rightarrow (1 - y \cos \theta) / y \sin \theta = 1 / \tan \phi$$

$$\Rightarrow (1 / y \sin \theta) - \cot \theta = \cot \phi \Rightarrow (1 / y \sin \theta) = \cot \phi + \cot \theta$$

$$\Rightarrow (1 / y \sin \theta) = (1 / x \sin \phi) \Rightarrow x/y = \sin \theta / \sin \phi$$

Trigonometric Ratios of Complementary Angles

13. The value of $\tan 1^\circ \times \tan 2^\circ \times \tan 3^\circ \times \dots \times \tan 89^\circ$ is :

(A) $\frac{1}{2}$

(B) 2

(C) 1

(D) 0

Answer: (C) 1

Solution: $\tan\theta\cot\theta=1$,

$$\tan(90-\theta)=\cot\theta$$

$$\text{and } \tan 45^\circ=1$$

$$\text{Given: } \tan 1^\circ \cdot \tan 2^\circ \cdot \tan 3^\circ \dots \tan 88^\circ \cdot \tan 89^\circ$$

$$= (\tan 1^\circ \cdot \tan 89^\circ) \cdot (\tan 2^\circ \cdot \tan 88^\circ) \dots (\tan 44^\circ \cdot \tan 46^\circ) (\tan 45^\circ)$$

$$= [(\tan 1^\circ \cdot \tan(90^\circ-1^\circ))] \cdot [(\tan 2^\circ \cdot \tan(90^\circ-2^\circ))] \dots [(\tan 44^\circ \cdot \tan(90^\circ-44^\circ))] \cdot 1$$

$$= (\tan 1^\circ \cdot \cot 1^\circ) \cdot (\tan 2^\circ \cdot \cot 2^\circ) \dots (\tan 44^\circ \cdot \cot 44^\circ)$$

$$= 1$$

14. If $\tan 2A = \cot(A-18^\circ)$, then value of A is:

(A) 27°

(B) 24°

(C) 36°

(D) 18°

Answer: (C) 36°

Solution: Given, $\tan 2A = \cot(A - 18^\circ)$

$$\Rightarrow \tan 2A = \tan(90 - (A - 18^\circ))$$

$$\Rightarrow \tan 2A = \tan(108^\circ - A)$$

$$\Rightarrow 2A = 108^\circ - A$$

$$\Rightarrow 3A = 108^\circ$$

$$\Rightarrow A = 36^\circ$$

15. If $\tan 4\theta = \cot(\theta - 10^\circ)$, where 4θ and $(\theta - 10^\circ)$ are acute angles then the value of θ in degrees is

- (A) 16°
- (B) 20°
- (C) 32°
- (D) 40°

Answer: (B) 20°

Solution: Given, $\tan 4\theta = \cot(\theta - 10^\circ)$

This can be written as

$$\cot(90^\circ - 4\theta) = \cot(\theta - 10^\circ) \text{ -----(i)}$$

$$(\because \tan \theta = \cot(90^\circ - \theta))$$


Hence, from (i) we have

$$\Rightarrow 90^\circ - 4\theta = \theta - 10^\circ$$

$$\Rightarrow 5\theta = 100^\circ$$

$$\Rightarrow \theta = 20^\circ$$

16. In the given triangle right angled at B, which pair of angles are complementary?



- (A) None of these
- (B) C and A
- (C) A and B
- (D) B and C

Answer: (B) C and A

Solution: Two angles are said to be complementary, if their sum is 90° . The triangle is right angled at B. With angle sum property of the triangle, $\angle A + \angle B + \angle C = 180^\circ$

$\angle A + \angle C = 90^\circ$, Hence angle A and C are complementary.

Trigonometric Ratios of Specific Angles

17. Which of the following is correct for some θ , such that $0^\circ \leq \theta < 90^\circ$

- (A) $1/\cos \theta < 1$
- (B) $\sec \theta = 0$
- (C) $1/\sec \theta < 1$
- (D) $1/\sec \theta > 1$

Answer: (C) $1/\sec \theta < 1$

Solution: $1/\sec \theta = \cos \theta$. And value of $\cos \theta$ ranges from 0 to 1

18. The value $\cot^2 30^\circ - 2\cos^2 60^\circ - 3/4 \sec^2 45^\circ - 4\sin^2 30^\circ$ is

- (A) 2
- (B) -1
- (C) 1
- (D) 0

Answer: (D) 0

Solution: $\cot^2 30^\circ - 2\cos^2 60^\circ - 3/4 (\sec^2 45^\circ) - 4\sin^2 30^\circ$

$$= (\sqrt{3})^2 - 2\left(\frac{1}{2}\right)^2 - \frac{3}{4}(\sqrt{2})^2 - 4\left(\frac{1}{2}\right)^2$$

$$= 3 - (1/2) - (3/2) - 1 = 0$$

19. If $\operatorname{Cosec}(A+B) = \frac{2}{\sqrt{3}}$ $\sec(A-B) = \frac{2}{\sqrt{3}}$

$0^\circ < A+B \leq 90^\circ$,
Find A and B.

- (A) $25^\circ, 35^\circ$
- (B) $30^\circ, 30^\circ$
- (C) $45^\circ, 15^\circ$
- (D) $10^\circ, 50^\circ$

Answer: (C) $45^\circ, 15^\circ$

Solution: If $A+B$ lies in this range $0^\circ < A+B \leq 90^\circ$

$$\operatorname{cosec}(A+B) = \frac{2}{\sqrt{3}} \quad \text{only when } A+B=60^\circ \dots\dots\dots (1)$$

$$\sec(A-B) = \frac{2}{\sqrt{3}} \quad \text{only when } A-B=30^\circ \dots\dots\dots (2)$$

By Solving equation 1 and equation 2
 $A=45^\circ$ and $B=15^\circ$

20. $\cos 1^\circ \times \cos 2^\circ \times \cos 3^\circ \times \dots \times \cos 180^\circ$ is equal to:

- (A) 0
- (B) 1
- (C) $\frac{1}{2}$
- (D) -1

Answer: (A) 0

Solution: Since $\cos 90^\circ = 0$
The given expression

$$\cos 1^\circ \times \cos 2^\circ \times \cos 3^\circ \times \dots \times \cos 90^\circ \times \dots \times \cos 180^\circ$$

reduces to zero as it contains $\cos 90^\circ$ which is equal to 0