

QUADRATIC EQUATION

1. SOLUTION OF QUADRATIC EQUATION & RELATION BETWEEN ROOTS & CO-EFFICIENTS :

(a) The solutions of the quadratic equation, $ax^2 + bx + c = 0$ is

$$\text{given by } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

(b) The expression $b^2 - 4ac \equiv D$ is called the discriminant of the quadratic equation.

(c) If α & β are the roots of the quadratic equation $ax^2 + bx + c = 0$, then ;

$$(i) \alpha + \beta = -b/a \quad (ii) \alpha\beta = c/a \quad (iii) |\alpha - \beta| = \sqrt{D}/|a|$$

(d) Quadratic equation whose roots are α & β is $(x - \alpha)(x - \beta) = 0$ i.e.

$$x^2 - (\alpha + \beta)x + \alpha\beta = 0 \text{ i.e. } x^2 - (\text{sum of roots})x + \text{product of roots} = 0.$$

2. NATURE OF ROOTS :

(a) Consider the quadratic equation $ax^2 + bx + c = 0$ where $a, b, c \in \mathbb{R}$ & $a \neq 0$ then ;

(i) $D > 0 \Leftrightarrow$ roots are real & distinct (unequal).

(ii) $D = 0 \Leftrightarrow$ roots are real & coincident (equal)

(iii) $D < 0 \Leftrightarrow$ roots are imaginary.

(iv) If $p + iq$ is one root of a quadratic equation, then the other root must be the conjugate $p - iq$ & vice versa.

$$(p, q \in \mathbb{R} \text{ \& } i = \sqrt{-1}).$$

(b) Consider the quadratic equation $ax^2 + bx + c = 0$

where $a, b, c \in \mathbb{Q}$ & $a \neq 0$ then ;

(i) If D is a perfect square, then roots are rational.

- (ii) If $\alpha = p + \sqrt{q}$ is one root in this case, (where p is rational & \sqrt{q} is a surd) then other root will be $p - \sqrt{q}$.

3. COMMON ROOTS OF TWO QUADRATIC EQUATIONS

- (a) Only one common root.

Let α be the common root of $ax^2 + bx + c = 0$ & $a'x^2 + b'x + c' = 0$ then $a\alpha^2 + b\alpha + c = 0$ & $a'\alpha^2 + b'\alpha + c' = 0$. By Cramer's

$$\text{Rule } \frac{\alpha^2}{bc' - b'c} = \frac{\alpha}{a'c - ac'} = \frac{1}{ab' - a'b}$$

$$\text{Therefore, } \alpha = \frac{ca' - c'a}{ab' - a'b} = \frac{bc' - b'c}{a'c - ac'}$$

So the condition for a common root is

$$(ca' - c'a)^2 = (ab' - a'b)(bc' - b'c)$$

- (b) If both roots are same then $\frac{a}{a'} = \frac{b}{b'} = \frac{c}{c'}$

4. ROOTS UNDER PARTICULAR CASES

Let the quadratic equation $ax^2 + bx + c = 0$ has real roots and

- (a) If $b = 0 \Rightarrow$ roots are of equal magnitude but of opposite sign

- (b) If $c = 0 \Rightarrow$ one root is zero other is $-b/a$

- (c) If $a = c \Rightarrow$ roots are reciprocal to each other

- (d) If $\left. \begin{matrix} a > 0, c < 0 \\ a < 0, c > 0 \end{matrix} \right\} \Rightarrow$ roots are of opposite signs

- (e) If $\left. \begin{matrix} a > 0, b > 0, c > 0 \\ a < 0, b < 0, c < 0 \end{matrix} \right\} \Rightarrow$ both roots are negative.

- (f) If $\left. \begin{matrix} a > 0, b < 0, c > 0 \\ a < 0, b > 0, c < 0 \end{matrix} \right\} \Rightarrow$ both roots are positive.

- (g) If sign of $a =$ sign of $b \neq$ sign of $c \Rightarrow$ Greater root in magnitude is negative.

- (h) If sign of $b =$ sign of $c \neq$ sign of $a \Rightarrow$ Greater root in magnitude is positive.

- (i) If $a + b + c = 0 \Rightarrow$ one root is 1 and second root is c/a .

5. MAXIMUM & MINIMUM VALUES OF QUADRATIC EXPRESSION :



Maximum & Minimum Values of expression $y = ax^2 + bx + c$ is $\frac{-D}{4a}$ which occurs at $x = -(b/2a)$ according as $a < 0$ or $a > 0$.

$$y \in \left[\frac{-D}{4a}, \infty \right) \text{ if } a > 0 \quad \& \quad y \in \left(-\infty, \frac{-D}{4a} \right] \text{ if } a < 0.$$



6. LOCATION OF ROOTS :

Let $f(x) = ax^2 + bx + c$, where $a, b, c \in \mathbb{R}$, $a \neq 0$



- (a) Conditions for both the roots of $f(x) = 0$ to be greater than a specified number 'd' are $D \geq 0$; $a \cdot f(d) > 0$ & $(-b/2a) > d$.



- (b) Conditions for the both roots of $f(x) = 0$ to lie on either side of the number 'd' in other words the number 'd' lies between the roots of $f(x) = 0$ is $a \cdot f(d) < 0$.





- (c) Conditions for exactly one root of $f(x) = 0$ to lie in the interval (d, e) i.e., $d < x < e$ is $f(d) \cdot f(e) < 0$



- (d) Conditions that both roots of $f(x) = 0$ to be confined between the numbers d & e are (here $d < e$).

$$D \geq 0; a \cdot f(d) > 0 \text{ \& } af(e) > 0; d < (-b/2a) < e$$



7. GENERAL QUADRATIC EXPRESSION IN TWO VARIABLES :

$f(x, y) = ax^2 + 2hxy + by^2 + 2gx + 2fy + c$ may be resolved into two linear factors if ;

$$\Delta = abc + 2fgh - af^2 - bg^2 - ch^2 = 0 \quad \text{OR} \quad \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = 0$$

8. THEORY OF EQUATIONS :

If $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$ are the roots of the equation ;

$f(x) = a_0 x^n + a_1 x^{n-1} + a_2 x^{n-2} + \dots + a_{n-1} x + a_n = 0$ where a_0, a_1, \dots, a_n are constants $a_0 \neq 0$ then,

$$\begin{aligned} \sum \alpha_1 &= -\frac{a_1}{a_0}, \quad \sum \alpha_1 \alpha_2 = +\frac{a_2}{a_0}, \quad \sum \alpha_1 \alpha_2 \alpha_3 \\ &= -\frac{a_3}{a_0}, \dots, \alpha_1 \alpha_2 \alpha_3 \dots \alpha_n = (-1)^n \frac{a_n}{a_0} \end{aligned}$$

Note :

(i) Every odd degree equation has at least one real root whose sign is opposite to that of its last term, when coefficient of highest degree term is (+)ve [If not then make it (+) ve].

$$\text{Ex. } x^3 - x^2 + x - 1 = 0$$

(ii) Even degree polynomial whose last term is (-)ve & coefficient of highest degree term is (+)ve has atleast two real roots, one (+)ve & one (-)ve.

(iii) If equation contains only even power of x & all coefficient are (+)ve, then all roots are imaginary.