

Topic : Gaseous State

Type of Questions		M.M., Min.
Single choice Objective ('-1' negative marking) Q.2 to Q.3	(3 marks, 3 min.)	[6, 6]
Multiple choice objective ('-1' negative marking) Q.1,4	(4 marks, 4 min.)	[8, 8]
Subjective Questions ('-1' negative marking) Q.5,9	(4 marks, 5 min.)	[8, 10]
Comprehension ('-1' negative marking) Q.6 to Q.8	(3 marks, 3 min.)	[9, 9]

- A gaseous organic compound has a density of 2.5 kg/m^3 at 2 atm and at 273°C . The molecular formula of the compound can be :
 (A) $\text{C}_3\text{H}_4\text{O}$ (B) $\text{C}_4\text{H}_6\text{O}$ (C) C_4H_8 (D) C_5H_{10}
- The ratio of rates of diffusion of SO_2 , O_2 and CH_4 under identical conditions is :
 (A) $1 : \sqrt{2} : 2$ (B) $1 : 2 : 4$ (C) $2 : \sqrt{2} : 1$ (D) $1 : 2 : \sqrt{2}$
- If the number of molecules of SO_2 (molecular weight = 64) effusing through an orifice of unit area of cross-section in unit time at 0°C and 1 atm pressure is n , the number of He molecules (atomic weight = 4) effusing under similar conditions at 273°C and 0.25 atm is :
 (A) $\frac{n}{\sqrt{2}}$ (B) $n\sqrt{2}$ (C) $2n$ (D) $\frac{n}{2}$
- The time taken for effusion of 32 mL of oxygen gas will be the same as the time taken for effusion of which gas sample under identical conditions : (Take $\sqrt{2} = 1.4$, $\sqrt{3} = 1.7$)
 (A) 64 mL of H_2 (B) 50 mL of N_2 (C) 44.8 mL of CH_4 (D) 22.4 mL of SO_2
- 5 mL of He gas diffuses out in 1 second from a hole. Find the volume of SO_2 that will diffuse out from the same hole under identical conditions in 2 seconds.

Comprehension # (Q.6 to Q.8)
Graham's Law :

"Under similar conditions of pressure (partial pressure), the rate of diffusion of different gases is inversely proportional to square root of the density of different gases."

$$\text{rate of diffusion } r \propto \frac{1}{\sqrt{d}} \quad (d = \text{density of gas})$$

$$r = \text{volume flow rate} = \frac{dV_{\text{out}}}{dt}$$

$$r = \text{moles flow rate} = \frac{dn_{\text{out}}}{dt}$$

$$r = \text{distance travelled by gaseous molecules per unit time} = \frac{dx}{dt}$$

The general form of the Grahams law of diffusion shows the variation of rate of diffusion of a gas with pressure of gas, temperature of gas, area of cross-section of orifice and molecular mass of the gas.

Now answer the following questions :

6. A bottle of dry NH_3 & a bottle of dry HCl connected through a long tube are opened simultaneously under identical conditions at both ends. The white ammonium chloride ring first formed will be:

(A) at the centre of the tube
(B) near the HCl bottle

(C) near the NH_3 bottle
(D) throughout the length of tube

7. At room temperature, A_2 gas (vapour density = 40) at 1 atm pressure and B_2 gas (vapour density = 10) at p atm pressure are allowed to diffuse through identical pinholes from opposite ends of a glass tube of 1m length and of uniform cross-section. The two gases first meet at a distance of 60 cm from the A_2 end. The value of p is :

(A) $\frac{4}{3}$ atm
(B) $\frac{1}{3}$ atm
(C) $\frac{3}{4}$ atm
(D) $\frac{1}{6}$ atm

8. A mixture containing 2 moles of He and 1 mole of CH_4 is taken in a closed container and made to effuse through a small orifice of container. Then, which is the correct effused volume percentage of He and CH_4 initially, respectively :

(A) 40%, 60%
(B) 20% , 80%
(C) 80% , 20%
(D) 60% , 40%

9. Pressure in a bulb dropped from 2000 to 1500 mm in 50 minute, when the contained oxygen leaked through a small hole. The bulb was then completely evacuated. A mixture of oxygen and another gas of molecular weight 72 in molar ratio 1 : 1 at a total pressure of 6000 mm was introduced. Find the molar ratio of two gases remaining in the bulb after a period of 70 minute.

Answer Key

DPP No. # 30

- | | | | | |
|------------|--------|--------|---------|---------|
| 1. (A) | 2. (A) | 3. (A) | 4. (CD) | |
| 5. 2.5 ml. | 6. (B) | 7. (B) | 8. (C) | 9. 9/46 |

Hints & Solutions

DPP No. # 30

$$1. \quad d = \frac{PM}{RT} \quad \therefore 2.5 = \frac{2 \times M_{\text{gas}}}{0.082 \times 546} \quad \therefore M_{\text{gas}} = 56$$

\therefore Both (A) & (C) options are correct.

$$5. \quad \text{Rate of diffusion of He} = \frac{5\text{mL}}{15} = 5\text{ml/s} = r_{\text{He}} \text{ (say)}$$

$$\therefore r_{\text{SO}_2} = r_{\text{He}} \times \frac{1}{4} = 5\text{ml/s} \times \frac{1}{4}$$

\therefore Volume of SO_2 diffused in 2.0 seconds

$$= \frac{5}{4} \times 2 \text{ ml} = 2.5 \text{ ml Ans.}$$

$$7. \quad \frac{r_{A_2}}{r_{B_2}} = \frac{P_{A_2}}{P_{B_2}} \sqrt{\frac{V_{D_{B_2}}}{V_{D_{A_2}}}}$$

$$\frac{60/\Delta t}{40/\Delta t} = \frac{1}{P} \sqrt{\frac{10}{40}}$$

$$P = \frac{1}{3} \text{ atm}$$

8. Rate of diffusion of He = r_1
rate of diffusion of CH_4 = r_2

$$\frac{r_1}{r_2} = \frac{n_1}{n_2} \sqrt{\frac{M_2}{M_1}} = \frac{2}{1} \sqrt{\frac{16}{4}} = \frac{4}{1}$$

$$\text{Diffused mole of He} = \frac{4}{5} \times 100 = 80\%$$

$$\text{Diffused mole of CH}_4 = \frac{1}{5} \times 100 = 20\%$$

9. 39/46