

# Electrochemistry

## Question1

The mass of silver (Molar mass of Ag :  $108\text{gmol}^{-1}$ ) displaced by a quantity of electricity which displaces 5600mL of  $\text{O}_2$  at S.T.P. will be \_\_\_\_g.

[27-Jan-2024 Shift 1]

**Answer: 108**

**Solution:**

Eq. of Ag = Eq. of  $\text{O}_2$

Let x gm silver displaced,

$$\frac{x \times 1}{108} = \frac{5.6}{22.7} \times 4$$

(Molar volume of gas at STP = 22.7 lit)

$$x = 106.57 \text{ gm}$$

Ans. 107

OR,

as per old STP data, molar volume = 22.4 lit

$$\frac{x \times 1}{108} = \frac{5.6}{22.4} \times 4, x = 108 \text{ gm}.$$

Ans. 108

---

## Question2

Which of the following statements is not correct about rusting of iron?

[27-Jan-2024 Shift 2]

**Options:**

A.

Coating of iron surface by tin prevents rusting, even if the tin coating is peeling off.

B.

When pH lies above 9 or 10, rusting of iron does not take place.

C.

Dissolved acidic oxides  $\text{SO}_2$ ,  $\text{NO}_2$  in water act as catalyst in the process of rusting.

D.

Rusting of iron is envisaged as setting up of electrochemical cell on the surface of iron object.

**Answer: A**

**Solution:**

As tin coating is peeled off, then iron is exposed to atmosphere.

---

## Question3

The hydrogen electrode is dipped in a solution of  $\text{pH} = 3$  at  $25^\circ\text{C}$ . The potential of the electrode will be  $\text{---} \times 10^{-2}\text{V}$ .


$$\left( \frac{2.303 RT}{F} = 0.059\text{V} \right)$$

**[27-Jan-2024 Shift 2]**

**Options:**

**Answer: 18**

**Solution:**



$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.059}{2} \log \frac{P_{\text{H}_2}}{[\text{H}^{+}]^2}$$

$$= 0 - 0.059 \times 3 = -0.177 \text{ volts.} = -17.7 \times 10^{-2}\text{V.}$$

---

## Question4

The mass of zinc produced by the electrolysis of zinc sulphate solution with a steady current of  $0.015 \text{ A}$  for  $15 \text{ minutes}$  is  $\text{---} \times 10^{-4}\text{g}$ .


(Atomic mass of zinc =  $65.4\text{amu}$ )

**[29-Jan-2024 Shift 1]**

**Options:**

**Answer: 45.75**

**Solution:**



$$W = Z \times i \times t$$

$$= \frac{65.4}{2 \times 96500} \times 0.015 \times 15 \times 60$$

$$= 45.75 \times 10^{-4} \text{ gm}$$


---

## Question5

A constant current was passed through a solution of  $\text{AuCl}_4^{-}$  ion between gold electrodes. After a period of 10.0 minutes, the increase in mass of cathode was 1.314g. The total charge passed through the solution is  $\text{---} \times 10^{-2}\text{F}$ .

(Given atomic mass of  $\overline{\text{Au}} = 197$  )

**[29-Jan-2024 Shift 2]**

**Options:**

**Answer: 2**

**Solution:**

$$\frac{W}{E} = \frac{\text{charge}}{1F}$$

$$\frac{1.314}{\frac{197}{3}} = \frac{Q}{1F}$$

$$Q = 2 \times 10^{-2}\text{F}$$


---

## Question6

**Reduction potential of ions are given below:**




$$E^{\circ} = 1.19\text{V} \quad E^{\circ} = 1.65\text{V} \quad E^{\circ} = 1.74\text{V}$$

**The correct order of their oxidising power is:**


**[30-Jan-2024 Shift 2]**

**Options:**

A.




B.



C.



D.



**Answer: B**

**Solution:**

Higher the value of  $\oplus$  ve SRP (Std. reduction potential) more is tendency to undergo reduction, so better is oxidising power of reactant.

Hence, ox. Power:-  $\text{BrO}_4^- > \text{IO}_4^- > \text{ClO}_4^-$

---

## Question7

**Identify the factor from the following that does not affect electrolytic conductance of a solution.**

**[31-Jan-2024 Shift 1]**

**Options:**

A.

The nature of the electrolyte added.

B.

The nature of the electrode used.

C.

Concentration of the electrolyte.

D.

The nature of solvent used.

**Answer: B**

**Solution:**

Conductivity of electrolytic cell is affected by concentration of electrolyte, nature of electrolyte and nature of solvent.

---

## Question8

The metals that are employed in the battery industries are

- A. Fe
- B. Mn
- C. Ni
- D. Cr
- E. Cd

Choose the correct answer from the options given below:

[31-Jan-2024 Shift 1]

Options:

A.

B, C and E only

B.

A, B, C, D and E

C.

A, B, C and D only

D.

B, D and E only

**Answer: A**

**Solution:**

Mn, Ni and Cd metals used in battery industries.

-----

## Question9




One Faraday of electricity liberates  $x \times 10^{-1}$  gram atom of copper from copper sulphate, x is \_\_\_\_\_

[31-Jan-2024 Shift 1]

Options:

**Answer: 5**

**Solution:**



$$0.5 \text{ mol} = 0.5 \text{ g atom} = 5 \times 10^{-1}$$

$$x = 5$$

## Question10

The values of conductivity of some materials at  $298.15\text{K}^{-1}\text{Sm}^{-1}$  are  $2.1 \times 10^3$ ,  $1.0 \times 10^{-16}$ ,  $1.2 \times 10$ ,  $3.91$ ,  $1.5 \times 10^{-2}$ ,  $1 \times 10^{-7}$ ,  $1.0 \times 10^3$ . The number of conductors among the materials is \_\_\_\_

[31-Jan-2024 Shift 2]

Answer: 4

Solution:

Conductivity ( $\text{Sm}^{-1}$ )

$$\left. \begin{array}{l} 2.1 \times 10^3 \\ 1.2 \times 10 \\ 3.91 \\ 1 \times 10^3 \end{array} \right\} \text{conductors at } 298.15\text{K}$$


$1 \times 10^{-16}$  Insulator at 298.15K

$$\left. \begin{array}{l} 1.5 \times 10^{-2} \\ 1 \times 10^{-7} \end{array} \right\} \text{Semiconductor at } 298.15\text{K}$$

Therefore number of conductors is 4 .

## Question11

The potential for the given half cell at 298K is  $\times 10^{-2}\text{V}$ .



$$[\text{H}^{+}] = 1\text{M}, P_{\text{H}_2} = 2 \text{ atm}$$

$$(\text{Given: } 2.303 \text{ RT}/F = 0.06\text{V}, \log 2 = 0.3 )$$

[1-Feb-2024 Shift 1]

Options:

**Answer: 1**

**Solution:**

$$E = E_{\text{H}^+/\text{H}_2}^\circ - \frac{0.06}{2} \log \frac{P_{\text{H}_2}}{[\text{H}^+]^2}$$

$$E = 0.00 - \frac{0.06}{2} \log \frac{2}{[1]^2}$$

$$E = -0.03 \times 0.3 = -0.9 \times 10^{-2} \text{V}$$

---


## Question12

The amount of electricity in Coulomb required for the oxidation of 1mol of  $\text{H}_2\text{O}$  to  $\text{O}_2$  is \_\_\_\_\_  $\times 10^5 \text{C}$ .

**[1-Feb-2024 Shift 2]**

**Answer: 2**

**Solution:**



$$\frac{W}{E} = \frac{Q}{96500}$$

$$\text{mole} \times \text{n-factor} = \frac{Q}{96500}$$

$$1 \times 2 = \frac{Q}{96500}$$

$$Q = 2 \times 96500 \text{C}$$

$$= 1.93 \times 10^5 \text{C}$$

---

## Question13


At 298K, a 1 litre solution containing 10 mmol of  $\text{Cr}_2\text{O}_7^{2-}$  and 100 mmol of  $\text{Cr}^{3+}$  shows a pH of 3.0.

Given :  $\text{Cr}_2\text{O}_7^{2-} \rightarrow \text{Cr}^{3+}$ ;  $E^0 = 1.330\text{V}$  and  $\frac{2.303RT}{F} = 0.059\text{V}$

The potential for the half cell reaction is  $x \times 10^{-3} \text{V}$ . The value of x is \_\_\_\_\_  
**[24-Jan-2023 Shift 1]**

**Answer: 917**

**Solution:**



$$E = 1.33 - \frac{0.059}{6} \log \frac{(0.1)^2}{(10^{-2})(10^{-3})^{14}}$$

$$E = 1.33 - \frac{0.059}{6} \times 42 = 0.917$$

$$E = 917 \times 10^{-3}$$

$$x = 917$$

---


## Question14

**Choose the correct representation of conductometric titration of benzoic acid vs sodium hydroxide.**


**[24-Jan-2023 Shift 2]**

**Options:**


A.



B.




C.



D.







**Answer: B**

**Solution:**

**Solution:**




(A) → (B) Free  $\text{H}^+$  ions are replaced by  $\text{Na}^+$  which decreases conductance.

(B) → (C) Un-dissociated benzoic acid reacts with NaOH and forms salt which increases ions & conductance increases.

(C) → (D) After equivalence point at (3), NaOH added further increases  $\text{Na}^+$  &  $\text{OH}^-$  ions which further increases the conductance.

## Question15

**Consider the cell**




**Given:**  $E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^\circ = 0.771\text{V}$  and  $E_{\text{H}^+ / \frac{1}{2}\text{H}_2}^\circ = 0\text{V}$ ,  $T = 298\text{K}$

**If the potential of the cell is 0.712V the ratio of concentration of  $\text{Fe}^{2+}$  to  $\text{Fe}^{3+}$  is \_\_\_\_\_ (Nearest integer)**

**[25-Jan-2023 Shift 1]**

**Answer: 10**

**Solution:**



$$E = E^\circ - \frac{0.059}{1} \log \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]}$$

$$\Rightarrow 0.712 = (0.771 - 0) - \frac{0.059}{1} \log \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]}$$

$$\Rightarrow \log \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]} = \frac{(0.771 - 0.712)}{0.059} = 1$$

$$\Rightarrow \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]} = 10$$


---

## Question 16

**Pt( s) | H<sub>2</sub>(g)(1 bar) | H<sup>+</sup>(aq)(1M) || M<sup>3+</sup>(aq), M<sup>+</sup>(aq) | Pt( s)**

**The E<sub>cell</sub> for the given cell is 0.1115V at 298K when  $\frac{[\text{M}^+(\text{aq})]}{[\text{M}^{3+}(\text{aq})]} = 10^a$**

**The value of a is \_\_\_\_\_**

**Given : E<sup>0</sup>M<sup>3+</sup> / M<sup>+</sup> = 0.2V**


$$\frac{2.303RT}{F} = 0.059V$$

**[25-Jan-2023 Shift 2]**

**Answer: 3**

**Solution:**

Overall reaction :-



$$E_{\text{Cell}} = E_{\text{Cathode}}^0 - E_{\text{anode}}^0 - \frac{0.059}{2} \log \frac{[\text{M}^+] \times 1^2}{[\text{M}^{3+}]1}$$

$$0.1115 = 0.2 - \frac{0.059}{2} \log \frac{[\text{M}^+]}{[\text{M}^{3+}]}$$

$$3 = \log \frac{[\text{M}^+]}{[\text{M}^{3+}]}$$

$$\therefore a = 3$$


---

## Question 17

**The standard electrode potential (M<sup>3+</sup> / M<sup>2+</sup>) for V, Cr, Mn & Co are -0.26V, -0.41V, +1.57V and +1.97V, respectively. The metal ions which can liberate H<sub>2</sub> from a dilute acid are**

**[29-Jan-2023 Shift 1]**

**Options:**

A. V<sup>2+</sup> and Mn<sup>2+</sup>

B. Cr<sup>2+</sup> and Co<sup>2+</sup>

C. V<sup>2+</sup> and Cr<sup>2+</sup>

D. Mn<sup>2+</sup> and Co<sup>2+</sup>

**Answer: C**

**Solution:**

Metal cation with  $(-)$  value of reduction potential ( $M^{+3} / M^{+2}$ ) or with  $(+)$  value of oxidation potential ( $M^{+2} / M^{+3}$ ) will liberate  $H_2$

Therefore they will reduce  $H^+$


i.e.  $eV^{+2}$  and  $Cr^{+2}$

---

## Question18

Following figure shows dependence of molar conductance of two electrolytes on concentration.

$\Lambda_m^0$  is the limiting molar conductivity



The number of Incorrect statement(s) from the following is \_\_\_\_\_

(A)  $\Lambda_m$  for electrolyte A is obtained by extrapolation

(B) For electrolyte B,  $\Lambda_m$  vs  $\sqrt{c}$  graph is a straight line with intercept equal to  $\Lambda_m^0$

(C) At infinite dilution, the value of degree of dissociation approach zero for electrolyte B.

(D)  $\Lambda_m$  for any electrolyte A or B can be calculated using  $\lambda^0$  for individual ions.

[29-Jan-2023 Shift 1]

**Answer: 2**

**Solution:**


**Solution:**

Statement (A) and Statement (C) are incorrect

---

## Question19

The equilibrium constant for the reaction




at 298K. The magnitude of standard electrode potential of Sn / Sn<sup>2+</sup> if  $E_{\text{Zn}^{2+} / \text{Zn}}^0 = -0.76\text{V}$  is \_\_\_\_\_  $\times 10^{-2}\text{V}$ . (Nearest integer)  
[29-Jan-2023 Shift 2]

**Answer: 17**

**Solution:**

$$\text{Given : } \frac{2.303 RT}{F} = 0.059\text{V}$$



$$\Delta G^\circ = -2.303 RT \log_{10} K_{\text{eq}}$$

$$-nF (E_{\text{cell}}^0) = -2.303 RT \log_{10} K_{\text{eq}}$$

$$0.76 + E_{\text{Sn}^{2+} / \text{Sn}}^0 = \frac{0.059}{2} \log_{10} 10^{20}$$

$$0.76 + E_{\text{Sn}^{2+} / \text{Sn}}^0 = \frac{0.059 \times 20}{2}$$


$$E_{\text{Sn}^{2+} / \text{Sn}}^0 = 0.59 - 0.76 = -0.17$$

$$E_{\text{Sn} / \text{Sn}^{2+}}^0 = 17 \times 10^{-2}\text{V}$$

$$\text{Ans.} = 17$$

## Question20

Consider the cell



When the potential of the cell is 0.712V at 298K, the ratio  $[\text{Fe}^{2+}] / [\text{Fe}^{3+}]$  is \_\_\_\_\_.

(Nearest integer)



Given:  $\text{Fe}^{3+} + \text{e}^- = \text{Fe}^{2+}$ ,  $E^\circ \text{Fe}^{3+}, \text{Fe}^{2+} | \text{Pt} = 0.771$

$$\frac{2.303 RT}{F} = 0.06\text{V}$$

[30-Jan-2023 Shift 1]

**Answer: 10**

**Solution:**



$$E^{\circ} = E_{\text{H}_2 | \text{H}^+}^{\circ} + E_{\text{Fe}^{3+} | \text{Fe}^{2+}}^{\circ} = 0.771\text{V}$$

$$E = E^{\circ} - \frac{0.06}{1} \log \frac{\text{Fe}^{2+}}{\text{Fe}^{3+}}$$


$$0.712 = (0 + 0.771) - \frac{0.06}{1} \log \frac{\text{Fe}^{2+}}{\text{Fe}^{3+}}$$

$$\log \frac{\text{Fe}^{2+}}{\text{Fe}^{3+}} = \frac{0.059}{0.06} \approx 1$$

$$\frac{\text{Fe}^{2+}}{\text{Fe}^{3+}} = 10$$

## Question21

The electrode potential of the following half cell at 298K .



is \_\_\_\_\_  $\times 10^{-2}\text{V}$  (Nearest integer).

Given:  $E_{\text{X}^{2+} | \text{X}}^{\circ} = -2.36\text{V}$


$E_{\text{Y}^{2+} | \text{Y}}^{\circ} = +0.36\text{V}$

$\frac{2.303RT}{F} = 0.06\text{V}$

[30-Jan-2023 Shift 2]

Answer: 275

Solution:



$E_{\text{Cell}}^{\circ} = 0.36 - (-2.36) = 2.72\text{V}$

$E_{\text{Cell}} = 2.72 - \frac{0.06}{2} \log \frac{0.001}{0.01}$

$= 2.72 + 0.03 = 2.75\text{V}$

$= 275 \times 10^{-2}\text{V}$

## Question22

Which one of the following statements is correct for electrolysis of brine solution?

[31-Jan-2023 Shift 1]

Options:

A.  $\text{Cl}_2$  is formed at cathode

B.  $\text{O}_2$  is formed at cathode

C.  $\text{H}_2$  is formed at anode






D.  $\text{OH}^-$  is formed at cathode

**Answer: D**

**Solution:**

**Solution:**

Electrolysis of brine solution



## Question23

The resistivity of a 0.8M solution of an electrolyte is  $5 \times 10^{-3} \Omega \text{ cm}$ . Its molar conductivity is  $\times 10^4 \Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$ . (Nearest integer)

[31-Jan-2023 Shift 2]

**Answer: 25**

**Solution:**

$$\Lambda_m = \frac{\kappa \times 1000}{M}$$

$$\Lambda_m = \frac{1}{\rho} \times \frac{1000}{M}$$

$$\frac{1}{5 \times 10^{-3}} \times \frac{1000}{0.8}$$

$$\text{Ans. } 25 \times 10^4 \Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

## Question24


At what pH, given half cell  $\text{MnO}_4^- (0.1\text{M}) \mid \text{Mn}^{2+} (0.001\text{M})$  will have electrode potential of 1.282 V ? \_\_\_\_\_ (Nearest Integer)

Given  $E_{\text{MnO}_4^- / \text{Mn}^{2+}}^\circ = 1.54\text{V}$ ,  $\frac{2.303RT}{F} = 0.059\text{V}$

[1-Feb-2023 Shift 1]

**Answer: 3**

**Solution:**



$$E = E^\circ - \frac{0.059}{5} \log \frac{[\text{Mn}^{2+}]}{[\text{MnO}_4^-][\text{H}^+]^8}$$

$$1.282 = 1.54 - \frac{0.059}{5} \log \frac{10^{-3}}{10^{-1} \times [\text{H}^+]^8}$$

$$\frac{0.258 \times 5}{0.059} = \log \frac{10^{-2}}{[\text{H}^+]^8}$$

$$\Rightarrow 21.86 = -2 + 8 \text{pH}$$

$$\therefore \text{pH} = 2.98$$

$$\approx 3$$

---

## Question25

$1 \times 10^{-5} \text{M AgNO}_3$  is added to 1L of saturated solution of AgBr. The conductivity of this solution at 298K is \_\_\_\_\_  $\times 10^{-8} \text{Sm}^{-1}$ .

[Given :  $K_{\text{sp}}(\text{AgBr}) = 4.9 \times 10^{-13}$  at 298K

$$\lambda_{\text{Ag}^+}^0 = 6 \times 10^{-3} \text{Sm}^2 \text{mol}^{-1}$$

$$\lambda_{\text{Br}^-}^0 = 8 \times 10^{-3} \text{Sm}^2 \text{mol}^{-1}$$

$$\lambda_{\text{NO}_3^-}^0 = 7 \times 10^{-3} \text{Sm}^2 \text{mol}^{-1} ]$$

[1-Feb-2023 Shift 2]

**Answer: 14**

**Solution:**

$$[\text{Ag}^+] = 10^{-5}$$

$$[\text{NO}_3^-] = 10^{-5}$$

$$[\text{Br}^-] = \frac{K_{\text{sp}}}{[\text{Ag}^+]} = 4.9 \times 10^{-8}$$

$$\Lambda_m = \frac{k}{1000 \times M}$$

For  $\text{Ag}^+$

$$6 \times 10^{-3} = \frac{K_{\text{Ag}^+}}{1000 \times 10^{-5}}$$

$$K_{\text{Ag}^+} = 6 \times 10^{-5}$$

$$\Rightarrow 6000 \times 10^{-8}$$

for  $\text{Br}^-$

$$8 \times 10^{-3} = \frac{K_{\text{Br}^-}}{1000 \times 4.9 \times 10^{-8}}$$

$$K_{\text{Br}^-} = 39.2 \times 10^{-8}$$

for  $\text{NO}_3^-$

$$7 \times 10^{-3} = \frac{K_{\text{NO}_3^-}}{1000 \times 10^{-5}}$$

$$K_{\text{NO}_3^-} = 7 \times 10^{-5}$$

$$= 7000 \times 10^{-8}$$

Conductivity of solution  
 $\Rightarrow (6000 + 7000 + 39.2) \times 10^{-8}$   
 $\Rightarrow 13039.2 \times 10^{-8} \text{Sm}^{-1}$

## Question26

**The standard electrode potential of  $\text{M}^+ / \text{M}$  in aqueous solution does not depend on**

**[6-Apr-2023 shift 1]**

**Options:**

- A. Ionisation of a solid metal atom
- B. Sublimation of a solid metal
- C. Ionisation of a gaseous metal atom
- D. Hydration of a gaseous metal ion

**Answer: A**

**Solution:**

**Solution:**

## Question27

**The product, which is not obtained during the electrolysis of brine solution is**

**[6-Apr-2023 shift 2]**

**Options:**


- A. NaOH
- B.  $\text{Cl}_2$
- C.  $\text{H}_2$
- D. HCl

**Answer: D**


**Solution:**




Brine is aq. Solution of NaCl



Cathode reaction



Anode reaction



So HCl will not form during electrolysis.

---

## Question28

The standard reduction potential at 298K for the following half cells are given below :-

$\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^- \rightarrow \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	$E^\circ = 0.97\text{V}$
$\text{V}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{V}(\text{s})$	$E^\circ = -1.19\text{V}$
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	$E^\circ = -0.04\text{V}$
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	$E^\circ = 0.80\text{V}$
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	$E^\circ = 1.40\text{V}$

The number of metal(s) which will be oxidized by  $\text{NO}_3^-$  in aqueous solution is  
[6-Apr-2023 shift 2]

**Answer: 3**

**Solution:**


**Solution:**

Metal having lower SRP than 0.97V will be oxidised by  $\text{NO}_3^-$ .

---

## Question29



The reaction




occurs in which of the given galvanic cell.

[8-Apr-2023 shift 1]

**Options:**







**Answer: B**

**Solution:**

**Solution:**



## Question30

The specific conductance of 0.0025M acetic acid is  $5 \times 10^{-5} \text{Scm}^{-1}$  at a certain temperature. The dissociation constant of acetic acid is \_\_\_\_\_  $\times 10^{-7}$ . (Nearest integer)

Consider limiting molar conductivity of  $\text{CH}_3\text{COOH}$  as  $400 \text{Scm}^2\text{mol}^{-1}$ .

[10-Apr-2023 shift 2]

**Answer: 66**

**Solution:**

**Solution:**

$$\Lambda_m = \frac{k}{C} \times 1000$$

Given  $k = 5 \times 10^{-5} \text{Scm}^{-1}$

$C = 0.0025 \text{M}$

$$\Lambda_m = \frac{5 \times 10^{-5} \times 10^3}{0.0025} = \frac{5 \times 10^{-2}}{2.5 \times 10^{-3}}$$

$$= 20 \text{Scm}^2\text{mol}^{-1}$$

$$\alpha = \frac{20}{400} = \frac{1}{20}$$

$$K_a = \frac{C\alpha^2}{1 - \alpha} = \frac{0.0025 \times \frac{1}{20} \times \frac{1}{20}}{\frac{19}{20}}$$

$$= \frac{0.0025}{19 \times 20} = 6.6 \times 10^{-6}$$

$$= 66 \times 10^{-7}$$


## Question31

In an electrochemical reaction of lead, at standard temperature, if  $E_{(\text{pb}^{2+} / \text{pb})}^\circ = m$  volt and  $E_{(\text{pb}^+ / \text{Pb}_b)}^\circ = n$  volt, then the value of  $E_{(\text{Pb}^{2+} / \text{Pb}^+)}^\circ$  is given by  $m - xn$ . The value of  $x$  is \_\_\_\_\_ (Nearest integer)

[11-Apr-2023 shift 1]

Answer: 2

Solution:



## Question32

The number of correct statements from the following is \_\_\_\_\_

A.  $E_{\text{cell}}$  is an intensive parameter

B. A negative  $E^\ominus$  means that the redox couple is a stronger reducing agent than the  $\text{H}^+ / \text{H}_2$  couple.

C. The amount of electricity required for oxidation or reduction depends on the stoichiometry of the electrode reaction.

D. The amount of chemical reaction which occurs at any electrode during electrolysis by a current is proportional to the quantity of electricity passed through the electrolyte.

[11-Apr-2023 shift 2]

Answer: 4

Solution:

Solution:

Given statements A, B, C and D are correct.

## Question33

For lead storage battery pick the correct statements

A. During charging of battery,  $\text{PbSO}_4$  on anode is converted into  $\text{PbO}_2$

**B. During charging of battery,  $\text{PbSO}_4$  on cathode is converted into  $\text{PbO}_2$**   
**C. Lead storage battery consists of grid of lead packed with  $\text{PbO}_2$  as anode**  
**D. Lead storage battery has ~38% solution of sulphuric acid as an electrolyte** Choose the correct answer from the options given below:  
**[12-Apr-2023 shift 1]**

**Options:**

A. B, D only

B. B, C only

C. B, C, D only

D. A,B,D only

**Answer: A**

**Solution:**

**Solution:**

Lead storage battery consists of lead anode and a grid of lead packed with lead oxide ( $\text{PbO}_2$ ) as cathode, a 38% solution of  $\text{H}_2\text{SO}_4$  is used as an electrolyte.

On charging the battery the reaction is reversed and  $\text{PbSO}_4$  (s) on anode and cathode is converted into Pb and  $\text{PbO}_2$  respectively.

## Question34

**A metal surface of  $100\text{cm}^2$  area has to be coated with nickel layer of thickness 0.001 mm. A current of 2A was passed through a solution of  $\text{Ni}(\text{NO}_3)_2$  for ' x ' seconds to coat the desired layer. The value of x is (Nearest integer)**

**(  $\rho_{\text{Ni}}$  ( .. density of Nickel) is  $10\text{gmL}^{-1}$ , Molar mass of Nickel is \_\_\_\_\_**

**$60\text{gmol}^{-1}$   $F = 96500\text{Cmol}^{-1}$  )**

**[13-Apr-2023 shift 1]**


**Answer: 161**

**Solution:**

$$\text{Volume of nickel required} = 100 \times 0.001 \times 10^{-3} \times 100 \\ = 0.01\text{cm}^3$$

$$\text{Mass of Nickel required} = 0.01 \times 10 \\ = 0.1 \text{ gm}$$

$$\text{Moles} = \frac{0.1}{60} = \frac{1}{600} \text{ mol}$$



for coating of 1 mol Ni, charge required =  $2 \times 96500\text{C}$

for coating of  $\frac{1}{600}$  mol, charge required =  $2 \times 96500 \times \frac{1}{600}\text{C}$

$$= \frac{965}{3}\text{C}$$

$$I = \frac{q}{t}$$

$$t = \frac{965/3}{2} = 160.83 \text{ sec} \approx 161$$

---

## Question35

At 298K, the standard reduction potential for  $\text{Cu}^{2+} / \text{Cu}$  electrode is 0.034V.

Given :  $K_{\text{sp}} \text{Cu(OH)}_2 = 1 \times 10^{-20}$

Take  $\frac{2.303RT}{F} = 0.059\text{V}$


The reduction potential at pH = 14 for the above couple is  $(-)\text{x} \times 10^{-2}\text{V}$ .

The value of x is \_\_\_\_\_.

[13-Apr-2023 shift 2]

**Answer: 25**


**Solution:**



$$K_{\text{sp}} = [\text{Cu}^{2+}][\text{OH}^{-}]^2$$

$$\text{pH} = 14; \text{pOH} = 0; [\text{OH}^{-}] = 1\text{M}$$

$$\therefore [\text{Cu}^{2+}] = \frac{K_{\text{sp}}}{[\text{OH}^{-}]^2} = 10^{-20}\text{M}$$



$$E = E^{\circ} - \frac{0.059}{2} \log_{10} \frac{1}{[\text{Cu}^{2+}]}$$

$$= 0.34 - \frac{0.059}{2} \log_{10} \frac{1}{10^{-20}}$$

$$= -0.25 = -25 \times 10^{-2}$$

---

## Question36

The number of correct statements from the following is \_\_\_\_\_.

(A) Conductivity always decreases with decrease in concentration for both strong and weak electrolysis.

(B) The number of ions per unit volume that carry current in a solution increases on dilution.

(C) Molar conductivity increases with decrease in concentration.

(D) The variation in molar conductivity is different for strong and weak electrolysis.

**(E) For weak electrolysis, the change in molar conductivity with dilution is due to decrease in degree of dissociation.**

**[15-Apr-2023 shift 1]**

**Answer: 3**

**Solution:**

**Solution:**

(A) Conductivity decreases with dilution for strong electrolyte as well as weak electrolyte.

(B) On dilution, The number of ions per unit volume that carry current in a solution decreases.

(C) Molar conductivity increases with dilution.

(D) Molar conductivity of strong electrolyte follows DHO equation but it is not applicable for weak electrolyte.

(E) On dilution degree of dissociation of weak electrolyte increases.

So answer is (A), (C) & (D)

---

## Question37

**The cell potential for the following cell**

**Pt | H<sub>2</sub>(g) | H<sup>+</sup>(aq) | Cu<sup>2+</sup>(0.01M) | Cu(s)**

**is 0.576V at 298K. The pH of the solution is \_\_\_\_ (Nearest integer)**

**(Given : E<sup>o</sup><sub>Cu<sup>2+</sup> / Cu</sub> = 0.34V and  $\frac{2.303RT}{F} = 0.06V$ )**

**[24-Jun-2022-Shift-1]**

**Answer: 5**

**Solution:**

$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.06}{2} \log \frac{[\text{H}^+]^2}{[\text{Cu}^{+2}]}$$

$$0.576 = 0.34 - 0.03 \log \frac{[\text{H}^+]^2}{[0.01]}$$

$$0.576 - 0.34 = -0.03 \log [\text{H}^+]^2 + 0.03 \log(0.01)$$

$$= 0.06 \text{pH} - 0.06$$

$$\text{pH} \approx 4.93 \approx 5$$

---

## Question38

**The resistance of a conductivity cell containing 0.01M KCl solution at 298K is 1750Ω. If the conductivity of 0.01M KCl solution at 298K is  $0.152 \times 10^{-3} \text{Scm}^{-1}$ , then the cell constant of the conductivity cell is**

$\underline{\hspace{1cm}} \times 10^{-3} \text{cm}^{-1}$   
[24-Jun-2022-Shift-2]

**Answer: 266**

**Solution:**

Molarity of KCl solution = 0.1M

Resistance = 1750 ohm

Conductivity =  $0.152 \times 10^{-3} \text{Scm}^{-1}$



Conductivity =  $\frac{\text{Cell constant}}{\text{Resistance}}$

$\therefore$  Cell constant =  $0.152 \times 10^{-3} \times 1750$   
=  $266 \times 10^{-3} \text{cm}^{-1}$

-----

## Question39


**In a cell, the following reactions take place**



**The standard electrode potential for the spontaneous reaction in the cell is  $x \times 10^{-2} \text{V}$  at 298K. The value of x is \_\_\_\_ (Nearest Integer)**  
[25-Jun-2022-Shift-1]

**Answer: 23**

**Solution:**



## Question40

**The correct order of reduction potentials of the following pairs is**

**A.  $\text{Cl}_2 / \text{Cl}^{-}$**

**B.  $\text{I}_2 / \text{I}^{-}$**

C.  $\text{Ag}^+ / \text{Ag}$

D.  $\text{Na}^+ / \text{Na}$

E.  $\text{Li}^+ / \text{Li}$

Choose the correct answer from the options given below.

[25-Jun-2022-Shift-2]

Options:

A.  $A > C > B > D > E$

B.  $A > B > C > D > E$

C.  $A > C > B > E > D$

D.  $A > B > C > E > D$

Answer: A

Solution:

$$E_{\text{C}_2 / \text{Cl}}^\circ = +1.36\text{V}$$

$$E_{\text{I}_2 / \text{I}^-}^\circ = +0.54\text{V}$$

$$E_{\text{Ag}^+ / \text{Ag}}^\circ = +0.80\text{V}$$

$$E_{\text{Na}^+ / \text{Na}}^\circ = -2.71\text{V}$$

$$E_{\text{L}^+ / \text{Li}}^\circ = -3.05\text{V}$$

---

## Question41

A solution of  $\text{Fe}_2(\text{SO}_4)_3$  is electrolyzed for ' x ' min with a current of 1.5A to deposit 0.3482g of Fe. The value of x is [nearest integer]

Given :  $1\text{F} = 96500\text{Cmol}^{-1}$

Atomic mass of Fe =  $56\text{gmol}^{-1}$

[25-Jun-2022-Shift-2]

Answer: 20

Solution:



$3\text{F} \rightarrow 1 \text{ mole Fe}$  is deposited

For 56g  $\rightarrow 3 \times 96500$  (required charge)

For 1g  $\rightarrow \frac{3 \times 96500}{56}$  (required charge)

For 0.3482g  $\rightarrow \frac{3 \times 96500}{56} \times 0.3482$

$$= 1800.06$$

$Q = it$

$$1800.06 = 1.5t$$

$$t = 20 \text{ min}$$



## Question42

The  $\left( \frac{\partial E}{\partial T} \right)_P$  of different types of half cells are as follows:

A	B	C	D
$1 \times 10^{-4}$	$2 \times 10^{-4}$	$0.1 \times 10^{-4}$	$0.2 \times 10^{-4}$

(Where E is the electromotive force)

Which of the above half cells would be preferred to be used as reference electrode?

[26-Jun-2022-Shift-1]

Options:

A. A

B. B

C. C

D. D

Answer: C


Solution:

**Solution:**

A cell with less variation in EMF with temperature is preferred as a reference electrode because it can be used for a wider range of temperatures without much derivation from standard value so a cell with less  $\left( \frac{\partial E}{\partial T} \right)_P$  is preferred.

-----

## Question43



The Gibbs free energy change for the above reaction at 298K is


$x \times 10^{-1} \text{ kJ mol}^{-1}$ . The value of x is \_\_\_\_\_ [nearest integer]

[Given :  $E_{\text{Cu}^{2+} / \text{Cu}}^{\ominus} = 0.34\text{V}$  ;  $E_{\text{Sn}^{2+} / \text{Sn}}^{\ominus} = -0.14\text{V}$  ;  $F = 96500 \text{ C mol}^{-1}$  ]

[26-Jun-2022-Shift-2]

Answer: 983

Solution:



$$\begin{aligned} E_{\text{cell}}^{\circ} &= E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ} \\ &= -0.14 - (0.34) \\ &= -0.48\text{V} \end{aligned}$$

$$\begin{aligned} E_{\text{cell}} &= E_{\text{cell}}^{\circ} - \frac{0.059}{2} \log \frac{[\text{Cu}^{2+}]}{[\text{Sn}^{2+}]} \\ &= -0.48 - \frac{0.059}{2} \log \frac{0.01}{0.001} \\ &= -0.509 \end{aligned}$$

$$\begin{aligned} \Delta G &= -nF E_{\text{cell}} \\ &= -2 \times 96500 \times (-0.5095) \\ &= 98333.5\text{J} / \text{mol} \\ &= 98.335\text{kJ} / \text{mol} \\ &= 983.35 \times 10^{-1}\text{kJ} / \text{mol} \end{aligned}$$

## Question44

**In 3d series, the metal having the highest  $\text{M}^{2+} / \text{M}$  standard electrode potential is**  
**[27-Jun-2022-Shift-2]**

**Options:**

A. Cr


B. Fe

C. Cu

D. Zn


**Answer: C**

**Solution:**



## Question45

**For the reaction taking place in the cell :**



**The value of  $\Delta_f G^{\circ}$  is \_\_\_\_  $\text{kJ mol}^{-1}$ . (in nearest integer)**

**[27-Jun-2022-Shift-2]**


**Answer: 103**

### Solution:




Anode    Cathode

# At anode, oxidation occur



\# At cathode, reduction occur



Adding equation (1) and (2), we get  $n = 2$ , where  $n =$  cancelled out electron

Now,

$$\begin{aligned}\Delta G^\circ &= -nF E_{\text{cell}}^\circ \\ &= -2 \times 96500 \times 0.5332 \\ &= -102907.6 \\ &= -102.9 \text{ kJ / mol} \\ &= -103 \text{ kJ / mol}\end{aligned}$$

---


## Question46

The solubility product of a sparingly soluble salt  $\text{A}_2\text{X}_3$  is  $1.1 \times 10^{-23}$ . If specific conductance of the solution is  $3 \times 10^{-5} \text{ Sm}^{-1}$ , the limiting molar conductivity of the solution is  $x \times 10^{-3} \text{ Sm}^2 \text{ mol}^{-1}$ . The value of  $x$  is \_\_\_\_\_

[28-Jun-2022-Shift-1]

Answer: 3

### Solution:



$$K_{\text{sp}} = (2s)^2(3s)^3 = 1.1 \times 10^{-23}$$

$$s \approx 10^{-5}$$

For sparingly soluble salts

$$\Lambda_m = \Lambda_m^0$$

$$\Lambda_m = \frac{k}{S \times 10^3}$$

$$= \frac{3 \times 10^{-5}}{10^{-5}} \times 10^{-3}$$

$$= 3 \times 10^{-3} \text{ Sm}^2 \text{ mol}^{-1}$$

---


## Question47

The quantity of electricity in Faraday needed to reduce 1 mol of  $\text{Cr}_2\text{O}_7^{2-}$  to  $\text{Cr}^{3+}$  is \_\_\_\_\_

[28-Jun-2022-Shift-1]

**Answer: 6**

**Solution:**




1 mol 6 mol

$\Rightarrow$  number of faradays = moles of electrons = 6

---

## Question48

For the given reactions






the electrode potentials are ;  $E_{\text{Sn}^{2+} / \text{Sn}}^0 = -0.140\text{V}$  and

$E_{\text{Sn}^{4+} / \text{Sn}}^0 = -0.010\text{V}$ . The magnitude of standard electrode potential for  $\text{Sn}^{4+} / \text{Sn}^{2+}$  i.e.  $E_{\text{Sn}^{4+} / \text{Sn}^{2+}}^0$  is \_\_\_\_\_  $\times 10^{-2}\text{V}$ . (Nearest integer)

[28-Jun-2022-Shift-2]

**Answer: 16**

**Solution:**



$$E_{\text{cell}}^0 = \frac{n_2 E_2^0 + n_1 E_1^0}{n} = \frac{4(0.010) + 2(0.140)}{2}$$

$$E_{\text{cell}}^0 = 0.16\text{V} = 16 \times 10^{-2}\text{V}$$

---

## Question49

A dilute solution of sulphuric acid is electrolysed using a current of 0.10A for 2 hours to produce hydrogen and oxygen gas. The total volume of gases produced at STP is \_\_\_\_\_  $\text{cm}^3$ . (Nearest integer)

[Given : Faraday constant  $F = 96500\text{Cmol}^{-1}$  at STP, molar volume of an ideal gas is  $22.7\text{Lmol}^{-1}$  ]


[29-Jun-2022-Shift-1]

**Answer: 127**


**Solution:**

**Solution:**

At anode



At cathode



$$\text{Now number of gm eq.} = \frac{i \times t}{96500}$$

$$= \frac{0.1 \times 2 \times 60 \times 60}{96500}$$

$$= 0.00746$$

$$V_{\text{O}_2} = \frac{0.00746}{4} \times 22.7 = 0.0423$$


$$V_{\text{H}_2} = \frac{0.00746}{2} \times 22.7 = 0.0846$$

$$V_{\text{Total}} \approx 127 \text{ ml or cc}$$

---

## Question50

The cell potential for the given cell at 298K



is 0.31V. The pH of the acidic solution is found to be 3 , whereas the concentration of  $\text{Cu}^{2+}$  is  $10^{-x}\text{M}$ . The value of x is \_\_\_\_\_

(Given :  $E_{\text{Cu}^{2+} / \text{Cu}}^\ominus = 0.34\text{V}$  and  $\frac{2.303RT}{F} = 0.06\text{V}$  )

[29-Jun-2022-Shift-2]

**Answer: 7**

**Solution:**

$$Q = \frac{[\text{H}^+]^2}{[\text{Cu}^{2+}]\text{pH}_2} = \frac{10^{-6}}{C} \quad \text{pH}_2 = 1$$

$$E = E_{\text{cell}}^\ominus - \frac{0.06}{n} \log Q$$

$$0.31 = 0.34 - \frac{0.06}{2} \log \frac{10^{-6}}{C}$$

$$\log \frac{10^{-6}}{C} = 1$$

$$C = 10^{-7}\text{M}$$

$$x = 7$$



---

## Question51

The cell potential for  $\text{Zn} | \text{Zn}^{2+}(\text{aq}) || \text{Sn}^{x+} | \text{Sn}$  is  $0.801\text{V}$  at  $298\text{K}$ . The reaction quotient for the above reaction is  $10^{-2}$ . The number of electrons involved in the given electrochemical cell reaction is ( Given :  $E_{\text{Zn}^{2+} | \text{Zn}}^{\circ} = -0.763\text{V}$ ,  $E_{\text{Sn}^{x+} | \text{Sn}}^{\circ} = +0.008\text{V}$  and  $\frac{2.303RT}{F} = 0.06\text{V}$  )  
[25-Jul-2022-Shift-1]

**Answer: 4**

**Solution:**



$$E_{\text{Cell}}^{\circ} = E_{\text{Zn} | \text{Zn}^{2+}}^{\circ} + E_{\text{Sn}^{x+} | \text{Sn}}^{\circ}$$

$$\Rightarrow 0.763 + 0.008 = 0.771\text{V}$$

From the Nernst equation,

$$E_{\text{Cell}} = E_{\text{Cell}}^{\circ} - \frac{2.303RT}{nF} \log Q$$

$$0.801 = 0.771 - \frac{0.06}{n} \log 10^{-2}$$

$$0.03 = \frac{0.06}{n} \times 2$$

$$n = 4$$

---

## Question52

The molar conductivity of a conductivity cell filled with 10 moles of  $20\text{mLN aCl}$  solution is  $\Lambda_{m1}$  and that of 20 moles another identical cell heaving  $80\text{mLN aCl}$  solution is  $\Lambda_{m2}$ . The conductivities exhibited by these two cells are same. The relationship between  $\Lambda_{m2}$  and  $\Lambda_{m1}$  is  
[25-Jul-2022-Shift-2]

**Options:**

A.  $\Lambda_{m2} = 2\Lambda_{m1}$

B.  $\Lambda_{m2} = \Lambda_{m1} / 2$

C.  $\Lambda_{m.2} = \Lambda_{m1}$

D.  $\Lambda_{m2} = 4\Lambda_{m1}$

**Answer: A**

**Solution:**

$$\Lambda_{m_1} = \frac{k_1 \times 1000}{M_1} = \frac{k \times 1000}{\frac{10}{0.02}}$$

$$\Lambda_{m_2} = \frac{k_2 \times 1000}{\frac{20}{0.08}}$$

It is given that  $k_1 = k_2$

$$k_1 = \frac{\Lambda_{m_1}}{2} \quad k_2 = \frac{\Lambda_{m_2}}{4}$$

Applying the given condition on conductivity.

$$\frac{\Lambda_{m_1}}{2} = \frac{\Lambda_{m_2}}{4}$$

$$\Lambda_{m_2} = 2\Lambda_{m_1}$$

## Question53

**The spin-only magnetic moment value of  $M^{3+}$  ion (in gaseous state) from the pairs  $Cr^{3+} / Cr^{2+}$ ,  $Mn^{3+} / Mn^{2+}$ ,  $Fe^{3+} / Fe^{2+}$  and  $Co^{3+} / Co^{2+}$  that has negative standard electrode potential, is \_\_\_\_ B.M. [Nearest integer]  
[25-Jul-2022-Shift-2]**

**Answer: 4**

**Solution:**

**Solution:**

Among the pairs given,  $Cr^{3+} / Cr^{2+}$  has negative reduction potential which is  $-0.41V$ .

$Cr(III) \Rightarrow d^3$

Number of unpaired electrons = 3

$\mu = \sqrt{3(3+2)} = \sqrt{15} \approx 4$  B.M.

## Question54

**The amount of charge in F (Faraday) required to obtain one mole of iron from  $Fe_3O_4$  is \_\_\_\_\_. (Nearest Integer)  
[26-Jul-2022-Shift-1]**


**Answer: 3**

**Solution:**

For  $\text{Fe}_3\text{O}_4$ ,

$$x = \frac{+8}{3}$$

where x is oxidation state of Fe.



$$\text{Charge required} = \frac{8}{3} \times F = \frac{8F}{3} \approx 3F$$

---

## Question55

Given below are two statements :

**Statement I : For KI, molar conductivity increases steeply with dilution**

**Statement II : For carbonic acid, molar conductivity increases slowly with dilution**

**In the light of the above statements, choose the correct answer from the options given below:**

**[27-Jul-2022-Shift-2]**

**Options:**

- A. Both Statement I and Statement II are true
- B. Both Statement I and Statement II are false
- C. Statement I is true but Statement II is false
- D. Statement I is false but Statement II is true


**Answer: B**

**Solution:**

**Solution:**

Statement I: KI is strong electrolyte thus almost constant on dilution.

Statement II: In weak electrolyte it increases, sharply.



---

## Question56

**Match List - I with List - II.**



List - I	List - II
(A) $\text{Cd}(s) + 2 \text{Ni}(\text{OH})_3(s) \rightarrow \text{CdO}(s) + 2 \text{Ni}(\text{OH})_2(s) + \text{H}_2\text{O}(l)$	(I) Primary battery
(B) $\text{Zn}(Hg) + \text{HgO}(s) \rightarrow \text{ZnO}(s) + \text{Hg}(l)$	(II) Discharging of secondary battery
(C) $2\text{PbSO}_4(s) + 2\text{H}_2\text{O}(l) \rightarrow \text{Pb}(s) + \text{PbO}_2(s) + 2\text{H}_2\text{SO}_4(aq)$	(III) Fuel cell
(D) $2\text{H}_2(g) + \text{O}_2(g) \rightarrow 2\text{H}_2\text{O}(l)$	(IV) Charging of secondary battery

**Choose the correct answer from the options given below:**  
**[28-Jul-2022-Shift-1]**

**Options:**

- A. (A) – (I), (B) – (II), (C) – (II), (D) – (IV)  
 B. (A) – (IV), (B) – (I), (C) – (I), (D) – (II)  
 C. (A) – (II), (B) – (I), (C) – (IV), (D) – ( III)  
 D. (A) – ( II) , (B) – (I), (C) – (II), (D) – (IV)

**Answer: C**

**Solution:**

**Solution:**

- (a)  $\text{Cd}(s) + 2 \text{Ni}(\text{OH})_3(s) \rightarrow \text{CdO}(s) + 2 \text{Ni}(\text{OH})_2(s) + \text{H}_2\text{O}(l)$  Discharge of secondary Battery  
 (b)  $\text{Zn}(Hg) + \text{HgO}(s) \rightarrow \text{ZnO}(s) + \text{Hg}(l)$  (Primary Battery Mercury cell)  
 (c)  $2\text{PbSO}_4(s) + 2\text{H}_2\text{O}(l) \rightarrow \text{Pb}(s) + \text{PbO}_2(s) + 2\text{H}_2\text{SO}_4(aq)$  (Charging of secondary Battery)  
 (d)  $2\text{H}_2(g) + \text{O}_2(g) \rightarrow 2\text{H}_2\text{O}(l)$  (Fuel cell)

## Question57

**Resistance of a conductivity cell (cell constant  $129\text{m}^{-1}$ ) filled with 74.5 ppm solution of KCl is  $100\Omega$  (labelled as solution 1). When the same cell is filled with KCl solution of 149 ppm, the resistance is  $50\Omega$  (labelled as solution 2). The ratio of molar conductivity of solution 1 and solution 2 is i.e.  $\frac{\Lambda_1}{\Lambda_2} = x \times 10^{-3}$ . The value of x is \_\_\_\_\_.(Nearest integer)**

**Given, molar mass of KCl is  $74.5\text{gmol}^{-1}$ .**

**[29-Jul-2022-Shift-1]**

**Answer: 1000**

**Solution:**

$$\frac{l}{A} = 129\text{m}^{-1}$$

$$\text{KCl solution 1} \Rightarrow 74.5 \text{ ppm, } R_1 = 100\Omega$$

$$\text{KCl solution 2} \Rightarrow 149 \text{ ppm, } R_2 = 50\Omega$$

$$\text{Here, } \frac{\text{ppm}_1}{\text{ppm}_2} = \frac{M_1}{M_2} = \left( \frac{w_1 / M_0}{V} \times \frac{V}{w_2 / M_0} \right)$$

$$\frac{\Lambda_1}{\Lambda_2} = \frac{k_1 \times \frac{1000}{M_1}}{k_2 \times \frac{1000}{M_2}}$$

$$= \frac{k_1}{k_2} \times \frac{M_1}{M_2}$$

$$= \frac{50}{100} \times 2$$

$$= \frac{\Lambda_1}{\Lambda_2} = 1000 \times 10^{-3}$$

$$= 1000$$

## Question58



For a cell,  $\text{Cu(s)} \mid \text{Cu}^{2+}(0.001\text{M}) \mid \text{Ag}^{+}(0.01\text{M}) \mid \text{Ag(s)}$  the cell potential is found to be  $0.43\text{V}$  at  $298\text{K}$ . The magnitude of standard electrode potential for  $\text{Cu}^{2+} / \text{Cu}$  is \_\_\_\_\_  $\times 10^{-2}\text{V}$

[Given :  $E_{\text{Ag}^{+} / \text{Ag}}^{\ominus} = 0.80\text{V}$  and  $\frac{2.303\text{RRT}}{F} = 0.06\text{V}$ ]

[29-Jul-2022-Shift-2]

Answer: 34

Solution:



$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.06}{2} \log \frac{[\text{Cu}^{2+}]}{[\text{Ag}^{+}]^2}$$

$$0.43 = E_{\text{cell}}^0 - \frac{0.06}{2} \log \left( \frac{10^{-3}}{(10^{-2})^2} \right)$$

$$0.43 = E_{\text{cell}}^0 - 0.03 \log 10$$

$$E_{\text{cell}}^0 = 0.46\text{V}$$

$$E_{\text{cell}}^0 = E_{\text{Ag}^{+} / \text{Ag}}^0 - E_{\text{Cu}^{2+} / \text{Cu}}^0$$

$$E_{\text{Cu}^{2+} / \text{Cu}}^0 = (0.80 - 0.46) = 0.34\text{V} = 34 \times 10^{-2}$$

## Question59

Emf of the following cell at  $298\text{K}$  in  $\text{V}$  is  $x \times 10^{-2}$ ,  $\text{Zn} \mid \text{Zn}^{2+}(0.1\text{M}) \mid \text{Ag}^{+}(0.01\text{M}) \mid \text{Ag}$  The value of  $x$  is .....



**(Rounded off to the nearest integer).**

**[Given,  $E_{\text{Zn}^{2+}/\text{Zn}}^{\circ} = -0.76\text{V}$ ,  $E_{\text{Ag}^{+}/\text{Ag}}^{\circ} = +0.80\text{V}$ ,  $\frac{2.303RT}{F} = 0.059$ ]**

**[26 Feb 2021 Shift 2]**

**Answer: 147**

**Solution:**



$$\Rightarrow Q = \frac{[\text{Zn}^{2+}][\text{Ag}]^2}{[\text{Zn}][\text{Ag}^{+}]^2} = \frac{0.1\text{M}}{1 \times (0.01)^2} = 10^3$$

$$n = 2 \text{ and } E_{\text{cell}}^{\circ} = E_{\text{Ag}^{+}/\text{Ag}}^{\circ} - E_{\text{Zn}^{2+}/\text{Zn}}^{\circ}$$
$$= 0.8 - (-0.76) = 1.56\text{V}$$

$$\Rightarrow E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{n} \log Q = 1.56 - \frac{0.059}{2} \log 10^3$$

$$= 1.4715\text{V}$$

$$= 147.15 \times 10^{-2}\text{V} = 147 \times 10^{-2}\text{V}$$

$$= x \times 10^{-2}\text{V}$$

$$x = 147$$

## Question60

**Copper reduces  $\text{NO}_3^-$  into  $\text{NO}$  and  $\text{NO}_2$  depending upon the concentration of  $\text{HNO}_3$  in solution. (Assuming fixed  $[\text{Cu}^{2+}]$  and  $p_{\text{NO}} = p_{\text{NO}_2}$ ), the  $\text{HNO}_3$  concentration at which the thermodynamic tendency for reduction of  $\text{NO}_3^-$  into  $\text{NO}$  and  $\text{NO}_2$  by copper is same is  $10^x\text{M}$ . The value of  $2x$  is ..... .(Rounded off to the nearest integer)**

**[ Given,  $E_{\text{Cu}^{2+}/\text{Cu}}^{\circ} = 0.34\text{V}$ ,  $E_{\text{NO}_3^-/\text{NO}}^{\circ} = 0.96\text{V}$**


**$E_{\text{NO}_3^-/\text{NO}_2}^{\circ} = 0.79\text{V}$  and at  $298\text{K}$ ,  $\frac{RT}{F}$**

**$(2.303) = 0.059$ ]**

**[25 Feb 2021 Shift 2]**

**Answer: 4**

**Solution:**




$$Q_1 = \frac{[\text{Cu}^{2+}]^3 \times (p_{\text{NO}})^2}{[\text{NO}_3^-]^2 \times [\text{H}^+]^8}$$

$$\because E_1^\circ = 0.96 - (-0.34) = 1.3V$$

$$E_1 = 1.3 - \frac{0.059}{6} \log Q_1$$

Cell-II ( $\text{HNO}_3 \rightarrow \text{NO}_2$ )



$$Q_2 = \frac{[\text{Cu}^{2+}] \times (p_{\text{NO}_2})^2}{[\text{NO}_3^-]^2 \times [\text{H}^+]^4}$$

$$E_2^\circ = 0.79 - (-0.34)V = 1.13V$$

$$E_2 = 1.13 - \frac{0.059}{2} \log Q_2$$

$$\because E_2$$

$$\text{Now, } E_1 = E_2$$

$$1.3 - \frac{0.059}{6} \log Q_1 = 1.13 - \frac{0.059}{2} \log Q_2$$

$$0.17 = \frac{0.059}{6} [\log Q_1 - 3 \log Q_2] = \frac{0.059}{6} \log \frac{Q_1}{Q_2}$$

$$= \frac{0.059}{6} \log \frac{[\text{Cu}^{2+}]^3 \times (p_{\text{NO}})^2}{[\text{NO}_3^-]^2 \times [\text{H}^+]^8} \times \frac{[\text{NO}_3^-]^6 \times [\text{H}^+]^{12}}{[\text{Cu}^{2+}]^3 \times (p_{\text{NO}_2})^6} = \frac{0.059}{6} \log \frac{[\text{H}^+]^4 \times [\text{NO}_3^-]^4}{(p_{\text{NO}_2})^4} \quad [\because p_{\text{NO}} = p_{\text{NO}_2}]$$

$$= \frac{0.059}{6} \log \frac{[\text{HNO}_3]^4}{(p_{\text{NO}_2})^4}$$

$$\text{Now, } p_{\text{NO}_2} \equiv [\text{HNO}_3]$$

$$\text{So, } 0.17 = \frac{0.059}{6} \log [\text{HNO}_3]^8$$

$$= \frac{0.059}{6} \times 8 \log [\text{HNO}_3]$$

$$\log [\text{HNO}_3] = 2.16$$

$$[\text{HNO}_3] = 10^{2.16} \text{M} = 10^x \text{M}$$

$$\therefore x = 2.16$$

$$\Rightarrow 2x = 2 \times 2.16 = 4.32 \sim \text{eq4}$$


## Question61

The magnitude of the change in oxidising power of the  $\text{MnO}_4^- / \text{Mn}^{2+}$  couple is  $x \times 10^{-4} \text{ V}$ , if the  $\text{H}^+$  concentration is decreased from 1M to  $10^{-4} \text{M}$  at  $25^\circ\text{C}$ . (Assume concentration of  $\text{MnO}_4^-$  and  $\text{Mn}^{2+}$  to be same on change in  $\text{H}^+$  concentration). The value of x is ..... (Rounded off to the nearest integer). [ Given,  $\frac{2.303RT}{F} = 0.059$  ]

[24 Feb 2021 Shift 2]

**Answer: 3776**

**Solution:**



$$\text{Applying Nernst equation, } E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{0.0591}{n} \log \frac{[\text{P}]}{[\text{R}]}$$

$$\text{or } E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{0.0591}{5} \log \frac{[\text{Mn}^{2+}]}{[\text{MnO}_4^-]} \left[ \frac{1}{[\text{H}^+]} \right]^8$$

$$(\text{I) Given, } [\text{H}^+] = 1\text{M}$$

$$E_1 = E^\circ - \frac{0.0591}{5} \log \frac{[Mn^{2+}]}{[MnO_4^-]}$$

(II) Now,  $[H^+] = 10^{-4}M$

$$E_2 = E^\circ - \frac{0.0591}{5} \log \frac{[Mn^{2+}]}{[MnO_4^-]} \times \frac{1}{(10^{-4})^8}$$

$$\therefore |E_1 - E_2|$$

$$|E_1 - E_2| = \frac{0.0591}{5} \times 32 = 0.3776V = 3776 \times 10^{-4}$$

$$x = 3776$$

## Question62

**Compound A used as a strong oxidising agent is amphoteric in nature. It is the part of lead storage batteries. Compound A is**  
**[26 Feb 2021 Shift 1]**

**Options:**

A.  $PbO_2$

B.  $PbO$

C.  $PbSO_4$

D.  $Pb_3O_4$

**Answer: A**

**Solution:**

**Solution:**

In the set of four lead compounds Pb (II) compounds are  $PbO$  and  $PbSO_4$ .  $PbO_2$  is a Pb (IV) compound whereas  $Pb_3O_4$  is a mixed oxide of Pb (II) and Pb (IV) i.e.  $2PbO \cdot PbO_2$ .

Pb is a member of group 14 and it shows +2 and +4 oxidation states. But due to inert pair effect,  $Pb^{2+}$  is more stable than  $Pb^{4+}$ . So, Pb (IV) compounds are strong oxidising agents as  $Pb^{4+}$  gets easily reduced to more stable  $Pb^{2+}$ .

$Pb^{+4} + 2e^- \rightarrow Pb^{2+}$ ,  $\Delta C^\circ < 0$  (spontaneous)

So,  $PbO_2$  or  $Pb_3O_4$  can be the compound A. But out of these two compounds only  $PbO_2$  is used in lead storage batteries where a grid of lead packed with  $PbO_2$  acts as cathode and also it is amphoteric in nature. It reacts with both acids and alkali.

(i)  $PbO_2 + 2HCl \rightarrow PbCl_2 + Cl_2 + H_2O$

Here,  $PbO_2$  acts as a basic oxide as well as an oxidising agent.

(ii)  $PbO_2 + 2NaOH \rightarrow Na_2PbO_2 + H_2O$

Here,  $PbO_2$  acts as an acidic oxide.

So, the compound A is  $PbO_2$  (option-a).

## Question63

**Consider the following reaction,**




**The quantity of electricity required in Faraday to reduce five moles of  $MnO_4^-$  is .....**

**[26 Feb 2021 Shift 1]**

**Answer: 25**

**Solution:**

In the reduction half-cell,



1mol 5F

5mol 25F

As one mole of  $\text{MnO}_4^-$  required 5F of charge,

5 moles of  $\text{MnO}_4^-$  will require charge,

$$Q = 5 \times 5 \text{ Faraday} = 25 \text{ Faraday}$$

---

## Question64

**A  $5.0 \text{ mmol dm}^{-3}$  aqueous solution of KCl has a conductance of  $0.55 \text{ mS}$  when measured in a cell constant  $1.3 \text{ cm}^{-1}$ .**

**The molar conductivity of this solution is  $\text{mSm}^2 \text{ mol}^{-1}$ .**

**(Round off to the nearest integer)**

**[16 Mar 2021 Shift 2]**

**Answer: 14**

**Solution:**

$$\Lambda_m = \frac{K}{1000 \times C}$$

where,  $\Lambda_m$  = molar conductivity  $\kappa$  = conductivity,  $C$  = molarity

$$\text{Conductivity } (\kappa) = G \frac{l}{A} = 0.55 \times (1.3 \times 100)$$

$$= 55 \times 1.3$$

$$\Lambda_m = \frac{55 \times 1.3}{1000 \times 5 \times 10^{-3}}$$

$$\Lambda_m = 14.3$$

Closest integer,  $\Lambda_m = 14$

---

## Question65

**The molar conductivities at infinite dilution of barium chloride, sulphuric acid and hydrochloric acid are  $280, 860$  and  $426 \text{ Scm}^2 \text{ mol}^{-1}$  respectively. The molar conductivity at infinite dilution of barium sulphate is .....  $\text{Scm}^2 \text{ mol}^{-1}$  (Round off to the nearest Integer).**

**[18 Mar 2021 Shift 2]**

**Answer: 288**

**Solution:**

Molar conductivity of  $\text{BaCl}_2 = 280 \text{ Scm}^2 \text{ mol}^{-1}$

Molar conductivity of  $\text{H}_2\text{SO}_4 = 860 \text{ Scm}^2 \text{ mol}^{-1}$

Molar conductivity of  $\text{HCl} = 426 \text{ Scm}^2 \text{ mol}^{-1}$

Molar conductivity of  $\text{BaSO}_4 = ?$

From Kohlrausch's law,

$$\Lambda_m^\infty(\text{BaSO}_4) = \lambda_m^\omega(\text{Ba}^{2+}) + \lambda_m^\omega(\text{SO}_4^{2-})$$

$$\Lambda_m^\infty(\text{BaSO}_4) = \Lambda_m^\infty(\text{BaCl}_2) + \Lambda_m^\infty(\text{H}_2\text{SO}_4) - 2\Lambda_m^\infty(\text{HCl})$$

On putting given values we get,


$$= 280 + 860 - 2(426)$$

$$= 288 \text{ Scm}^2 \text{ mol}^{-1}$$

---

## Question 66

**For the reaction,**



**The magnitude of the standard molar Gibbs free energy change,**

$$\Delta_r G_m^\circ = -\dots\dots\dots \text{ kJ}$$

**(Round off to the nearest integer).**



$$\left[ \begin{array}{ll} E_{\text{Fe}^{2+}/\text{Fe(s)}}^\circ = -0.440\text{V}; & E_{\text{Fe}^{3+}/\text{Fe(s)}}^\circ = -0.036\text{V} \\ E_{\text{I}_2/2\text{I}^{-}} = 0.539\text{V}; & F = 96500\text{C} \end{array} \right]$$

**[18 Mar 2021 Shift 1]**

**Answer: 45**

**Solution:**

Given reaction is



$$\begin{array}{c} E_3^\circ \\ | \text{-----} | \\ nE_1^\circ + nE_2^\circ = nE_3^\circ \quad [n = \text{number of electron transferred}] \end{array}$$

$$E_1^\circ + 2E_2^\circ = 3E_3^\circ$$

$$E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^\circ + 2E_{\text{Fe}^{2+}/\text{Fe(s)}}^\circ = 3E_{\text{Fe}^{3+}/\text{Fe(s)}}^\circ$$

$$E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^\circ + (-0.440) \times 2 = (-0.036) \times 3$$

$$E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^\circ = 0.772\text{V}$$

$$E_{\text{cell}}^{\circ} = E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ}$$

$$= 0.772 - 0.539 = 0.233\text{V}$$

Using standard Gibb's free energy,

$$\Delta G^{\circ} = nF E_{\text{cell}}^{\circ} = +2 \times 96500 \times 0.233$$

$$\Delta G^{\circ} = 44969\text{J} = 44.9\text{kJ} = 45\text{kJ}$$


---

## Question67

Potassium chlorate is prepared by electrolysis of KCl in basic solution as shown by following equation.



A current of x A has to be passed for 10h to produce 10.0g of potassium chlorate. the value of x is \_\_\_\_\_. (Nearest integer)

(Molar mass of KClO<sub>3</sub> = 122.6g mol<sup>-1</sup>, F = 96500C)

[20 Jul 2021 Shift 2]

**Answer: 1**

**Solution:**

$$W = \frac{E}{F} \times I \times t$$

$$10 = \frac{122.6}{96500 \times 6} \times x \times 10 \times 3600$$

$$X = 1.311$$


---

## Question68

For the cell



the cell potential  $E_1 = 0.3095\text{V}$  For the cell



the cell potential = \_\_\_\_  $10^{-2}\text{V}$ . (Round off the Nearest Integer).

$$\left[ \text{Use : } \frac{2.303RT}{F} = 0.059 \right]$$


[27 Jul 2021 Shift 2]

**Answer: 28**

**Solution:**



Cell reaction is :



$$\text{Now, } E_{\text{cell}} = E^{\circ}_{\text{Cell}} - \frac{0.059}{2} \log \frac{[\text{Cu}^{2+}]}{[\text{Ag}^+]^2} \dots\dots(1)$$

$$\therefore E_1 = 0.3095 = E^{\circ}_{\text{Cell}} - \frac{0.059}{2} \cdot \log \frac{0.01}{(0.001)^2}$$

$$\text{From (1) and (2), } E_2 = 0.28\text{V} = 28 \times 10^{-2}\text{V}$$

---

## Question69

Consider the cell at 25°C






The fraction of total iron present as  $\text{Fe}^{3+}$  ion at the cell potential of 1.500V is  $x \times 10^{-2}$ . The value of x is \_\_\_\_\_. (Nearest integer)

(Given :  $E^{\circ}_{\text{Fe}^{3+}/\text{Fe}^{2+}} = 0.77\text{V}$ ,  $E^{\circ}_{\text{Zn}^{2+}/\text{Zn}} = -0.76\text{V}$  )

[25 Jul 2021 Shift 1]

**Answer: 24**

**Solution:**



$$E^{\circ}_{\text{cell}} = 0.77 - (0.76)$$

$$= 1.53\text{V}$$

$$1.50 = 1.53 - \frac{0.06}{2} \log \left( \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]} \right)^2$$

$$\log \left( \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]} \right) = \frac{0.03}{0.06} = \frac{1}{2}$$

$$\frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]} = 10^{1/2} = \sqrt{10}$$

$$\frac{[\text{Fe}^{3+}]}{[\text{Fe}^{2+}]} = \frac{1}{\sqrt{10}}$$

$$\frac{[\text{Fe}^{3+}]}{[\text{Fe}^{2+}] + [\text{Fe}^{3+}]} = \frac{1}{1 + \sqrt{10}} = \frac{1}{4.16}$$


$$= 0.2402$$

$$= 24 \times 10^{-2}$$

---

## Question70

Assume a cell with the following reaction



$$E^{\circ}_{\text{cell}} = 2.97\text{V}$$

$E_{\text{cell}}$  for the above reaction is \_\_\_\_\_ V.

(Nearest integer)

[ Given :  $\log 2.5 = 0.3979$ ,  $T = 298\text{K}$  ]

[22 Jul 2021 Shift 2]

**Answer: 3**

**Solution:**

$$\begin{aligned} E &= E^\circ - \frac{0.059}{2} \log \frac{[\text{Cu}^{+2}]}{[\text{Ag}^+]^2} \\ &= 2.97 - \frac{0.059}{2} \log \frac{0.25}{(10^{-3})^2} = 2.81\text{V} \end{aligned}$$

---

## Question71

The conductivity of a weak acid HA of concentration  $0.001\text{mol L}^{-1}$  is  $2.0 \times 10^{-5}\text{Scm}^{-1}$ . If  $\Lambda_m^\circ(\text{HA}) = 190\text{Scm}^2\text{mol}^{-1}$ , the ionization constant ( $K_a$ ) of HA is equal to \_\_\_\_\_  $\times 10^{-6}$ .

(Round off to the Nearest Integer)


[27 Jul 2021 Shift 1]

**Answer: 12**

**Solution:**

$$\begin{aligned} \Lambda_m &= 1000 \times \frac{\kappa}{M} \\ &= 1000 \times \frac{2 \times 10^{-5}}{0.001} = 20\text{Scm}^2\text{mol}^{-1} \end{aligned}$$

$$\Rightarrow \alpha = \frac{\Lambda_m}{\Lambda_m^\circ} = \frac{20}{190} = \left( \frac{2}{19} \right)$$



$$0.001(1 - \alpha) \quad 0.001\alpha \quad 0.001\alpha$$

$$\Rightarrow K_a = 0.001 \left( \frac{\alpha^2}{1 - \alpha} \right) = \frac{0.001 \times \left( \frac{2}{19} \right)^2}{1 - \left( \frac{2}{19} \right)}$$

$$= 12.3 \times 10^{-6}$$

---

## Question72

**Given below are two statements.**

**Statement I** The limiting molar conductivity of KCl (strong electrolyte) is higher compared to that of  $\text{CH}_3\text{COOH}$  (weak electrolyte).

**Statement II** Molar conductivity decreases with decrease in concentration of electrolyte. In the light of the above statements, choose the most appropriate answer from the options given below  
[26 Aug 2021 Shift 1]

**Options:**

- A. Statement I is true but statement II is false.
- B. Statement I is false but statement II is true.
- C. Both statement I and statement II are true.
- D. Both statement I and statement II are false.

**Answer: D**

**Solution:**

**Solution:**

Limiting molar conductivity of KCl

$$\Lambda_m^\circ \text{KCl} = \Lambda_m^\circ (\text{K}^+) + \Lambda_m^\circ (\text{Cl}^-)$$

$$= 73.55 \text{ S cm}^2 / \text{mol} + 76.3 \text{ S cm}^2 / \text{mol}$$

$$= 149.3 \text{ S cm}^2 / \text{mol}$$

Limiting molar conductivity of  $\text{CH}_3\text{COOH}$

$$\Lambda_m^\circ (\text{CH}_3\text{COOH}) = \Lambda_m^\circ (\text{CH}_3\text{COO}^-) + \Lambda_m^\circ (\text{H}^+)$$

$$= 349.8 \text{ S cm}^2 / \text{mol} + 40.9 \text{ S cm}^2 / \text{mol} = 390.7 \text{ S cm}^2 / \text{mol}$$

$\therefore$  Limiting molar conductivity of  $\text{CH}_3\text{COOH}$  is more than KCl.

Hence, statement I is false.

Also, as the concentration decreases, dilution of the electrolyte increases which will increase the degree of dissociation of weak electrolyte. Thus, the number of ions in the solution increases and hence, the molar conductance of electrolyte increases. Therefore, we can see that with decrease in concentration of electrolyte, molar conductance increases.

Hence, statement II is also false.

---

## Question73

**These are physical properties of an element.**

**A. Sublimation enthalpy**

**B. Ionisation enthalpy**

**C. Hydration enthalpy**

**D. Electron gain enthalpy**

**The total number of above properties that affect the reduction potential is..... (Integer answer)**

**[26 Aug 2021 Shift 1]**

**Answer: 3**

**Solution:**


**Solution:**

The physical properties of an element is affected by sublimation enthalpy, ionisation enthalpy and hydration enthalpy as these 3 enthalpies will affect the reduction potential.

---

## Question74

For the galvanic cell,



$$E_{\text{cell}} = \dots \times 10^{-2} \text{V. (Nearest integer)}$$

$$[\text{Use } E^{\circ}_{\text{Cu} / \text{Cu}^{2+}} = -0.34\text{V}, E^{\circ}_{\text{Zn} / \text{Zn}^{2+}} = +0.76\text{V},$$

$$\frac{2.303RT}{F} = 0.059\text{V}]$$


[26 Aug 2021 Shift 2]

**Answer: 109**

**Solution:**

**Solution:**

Cell reaction,



$$E^{\circ}_{\text{cell}} = E^{\circ}_{\text{Cu}^{2+} / \text{Cu}} - E^{\circ}_{\text{Zn}^{2+} / \text{Zn}}$$

$$E_{\text{cell}} = E^{\circ}_{\text{cell}} - \frac{0.059}{2} \log \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]}$$

$$= 1.1 - \frac{0.059}{2} \log \frac{[0.04]}{[0.02]}$$

$$= 1.1 - 0.03 \log 2 = 1.1 - 0.03 \times 0.30$$

$$= 1.09\text{V} = 109 \times 10^{-2}\text{V}$$

∴ Answer is 109.

---

## Question75

The resistance of a conductivity cell with cell constant  $1.14\text{cm}^{-1}$ , containing  $0.001\text{ M KCl}$  at  $298\text{K}$  is  $1500\Omega$ . The molar conductivity of  $0.001\text{ M KCl}$  solution at  $298\text{K}$  in  $\text{Scm}^2\text{mol}^{-1}$  is ..... .(Integer answer)  
[27 Aug 2021 Shift 2]

**Answer: 760**

**Solution:**

**Solution:**

$$\text{Given, } C^* = 1.14\text{cm}^{-1} = 1.14 \times 10^3\text{m}^{-1}$$

$$R = 1500 \, \Omega$$

$$C = 0.001 \, \text{M}$$

$$G^* = R \times \kappa$$

[where,  $R$  = resistance

$G^*$  = cell constant

$\kappa$  = conductivity

$C$  = concentration]

$$\therefore \kappa = \frac{G^*}{R} = \frac{1.14 \times 10^3 \text{m}^{-1}}{1500}$$

$$\Lambda_m = \frac{\kappa}{C}$$


[where,  $\Lambda_m$  = molar conductivity]

$$= \frac{1.14 \times 10^3}{1500 \times 0.001} = 760 \text{Scm}^2 \text{mol}^{-1}.$$

## Question76

### Section B : Numerical Type Questions

Consider the following cell reaction,



The value of  $E_{\text{cell}}^\circ$  is 4.315V at 25°C. If  $\Delta H^\circ = -825.2 \text{kJ mol}^{-1}$ , the standard entropy change  $\Delta S^\circ$  in  $\text{JK}^{-1}$  is.... (Nearest integer)

[Given, Faraday constant =  $96487 \text{Cmol}^{-1}$ ]

[31 Aug 2021 Shift 1]

**Answer: 25**

**Solution:**

**Solution:**

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ \dots(i)$$

where,  $\Delta G^\circ$  = standard Gibb's free energy change

$\Delta H^\circ$  = standard change in enthalpy

$\Delta S^\circ$  = standard entropy change

$$\text{Also, } \Delta G^\circ = -nFE^\circ$$

where,  $n$  = number of electrons

$F$  = Faraday constant

$E^\circ$  = standard cell potential

$\therefore$  Equation (i) is,

$$-nFE^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$-nFE^\circ - \Delta H^\circ = T\Delta S^\circ$$

$$\Rightarrow \Delta S^\circ = \frac{nFE^\circ + \Delta H^\circ}{T} \quad (2 \times 96487 \text{Cmol}^{-1} \times 4.315\text{V})$$

$$= \frac{+(-825.2 \text{kJ} / \text{mol})}{298\text{K}}$$

$$= \frac{832.682 \times 10^3 \text{J mol}^{-1} - 825.2 \times 10^3 \text{J} / \text{mol}}{298\text{K}}$$

$$= 25.11 \text{J} / \text{Kmol}$$

## Question77

**Match List-I with List-II.**

List-I (Parameter)	List-II (Unit)
A. Cell constant	1. $\text{Scm}^2\text{mol}^{-1}$
B. Molar conductivity	2. Dimensionless
C. Conductivity	3. $\text{m}^{-1}$
D. Degree of dissociation of electrolyte	4. $\Omega^{-1}\text{m}^{-1}$

**Choose the most appropriate answer from the options given below  
[31 Aug 2021 Shift 2]**

**Options:**

A. A-3 B-1 C-4 D-2

B. A-3 B-1 C-2 D-4

C. A-1 B-4 C-3 D-2

D. A-2 B-1 C-3 D-4

**Answer: A**

**Solution:**

**Solution:**

(A) Cell constant =  $\frac{l}{A}\text{m}^{-1}$

(B) Molar conductivity ( $\lambda_m$ ) =  $\frac{\kappa \times 1000}{\text{molarity}}\text{Scm}^2\text{mol}^{-1}$

(C) Conductivity ( $\kappa$ ) =  $\frac{1}{\rho} = \frac{1}{RA}\Omega^{-1}\text{m}^{-1}$

(D) Degree of dissociation of electrolyte = Number of moles dissociated out of one mole.

It is a ratio. Hence, it is dimensionless.

Thus, the correct match is

A  $\rightarrow$  3, B  $\rightarrow$  1, C  $\rightarrow$  4, D  $\rightarrow$  2.

## Question78

**If the conductivity of mercury at  $0^\circ\text{C}$  is  $1.07 \times 10^6\text{Sm}^{-1}$  and the resistance of a cell containing mercury is  $0.243\Omega$ , then the cell constant of the cell is  $x \times 10^4\text{m}^{-1}$ . The value of x is..... .(Nearest integer)**

**[1 Sep 2021 Shift 2]**

**Answer: 26**

**Solution:**

Conductance (G) is reciprocal of resistance (R).

$$R = \frac{1}{G} \text{ or } G = \frac{1}{R} = \frac{1}{0.243\Omega} = 4.115\Omega^{-1}$$

Relation between conductance (G),

conductivity (kappa) and cell constant  $\left(\frac{l}{A}\right)$  is given as

$$k = \frac{Gl}{A}$$

$$\Rightarrow \frac{l}{A} = \frac{k}{G} = \frac{1.07 \times 10^6 \text{Sm}^{-1}}{4.115\Omega^{-1}} = 26 \times 10^4 \text{m}^{-1}$$

$$\Rightarrow x = 26$$

$\therefore$  Answer is 26.

## Question79

**The equation that is incorrect is:  
[Jan. 07, 2020 (II)]**

**Options:**

A.  $(\Lambda_m^0)_{\text{NaBr}} - (\Lambda_m^0)_{\text{NaCl}} = (\Lambda_m^0)_{\text{KBr}} - (\Lambda_m^0)_{\text{KCl}}$

B.  $(\Lambda_m^0)_{\text{KCl}} - (\Lambda_m^0)_{\text{NaCl}} = (\Lambda_m^0)_{\text{KBr}} - (\Lambda_m^0)_{\text{NaBr}}$

C.  $(\Lambda_m^0)_{\text{H}_2\text{O}} = (\Lambda_m^0)_{\text{HCl}} + (\Lambda_m^0)_{\text{NaOH}} - (\Lambda_m^0)_{\text{NaCl}}$

D.  $(\Lambda_m^0)_{\text{NaBr}} - (\Lambda_m^0)_{\text{NaI}} = (\Lambda_m^0)_{\text{KBr}} - (\Lambda_m^0)_{\text{NaBr}}$

**Answer: D**

**Solution:**

**Solution:**

$$\begin{aligned} &(\Lambda_m^0)_{\text{NaBr}} - (\Lambda_m^0)_{\text{NaI}} \\ &= \Lambda_m^0 \text{Na}^+ + \Lambda_m^0 \text{Br}^- - (\Lambda_m^0 \text{Na}^+ + \Lambda_m^0 \text{I}^-) \\ &= \Lambda_m^0 \text{Na}^+ + \Lambda_m^0 \text{Br}^- - \Lambda_m^0 \text{Na}^+ - \Lambda_m^0 \text{I}^- \\ &= \Lambda_m^0 \text{Br}^- + \Lambda_m^0 \text{I}^- \end{aligned}$$

$$\begin{aligned} &(\Lambda_m^0)_{\text{KBr}} - (\Lambda_m^0)_{\text{NaBr}} \\ &= \Lambda_m^0 \text{K}^+ + \Lambda_m^0 \text{Br}^- - (\Lambda_m^0 \text{Na}^+ + \Lambda_m^0 \text{Br}^-) \\ &= \Lambda_m^0 \text{K}^+ + \Lambda_m^0 \text{Br}^- - \Lambda_m^0 \text{Na}^+ - \Lambda_m^0 \text{Br}^- \\ &= \Lambda_m^0 \text{K}^+ + \Lambda_m^0 \text{Na}^+ \end{aligned}$$

$$\therefore (\Lambda_m^0)_{\text{NaBr}} - (\Lambda_m^0)_{\text{NaI}} \neq (\Lambda_m^0)_{\text{KBr}} - (\Lambda_m^0)_{\text{NaBr}}$$

## Question80

**108g of silver (molar mass  $108\text{gmol}^{-1}$ ) is deposited at cathode from  $\text{AgNO}_3(\text{aq})$  solution by a certain quantity of electricity. The volume (in L) of oxygen gas produced at 273K and 1 bar pressure from water by the same quantity of electricity is  
[NV, Jan. 09, 2020(I)]**


**Answer: 5.68**

**Solution:**


**Solution:**

No. of moles of silver deposited.

$$= \frac{108}{108} = 1 \text{ mol}$$



1F charge is required to deposit 1mol e of Ag



2F charge deposit  $\rightarrow \frac{1}{2}$  moles of oxygen

1F charge will deposit  $\rightarrow \frac{1}{4}$  moles of oxygen

$$V_{\text{O}_2} = \frac{nRT}{P}$$

$$= \frac{1}{4} \times \frac{0.08314 \text{ L bar K}^{-1} \text{ mol}^{-1} \times 273 \text{ K}}{1 \text{ bar}}$$


$$= \frac{1}{4} \times 22.7$$

$$V_{\text{O}_2} = 5.675 \text{ L}$$

---

## Question81

**What would be the electrode potential for the given half cell reaction at pH = 5 ?**



**( R =  $8.314 \text{ J mol}^{-1} \text{ K}^{-1}$ ; T emp =  $298 \text{ K}$  ; oxygen under std. atm. pressure of 1 bar )**  
**[NV, Jan. 08, 2020 (I)]**

**Answer: 1.52**

**Solution:**

$$E = 1.23 - \frac{0.0591}{4} \log[\text{H}^+]^4$$

$$= 1.23 + 0.0591 \times \text{pH}$$

$$= 1.23 + 0.0591 \times 5$$

$$= 1.23 + 0.2955$$

$$= 1.52 \text{ V}$$

---

## Question82



**For an electrochemical cell**

**$\text{Sn(s)} \mid \text{Sn}^{2+}(\text{aq}, 1\text{M}) \mid \text{Pb}^{2+}(\text{aq}, 1\text{M}) \mid \text{Pb(s)}$  the ratio  $\frac{[\text{Sn}^{2+}]}{[\text{Pb}^{2+}]}$**


**when this cell attains equilibrium is**

**[NV, Jan. 08, 2020 (II)]**

**Answer: 2.15**

**Solution:**

At equilibrium state  $E_{\text{cell}} = 0$ ;  $E_{\text{cell}}^0 = 0.01\text{V}$



$$E = E_{\text{cell}}^0 - \frac{0.06}{n} \log \frac{[\text{P}]}{[\text{R}]}$$

$$0 = 0.01 - \frac{0.06}{2} \log \frac{[\text{Sn}^{2+}]}{[\text{Pb}^{2+}]}$$

$$-0.01 = -\frac{0.06}{2} \log \frac{[\text{Sn}^{2+}]}{[\text{Pb}^{2+}]}$$

$$\frac{1}{3} = \log \frac{[\text{Sn}^{2+}]}{[\text{Pb}^{2+}]}$$

$$\frac{[\text{Sn}^{2+}]}{[\text{Pb}^{2+}]} = 10^{1/3} = 2.15$$

---

## Question83

**Given that the standard potentials (E) of  $\text{Cu}^{2+} / \text{Cu}$  and  $\text{Cu}^+ / \text{Cu}$  are 0.34V and 0.522V respectively, the  $E^0$  of  $\text{Cu}^{2+} / \text{Cu}$  is:**

**[Jan. 07,2020(I)]**

**Options:**

A. +0.182V

B. +0.158V


C. -0.182V

D. -0.158V


**Answer: B**

**Solution:**

**Solution:**



Subtract (ii) from (i)




$$\therefore \Delta G_1^\circ - \Delta G_2^\circ = \Delta G_3^\circ$$

$$\Rightarrow -F E^0 = -2F(0.34) + F(0.522)$$

$$\Rightarrow E^0 = 0.68 - 0.522 = 0.158\text{V}$$

## Question84

The variation of molar conductivity with concentration of an electrolyte (X) in aqueous solution is shown in the given figure.



The electrolyte X is :  
[Sep. 05, 2020 (II)]


Options:

- A. HCl
- B. NaCl
- C. KNO<sub>3</sub>
- D. CH<sub>3</sub>COOH

Answer: D

Solution:

Solution:



Among given electrolytes, CH<sub>3</sub>COOH is weak electrolyte.

## Question85

Let  $C_{\text{NaCl}}$  and  $C_{\text{BaSO}_4}$  be the conductances (in S) measured for saturated aqueous solutions of NaCl and BaSO<sub>4</sub>, respectively, at a temperature T. Which of the following is false?  
[Sep. 03, 2020 (I)]

Options:

- A. Ionic mobilities of ions from both salts increase with T
- B.  $C_{\text{BaSO}_4}(T_2) > C_{\text{BaSO}_4}(T_1)$  for  $T_2 > T_1$

C.  $C_{\text{NaCl}}(T_2) > C_{\text{NaCl}}(T_1)$  for  $T_2 > T_1$

D.  $C_{\text{NaCl}} \gg C_{\text{BaSO}_4}$  at a given T

E. None of above

**Answer: E**

**Solution:**

**Solution:**

(N) Ionic mobility increases with increase in temperature, which increases the conductance of the solution, while conductance of NaCl solution is independent of temperature above 400°C. Because the temperature is not given, so none of the option is correct.

-----

## Question86

Potassium chlorate is prepared by the electrolysis of KCl in basic solution  $6\text{OH}^- + \text{Cl}^- \rightarrow \text{ClO}_3^- + 3\text{H}_2\text{O} + 6\text{e}^-$  If only 60% of the current is utilized in the reaction, the time (rounded to the nearest hour) required to produce 10 g of  $\text{KClO}_3$  using a current of 2A is \_\_\_\_\_ .


(Given :  $F = 96,500 \text{ C mol}^{-1}$ ; molar mass of  $\text{KClO}_3 = 122 \text{ g mol}^{-1}$  )

[NV, Sep. 06, 2020 (I)]

**Answer: 11**

**Solution:**

**Solution:**



For synthesis of 1 mole of  $\text{ClO}_3^-$ , 6F of charge is required.

∴ Current efficiency = 60%

∴ To synthesis 1 mole of  $\text{ClO}_3^-$ , 10F of charge is required.

To synthesis  $\frac{10}{122}$  moles of  $\text{KClO}_3$ , charge =  $\frac{10 \times 10}{122} F$   $Q = I \cdot t$

$$t = \frac{100 \times 96500}{122 \times 2} = 39549.18 \text{ s}$$

$$= \frac{79098.365}{3600 \text{ s}} = 10.99 \text{ h}$$

$$\therefore t = 11 \text{ h.}$$

-----

## Question87

250mL of a waste solution obtained from the workshop of a goldsmith contains 0.1M  $\text{AgNO}_3$  and 0.1M  $\text{AuCl}$ . The solution was electrolyzed at 2V by passing a current of 1 A for 15 minutes. The metal/metals

**electrodeposited will be:**  
**[Sep. 04, 2020(II)]**

$$(E_{\text{Ag}^+ / \text{Ag}}^0 = 0.80\text{V}, E_{\text{Au}^+ / \text{Au}}^0 = 1.69\text{V})$$

**Options:**

- A. only gold
- B. silver and gold in proportion to their atomic weights
- C. only silver
- D. silver and gold in equal mass proportion

**Answer: A**

**Solution:**

**Solution:**

$$\text{Millimoles of Au}^+ = 0.1 \times 250 = 25$$

$$\text{Mole of Au}^+ = \frac{25}{1000} = \frac{1}{40} = 0.025$$

$$\text{Similarly, moles of Ag}^+ = 0.025$$


$$\text{Charge passed} = I \times t = 1 \times 15 \times 60 = 900\text{C}$$

$$\text{Moles of e}^- \text{ passed} = \frac{900}{96500} = 0.0093\text{mol}$$

Species with higher value of SRP will get deposited first at cathode.

## Question 88

**An acidic solution of dichromate is electrolyzed for 8 minutes using 2 A current. As per the following equation**



**The amount of  $\text{Cr}^{3+}$  obtained was 0.104g. The efficiency of the process (in %) is (Take :  $F = 96000\text{C}$ , At. mass of chromium = 52 ) \_\_\_\_\_.**  
**[NV, Sep. 03, 2020 (II)]**


**Answer: 60**

**Solution:**

**Solution:**

$$\text{Charge (Q)} = I t = 2 \times 8 \times 60 = 960\text{C}$$

$$\Rightarrow \frac{960}{96000} = 0.01F$$




$$\text{Theoretical mass of Cr}^{3+} = \frac{1}{3} \times \frac{960}{96000} \times 52 = 0.173\text{g}$$

$$\text{So, efficiency} = \frac{W_{\text{actual}}}{W_{\text{Theoretical}}} \times 100 = \frac{0.104}{0.173} \times 100 = 60\%$$


---

## Question89

**For the given cell;**



**change in Gibbs energy ( $\Delta G$ ) is negative, if:**

**[Sep. 06, 2020 (II)]**

**Options:**

A.  $C_1 = C_2$

B.  $C_2 = \frac{C_1}{\sqrt{2}}$




C.  $C_1 = 2C_2$

D.  $C_2 = \sqrt{2}C_1$

**Answer: D**

**Solution:**

For the concentration cell,  $E_{\text{cell}}^0 = 0$



As  $\Delta G = -nF E$

If  $\Delta G = -ve$ , then  $E_{\text{cell}}$  is  $+ve$ .

$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{RT}{2F} \ln \frac{C_1}{C_2}$$

$$E_{\text{cell}} = 0 - \frac{RT}{2F} \ln \frac{C_1}{C_2}$$

$$E_{\text{cell}} = \frac{RT}{2F} \ln \frac{C_2}{C_1}$$

So,  $C_2 > C_1$ .

Thus,  $C_2 = \sqrt{2}C_1$  relation is correct.

---

## Question90

**An oxidation-reduction reaction in which 3 electrons are transferred has a  $\Delta G^0$  of  $17.37\text{kJ mol}^{-1}$  at  $25^\circ\text{C}$ . The value of  $E_{\text{cell}}^0(\text{inV})$  is \_\_\_\_\_  $\times 10^{-2}$**

**( $1F = 96,500\text{Cmol}^{-1}$ )**

**[NV, Sep. 05, 2020(I)]**


**Answer: -6**

**Solution:**

$$\Delta G^0 = -nF E_{\text{cell}}^0$$
$$17.37 \times 10^3 = -3 \times 96500 \times E_{\text{cell}}^0$$
$$E_{\text{cell}}^0 = -0.06V \sim \text{eq} - 6.0 \times 10^{-2}V$$

---

## Question91



$$E_{\text{Cu}^{2+} | \text{Cu}}^0 = +0.34V ; E_{\text{Zn}^{2+} | \text{Zn}}^0 = -0.76V$$

**Identify the incorrect statement from the options below for the above cell :**

**[Sep. 04, 2020(I)]**

**Options:**

- A. If  $E_{\text{ext}} > 1.1V$ ,  $e^-$  flows from Cu to Zn
- B. If  $E_{\text{ext}} > 1.1V$ , Zn dissolves at Zn electrode and Cu deposits at Cu electrode
- C. If  $E_{\text{ext}} < 1.1V$ , Zn dissolves at anode and Cu deposits at cathode
- D. If  $E_{\text{ext}} = 1.1V$ , no flow of  $e^-$  or current occurs

**Answer: B**

**Solution:**

**Solution:**

$$E_{\text{cell}}^0 = E_{\text{Cu}^{2+} | \text{Cu}}^0 - E_{\text{Zn}^{2+} | \text{Zn}}^0 = 1.1V$$

So, if  $E_{\text{ext.}} = 1.1V$  no electron will flow

At,  $E_{\text{ext.}} > 1.1V$  cell act as electrolytic cell and electron will flow from Cu to Zn.

At,  $E_{\text{ext.}} < 1.1V$  cell act as electrochemical cells so Zn dissolve and Cu deposit.

---

## Question92

The photoelectric current from Na (work function,  $w_0 = 2.3\text{eV}$  ) is stopped by the output voltage of the cell  
 $\text{Pt(s)} \mid \text{H}_2(\text{g}, 1\text{bar}) \mid \text{HCl}(\text{aq.}, \text{pH} = 1) \mid \text{AgCl(s)} \mid \text{Ag(s)}$  The pH of aq. HCl required to stop the photoelectric current from K ( $w_0 = 2.25\text{eV}$  ), all other conditions remaining the same, is \_\_\_\_\_  $\times 10^{-2}$  (to the nearest integer).

Given,

$$2.303 \frac{RT}{F} = 0.06\text{V}; E_{\text{AgCl} \mid \text{Ag} \mid \text{Cl}^-}^0 = 0.22\text{V}$$

[NV, Sep. 03 , 2020(I)]

**Answer: 142**




**Solution:**

**Solution:**

**Sodium metal :**

$$E = E_0 + (KE)_{\text{max}}; E_{\text{cell}}^0 = 0.22\text{V}$$

Cell reaction



$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.06}{1} \log[\text{H}^+][\text{Cl}^-]$$

$$E_{\text{cell}} = 0.22 - \frac{0.06}{1} \log[10^{-1}][10^{-1}]$$

$$= 0.22 + 0.12 = 0.34\text{V}$$

$$(KE)_{\text{max}} = E_{\text{cell}} = 0.34\text{eV}$$

$$\text{So, } E = 2.3 + 0.34 = 2.64\text{eV} = \text{Energy of photon incident}$$




**For potassium metal :**

$$E = E_0 + (KE)_{\text{max}}$$

$$2.64 = 2.25 + (KE)_{\text{max}}$$

$$(KE)_{\text{max}} = 0.39 = E_{\text{cell}}$$

Cell reaction



$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.06}{1} \log[\text{H}^+][\text{Cl}^-]$$

$$0.39 = 0.22 - 0.06 \log[\text{H}^+]^2$$


$$0.39 = 0.22 - 0.12 \log[\text{H}^+]$$

$$0.17 = 0.12 \times \text{pH}$$

$$\text{pH} = 17 / 12 = 1.4166 \sim \text{eq} 1.42 = 142 \times 10^{-2}$$

## Question93

The Gibbs energy change (in J) for the given reaction at  $[\text{Cu}^{2+}] = [\text{Sn}^{2+}] = 1\text{M}$  and 298K is :



$$\left( E^0_{\text{Sn}^{2+} | \text{Sn}} = -0.16\text{V}, E^0_{\text{Cu}^{2+} | \text{Cu}} = 0.34\text{V}, \text{ Take } F = 96500\text{cmol}^{-1} \right)$$

[NV, Sep. 02, 2020(I)]

**Answer: 96500**

**Solution:**

**Solution:**

$$\text{Sol. } E_{\text{cell}}^0 = E_{\text{Sn}^{2+} / \text{Sn}}^0 - E_{\text{Cu}^{2+} / \text{Cu}}^0$$

$$= -0.16 - 0.34$$

$$= -0.50\text{V}$$

$$\Delta G^0 = -nF E_{\text{cell}}^0$$

$$= -2 \times 96500 \times (-0.5)$$

$$= 96500\text{J}$$

$$= 96.5\text{KJ} = 96500\text{J}$$

## Question94

For the disproportionation reaction  $2\text{Cu}^+(\text{aq}) \rightleftharpoons \text{Cu(s)} + \text{Cu}^{2+}(\text{aq})$  at 298K,  $\ln K$

(where K is the equilibrium constant) is  $\times 10^{-1}$ .

Given


[NV, Sep. 02,2020(II)]

$$\left( E_{\text{Cu}^{2+} / \text{Cu}^+}^0 = 0.16\text{V}; E_{\text{Cu}^+ / \text{Cu}}^0 = 0.52\text{V}; \frac{RT}{F} = 0.025 \right)$$

**Answer: 144**

**Solution:**

**Solution:**



$$E_{\text{cell}}^0 = E_{\text{Cu}^+ / \text{Cu}}^0 - E_{\text{Cu}^{2+} / \text{Cu}^+}^0 = 0.52 - 0.16 = 0.36\text{V}$$

$$E_{\text{cell}}^0 = \frac{RT}{nF} \ln K_{\text{eq}}$$

$$\Rightarrow 0.36 = \frac{0.025}{1} \ln K$$

$$\Rightarrow \ln K = 14.4 = 144 \times 10^{-1}$$

## Question95



$\Lambda_m^\circ$  for NaCl, HCl and NaA are 126.4, 425.9 and 100.5  $\text{Scm}^2\text{mol}^{-1}$ , respectively. If the conductivity of 0.001 M HA is  $5 \times 10^{-5} \text{Scm}^{-1}$ , degree of dissociation of HA is :  
[Jan. 12, 2019 (II)]

**Options:**

- A. 0.50
- B. 0.25
- C. 0.125
- D. 0.75

**Answer: C**

**Solution:**

**Solution:**

Given:  $\Lambda_m^\circ(\text{NaCl}) = 126.4 \text{Scm}^2\text{mol}^{-1}$ ,  $\Lambda_m^\circ(\text{HCl}) = 425.9 \text{Scm}^2\text{mol}^{-1}$  and  $\Lambda_m^\circ(\text{NaA}) = 100.5 \text{Scm}^2\text{mol}^{-1}$

$$\Lambda_m^\circ(\text{HA}) = \lambda_{\text{H}^+}^\circ + \lambda_{\text{A}^-}^\circ$$

$$= \lambda_{\text{H}^+}^\circ + \lambda_{\text{Cl}^-}^\circ + \lambda_{\text{A}^+}^\circ + \lambda_{\text{Na}^+}^\circ - \lambda_{\text{Cl}^-}^\circ - \lambda_{\text{Na}^+}^\circ$$

$$= \Lambda_m^\circ(\text{HCl}) + \Lambda_m^\circ(\text{NaA}) - \Lambda_m^\circ(\text{NaCl})$$

$$\Lambda_m^\circ(\text{HA}) = 425.9 - 126.4 + 100.5 = 400 \text{Scm}^2\text{mol}^{-1}$$

$$\kappa(\text{HA}) = 5 \times 10^{-5} \text{Scm}^{-1}$$

$$(\text{HA}) = \frac{\kappa(\text{HA}) \times 1000}{\text{Molality of HA}} = \frac{5 \times 10^{-5} \times 1000}{0.001} = 50 \text{Scm}^2\text{mol}^{-1}$$

$$\alpha = \frac{\Lambda_m(\text{HA})}{\Lambda_m^\circ(\text{HA})} = \frac{50}{400} = 0.125$$

## Question96

The anodic half-cell of lead-acid battery is recharged using electricity of 0.05 Faraday. The amount of  $\text{PbSO}_4$  electrolyzed in g during the process is :

(Molar mass of  $\text{PbSO}_4 = 303 \text{gmol}^{-1}$ )

[Jan. 9, 2019 (I)]

**Options:**

- A. 22.8
- B. 15.2
- C. 7.6
- D. 11.4

**Answer: A**

**Solution:**


According to the Faraday's law of electrolysis,  $nF$  of current is required for the deposition of 1mol  
 According to the reaction,  

$$\text{Ni}(\text{NO}_3)_2 \rightarrow \text{Ni}^{2+} + 2\text{NO}_3^-$$
  
 $2F$  of current deposits = 1mol  
 $\therefore 0.1F$  of current deposits =  $\frac{0.1}{2} = 0.05\text{mol}$

---

## Question97

The standard electrode potential  $E^\circ$  and its temperature coefficient  $\left(\frac{dE^\circ}{dT}\right)$  for a cell are 2V and  $-5 \times 10^{-4} \text{V K}^{-1}$  at 300K respectively. The cell reaction is:



The standard reaction enthalpy ( $\Delta_r H^\circ$ ) at 300K in  $\text{kJ mol}^{-1}$  is,

[ Use  $R = 8 \text{J K}^{-1} \text{mol}^{-1}$  and  $F = 96,000 \text{Cmol}^{-1}$  ]

[Jan. 12, 2019 (I)]

Options:

A. -412.8

B. -384.0

C. 192.0

D. 206.4

Answer: A


Solution:

$$\Delta G^\circ = \Delta_r H^\circ - T \Delta S^\circ$$

$$\Delta_r H^\circ = \Delta G^\circ + T \Delta S^\circ$$

$$\Delta_r H^\circ = -nF E^\circ + T nF \frac{dE^\circ}{dT}$$

$$\Delta_r H^\circ = -nF E^\circ + nF T \frac{dE^\circ}{dT}$$



$$\Delta_r H^\circ = -nF \left( E^\circ - T \frac{dE^\circ}{dT} \right)$$

$$\Delta_r H^\circ = -2 \times 96000(2 - 300 \times -5 \times 10^{-4})$$

$$\Delta_r H^\circ = -2 \times 96000(2 + 300 \times 5 \times 10^{-4})$$

$$= -2 \times 96000(2 + 0.15)$$

$$= -412.8 \times 10^3 \text{J / mol} ; = -412.8 \text{kJ / mol}$$


---

## Question98

For the cell  $\text{Zn(s)} | \text{Zn}^{2+}(\text{aq}) | \text{M}^{x+}(\text{aq}) | \text{M(s)}$ , different half cells and their standard electrode potentials are given below:

$M^{x+}(aq) \backslash$	$Au^{3+}(aq)$	$Ag^+(aq) \backslash$	$Fe^{3+}(aq)$	$Fe^{2+}(aq)$
$M(s)$	$Au(s)$	$Ag(s)$	$Fe^{2+}(aq)$	$Fe(s)$
$E^\circ M^{x+/M} (V)$	1.40	0.80	0.77	0.44

If  $E_{Zn^{2+}/Zn}^\circ = -0.76V$ , which cathode will give a maximum value of  $E_{cell}^\circ$  per electron transferred?  
**[Jan. 11, 2019 (I)]**

**Options:**

- A.  $Ag^+ / Ag$
- B.  $Fe^{3+} / Fe^{2+}$
- C.  $Au^{3+} / Au$
- D.  $Fe^{2+} / Fe$

**Answer: A**

**Solution:**

$$Zn + 2Ag^+ \longrightarrow 2Ag + Zn^{2+} (n = 2e^-)$$

$$E_{cell}^\circ = (E_{R.P}^\circ)_{cathode} - (E_{R.P}^\circ)_{anode}$$


$$= 0.80 - (-0.76) = 1.56V \text{ for } 2e^-$$

$$\therefore E_{cell}^\circ \text{ for } 1e^- = \frac{1.56}{2} = 0.78V$$

## Question99

**Given the equilibrium constant:**

**$K_c$  of the reaction:**



**$10 \times 10^{15}$  calculate the  $E_{cell}^\circ$  of this reaction at 298K**

$$\left[ 2.303 \frac{RT}{F} \text{ at } 298K = 0.059V \right]$$

**[Jan. 11, 2019 (II)]**

**Options:**

- A. 0.04736mV
- B. 0.4736mV
- C. 0.4736V
- D. 0.04736V

**Answer: C**

## Solution:

### Solution:

$$E_{\text{cell}}^0 = \frac{2.303RT}{nF} \log K_C \text{ or } E_{\text{cell}}^0 = \frac{0.059V}{n} \log K_C$$
$$= \frac{0.059V}{2} \log 10^{16} = 0.4736V$$

---

## Question 100

### In the cell

**Pt(s) | H<sub>2</sub>(g, 1 bar) / HCl(aq) | AgCl(s) / Ag(s) | Pt(s), the cell potential is 0.92V when a 10<sup>-6</sup> molal HCl solution is used. The standard electrode potential of (AgCl / Ag, Cl<sup>-</sup>) electrode is:**

**{ Given :  $\frac{2.303RT}{F} = 0.06V$  at 298K }**  
**[Jan. 10, 2019 (II)]**

### Options:


- A. 0.94V
- B. 0.76V
- C. 0.40V
- D. 0.20V

**Answer: D**

## Solution:



### Solution:

Given that:




$$E_{\text{cell}} = 0.92V$$

$$\text{Now, } E_{\text{cell}} = E_{\text{H}_2(\text{g}) / \text{H}^+(\text{aq})}^{\circ} + E_{\text{AgCl(s)} / \text{Ag(s)} / \text{Cl}^-}^{\circ} - \frac{0.06}{n} \log Q$$



Net cell reaction:



$$\therefore Q = \frac{[\text{H}^+][\text{Cl}^-]}{(P_{\text{H}_2})^{1/2}}$$

Here, 10<sup>-6</sup> molal HCl solution is used

$$\text{So } Q = \frac{10^{-6} \times 10^{-6}}{1} = 10^{-12}$$

(assuming molality = molarity)


$$\text{Now, } 0.92 = E_{\text{AgCl(s)} / \text{Ag(s)} / \text{Cl}^-}^{\circ} - \frac{0.06}{1} \log 10^{-12}$$

$$E_{\text{AgCl(s)} / \text{Ag(s)} / \text{Cl}^-}^{\circ} = 0.92 + [0.06 \times (-12)]$$
$$= 0.92 - 0.72 = 0.20V$$

---

## Question101

If the standard electrode potential for a cell is 2V at 300K, the equilibrium constant (K) for the reaction



at 300K is approximately ( $R = 8\text{ J K}^{-1}\text{mol}^{-1}$ ,  $F = 96000\text{ Cmol}^{-1}$ )  
[Jan. 9, 2019 (II)]

Options:

A.  $e^{-80}$

B.  $e^{-160}$

C.  $e^{320}$

D.  $e^{160}$

Answer: D

Solution:

Solution:

We know that,

$$\ln K = \frac{nF E_{\text{cell}}^{\circ}}{RT}$$

$$\Delta G = -nF E_{\text{cell}}^{\circ}$$

$$\Delta G = -RT \ln k$$

Now,

$$\text{After putting the given values, we get } \ln K = \frac{2 \times 96000 \times 2}{8 \times 300} = 160$$

$$\therefore K = e^{160}$$

---

## Question102

The decreasing order of electrical conductivity of the following aqueous solutions is:

0.1 M Formic acid (A),

0.1 M Acetic acid (B),

0.1 M Benzoic acid (C).

[April 12, 2019 (II)]

Options:

A.  $A > C > B$

B.  $C > B > A$

C.  $A > B > C$

D.  $C > A > B$

Answer: A


## Solution:

### Solution:

Order of acidic strength is  $\text{HCOOH} > \text{C}_6\text{H}_5\text{COOH} > \text{CH}_3\text{COOH}$

More the acidic strength more will be the dissociation of acid into ions and more will be the conductivity.

Thus, order of conductivity will be,



---

## Question 103

Consider the statements S1 and S2 :

**S1: Conductivity always increases with decrease in the concentration of electrolyte. S2: Molar conductivity always increases with decrease in the concentration of electrolyte. The correct option among the following is: [April 10, 2019 (I)]**

### Options:

- A. Both S1 and S2 are wrong
- B. S1 is wrong and S2 is correct
- C. Both S1 and S2 are correct
- D. S1 is correct and S2 is wrong

**Answer: B**

## Solution:

### Solution:

Conductivity of an electrolyte is the conductance of  $1\text{cm}^3$  of the given electrolyte. It increases with the increase in concentration of electrolyte due to increase in the number of ions per unit volume. Molar conductivity ( $\lambda_m$ ) is the conductance of a solution containing 1 mole of the electrolyte. It increases with the decrease of concentration due to increase in the total volume having one mole of electrolyte. Thus, interionic attraction increases and degree of ionisation decreases. Therefore, (S<sub>1</sub>) is wrong and (S<sub>2</sub>) is correct.


---

## Question 104


**Which one of the following graphs between molar conductivity ( $\Lambda_m$ ) versus  $\sqrt{C}$  is correct? [April 10, 2019 (II)]**

### Options:


- A.




B.



C.



D.



**Answer: A**

**Solution:**

**Solution:**

Since, KCl is more conducting than NaCl, therefore, graph (a) is correct.

## Question 105

A solution of  $\text{Ni}(\text{NO}_3)_2$  is electrolysed between platinum electrodes using 0.1 Faraday electricity. How many mole of Ni will be deposited at the cathode?

[April 9, 2019 (II)]


**Options:**

- A. 0.05
- B. 0.20
- C. 0.15
- D. 0.10


**Answer: C**

**Solution:**


The reaction involved in the process is given below



So, half cell reaction is



According to the reaction:



We require 2F for the electrolysis of 1mol or 303g of  $\text{PbSO}_4$


$\therefore$  Amount of  $\text{PbSO}_4$  electrolysed by 0.05F

$$= \frac{303}{2} \times 0.05 = 7.575\text{g} \approx 7.6\text{g}$$

---

## Question 106

**The standard Gibbs energy for the given cell reaction in  $\text{kJ mol}^{-1}$  at 298K is:**



**$E^\circ = 2\text{V}$  at 298K**

**(Faraday's constant,  $F = 96000\text{Cmol}^{-1}$ )**

**[April 9, 2019(I)]**

**Options:**

- A.  $-384$
- B. 384
- C. 192
- D.  $-192$

**Answer: A**

**Solution:**

**Solution:**


$$\Delta G^\circ = -nF E_{\text{cell}}^\circ$$

$$= -2 \times (96000) \times 2\text{V} = -384000\text{J / mol} = -384\text{kJ / mol}$$



## Question107

Calculate the standard cell potential (in V ) of the cell in which following reaction takes place:



Given that

$$E_{\text{Ag}^+ / \text{Ag}}^0 = x\text{V}; E_{\text{Fe}^{2+} / \text{Fe}}^0 = y\text{V}; E_{\text{Fe}^{3+} / \text{Fe}}^0 = z\text{V}$$


[April 8, 2019 (II)]

Options:


- A.  $x - z$
- B.  $x - y$
- C.  $x + 2y - 3z$
- D.  $x + y - z$

Answer: C

Solution:



$$E_{\text{cell}}^0 = E_{\text{Ag}^+ / \text{Ag}}^0 - E_{\text{Fe}^{3+} / \text{Fe}^{2+}}^0$$



$$E_{\text{Fe}^{3+} / \text{Fe}^{2+}}^0 = 3z - 2y$$

$$E_{\text{Ag}^+ / \text{Ag}}^0 = x \quad (\text{given})$$

$$\therefore E_{\text{cell}}^0 = x - 3z + 2y$$

## Question108

How long (approximate) should water be electrolysed by passing through 100 amperes current so that the oxygen released can completely burn 27.66g of diborane? (Atomic weight of B = 10.8u ) [2018]


Options:

- A. 6.4 hours
- B. 0.8 hours
- C. 3.2 hours
- D. 1.6 hours

**Answer: C**


**Solution:**

**Solution:**



27.66g of  $\text{B}_2\text{H}_6$  (1mol e) requires 3 moles of oxygen ( $\text{O}_2$ ) for complete burning.

Now the oxygen is produced by the electrolysis of  $\text{H}_2\text{O}$ .



1 mole  $\text{O}_2$  is produced by  $4\text{F}$  charge

$\therefore$  3mol  $\text{eO}_2$  will be produced by  $12\text{F}$  charge.

$$\therefore Q = It$$

$$12 \times 96500\text{C} = I \times t$$

$$12 \times 96500\text{C} = 100 \times t$$

$$t = \frac{12 \times 96500}{100} \text{second}$$

$$= \frac{12 \times 96500}{100 \times 3600} \text{hour}$$

$$= 3.2 \text{ hours}$$

## Question 109

**When 9.65 ampere current was passed for 1.0 hour into nitrobenzene in acidic medium, the amount of p -aminophenol produced is [Online April 16, 2018]**

**Options:**

A. 109.0g

B. 98.1g

C. 9.81g

D. 10.9g

**Answer: C**



**Solution:**

**Solution:**

9.65 ampere current was passed for 1.0 hour ( 3600 seconds)

Number of moles of electrons passed

$$= \frac{I (\text{A}) \times t (\text{s})}{96500} = \frac{9.65\text{A} \times 3600\text{s}}{96500} = 0.36\text{mol es}$$



$\therefore$  4 moles of electrons reduces 1 mole of nitrobenzene to p -aminophenol.  $\therefore$  0.36 moles of electrons will reduce

$$\frac{0.36}{4} = 0.09 \text{ moles of nitrobenzene to p -aminophenol.}$$

p -aminophenol molar mass = 109.14g / mol

Mass of p -aminophenol obtained = 109.14g / mol  $\times$  0.09mol = 9.81g

## Question110

**When an electric current is passes through acidified water, 112mL of hydrogen gas at N.T.P. was collected at the cathode in 965 seconds. The current passed, in ampere, is :  
[Online April 15,2018 (I)]**

**Options:**

A. 2.0

B. 0.1

C. 0.5


D. 1.0

**Answer: D**

**Solution:**

**Solution:**

Reduction at cathode:



At NTP 22400mL of  $H_2$  = 1mol e of  $H_2$

$$112\text{mL of } H_2 = \frac{1}{22400} \times 112 = 0.005\text{mol e of } H_2$$

$$\text{Moles of } H_2 \text{ produced} = \frac{I \times t}{96500} \times \text{molar ratio}$$

$$0.005 = \frac{I \times 965}{96500} \times \frac{1\text{mol e of } H_2}{2\text{mol e of } e^-} \quad I = 1.0\text{A}$$

## Question111

**Given**

$$E_{Cl_2 / Cl^-}^0 = 1.36V, E_{Cr^{3+} / Cr}^0 = -0.74V$$

$$E_{Cr_2 / O_7^{2-} / Cr^{3+}} = 1.33V, E_{MnO_4^- / Mn^{2+}}^0 = 1.51V$$

**Among the following, the strongest reducing agent is  
[2017]**

**Options:**

A. Cr

B.  $Mn^{2+}$

C.  $Cr^{3+}$

D. Cl

**Answer: A**

**Solution:**

**Solution:**

$$E_{\text{MnO}_4^- / \text{Mn}^{2+}}^\circ = 1.51\text{V}$$

$$E_{\text{Cl}_2 / \text{Cl}^-}^\circ = 1.36\text{V}$$

$$E_{\text{C}_2\text{O}_7^{2-} / \text{Cr}^{3+}} = 1.33\text{V}$$

$$E_{\text{Cr}^{3+} / \text{Cr}}^\circ = -0.74$$

Since  $\text{Cr}^{3+}$  is having least reduction potential, so Cr is the best reducing agent.

## Question 112

Consider the following standard electrode potentials ( $E^\circ$  in volts) in aqueous solution:

Element	$M^{3+}/M$	$M^+/M$
Al	-1.66	+0.55
Tl	+1.26	-0.34

Based on these data, which of the following statements is correct?  
[Online April 8, 2017]



**Options:**

- A.  $\text{Tl}^+$  is more stable than  $\text{Al}^{3+}$
- B.  $\text{Al}^+$  is more stable than  $\text{Al}^{3+}$
- C.  $\text{Tl}^+$  is more stable than  $\text{Al}^+$
- D.  $\text{Tl}^{3+}$  is more stable than  $\text{Al}^{3+}$

**Answer: C**

**Solution:**

**Solution:**





$\text{Tl}^+$  has negative electrode potential ( $E^\circ = -0.34$ ) means, it does not prefer to convert into Tl but reverse must be preferred that's why it is more stable than  $\text{Tl}^{3+}$  ( $E^\circ = +1.26$ ). In Al,  $\text{Al}^{3+}$  is more stable ( $E^\circ = -1.66$ ) than  $\text{Al}^+$  ( $E^\circ = +0.55$ ) and also from Tl due to more negative value of  $E^\circ$ . Therefore, by comparison it confirms that  $\text{Tl}^+$  is more stable than  $\text{Al}^+$ .

## Question 113

**What is the standard reduction potential ( $E^\circ$ ) for  $\text{Fe}^{3+} \rightarrow \text{Fe}$  ?**

**Given that :**



**[Online April 8, 2017]**

**Options:**

A.  $-0.057\text{V}$

B.  $+0.057\text{V}$

C.  $+0.30\text{V}$




D.  $-0.30\text{V}$

**Answer: A**

**Solution:**

**Solution:**

$$\Delta G^\circ = -nFE^\circ$$



(i)  $\Delta G^\circ = -nFE^\circ = -2(-0.47)F = 0.94F$

(ii)  $\Delta G^\circ = -nFE^\circ = -1(+0.77)F = -0.77F$

(iii) On adding :  $\Delta G^\circ = +0.17F$

$$\Delta G^\circ = -nFE^\circ$$

$$E^\circ \text{ for } (\text{Fe}^{3+} \rightarrow \text{Fe}) = \frac{\Delta G^\circ}{-nF} = \frac{0.17F}{-3F} = -0.057\text{V}$$

---

## Question114

**To find the standard potential of  $\text{M}^{3+} / \text{M}$  electrode, the following cell is constituted :  $\text{Pt} / \text{M} / \text{M}^{3+}(0.001\text{mol L}^{-1}) \vee \text{Ag}^+(0.01\text{mol L}^{-1}) / \text{Ag}$**

**The emf of the cell is found to be 0.421 volt at 298K . The standard potential of half reaction  $\text{M}^{3+} + 3\text{e}^- \rightarrow \text{M}$  at 298K will be:**

**(Given  $E^\circ_{\text{Ag}^+ / \text{Ag}}$  at 298K = 0.80V ol t )**

**[Online April 9, 2017]**

**Options:**

A. 0.38 Volt

B. 0.32 Volt


C. 1.28 Volt

D. 0.66 Volt

**Answer: B**

**Solution:**

Cell reaction:



$$E = E^0 - \frac{0.0591}{n} \log \left[ \frac{\text{Reduced state}}{\text{Oxidised state}} \right]$$

$$0.421 = E^0 - \frac{0.0591}{3} \log \frac{0.001}{(0.01)^3}$$

$$E^0 = 0.48$$

$$E^0 = E_{Ag^+/Ag}^0 - E_{M^{3+}/M}^0$$

$$E_{M^{3+}/M}^0 = 0.8V - 0.48V = 0.32 \text{ volt}$$

---

## Question 115

**What will occur if a block of copper metal is dropped into a beaker containing a solution of 1M ZnSO<sub>4</sub> ?**

**[Online April 9, 2016]**

**Options:**

- A. The copper metal will dissolve with evolution of oxygen gas
- B. The copper metal will dissolve with evolution of hydrogen gas
- C. Noreaction will occur
- D. The copper metal will dissolve and zinc metal will be deposited.

**Answer: C**

**Solution:**

**Solution:**

No reaction will occur as the Zn is placed above Cu in electrochemical series. Hence there will be no displacement reaction.

---

## Question 116

**Galvanization is applying a coating of:**  
**[2016]**

**Options:**

- A. Cu
- B. Zn
- C. Pb
- D. Cr

**Answer: B**

**Solution:**

**Solution:**

Galvanization is the process by which zinc is coated over corrosive (easily rusted) metals to prevent them from corrosion.

---

## Question 117

**Identify the correct statement:**

**[Online April 10, 2016]**

**Options:**

- A. Corrosion of iron can be minimized by forming a contact with another metal with a higher reduction potential
- B. Iron corrodes in oxygen free water
- C. Corrosion of iron can be minimized by forming an impermeable barrier at its surface
- D. Iron corrodes more rapidly in salt water because its electrochemical potential is higher

**Answer: C**

**Solution:**

**Solution:**

Corrosion of iron can be minimized by forming an impermeable barrier at its surface.

---

## Question 118


**Two Faraday of electricity is passed through a solution of  $\text{CuSO}_4$ . The mass of copper deposited at the cathode is (at. mass of Cu = 63.5 amu ) [2015]**

**Options:**

- A. 2g
- B. 127g
- C. 0g
- D. 63.5g

**Answer: D**

**Solution:**

**Solution:**

2F i.e.  $2 \times 96500\text{C}$  deposit Cu = 1mol  
= 63.5g

---

## Question119

A variable, opposite external potential ( $E_{\text{ext}}$ ) is applied to the cell  $\text{Zn} | \text{Zn}^{2+}(1\text{M}) | \text{Cu}^{2+}(1\text{M}) | \text{Cu}$ , of potential 1.1V. When  $E_{\text{ext}} < 1.1\text{V}$  and  $E_{\text{ext}} > 1.1\text{V}$ , respectively electrons flow from:  
[Online April 10, 2015]

**Options:**

- A. anode to cathode in both cases
- B. cathode to anode and anode to cathode
- C. anode to cathode and cathode to anode
- D. cathode to anode in both cases

**Answer: C**

**Solution:**

**Solution:**

EMF of galvanic cell = 1.1 volt

If  $E_{\text{ext}} < E_{\text{MF}}$  then electrons flow steadily from anode to cathode, while if  $E_{\text{ext}} > E_{\text{MF}}$  then electrons flow from cathode to anode as polarity is changed.

-----

## Question120

At 298K, the standard reduction potentials are 1.51V for  $\text{MnO}_4^- | \text{Mn}^{2+}$ , 1.36V for  $\text{Cl}_2 | \text{Cl}^-$ , 1.07V for  $\text{Br}_2 | \text{Br}^-$  and 0.54V for  $\text{I}_2 | \text{I}^-$ . At pH = 3, permanganate is expected to oxidize :  $\left( \frac{RT}{F} = 0.059\text{V} \right)$   
[Online April 11, 2015]


**Options:**

- A. Cl, Br and I
- B. Br and  $\text{I}^-$
- C.  $\text{Cl}^-$  and  $\text{Br}^-$
- D.  $\text{I}^-$  only

**Answer: B**

**Solution:**

**Solution:**



$$E = 1.51 - \frac{0.059}{5} \log \frac{[\text{Mn}^{2+}]}{[\text{MnO}_4^-][\text{H}^+]^8}$$

Taking  $\text{Mn}^{2+}$  and  $\text{MnO}_4^-$  in standard state i.e. 1M,



$$E = 1.51 - \frac{0.059}{5} \times 8 \log \frac{1}{[H^+]}$$

$$= 1.51 - \frac{0.059}{5} \times 8 \times 3 = 1.2268V$$

Hence at this pH,  $MnO_4^-$  will oxidise only  $Br^-$  and  $I^-$  as SRP of  $Cl_2 / Cl^-$  is 1.36V which is greater than that for  $MnO_4^- / Mn^{2+}$

## Question121

**Resistance of 0.2M solution of an electrolyte is  $50\Omega$ . The specific conductance of the solution is  $1.4Sm^{-1}$ . The resistance of 0.5M solution of the same electrolyte is  $280\Omega$ . The molar conductivity of 0.5M solution of the electrolyte in  $Sm^2mol^{-1}$  is:**  
**[2014]**

**Options:**

- A.  $5 \times 10^{-4}$
- B.  $5 \times 10^{-3}$
- C.  $5 \times 10^3$
- D.  $5 \times 10^2$

**Answer: A**

**Solution:**

**Solution:**

Given for 0.2M solution

$R = 50\Omega$

$\kappa = 1.4Sm^{-1} = 1.4 \times 10^{-2}Scm^{-1}$

Now,  $R = \rho \frac{l}{a} = \frac{1}{\kappa} \times \frac{l}{a}$

$\Rightarrow \frac{l}{a} = R \times \kappa = 50 \times 1.4 \times 10^{-2}cm^{-1}$

For 0.5 M solution

$R = 280\Omega$

$\kappa = ?$

$\frac{l}{a} = 50 \times 1.4 \times 10^{-2}cm^{-1}$

$\Rightarrow R = \rho \frac{l}{a} = \frac{1}{\kappa} \times \frac{l}{a}$

$\Rightarrow \kappa = \frac{1}{280} \times 50 \times 1.4 \times 10^{-2}$

$= \frac{1}{280} \times 70 \times 10^{-2}$

$= 2.5 \times 10^{-3}Scm^{-1}$

Now,  $\Lambda_m = \frac{\kappa \times 1000}{M}$

$= \frac{2.5 \times 10^{-3}Scm^{-1} \times 1000cm^3 / L}{0.5mol / L}$

$= 5Scm^2mol^{-1} = 5 \times 10^{-4}Sm^2mol^{-1}$

## Question122

The equivalent conductance of NaCl at concentration C and at infinite dilution are  $\lambda_C$  and  $\lambda_\infty$ , respectively. The correct relationship between  $\lambda_C$  and  $\lambda_\infty$  is given as:

(Where the constant B is positive)  
[2014]

Options:

- A.  $\lambda_C = \lambda_\infty + (B)C$
- B.  $\lambda_C = \lambda_\infty - (B)C$
- C.  $\lambda_C = \lambda_\infty - (B)\sqrt{C}$
- D.  $\lambda_C = \lambda_\infty + (B)\sqrt{C}$

Answer: C

Solution:

Solution:

According to Debye Huckle onsager equation,

$$\lambda_C = \lambda_\infty - B\sqrt{C}$$

---

## Question123

The standard electrode potentials ( $E_{M^+/M}^\circ$ ) of four metals A, B, C and D are  $-1.2V$ ,  $0.6V$ ,  $0.85V$  and  $-0.76V$ , respectively. The sequence of deposition of metals on applying potential is:  
[Online April 9, 2014]

Options:

- A. A, C, B, D
- B. B, D, C, A
- C. C, B, D, A
- D. D, A, B, C

Answer: C

Solution:

Solution:

As the value of reduction potential decreases the reducing power increases i.e.

$$C < B < D < A$$

$$(0.85) \cdot (0.6) \cdot (-0.76)(-1.2)$$

---

## Question124

A current of 10.0A flows for 2.00h through an electrolytic cell containing a molten salt of metal X. This results in the decomposition of 0.250 mol of metal X at the cathode. The oxidation state of X in the molten salt is: (F = 96, 500C)  
[Online April 9, 2014]

Options:

- A. 1+
- B. 2+
- C. 3+
- D. 4

Answer: C

Solution:

**Solution:**

According to Faraday's first law of electrolysis

$$W = \frac{E \times i \times t}{96500}$$

Where E = equivalent weight

$$= \frac{\text{mol. mass of metal (M)}}{\text{oxidation state of metal (x)}}$$

Substituting the value in the formula

$$W = \frac{M}{x} \times \frac{i \times t}{96500}$$

$$\text{or } x = \frac{M}{W} \times \frac{i \times t}{96500} = \frac{10 \times 2 \times 60 \times 60}{96500 \times 0.250} = 3$$


$$[ \text{Given : no. of moles} = \frac{W}{M} = 0.250 ]$$

Hence oxidation state of metal is (+3)

-----

## Question125

Given



Considering the electrode potentials, which of the following represents the correct order of reducing power?

[Online April 11, 2014]

Options:

- A.  $\text{F e}^{2+} < \text{Al} < \text{Br}^{-}$
- B.  $\text{Br}^{-} < \text{F e}^{2+} < \text{Al}$
- C.  $\text{Al} < \text{Br}^{-} < \text{F e}^{2+}$

D.  $\text{Al} < \text{Fe}^{2+} < \text{Br}$

**Answer: D**

**Solution:**

**Solution:**

Reducing character decreases down the series. Hence the correct order is

$\text{Al} < \text{Fe}^{2+} < \text{Br}^-$

---

## Question126

**How many electrons would be required to deposit 6.35g of copper at the cathode during the electrolysis of an aqueous solution of copper sulphate? (Atomic mass of copper = 63.5u,  $N_A$  = Avogadro's constant ) [Online April 12, 2014]**

**Options:**

A.  $\frac{N_A}{20}$

B.  $\frac{N_A}{10}$


C.  $\frac{N_A}{5}$

D.  $\frac{N_A}{2}$

**Answer: C**

**Solution:**

**Solution:**



i.e, to deposit 1 mole of Cu at cathode from  $\text{Cu}^{2+}\text{SO}_4^{2-}$  solution = 2 moles of electrons are required

$$\text{i.e, To deposit 6.35g} = \frac{6.35}{63.5} \times 2 = \frac{2}{10} = \frac{1}{5} \text{mol}$$

$$\text{Thus total no. of electrons required} = \frac{N_A}{5}$$

---

## Question127

**Given below are the half-cell reactions:**



**The E for  $3\text{Mn}^{2+} \rightarrow \text{Mn} + 2\text{Mn}^{3+}$  will be: [2014]**

**Options:**

- A.  $-2.69\text{V}$  ; the reaction will not occur
- B.  $-2.69\text{V}$  ; the reaction will occur
- C.  $-0.33\text{V}$  ; the reaction will not occur
- D.  $-0.33\text{V}$  ; the reaction will occur

**Answer: A**

**Solution:**

**Solution:**

(a)  $\text{Mn}^{2+} + 2\text{e}^- \rightarrow \text{Mn}$ ;  $E^\circ = -1.18\text{V}$  ;...(i)

(b)  $\text{Mn}^{3+} + \text{e}^- \rightarrow \text{Mn}^{2+}$ ;  $E^\circ = -1.51\text{V}$  ; ... (ii)

Now multiplying equation (ii) by two and subtracting from equation (i)

$3\text{Mn}^{2+} \rightarrow \text{Mn} + 2\text{Mn}^{3+}$

$E^\circ = E_{\text{Ox.}} + E_{\text{Red}}$

$= -1.18 + (-1.51) = -2.69\text{V}$

(-ve value of EMF (i.e.  $\Delta G = +ve$ ) shows that the reaction is non-spontaneous)

## Question128

**Electrode potentials ( $E^\circ$ ) are given below:**

**$\text{Cu}^+ / \text{Cu} = +0.52\text{V}$**

**$\text{Fe}^{3+} / \text{Fe}^{2+} = +0.77\text{V}$**

**$\frac{1}{2}\text{I}_2(\text{s}) / \text{I}^- = +0.54\text{V}$**

**$\text{Ag}^+ / \text{Ag} = +0.88\text{V}$**

**Based on the above potentials, strongest oxidizing agent will be:**

**[Online April 9, 2013]**

**Options:**

- A.  $\text{Cu}^+$
- B.  $\text{Fe}^{3+}$
- C.  $\text{Ag}^+$
- D.  $\text{I}_2$

**Answer: C**

**Solution:**

**Solution:**

Higher the value of reduction potential stronger will be the oxidising agent, hence based on the given values  $\text{Ag}^+$  will be strongest oxidizing agent.

## Question129

**Given:**

$$E_{\frac{1}{2}\text{Cl}_2 / \text{Cl}^-} = 1.36\text{V}, E_{\text{Cr}^{3+} / \text{Cr}}^\circ = -0.74\text{V}$$

$$E^\circ_{\text{Cr}_2\text{O}_7^{2-} / \text{Cr}^{3+}} = 1.33\text{V}, E^\circ_{\text{MnO}_4^- / \text{Mn}^{2+}} = 1.51\text{V}$$

The correct order of reducing power of the species( $\text{Cr}$ ,  $\text{Cr}^{3+}$ ,  $\text{Mn}^{2+}$  and  $\text{Cl}^-$ ) will be:

[Online April 23, 2013]

Options:

A.  $\text{Mn}^{2+} < \text{Cl}^- < \text{Cr}^{3+} < \text{Cr}$

B.  $\text{Mn}^{2+} < \text{Cl}^{3+} < \text{Cl}^- < \text{Cr}$

C.  $\text{Cr}^{3+} < \text{Cl}^- < \text{Mn}^{2+} < \text{Cr}$

D.  $\text{Cr}^{3+} < \text{Cl}^- < \text{Cr} < \text{Mn}^{2+}$

Answer: A

Solution:

Solution:

Lower the value of reduction potential higher will be reducing power hence the correct order will be  $\text{Mn}^{2+} < \text{Cl}^- < \text{Cr}^{3+} < \text{Cr}$

## Question130

A solution of copper sulphate ( $\text{CuSO}_4$ ) is electrolysed for 10 minutes with a current of 1.5 amperes. The mass of copper deposited at the cathode (at. mass of  $\text{Cu} = 63\text{u}$ ) is:

[Online April 25, 2013]

Options:

A. 0.3892g

B. 0.2938g

C. 0.2398g

D. 0.3928g

Answer: B

Solution:

Solution:

$$W = Z \text{ it}$$

where  $Z$  = Electrochemical equivalent

$$\text{Eq. wt. of copper} = \frac{63}{2} = 31.5$$

$$Z = \frac{31.5}{96500}$$

$$W = Z \text{ it} = \frac{31.5}{96500} \times 1.5 \times 10 \times 60 = 0.2938\text{g}$$

## Question131

**Given:**

$$E_{\text{Cr}^{3+} / \text{Cr}}^{\circ} = -0.74\text{V}; E_{\text{MnO}_4^{-} / \text{Mn}^{2+}}^{\circ} = 1.51\text{V}$$

$$E_{\text{Cr}_2\text{O}_7^{2-} / \text{Cr}^{3+}} = 1.33\text{V}; E_{\text{Cl} / \text{Cl}^{-}}^{\circ} = 1.36\text{V}$$

**Based on the data given above, strongest oxidising agent will be:  
[2013]**

**Options:**

A. Cl

B.  $\text{Cr}^{3+}$

C.  $\text{Mn}^{2+}$

D.  $\text{MnO}_4^{-}$

**Answer: D**

**Solution:**

**Solution:**

Higher the value of standard reduction potential, stronger is the oxidising agent, hence  $\text{MnO}_4^{-}$  is the strongest oxidising agent.

-----

## Question132

**The standard reduction potentials for  $\text{Zn}^{2+} / \text{Zn}$ ,  $\text{Ni}^{2+} / \text{Ni}$  and  $\text{Fe}^{2+} / \text{Fe}$  are  $-0.76$ ,  $-0.23$  and  $-0.44\text{V}$  respectively.**

**The reaction  $\text{X} + \text{Y}^{2+} \rightarrow \text{X}^{2+} + \text{Y}$  will be spontaneous when :  
[2012]**

**Options:**

A.  $\text{X} = \text{Ni}$ ,  $\text{Y} = \text{Fe}$

B.  $\text{X} = \text{Ni}$ ,  $\text{Y} = \text{Zn}$

C.  $\text{X} = \text{Fe}$ ,  $\text{Y} = \text{Zn}$

D.  $\text{X} = \text{Zn}$ ,  $\text{Y} = \text{Ni}$

**Answer: D**


**Solution:**

**Solution:**

For a spontaneous reaction  $\Delta G^{\circ}$  must be -ve

Since  $\Delta G^{\circ} = -nF E^{\circ}$

Hence for  $\Delta G^\circ$  to be  $-ve$ ,  $\Delta E^\circ$  has to be positive. Which is possible when  $X = Zn$ ,  $Y = Ni$




$$E_{\text{cell}}^\circ = E_{Ni^{2+}/Ni}^\circ - E_{Zn^{2+}/Zn}^\circ \\ = -0.23 - (-0.76) = +0.53 \text{ (positive)}$$

---

## Question133

A battery is constructed of Cr and  $Na_2Cr_2O_7$ . The unbalanced chemical equation when such a battery discharges is following:



If one Faraday of electricity is passed through the battery during the charging, the number of moles of  $Cr^{3+}$  removed from the solution is [Online May 7, 2012]

Options:

A.  $\frac{4}{3}$

B.  $\frac{1}{3}$

C.  $\frac{3}{3}$


D.  $\frac{2}{3}$

**Answer: C**


**Solution:**

**Solution:**


Reduction half reaction:



Oxidation half reaction:



Overall reaction :



$$3F \text{ of electricity} = 3 \text{ moles of } Cr^{3+}$$

$$1F \text{ of electricity} = \frac{3}{3} \text{ moles of } Cr^{3+}$$

---

## Question134

The standard potentials of  $Ag^+ / Ag$ ,  $Hg_2^{2+} / 2Hg$ ,  $Cu^{2+} / Cu$  and  $Mg^{2+} / Mg$  electrodes are 0.80, 0.79, 0.34 and  $-2.37V$ , respectively. An aqueous solution which contains one mole per litre of the salts of each of the four metals is electrolyzed. With increasing voltage, the correct sequence of deposition of the metals at the cathode is [Online May 19, 2012]

Options:



- A. Ag, H g, Cu, M g
- B. Cu, H g, Ag only
- C. Ag, Hg, Cu only
- D. M g, Cu, H g, Ag

**Answer: C**

**Solution:**

**Solution:**

More the reduction potential, more is the deposition of metals at cathode. Cation having  $E^\circ$  value less than  $-0.83V$  (reduction potential of  $H_2O$ ) will not deposit from aqueous solution. Hence correct order of deposition of the metal at the cathode is

Ag > H g > Cu

## Question135

**Given**

$$E_{Cu^{2+}/Cu}^\circ = 0.34V, E_{Cu^{2+}/Cu^+}^\circ = 0.15V$$



**Standard electrode potential for the half cell  $Cu^+ / Cu$  is [Online May 26,2012]**

**Options:**


- A. 0.38V
- B. 0.53V
- C. 0.19V
- D. 0.49V

**Answer: B**

**Solution:**



On subtracting eq.(i) from eq. (ii) we get



$$-nE^\circ F = -(n_2 E_2^\circ F - n_1 E_1^\circ F)$$

$$E^\circ = \frac{n_2 E_2^\circ F - n_1 E_1^\circ F}{nF}$$

$$= \frac{2 \times 0.34 - 0.15}{1} = 0.53V$$

## Question136

**Resistance of 0.2M solution of an electrolyte is  $50\Omega$ . The specific**

**conductance of the solution is  $1.3\text{Sm}^{-1}$ . If resistance of the  $0.4\text{M}$  solution of the same electrolyte is  $260\ \Omega$ , its molar conductivity is : [2011RS]**

**Options:**

A.  $6.25 \times 10^{-4}\text{Sm}^2\text{mol}^{-1}$

B.  $625 \times 10^{-4}\text{Sm}^2\text{mol}^{-1}$

C.  $62.5\text{Sm}^2\text{mol}^{-1}$

D.  $6250\text{Sm}^2\text{mol}$

**Answer: A**

**Solution:**

$$k = \frac{1}{R} \times \frac{1}{a}$$

$$1.3 = \frac{1}{50} \times \frac{1}{a}$$

$$\frac{1}{a} = 65\text{m}^{-1}$$

$$\Lambda_m = \frac{k \times 1000}{\text{molarity}}$$

[molarity is in moles/litre but 1000 is used to convert litre into  $\text{cm}^3$ ]

$$= \frac{\left( \frac{1}{260} \times 65\text{m}^{-1} \right) \times 1000\text{cm}^3}{0.4\text{mol}}$$


$$= \frac{65\text{m}^{-1}}{260 \times 0.4\text{mol}} \times \frac{1}{1000}\text{m}^3$$

$$= 6.25 \times 10^{-4}\text{Sm}^2\text{mol}^{-1}$$

---

## Question137

**The Gibbs energy for the decomposition of  $\text{Al}_2\text{O}_3$  at  $500^\circ\text{C}$  is as follows:**



**The potential difference needed for electrolytic reduction of  $\text{Al}_2\text{O}_3$  at  $500^\circ\text{C}$  is at least [2010]**

**Options:**

A.  $4.5\text{V}$


B.  $3.0\text{V}$

C.  $2.5\text{V}$



D.  $5.0\text{V}$

**Answer: C**

**Solution:**



The ionic reactions are



Thus, no. of electron transferred  $\Rightarrow n = 4$

$$\Delta G = -nF E$$

$$\text{or } E = \frac{\Delta G}{-nF} = \frac{966 \times 10^3}{4 \times 96500} = -2.5\text{V}$$

$\therefore$  The potential difference needed for the reduction = 2.5V .

---


## Question138

**Given:**

$$E_{\text{Fe}^{3+}/\text{Fe}}^\circ = -0.036\text{V}$$

$$E_{\text{Fe}^{2+}/\text{Fe}}^\circ = -0.439\text{V}$$

**The value of standard electrode potential for the change,**



**[2009]**

**Options:**

A. 0.385V

B. 0.770V


C. -0.270V

D. -0.072V


**Answer: B**

**Solution:**

Given




$$E_{\text{Fe}^{3+}/\text{Fe}}^\circ = -0.036\text{V} \dots(\text{i})$$




$$E_{\text{Fe}^{2+}/\text{Fe}}^\circ = -0.439\text{V} \dots(\text{ii})$$

we have to calculate



To obtain this equationsubtract equ. (ii) from (i) we get



$$\text{As we know that } \Delta G^\circ = -nF E^\circ$$

Thus for reaction (iii)

$$\Delta G_3 = \Delta G_1 - \Delta G_2$$

$$-nF E_3^\circ = -nF E_1^\circ - (-nF E_2^\circ)$$

$$-nF E_3^\circ = nF E_2^\circ - nF E_1^\circ$$

$$-1F E_3^\circ = 2 \times 0.439F - 3 \times 0.036F$$


$$-1F E_3^\circ = 0.770F$$

$$\therefore E_3^\circ = -0.770\text{V}$$


---

## Question139

In a fuel cell methanol is used as fuel and oxygen gas is used as an oxidizer. The reaction is




At 298K standard Gibb's energies of formation for  $\text{CH}_3\text{OH}(\text{l})$ ,  $\text{H}_2\text{O}(\text{l})$  and  $\text{CO}_2(\text{g})$  are  $-166.2$ ,  $-237.2$  and  $-394.4\text{kJ mol}^{-1}$  respectively. If standard enthalpy of combustion of methanol is  $-726\text{kJ mol}^{-1}$ , efficiency of the fuel cell will be:  
[2009]

Options:

- A. 87%
- B. 90%
- C. 97%
- D. 80%

Answer: C

Solution:



$$\Delta G_r = \Delta G_f(\text{CO}_2, \text{g}) + 2 \Delta G_f(\text{H}_2\text{O}, \text{l}) -$$

$$\Delta G_f(\text{CH}_3\text{OH}, \text{l}) - \frac{3}{2} \Delta G_f(\text{O}_2, \text{g})$$

$$= -394.4 + 2(-237.2) - (-166.2) - 0$$

$$= -394.4 - 474.4 + 166.2 = -702.6\text{kJ}$$

$$\% \text{ efficiency} = \frac{702.6}{726} \times 100 = 97\%$$

---

## Question140

Given  $E^\circ \text{Cr}^{3+} / \text{Cr} = -0.72\text{V}$

$E^\circ \text{Fe}^{2+} / \text{Fe} = -0.42\text{V}$ .

The potential for the cell

$\text{Cr} | \text{Cr}^{3+}(0.1\text{M}) | \text{Fe}^{2+}(0.01\text{M}) | \text{Fe}$  is  
[2008]

Options:

- A. 0.26V
- B. 0.336V
- C.  $-0.339$
- D. 0.26V

**Answer: D**

**Solution:**

**Solution:**

From the given representation of the cell,  $E_{\text{cell}}$  can be found as follows.

$$E_{\text{cell}} = E_{\text{Fe}^{2+}/\text{Fe}}^{\circ} - E_{\text{Cr}^{3+}/\text{Cr}}^{\circ} - \frac{0.059}{6} \log \frac{[\text{Cr}^{3+}]^2}{[\text{Fe}^{2+}]^3}$$

[Nernst -Equ.]

$$= -0.42 - (-0.72) - \frac{0.059}{6} \log \frac{(0.1)^2}{(0.01)^3}$$

$$= -0.42 + 0.72 - \frac{0.059}{6} \log \frac{0.1 \times 0.1}{0.01 \times 0.01 \times 0.01}$$

$$= 0.3 - \frac{0.059}{6} \log \frac{10^{-2}}{10^{-6}} = 0.3 - \frac{0.059}{6} \times 4$$

$$= 0.30 - 0.0393 = 0.26\text{V}$$

## Question141

The equivalent conductances of two strong electrolytes at infinite dilution in  $\text{H}_2\text{O}$  (where ions move freely through a solution) at  $25^\circ\text{C}$  are given below:

$$\Lambda_{\text{CH}_3\text{COONa}} = 91.0 \text{Scm}^2 / \text{equiv.}$$

$$\Lambda_{\text{HCl}}^{\circ} = 426.2 \text{Scm}^2 / \text{equiv.}$$

What additional information/ quantity one needs to calculate  $\Lambda^{\circ}$  of an aqueous solution of acetic acid?

[2007]

**Options:**

A.  $\Lambda$  of chloroacetic acid ( $\text{ClCH}_2\text{COOH}$ )

B.  $\Lambda$  of  $\text{NaCl}$

C.  $\Lambda$  of  $\text{CH}_3\text{COOK}$

D. the limiting equivalent conductance of  $\text{H}^+$  ( $\lambda_{\text{H}^+}^{\circ}$ ).

**Answer: B**

**Solution:**

**Solution:**

**Note:** According to Kohlrausch's law, equivalent conductivity of weak electrolyte, acetic acid ( $\text{CH}_3\text{COOH}$ ) can be calculated as follows:

$$\Lambda_{\text{CH}_3\text{COOH}}^{\circ} = (\Lambda_{\text{CH}_3\text{COONa}}^{\circ} + \Lambda_{\text{HCl}}^{\circ}) - \Lambda_{\text{NaCl}}^{\circ}$$

$\therefore$  Value of  $\Lambda_{\text{NaCl}}^{\circ}$  should also be known for calculating value of  $\Lambda_{\text{CH}_3\text{COOH}}^{\circ}$

## Question142

The cell,  $\text{Zn} | \text{Zn}^{2+}(1\text{M}) | \text{Cu}^{2+}(1\text{M}) | \text{Cu}$  ( $E_{\text{cell}}^{\circ} = 1.10\text{V}$ ) was allowed to be completely discharged at  $298\text{K}$ .

The relative concentration of  $\text{Zn}^{2+}$  to  $\text{Cu}^{2+}$   $\left( \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]} \right)$  is

[2007]

Options:

A.  $9.65 \times 10^4$

B.  $\text{antilog}(24.08)$

C. 37.3

D.  $10^{373}$ .

Answer: D

Solution:

Solution:

$E_{\text{cell}} = 0$ ; when cell is completely discharged.

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{2} \log \left( \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]} \right)$$

$$\text{or } 0 = 1.1 - \frac{0.059}{2} \log \left( \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]} \right)$$

$$\log \left( \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]} \right) = \frac{2 \times 1.1}{0.059} = 37.3$$

$$\therefore \left( \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]} \right) = 10^{37.3}$$

## Question143

The molar conductivities  $\Lambda_{\text{NaOAc}}^{\circ}$  and  $\Lambda_{\text{HCl}}^{\circ}$  at infinite dilution in water at  $25^{\circ}\text{C}$  are  $91.0$  and  $426.2\text{Scm}^2/\text{mol}$  respectively. To calculate  $\Lambda_{\text{H OAc}}^{\circ}$ , the additional value required is

[2006]

Options:

A.  $\Lambda_{\text{NaOH}}^{\circ}$

B.  $\Lambda_{\text{NaCl}}^{\circ}$

C.  $\Lambda_{\text{H}_2\text{O}}^{\circ}$

D.  $\Lambda_{\text{KCl}}^{\circ}$

Answer: B

Solution:

$\Lambda_{\text{CH}_3\text{COOH}}^\circ$  is given by the following equation

$$\Lambda_{\text{CH}_3\text{COOH}}^\circ = (\Lambda_{\text{CH}_3\text{COONa}}^\circ + \Lambda_{\text{HCl}}^\circ) - (\Lambda_{\text{NaCl}}^\circ)$$

Hence  $\Lambda_{\text{NaCl}}^\circ$  is required.

---

## Question 144

**Resistance of a conductivity cell filled with a solution of an electrolyte of concentration 0.1M is  $100\Omega$ . The conductivity of this solution is  $1.29\text{Sm}^{-1}$ . Resistance of the same cell when filled with 0.2M of the same solution is  $520\Omega$ . The molar conductivity of 0.2M solution of electrolyte will be [2006]**

**Options:**

- A.  $1.24 \times 10^{-4}\text{Sm}^2\text{mol}^{-1}$
- B.  $12.4 \times 10^{-4}\text{Sm}^2\text{mol}^{-1}$
- C.  $124 \times 10^{-4}\text{Sm}^2\text{mol}^{-1}$
- D.  $1240 \times 10^{-4}\text{Sm}^2\text{mol}^{-1}$

**Answer: B**

**Solution:**

$$R = 100\Omega, K = \frac{1}{R} \left( \frac{l}{a} \right)$$

$$\frac{1}{a} \left( \text{cell constant} \right) = 1.29 \times 100\text{m}^{-1}$$

Given,  $R = 520\Omega$

$$C = 0.2\text{M} = 0.2 \times 1000\text{mol m}^{-3}$$

$\Lambda_m$  (molar conductivity) = ?

$\Lambda_m = K \times V$  (K can be calculated as cell constant is known. )

$$K = \frac{1}{R} \left( \frac{l}{a} \right)$$

Hence,

$$K = \frac{129\text{m}^{-1}}{520\Omega} = 0.248\text{Sm}^{-1}$$

$$\begin{aligned}\Lambda_m &= \frac{K}{C} = \frac{0.248\text{Sm}^{-1}}{0.2 \times 1000\text{mol m}^{-3}} \\ &= 12.4 \times 10^{-4}\text{Sm}^2\text{mol}^{-1}\end{aligned}$$

---

## Question 145

**The highest electrical conductivity of the following aqueous solutions is of [2005]**

**Options:**

- A. 0.1M difluoroacetic acid
- B. 0.1M fluoroacetic acid
- C. 0.1M chloroacetic acid
- D. 0.1M acetic acid

**Answer: A**

**Solution:**

**Solution:**

Difluoro acetic acid being strongest acid will furnish maximum number of ions showing highest electrical conductivity. The decreasing acidic strength of the carboxylic acids given is difluoro acetic acid > fluoro acetic acid > chloro acetic acid > acetic acid.

## Question146

Electrolyte:	$KCl$	$KNO_3$	$HCl$	$NaOAc$	$NaCl$
$\Lambda^\infty (S\,cm^2\,mol^{-1})$ :	149.9	145	426.2	91	126.5

**Calculate  $\Lambda_{HOAc}^\infty$  using appropriate molar conductance's of the electrolytes listed above at infinite dilution in  $H_2O$  at  $25^\circ C$  [2005]**

**Options:**

- A. 217.5
- B. 390.7
- C. 552.7
- D. 517.2

**Answer: B**

**Solution:**

**Solution:**

$$\Lambda_{HCl}^\infty = 426.2 \dots (i)$$

$$\Lambda_{ACONa}^\infty = 91.0 \dots (ii)$$

$$\Lambda_{NaCl}^\infty = 126.5 \dots (iii)$$

$$\Lambda_{AcOH}^\infty = (i) + (ii) - (iii)$$

$$= [426.2 + 91.0 - 126.5] = 390.7$$

## Question147

**Aluminium oxide may be electrolysed at  $1000^\circ C$  to furnish aluminium**



**metal (At. Mass = 27 amu; 1 Faraday = 96,500 Coulombs). The cathode reaction is-  $\text{Al}^{3+} + 3\text{e}^- \rightarrow \text{Al}^\circ$  To prepare 5.12kg of aluminium metal by this method we require [2005]**

**Options:**

- A.  $5.49 \times 10^1 \text{C}$  of electricity
- B.  $5.49 \times 10^4 \text{C}$  of electricity
- C.  $1.83 \times 10^7 \text{C}$  of electricity
- D.  $5.49 \times 10^7 \text{C}$  of electricity

**Answer: D**

**Solution:**

**Solution:**

1 mole of  $\text{e}^- = 1\text{F} = 96500\text{C}$

27g of Al is deposited by  $3 \times 96500\text{C}$

5120g of Al will be deposited by

$$= \frac{3 \times 96500 \times 5120}{27} = 5.49 \times 10^7 \text{C}$$

We know,  $Q = \frac{mFz}{M}$

$$\therefore Q = \frac{5120 \times 96500 \times 3}{27} = 5.49 \times 10^7 \text{C}$$

## Question 148

**For a spontaneous reaction the  $\Delta G$ , equilibrium constant ( $K$ ) and  $E_{\text{cell}}^\circ$  will be respectively [2005]**

**Options:**

- A.  $-ve, >1, -ve$
- B.  $-ve, <1, -ve$
- C.  $+ve, >1, -ve$
- D.  $-ve, >1, +ve$

**Answer: D**

**Solution:**

**Solution:**

**Note:** For spontaneous reaction  $\Delta G$  should be negative. Equilibrium constant should be more than one

( $\Delta G = -2.303RT \log K_c$ , If  $K_c = 1$  then

$\Delta G = 0$ ; If  $K_c > 1$  then  $\Delta G = -ve$ ).

Again  $\Delta G = -nF E_{\text{cell}}^\circ$

$E_{\text{cell}}^\circ$  must be  $+ve$  to have  $\Delta G -ve$ .

## Question149

The limiting molar conductivities  $\Lambda^\circ$  for NaCl, KBr and KCl are 126, 152 and 150  $\text{Scm}^2\text{mol}^{-1}$  respectively. The  $\Lambda$  for NaBr is [2004]

Options:

- A. 278  $\text{Scm}^2\text{mol}^{-1}$
- B. 176  $\text{Scm}^2\text{mol}^{-1}$
- C. 128  $\text{Scm}^2\text{mol}^{-1}$
- D. 302  $\text{Scm}^2\text{mol}^{-1}$

Answer: C

Solution:

Solution:

$$\Lambda^\circ_{\text{NaCl}} = \lambda^\circ_{\text{Na}^+} + \lambda^\circ_{\text{Cl}^-} \dots (i)$$

$$\Lambda^\circ_{\text{KBr}} = \lambda^\circ_{\text{K}^+} + \lambda^\circ_{\text{Br}^-} \dots (ii)$$

$$\Lambda^\circ_{\text{KCl}} = \lambda^\circ_{\text{K}^+} + \lambda^\circ_{\text{Cl}^-} \dots (iii)$$

operating (i) + (ii) - (iii)

$$\begin{aligned}\Lambda^\circ_{\text{NaBr}} &= \lambda^\circ_{\text{Na}^+} + \lambda^\circ_{\text{Br}^-} \\ &= 126 + 152 - 150 = 128 \text{Scm}^2\text{mol}^{-1}\end{aligned}$$

---

## Question150

Consider the following

$E^\circ$  values  $E_{\text{Fe}^{3+}/\text{Fe}^{2+}} = +0.77\text{V}$ ;  $E_{\text{Sn}^{2+}/\text{Sn}} = -0.14\text{V}$

Under standard conditions the potential for the reaction

$\text{Sn(s)} + 2\text{Fe}^{3+}(\text{aq}) \rightarrow 2\text{Fe}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq})$  is

[2004]

Options:

- A. 0.91V
- B. 1.40V
- C. 1.68V
- D. 0.63V

Answer: A

Solution:

$\text{Fe}^{3+} + \text{e}^{-} \rightarrow \text{Fe}^{2+}$ ;  $\text{Sn}^{2+} + 2\text{e}^{-} \rightarrow \text{Sn(s)}$ ;  
 for  $\text{Sn(s)} + 2\text{Fe}^{3+}(\text{aq}) \rightarrow 2\text{Fe}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq})$   
 $\therefore$  Standard potential for the given reaction or  
 $E_{\text{cell}}^{\circ} = E_{\text{Sn} / \text{Sn}^{2+}}^{\circ} + E_{\text{Fe}^{3+} / \text{Fe}^{2+}}$   
 $= 0.14 + 0.77 = 0.91\text{V}$

---

## Question151

The  $E^{\circ} \text{M}^{3+} / \text{M}^{2+}$  values for Cr, Mn, Fe and Co are  $-0.41$ ,  $+1.57$ ,  $+0.77$  and  $+1.97\text{V}$  respectively. For which one of these metals the change in oxidation state from +2 to +3 is easiest?  
 [2004]

**Options:**

- A. Fe
- B. Mn
- C. Cr
- D. Co

**Answer: C**

**Solution:**

**Solution:**

The given values show that Cr has maximum oxidation potential, therefore its oxidation will be easiest. (Change the sign to get the oxidation values)

---

## Question152

The standard e.m.f. of a cell involving one electron change is found to be  $0.591\text{V}$  at  $25^{\circ}\text{C}$ . The equilibrium constant of the reaction is  
 ( $F = 96,500\text{Cmol}^{-1}$ ;  $R = 8.314\text{J K}^{-1}\text{mol}^{-1}$ )  
 [2004]

**Options:**

- A.  $1.0 \times 10^{10}$
- B.  $1.0 \times 10^5$
- C.  $1.0 \times 10^1$
- D.  $1.0 \times 10^{30}$

**Answer: A**

**Solution:**

$$(a) E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{n} \log K_c$$

$$\text{or } 0 = 0.591 - \frac{0.0591}{1} \log K_c$$

$$\text{or } \log K_c = \frac{0.591}{0.0591} = 10 \text{ or } K_c = 1 \times 10^{10}$$


---

## Question153

**In a cell that utilises the reaction**  
 $\text{Zn(s)} + 2\text{H}^+(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + \text{H}_2(\text{g})$  **addition of  $\text{H}_2\text{SO}_4$  to cathode compartment, will**  
**[2004]**

**Options:**

- A. increase the  $E$  and shift equilibrium to the right
- B. lower the  $E$  and shift equilibrium to the right
- C. lower the  $E$  and shift equilibrium to the left
- D. increase the  $E$  and shift equilibrium to the left

**Answer: A**

**Solution:**

**Solution:**  
 $\text{Zn(s)} + 2\text{H}^+(\text{aq}) \rightleftharpoons \text{Zn}^{2+}(\text{aq}) + \text{H}_2(\text{g})$   

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{2} \log \frac{[\text{Zn}^{2+}][\text{H}_2]}{[\text{H}^+]^2}$$
 Addition of  $\text{H}_2\text{SO}_4$  will increase  $[\text{H}^+]$  and  $E_{\text{cell}}$  will also increase and thus the equilibrium shifts towards RHS.

---

## Question154

**In a hydrogen-oxygen fuel cell, combustion of hydrogen occurs to**  
**[2004]**

**Options:**

- A. produce high purity water
- B. create potential difference between two electrodes
- C. generate heat
- D. remove adsorbed oxygen from electrode surfaces

**Answer: B**

**Solution:**

In  $\text{H}_2 - \text{O}_2$  fuel cell, the combustion of  $\text{H}_2$  occurs to create potential difference between the two electrodes

---

## Question 155

Standard reduction electrode potentials of three metals A, B & C are respectively  $+0.5\text{V}$ ,  $-3.0\text{V}$  &  $-1.2\text{V}$ .

The reducing powers of these metals are  
[2003]

Options:

- A.  $A > B > C$
- B.  $C > B > A$
- C.  $A > C > B$
- D.  $B > C > A$

**Answer: D**

**Solution:**

**Solution:**

**Note:** The higher the negative value of reduction potential, the more is the reducing power.

Hence  $B > C > A$ .

---

## Question 156

When during electrolysis of a solution of  $\text{AgNO}_3$  9650 coulombs of charge pass through the electroplating bath, the mass of silver deposited on the cathode will be  
[2003]

Options:

- A. 10.8g
- B. 21.6g
- C. 108g
- D. 1.08g

**Answer: A**

**Solution:**

**Solution:**

When 96500 coulomb of electricity is passed through the electroplating bath the amount of Ag deposited = 108g  
 $\therefore$  when 9650 coulomb of electricity is passed deposited Ag.

$$= \frac{108}{96500} \times 9650 = 10.8\text{g}$$


---

## Question157

For a cell reaction involving a two-electron change, the standard e.m.f. of the cell is found to be 0.295V at 25°C. The equilibrium constant of the reaction at 25°C will be [2003]

Options:

- A.  $29.5 \times 10^{-2}$
- B. 10
- C.  $1 \times 10^{10}$
- D.  $1 \times 10^{-10}$

Answer: C

Solution:

**Solution:**

The equilibrium constant is related to the standard emf of cell by the expression


$$\log K = \frac{n}{0.059} \times E_{\text{cell}}^{\circ} = 0.295 \times \frac{2}{0.059}$$

$$\log K = \frac{590}{59} = 10 \text{ or } K = 1 \times 10^{10}$$


---

## Question158

For the redox reaction : [2003]



taking place in a cell,  $E_{\text{cell}}^{\circ}$  is 1.10 volt.  $E_{\text{cell}}$  for the cell will be

$$\left( 2.303 \frac{RT}{F} = 0.0591 \right)$$

Options:

- A. 1.80 volt
- B. 1.07 volt
- C. 0.82 volt
- D. 2.14 volt

Answer: B

## Solution:

### Solution:

$$\begin{aligned}E_{\text{cell}} &= E_{\text{cell}}^{\circ} + \frac{0.059}{n} \log \frac{[\text{Cu}^{2+}]}{[\text{Zn}^{2+}]} \\&= 1.10 + \frac{0.059}{2} \log[0.1] \\&= 1.10 - 0.0295 = 1.07\text{V}\end{aligned}$$

---

## Question159

**Conductivity (unit Siemen's S ) is directly proportional to area of the vessel and the concentration of the solution in it and is inversely proportional to the length of the vessel then the unit of the constant of proportionality is [2002]**

### Options:

- A.  $\text{Smmol}^{-1}$
- B.  $\text{Sm}^2\text{mol}^{-1}$
- C.  $\text{S}^{-2}\text{m}^2\text{mol}$
- D.  $\text{S}^2\text{m}^2\text{mol}^{-2}$ .

**Answer: B**

## Solution:

### Solution:

$$\begin{aligned}\text{Given conductivity} &\propto \frac{\text{area} \times \text{conc.}}{\text{length}} \\&= \frac{\text{K m}^2\text{mol}}{\text{m} \times \text{m}^3} \\\therefore \text{K} &= \text{Sm}^2\text{mol}^{-1}\end{aligned}$$

---

## Question160

**Which of the following reaction is possible at anode? [2002]**


### Options:

- A.  $2\text{Cr}^{3+} + 7\text{H}_2\text{O} \rightarrow \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+$  [2002]
- B.  $\text{F}_2 \rightarrow 2\text{F}$
- C.  $(1/2)\text{O}_2 + 2\text{H}^+ \rightarrow \text{H}_2\text{O}$
- D. none of these.

**Answer: A**

**Solution:**

**Solution:**



O.S. of Cr changes from +3 to +6 by loss of electrons. At anode oxidation takes place.

---

## Question161

**When the sample of copper with zinc impurity is to be purified by electrolysis, the appropriate electrodes are Cathode    Anode [2002]**

**Options:**

A. Cathode - pure zinc  
Anode - pure copper

B. Cathode - impure sample  
Anode - pure copper

C. Cathode - impure zinc  
Anode - impure sample

D. Cathode - pure copper  
Anode - impure sample.

**Answer: D**

**Solution:**

**Solution:**

Pure metal always deposits at cathode.

---

## Question162

**EMF of a cell in terms of reduction potential of its left and right electrodes is [2002]**

**Options:**

A.  $E = E_{\text{left}} - E_{\text{right}}$

B.  $E = E_{\text{left}} + E_{\text{right}}$

C.  $E = E_{\text{right}} - E_{\text{left}}$

D.  $E = -(E_{\text{righ}} + E_{\text{lef}})$ .

**Answer: C**



## Solution:


### Solution:

$$\begin{aligned} E_{\text{cell}} &= \text{Reduction potential of cathode (right)} \\ &\quad - \text{reduction potential of anode (left)} \\ &= E_{\text{right}} - E_{\text{left}} \end{aligned}$$

---

## Question 163

What will be the emf for the given cell



**Options:**

A.  $\frac{RT}{F} \log_e \frac{P_1}{P_2}$

B.  $\frac{RT}{2F} \log_e \frac{P_2}{P_1}$


C.  $\frac{RT}{F} \log_e \frac{P_2}{P_1}$

D. none of these.


**Answer: C**

### Solution:


Oxidation half cell:-



P<sub>1</sub> Reduction half cell



P<sub>2</sub> The net cell reaction



P<sub>t</sub>

$$E_{\text{cell}}^\circ = 0.00\text{V}; n = 2$$

$$\therefore E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{nF} \log_e K$$

$$= 0 - \frac{RT}{nF} \log_e \frac{P_2}{P_1}$$

$$\text{or } E_{\text{cell}} = \frac{RT}{2F} \log_e \frac{P_1}{P_2}$$

---