
Chapter 6. Triangle

Question-1

The bisector of the exterior angle $\angle A$ of \triangle ABC intersects side BC produced at D. Prove that $\frac{AB}{AC} = \frac{BD}{DC}$.



Solution:

Given: ABC is a triangle; AD is the exterior bisector of $\angle A$ and meets BC produced at D; BA is produced to F.

To prove: $\frac{AB}{AC} = \frac{BD}{DC}$

Construction: Draw CE||DA to meet AB at E.

Proof: In \triangle ABC, CE || AD cut by AC. \angle CAD = \angle ACE (Alternate angles)

Similarly CE || AD cut by AB

∠FAD = ∠AEC (corresponding angles)

Since ∠FAD = ∠CAD (given) ∴ ∠ACE = ∠AEC

 $\ AC = AE (by isosceles \Delta theorem)$


Now in \triangle BAD, CE || DA <u>AE</u> = <u>DC</u>(BPT) AB BD

But AC = AE (proved above)

\ <u>AC</u> = <u>DC</u> AB BD or

$$\frac{AB}{AC} = \frac{BD}{DC}$$
 (proved).

State and prove the converse of angle bisector theorem.



Solution:

Given: ABC is a Δ ; AD divides BC in the ratio of the sides containing the angles $\angle A$ to meet BC at D.

i.e.
$$\frac{AB}{AC} = \frac{BD}{DC}$$

To prove: AD bisects $\angle A$.

Construction: Draw CE || DA to meet BA produced at E.

Proof: In Δ ABC, CE || DA cut by AE.

\ ∠BAD = ∠AEC (corresponding angle) ----(i)

Similarly CE || DA cut by AC

\ ∠ DAC = ∠ACE (alternate angles) ----(ii)

In DBEC; CE | AD

 $\ \ AB = BD (BPT)$

AE DC

But AB = BD (given)

AC DC

 \land AB = AB

AE AC

 $\AE = AC$

⇒ ∠AEC = ∠ACE (isosceles property) ----(iii)


According to equation (i), (ii) and (iii) $\angle BAD = \angle DAC$

 \Rightarrow AD bisects $\angle A$.

Question-3

If a parallelogram has all its sides equal and one of its diagonal is equal to a side, show that its diagonals are in the ratio $\sqrt{3}$: 1.

Solution:



Given: ABCD is a parallelogram, where AC and BD are the diagonals

meeting at O. AB = BC = AC.

To Prove: BD : AC :: √3 : 1

Proof: In \triangle ABC, AB = BC = CA (given).

= a (say)

Hence ABC is an equilateral triangle. (Definition of equilateral triangle)

AC and BD are the diagonals of parallelogram ABCD,

⇒ AC = BD (Diagonals of a parallelogram bisect each other)

or AO = OC.

i.e BO is the median of the equilateral ABC.

Hence BO = $\frac{\sqrt{3}}{3}$ a


∴ BD = √3 a

⇒ BD : AC :: √3 a : a

⇒ BD : AC :: √3 : 1.

Question-4

In \triangle ABC, P, Q are points on AB and AC respectively and PQ || BC. Prove that the median AD bisects PQ.



Solution:

Given: ABC is a triangle, PQ || BC; AD is the median which cuts PQ at R.

To prove: AD bisects PQ at R.

Proof: In ∆ ABD; PR || BD

 $\underline{AP} = \underline{AR} (BPT)$

PB RD

In Δ ACD, RQ || DC

 \therefore AR = AQ (BPT)

RD QC

In $\triangle APR$ and $\triangle ABD$,

 $\angle APR = \angle ABD$ (corresponding angles.)

 \angle ARP = \angle ADB (corresponding angles.)

 $\Delta \Delta$ APR is similar to $\Delta \Delta$ ABD (AA similarity)

 \therefore <u>AP</u> = <u>AR</u> = <u>PR</u> (corresponding sides of similar triangles are proportional)---

-(i)

AB AD BD

Similarly \triangle ARQ is similar to \triangle ADC \ $\underline{AQ} = \underline{AR} = \underline{RQ}$ ----(ii) AC AD DC

According to equation (i) and (ii),

AR = PR = RQ

AD BD DC


but BD = DC (given)

∴ PR = RQ

or AD bisects PQ at R (proved).


Question-5

Prove that the line joining the midpoints of the sides of the triangle form four triangles, each of which is similar to the original triangle.



Solution:

Given: In \triangle ABC, D, E, F are the midpoints of AB, BC and AC respectively.



To prove: ΔABC ~ ΔDEF

ΔABC ~ ΔADF

ΔABC ~ ΔBDE

ΔABC ~ ΔEFC

Proof: In \triangle ABC, D and F are mid points of AB and AC respectively.

.. DF || BC (midpoint theorem)

In \triangle ABC and \triangle ADF

 $\angle A$ is common; $\angle ADF = \angle ABC$ (corresponding angles)

 \triangle ABC ~ \triangle DF (AA similarity) ----(1)

Similarly we can prove $\triangle ABC \sim \triangle BDE$ (AA similarity)----(2)

 \triangle ABC ~ \triangle EFC (AA similarity)----(3)

In \triangle ABC and \triangle DEF;


since D,E, F are the midpoints of AB, BC and AC respectively,

DF= $(1/2) \times$ BC; DE = $(1/2) \times$ AC; EF = $(1/2) \times$ AB; (midpoint theorem) \therefore AB = BC = CA = 2 EF DF DE \therefore \triangle ABC \sim \triangle EFD (SSS similarity)------(4) From (1), (2), (3) and (4) \triangle ABC \sim \triangle DEF \triangle ABC \sim \triangle ADF \triangle ABC \sim \triangle BDE \triangle ABC \sim \triangle EFC.

Question-6

In triangle ABC, DE||BC and AD : DB = 2 : 3. Determine the ratio of the area triangle ADE to the area triangle ABC.


Solution:




$$\frac{\text{Area }\triangle ADE}{\text{Area }\triangle ABC} = \frac{2^2}{5^2} = \frac{4}{25}$$

Question-7

One angle of a triangle is equal to one angle of another triangle and the bisectors of these equal angles divide the opposite sides in the same ratio. Prove that the triangles are similar.





Solution:

Given: ΔABC and ΔPQR

$$\angle A = \angle P$$

AD and PS bisects $\angle A$ and $\angle P$ respectively.

$$BD = QS$$
 $DC SR$

To prove: $\triangle ABC \sim \triangle PQR$ **Proof**: In $\triangle ABC$ and $\triangle PQR$

 \therefore <u>AB</u> = <u>BD</u> (Angle bisector theorem) ----(1)

AC DC

Similarly in Δ PQR,

 $\underline{PQ} = \underline{QS}$ (Angle bisector theorem) ----(2)

PR SR

But BD = QS (given)

DC SR

: According to equation (1) and (2)

$$\underline{AB} = \underline{PQ} \triangleright \underline{AB} = \underline{AC}$$


AC PR PQ PR

$$\angle A = \angle P$$
 (given)

∴ \triangle ABC ~ \triangle PQR (SAS similarity).

Question-8

The bisector of interior angle A of a triangle AB meets BC in D and the bisector of exterior angle A meets BC produced in E. Prove that $\frac{BD}{BE} = \frac{CD}{CE}$



Solution:

Given: $\triangle ABC$, AD bisects interior $\angle A$ and AE bisects exterior $\angle A$ meeting BC at D and BC produced at E.

To prove: $\frac{BD}{BE} = \frac{CD}{CE}$

Proof: In \triangle ABC, AD bisects interior \angle A

$$\therefore \frac{AB}{AC} = \frac{BD}{DC} \text{ (Angle Bisector theorem)}....(1)$$

Similarly in DABC, AE bisects exterior ∠A

$$\therefore \frac{AB}{AC} = \frac{BE}{CE} \dots (2)$$


From equation (1) and (2),

$$\frac{AB}{AC} = \frac{BD}{CD} = \frac{BE}{CE} \implies \frac{BD}{BE} = \frac{CD}{CE}$$

Hence Proved.

Question-9

In a triangle ABC, XY || AC divides the triangle into two parts equal in areas. Determine $\frac{A\times}{AB}$.



Solution:

Given: ABC is a triangle with XY || AC divides the triangle into two parts equal in areas.

To find: $\frac{A\times}{AB}$

Proof:

ar ΔBXY = ar trap. XYCA (Given) \therefore ar ΔBXY = $\frac{1}{2}$ ar ΔABC

In ΔBXY andBAC,

 \angle BXY = \angle BAC (Corresponding angles)

∠BYX = ∠BCA (Corresponding angles)

ΔBXY ~ ΔBAC (AA similarity)

$$\therefore \frac{\text{Area } \Delta B \times Y}{\text{Area } \Delta B \wedge C} = \frac{B \times^2}{AB^2}$$
 (Areas of similar triangle)

$$\therefore \frac{1}{2} = \frac{B \times ^2}{AB^2}$$

$$\therefore \frac{1}{\sqrt{2}} = \frac{B \times}{AB}$$


$$\therefore$$
 AB - BX = $\sqrt{2}$ BX - BX


$$\therefore AX = (\sqrt{2} - 1)BX$$

$$\frac{A\times}{AB} = \frac{(\sqrt{2}-1)B\times}{\sqrt{2}B\times} = \frac{\sqrt{2}-1}{\sqrt{2}}.$$

Question-10

If two sides and a median bisecting the third side of a Δ are respectively proportional to the corresponding sides and the median of another triangle, then prove that the two triangles are similar.





Solution:

Given: \triangle ABC and \triangle PQR where BD and QS are the medians and $\underline{AB} = \underline{BC} = \underline{BD}$

PQ QR QS

To prove: $\triangle ABC \sim \triangle PQR$

Construction: Produce BD and QS to E and T respectively such that BD = DE and QS = ST. CE and TR are joined.

Proof: In ΔADB and DCDE,

AD = DC (given)


 $\angle ADB = \angle CDE$ (Vertically opposite angles)

BD = DE

 $\Delta ADB \cong \Delta CDE (SAS \cong axiom)$

```
Hence AB = CE and \angle ABD = \angle DEC.
Similarly \triangle PQS \cong \triangle RST,
hence PQ = TR and \anglePQS = \angleSTR.
Consider \triangle EBC and \triangle TQR,
BD = 2 BD = BE (from given and construction)-----(1)
QS 2QS QT
AB = CE and PQ = RT (proved),
    \underline{AB} = \underline{CE}----(2)
    PQ RT
    AB = BC = BD (Given)
                                               ----(3)
    PQ QR QS
From (1),(2) and (3),
    BE = CE = BC
    QT RT QR
∴ ΔEBC ~ ΔTQR(SSS similarity axiom).
⇒ ∠DBC = ∠SQR and ∠DEC = ∠STR ---(4) (corresponding angles of similar
triangles are proportional)
 But \angle ABD = \angle DEC and \angle PQS = \angle STR (proved)----(5)
 \therefore \angle ABD = \angle PQS \text{ (from (4) and (5)) ----(6)}
 From (5) and (6),
 \angle ABC = \angle PQR ----I
 In \triangle ABC and \triangle PQR,
    AB = BC (given)
    PQ QR
 And \angle ABC = \angle PQR (from I)
 .: ΔABC ~ ΔPQR (SAS Similarity).
```

Two right triangles ABC and DBC are drawn on the same hypotenuse BC and on the same sides of BC. If AC and DB intersect at P, prove that AP × PC = BP × PD.



Solution:

Given: Two right triangles ABC and BDC on the same hypotenuse BC. AC and BD intersect at P.

To prove: AP × PC = BP × PD

Proof: In ΔABP and ΔDCP

∠A = ∠D (= 90°) (given)

∠APB = ∠DPC (vertically opposite angles)

∴ ΔABP ~ ΔDCP (AA similarity axiom)

∴ AB = BP = AP (corresponding sides of similar Δ s are proportional)

.......(1)

DC CP DP

From (1) BP = AP


CP DP

By cross multiplication,


BP × DP = AP × PC (proved).

Question-12

If ABC is an equilateral triangle of side 2a prove that the altitude AD = 3a and $3AB^2 = 4AD^2$.



Solution:



Given: \triangle ABC is an equilateral triangle of side 2a. AD is the altitude of triangle.

To Prove: $AD = a\sqrt{3}$ and $3AB^2 = 4AD^2$


Proof:

In rt. \triangle ADC, AD² = AC² - DC² = (2a)² - a² = 4a² - a² = 3a²

∴ AD =
$$a\sqrt{3}$$

 $3AB^2 = 3(2a)^2$
= $3(2a)^2$
= 3^4a^2
= $4(a\sqrt{3})^2$
= $4AD^2$
∴ $3AB^2 = 4AD^2$

Two isosceles Δ s have equal vertical angles and their areas are in the ratio 9:16. Find the ratio of their corresponding heights (altitudes).



Solution:

Given: \triangle ABC and \triangle PQR are isosceles and \angle A = \angle P. AD, PS are the altitudes and \triangle ABC = $\frac{9}{2}$

and $\frac{\Delta ABC}{\Delta PQR} = \frac{9}{16}$. **To find:** AD PS

Proof: In \triangle ABC, \angle B = \angle C (isosceles \triangle property)

Similarly in \triangle PQR, \angle Q = \angle R.

$$\angle A = \angle P$$
 (given)

$$\therefore \angle B = \angle C = \frac{180^{\circ} - \angle A}{2}$$

Since $\angle A = \angle P$

$$\angle$$
 B = \angle C = \angle Q = \angle R


$$\therefore \Delta ABC \sim \Delta PQR (AA)$$

If 2 triangles are similar then the ratio of areas will be equal to the square of the corresponding sides,

... The ratio of their corresponding heights is 3:4.

The diagonals of a quadrilateral ABCD intersect each other at the point 0 such that $\frac{AO}{BO} = \frac{CO}{DO}$. Show that ABCD is a trapezium.

Solution:



Given: ABCD is a trapezium with AB || CD and the diagonals AC and BD intersect at 'O'.

To prove: $\frac{OA}{OC} = \frac{OB}{OD}$

Proof:

In the figure consider the triangle OAB and OCD

∠DOC = ∠AOB (Vertically opposite angles are equal)

since AB || DC,

 \angle DCO = \angle OAB (Alternate angles are equal)

.. By AA corollary of similar triangles.


.. Δ OAB \sim Δ OCB When the two triangle are similar, the side are proportionally.

$$\Rightarrow \frac{OA}{OC} = \frac{OB}{OD}$$

Hence proved.

Question-15

P and Q are the points on the sides AB and AC respectively of a \triangle ABC. If AP = 2 cm, PB = 4 cm, AQ = 3 cms, QC = 6 cm, prove that BC = 3PQ.



Solution:

Given: \triangle ABC, PQ are points on AB and AC such that AP = 2 cm, BP = 4

cm, AQ = 3 cm, QC = 6 cm

To prove: BC = 3PQ

Proof: In \triangle ABC, $\frac{AP}{PB} = \frac{2}{4} = \frac{1}{2} \cdot \frac{AQ}{QC} = \frac{3}{6} = \frac{1}{2}$

As
$$\underline{AP} = \underline{AQ}$$

PB QC

According to converse of BPT, PQ || BC

In \triangle APQ and \triangle ABC

$$\therefore \angle APQ = \angle ABC$$
 (Corresponding angles)

∠ A is Common

 $\Delta APQ \sim \Delta ABC$ (AAS similarity)

 $\frac{\Delta P}{\Delta B} = \frac{AQ}{QC}$ (corresponding sides of similar Δ s are proportional)

But
$$\frac{AP}{AB} = \frac{PQ}{BC}$$


$$PQ = 2 = 1$$

BC 6 3

: 3PQ = BC (Proved).

Question-16

Through the midpoint of M of the side CD of a parallelogram ABCD, the line BM is drawn intersecting AC in L and AD produced in E. Prove that EL = 2BL.



Solution:

Given: ABCD is a parallelogram, M is the midpoint of CD. BM intersects AC at L and AD produced at E.

To prove: EL = 2BL

Proof: In \triangle BMC and \triangle EDM

 \angle DME = \angle BMC (Vetically opposite angles)

DM = MC (given)

 $\angle DEM = \angle MBC$ (alternate angles)

 $\therefore \triangle$ BMC $\cong \triangle$ EDM (ASA congruence)

 \therefore DE = BC (c.p.c.t)

But BC = AD (opposite sides of parallelogram ABCD)

 \therefore AD = DE \Rightarrow AE = 2AD = 2BC

In \triangle AEL and \triangle CBL

 \angle ALE = \angle BLC (Vertically opposite angles)

∠ AEL = ∠ LBC (alternate angles)

 $\Delta \Delta AEL \sim \Delta CBL$ (AA similarity axiom)

$$\Rightarrow \frac{AE}{BC} = \frac{AL}{LC} = \frac{EL}{BL}$$

$$\Rightarrow \frac{EL}{BL} = \frac{AE}{BC}$$

$$\Rightarrow \frac{EL}{BL} = \frac{AD + DE}{BC}$$

$$\Rightarrow \frac{EL}{BL} = \frac{BC + BC}{BC}$$

$$\Rightarrow \frac{EL}{BL} = \frac{2BC}{BC}$$


$$\Rightarrow \frac{EL}{BL} = 2$$

∴ EL = 2 BL

Question-17

The side BC of a triangle ABC is bisected at D; O is any point in AD. BO, CO produced meet AC, AB in E,F respectively, and AD is produced to X so that D is the mid point of OX. Prove that AO: AX = AF: AB and show that EF is parallel to BC.

Solution:



Given: The side BC of a triangle ABC is bisected at D; O is any point in AD. BO, CO produced meet AC, AB in E, F respectively, and AD is produced to X so that D is the mid point of OX.

To Prove : AO : AX = AF : AB and show that EF is parallel to BC.

Construction: Join BX and CX.

Proof: In quadrilateral BOCX, BD = DC and DO = DX (given)

.. BOCX is a parallelogram (When the diagonals of a quadrilateral bisect each other, then the quad. is a parallelogram)

∴ BX || CO (Definition of a parallelogram) or BX || FO. In \triangle ABX, BX || FO(proved).

∴ AO : AX = AF : AB (using B.P.T) -----(i)

Similarly, AO : AX = AE : AC -----(ii)


From (i) and (ii), AF: AB = AE: AC

By corollary to B.P.T, EF is parallel to BC.

Question-18

ABC is a triangle in which \angle BAC = 90° and DEFG is a square, prove that DE² = BD × EC.

Solution:



Given: ABC is a triangle in which ∠BAC = 90° and DEFG is a square.

To prove: $DE^2 = BD \times EC$.

Proof: In \triangle AGF and \triangle DBG,

∠AGF = ∠GBD (corresponding angles)

 \angle GAF = \angle BDG (each = 90°)

 $\cdot \cdot \cdot \Delta AGF \sim \Delta DBG.$ -----(i)

Similarly, ΔAFG ~ ΔECF (AA Similarity)-----(ii)

From (i) and (ii), $\Delta DBG \sim \Delta ECF$.

$$\frac{BD}{EF} = \frac{BG}{FC} = \frac{DG}{EC}$$

$$\frac{BD}{EE} = \frac{DG}{EC}$$


EF × DG = BD × EC. ----(iii)

Also DEFG is a square ⇒ DE = EF = FG = DG -----(iv)

From (iii) and (iv), $DE^2 = BD \times EC$.

Question-19

In fig. DE || BC. If AD = x, DB = x - 2, AE = x + 2 and EC = x - 1, find the value x.



Solution:

Given: ABC is a triangle, DE || BC, AD = x, DB = x - 2, AE = x + 2 and EC = x - 1.

To find: x

In \triangle ABC, we have

DE || BC

Therefore [By Thale's theorem]

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$AD \times EC = AE \times DB$$

$$x(x-1) = (x-2)(x+2)$$

 $x^2 - x = x^2 - 4$

$$x^2 - x = x^2 - 4$$

 $x = 4$

Question-20

In a \triangle ABC, D and E are points on the sides AB and AC respectively such that DE || BC. If AD = 4 cm, AE = 8 cm, DB = x - 4 and EC = 3x - 19, find x.

Solution:

Given: In \triangle ABC, D and E are points on the sides AB and AC respectively such that DE || BC. AD = 4 cm, AE = 8 cm, DB = x - 4 and EC = 3x - 19.

To find: x.

In Δ ABC, we have DE || BC

Therefore $\frac{AD}{DB} = \frac{AE}{EC}$ [By Thale's theorem]


$$\frac{4}{x-4} = \frac{8}{3x-19}$$

$$4(3x - 19) = 8(x - 4)$$

$$12x - 76 = 8x - 32$$

$$4x = 44$$

$$x = 11$$




In a \triangle ABC, AD is the bisector of \angle A, meeting side BC at D. If AC = 4.2 cm, DC = 6 cm, BC = 10 cm, find AB.

Solution:

Given: In a \triangle ABC, AD is the bisector of \angle A, meeting side BC at D. AC = 4.2 cm, DC = 6 cm and BC = 10 cm.

To find: AB.

In \triangle ABC, $\frac{AB}{AC} = \frac{BD}{DC}$ [By internal bisector theorem] $AB = 4 \times 4.2/6 = 2.8 \text{ cm}$ \therefore AB = 2.8 cm



Question-22

In \triangle ABC, \angle B = 90° and D is the mid-point of BC. Prove that AC² = AD² + 3CD².



Solution:

Given: In \triangle ABC, \angle B = 90° and D is the mid-point of BC.

To Prove: $AC^2 = AD^2 + 3CD^2$

Proof: In \triangle ABD, $AD^2 = AB^2 + BD^2$ $AB^2 = AD^2 - BD^2$ (i)

In \triangle ABC, AC² = AB² + BC² AB² = AC² - BC²(ii)

Equating (i) and (ii) $AD^2 - BD^2 = AC^2 - BC^2$ $AD^2 - BD^2 = AC^2 - (BD + DC)^2$

$$AD^2 - BD^2 = AC^2 - BD^2 - DC^2 - 2BD \times DC$$


 $AD^2 = AC^2 - DC^2 - 2DC^2$ (DC = BD)

 $AD^2 = AC^2 - 3DC^2$

ABC is a triangle in which AB = AC and D is a point on the side AC such that $BC^2 = AC \times CD$. Prove that BD = BC.

Solution:

Given: A \triangle ABC in which AB = AC. D is a point on AC such that BC² = AC × CD.



To prove: BD = BC

Proof: Since $BC^2 = AC \times CD$ Therefore $BC \times BC = AC \times CD$ AC/BC = BC/CD....(i)

Also ∠ACB = ∠BCD

Since $\triangle ABC \sim \triangle BDC$ [By SAS Axiom of similar triangles]

AB/AC = BD/BC(ii)


But AB = AC (Given)(iii)

From (i), (ii) and (iii) we get BD = BC.

Question-24

The diagonal BD of a parallelogram ABCD intersects the segment AE at the point F, where E is any point on the side BC. Prove that $DF \times EF = FB \times FA$.

Solution:



Given: The diagonal BD of parallelogram ABCD intersects the segment AE at F, where E is any point on BC.

To prove: DF × EF = FB × FA

Proof: In triangles AFD and BFE,

 \angle FAD = \angle FEB (Alternate angles)

∠AFD = ∠BFE (Vertically opposite

angles)

Therefore $\triangle ADF \sim \triangle BFE$ (AA similarity)

 $\frac{DF}{FA} = \frac{FB}{EF}$


Hence DF \times EF = FB \times FA

Question-25

ABC is a triangle, right-angled at C and AC = $\sqrt{3}$ BC. Prove that \angle ABC = 60°.

Solution:

Given: \triangle ABC is right angled at C and AC = $\sqrt{3}$ BC.



To prove: \angle ABC = 60°.

Proof:

Let D be the midpoint of AB. Join CD.

Now, $AB^2 = BC^2 + AC^2 = BC^2 + (\sqrt{3}BC)^2 = 4BC^2$

Therefore AB = 2BC.

Now, BD = $\frac{1}{2}$ AB = $\frac{1}{2}$ (2BC) = BC.

But, D being the midpoint of hypotenuse AB, it is equidistant from all the

three vertices.

Therefore CD = BD = DA or CD = $\frac{1}{2}$ AB = BC.


Thus, BC = BD = CD,

i.e., ΔBCD is a equilateral triangle.

Hence, ∠ABC = 60°.

Question-27

Let ABC be a triangle, right-angled at C. If D is the mid-point of BC, prove that $AB^2 = 4AD^2 - 3AC^2$.



Solution:

Given: ABC be a triangle, right-angled at C and D is the mid-point of BC.

To Prove: $AB^2 = 4AD^2 - 3AC^2$.

Proof:


From right triangle ACB, we have,

$$AB^2 = AC^2 + BC^2$$

= $AC^2 + (2CD)^2 = AC^2 + 4CD^2$ [Since BC = 2CD]
= $AC^2 + 4(AD^2 - AC^2)$ [From right \triangle ACD]
= $4AD^2 - 3AC^2$.

Question-28

In the given figure, points D and E trisect BC and $\angle B = 90^{\circ}$. Prove that $8AE^2 = 3AC^2 + 5AD^2$.



Solution:

Given: In \triangle ABC, points D and E trisect on BC and \angle B = 90°.

To Prove: $8AE^2 = 3AC^2 + 5AD^2$.

Proof: Let ABC be the triangle in which $h \angle B = 90^{\circ}$. Let the points D and E

trisect BC.

Join AD and AE. Then,

$$AC^2 = AB^2 + BC^2$$

$$3AC^2 = 3AB^2 + 3BC^2$$
(i)


$$AD^2 = AB^2 + BD^2$$

$$5AD^2 = 5AB^2 + 5BD^2$$
(ii)

Therefore
$$3AC^2 + 5AD^2 = 8AB^2 + 3BC^2 + 5BD^2$$

 $= 8AB^2 + 3 \cdot \left(\frac{3}{2}BE\right)^2 + 5\left(\frac{1}{2}BE\right)^2$
 $= 8AB^2 + \left(\frac{27}{4} + \frac{5}{4}\right)BE^2$
 $= 8AB^2 + 8BE^2$
 $= 8(AB^2 + BE^2)$
 $= 8AE^2$

In fig., ABC is a triangle in which AB = AC. D and E are points on the sides AB and AC respectively such that AD = AE. Show that the points B, C, E and D are concyclic.



Solution:

Given: In \triangle ABC, AB = AC. D and E are points on the sides AB and AC respectively such that AD = AE.

To Prove: Points B, C, E and D are concyclic.

Proof: In order to prove that the points B, C, E and D are concyclic, it is sufficient to show that \angle ABC + \angle CED = 180° and

 \angle ACB + \angle BDE = 180°.

In \triangle ABC, we have

AB = AC and AD = AE

AB - AD = AC - AE

DB = EC

Thus, we have

AD = AE and DB = EC.

 $\frac{AD}{DB} = \frac{AE}{EC}$

DE || BC [By the converse of Thale's Theorem]

 $\angle ABC = \angle ADE$ [Corresponding angles]

 $\angle ABC + \angle BDE = \angle ADE + \angle BDE$ [adding $\angle BDE$ on both sides]

 \angle ABC + \angle BDE = 180°

 \angle ACB + \angle BDE = 180° [Since AB = AC Therefore \angle ABC =

∠ACB]

Again DE||BC

∠ACB = ∠AED

 $\angle ACB + \angle CED = \angle AED + \angle CED$ [Adding $\angle CED$ on both sides]


 \angle ACB + \angle CED = 180°

 $\angle ABC + \angle CED = 180^{\circ}$ [Since $\angle ABC = \angle ACB$]

Therefore BDEC is a cyclic quadrilateral.

Hence, B, C, E and D are concyclic points.

In fig, \angle B < 90° and segment AD ^ BC, show that $b^2 = h^2 + a^2 + x^2 - 2ax$



Solution:

Given: In \triangle ABC, \angle B < 90° and segment AD \bot BC.

To prove: $b^2 = h^2 + a^2 + x^2 - 2ax$


Proof:

$$b^2 = h^2 + (a - x)^2$$

$$b^2 = h^2 + a^2 + x^2 - 2ax$$
.

Question-31

In the given figure, ABCD is a parallelogram P is a point on BC, such that BP : PC = 1 : 2. DP produced meets AB produced at Q. Given area of triangle CPQ = 20 m^2 , calculate the area of triangle DCP.




Solution:

Given: ABCD is a parallelogram P is a point on BC, such that BP : PC = 1 : 2. DP produced meets AB produced at Q.

Area of triangle CPQ = 20 m^2 .

To Find: Area of triangle DCP.

Construction: Join DB.



 \angle BPQ = \angle DPC (Vertically opposite angles)

```
∠ BQP = ∠ PDC (alternate angles, BQ || DC, DQ meets them)
∴ ∆ BPQ \sim ∆ CPD (AA similarity)

BP/CP = \frac{1}{2} (Given)
\frac{\text{are aΔBPQ}}{\text{are aΔCPD}} = \left(\frac{\text{BP}}{\text{CP}}\right)^2 = \frac{1}{4}
∴ Area D CPD = Area D BPQ
\frac{\text{are aΔBPQ}}{\text{are aΔCPD}} = \left(\frac{\text{BP}}{\text{CP}}\right) = \frac{1}{2} \text{ and area } Δ\text{CPQ} = 20 \text{ cm}^2 \text{ (Given)}

Area ∆BPQ = 10 cm²
Area ∆CPD = 40 cm²
\frac{\text{are aΔDBC}}{\text{are aΔDPC}} = \frac{3}{2} \text{ (Proportional to bases BC and PC)}
```

 $\frac{\text{area\DeltaDBC}}{40} = \frac{3}{2}$:: Area \triangle DBC = $40 \times 3/2 = 60 \text{ cm}^2$.