


PHYSICS

1. At $t = 0$ particle is at $\frac{A}{2}$ from mean position and moving in +ve x-direction. At general time its equation is $A \sin (\omega t + \phi)$. Value of ϕ is?

- (1) $\frac{\pi}{3}$ (2) $\frac{\pi}{6}$ (3) $\frac{5\pi}{6}$ (4) $\frac{\pi}{2}$

Ans. (2)



$$x = A \sin (\omega t + \phi)$$

at $t = 0$ $x = \frac{A}{2}$

$$\frac{A}{2} = A \sin [\omega(0) + \phi]$$


$$\sin \phi = \frac{1}{2}$$

$$\phi = \frac{\pi}{6}$$

2. A ball of mass ' m ' moving with velocity ' v ' collides and sticks to the body of mass ' $2m$ ', initially at rest. Find the final velocity of combined mass.

- (1) $\frac{v}{3}$ (2) $\frac{v}{4}$ (3) $\frac{v}{8}$ (4) $\frac{v}{10}$

Ans. (1)



$$mv = 3m v'$$

$$\frac{v}{3} m / s = v'$$

3. $y = A \sin (6t + 0.003 x)$. Find speed of wave 'x' is in centimeter :

- (1) 10 m/s (2) 20 m/s (3) 30 m/s (4) 40 m/s


Ans. (2)

Sol. $\omega = 6 \text{ rad/sec}$

$k = 0.003 \text{ rad/cm}$

$$v = \frac{\omega}{k} = \frac{6}{0.3} = 20 \text{ m / s}$$


4. Find equivalent capacitance between A and B



- (1) $4C$ (2) $2C$ (3) $C/2$ (4) $5C/3$

Ans. (2)

Sol. Circuit is reduced to



5. The de-Broglie wavelength of gas particle is λ for temperature 300 k, find the de-Broglie wavelength when temperature is 600 k?

- (1) $\frac{\lambda}{\sqrt{2}}$ (2) $\frac{\lambda}{\sqrt{3}}$ (3) $\frac{\lambda}{2}$ (4) $\frac{\lambda}{5}$

Ans. (1)

Sol. $\lambda = \frac{h}{\sqrt{2mk}}$ ($\because k = \frac{3}{2}kT$)

$$\lambda \propto \frac{1}{\sqrt{T}}$$

$$\lambda_1 \sqrt{T_1} = \lambda_2 \sqrt{T_2}$$

$$\lambda \sqrt{\frac{300}{600}} = \lambda'$$

$$\frac{\lambda}{\sqrt{2}} = \lambda' (\text{new wavelength})$$

6. If the weight on the surface of a planet of mass, radius R is 200 N. Find weight at depth R/2 from surface of planet.


(1) 200 N

(2) 300 N

(3) 100 N

(4) 400 N

Ans. (3)



$$200 = \frac{GM}{R^2} m$$


$$M = \frac{4}{3} \pi R^3 \rho$$

$$200 = \frac{G}{3} \pi \rho R m$$

$$\text{weight} \propto R$$

$$\text{Hence at } \frac{R}{2} \text{ from centre weight} = 100 \text{ N}$$

7. Force acting on rod is :



(1) 0.18 N

(2) 0.018 N

(3) 1.8 N

(4) 18 N

Sol. $F = i \ell B$

$$= \left(\frac{\varepsilon}{R} \right) \ell B = \left(\frac{v B \ell}{R} \right) \ell B = \frac{v B^2 \ell^2}{R} = \frac{4}{5} \times \left(\frac{15}{100} \right)^2 \times 1^2$$


$$= \frac{4}{5} \times \frac{225}{10^4}$$

$$= \frac{180}{10^4} = 0.018 \text{ N}$$

8. If a projectile is thrown with speed u at an angle 15° , the range obtained is 50 m. What will be range obtained if the same particle is thrown at an angle of 45° with same speed u .

(1) 50 m (2) 100 m (3) 200 m (4) 150 m

Ans. (2)



Sol.


$$50 = \frac{u^2 \sin 30}{g}$$

$$R_1 = \frac{u^2 \sin 90}{g}$$

$$\frac{50}{R_1} = \frac{1}{2}$$

$$R' = 100 \text{ m}$$


9. Find R_{eq} across A and B



(1) $\frac{16}{5} \Omega$ (2) $\frac{18}{5} \Omega$ (3) 4Ω (4) 6Ω

Ans. (1)

Sol. The Circuit can be required to



$$\Rightarrow R_{eq} = \frac{16 \times 4}{16 + 4} = \frac{16}{5} \Omega$$

10. If frequency of electromagnetic wave is f then frequency of energy density of electromagnetic wave is


- (1) $1.2 f$ (2) f (3) $\frac{f}{2}$ (4) $2f$

Ans. (4)

Sol. $E = E_0 \sin(\omega t - kx)$

$$\frac{du}{dv} = \epsilon_0 E_0^2 \sin^2(\omega t - kx)$$

11.



- A. A takes less time to reach home.
 B. B takes less time to reach home.
 C. A is faster.
 D. B is faster.
 E. A's home is farther than B.

Correct statements are :


- (1) B, C (2) B, D (3) A, C (4) A, D, E

Ans. (2)

12. The volume of Earth shrinks to $1/64$ of its initial value, mass staying the same then ratio of initial and final value of time periods of rotation of Earth about its axis is t_1/x where $t_1 = 24$. Find x :

Ans. 16

Sol. From conservation of angular momentum



$$MR^2\omega_1 = M\left(\frac{R}{4}\right)^2\omega_2$$

$$\Rightarrow MR^2\omega_1 = \frac{MR^2}{16}\omega_2$$

$$\Rightarrow \frac{\omega_1}{\omega_2} = \frac{1}{16} \Rightarrow \frac{T_2}{T_1} = \frac{1}{16} \Rightarrow \frac{T_1}{T_2} = \frac{16}{1} = \frac{t_1}{x}$$

$$\therefore t_1 = 24 \Rightarrow \frac{16}{1} = \frac{24}{t_2} \Rightarrow x = 16$$

13. Statement 1: Current sensitivity doubles when number of turns is doubled

Statement 2: Both voltage sensitivity and current sensitivity increases equally an increasing no of turns.

- (1) Statement-1 and statement-1 both are correct.
- (2) Statement-1 and statement-1 both are wrong.
- (3) Statement-1 is wrong and statement-2 is correct.
- (4) Statement-1 is correct and statement-2 is wrong.

Ans. (4)

Sol. $BINA = C\phi \rightarrow \frac{\phi}{I} = \frac{BNA}{C}$: Current sensitivity voltage sensitivity = $\frac{\phi}{V} = \frac{BNA}{CR}$

as $N \uparrow \Rightarrow R \uparrow \Rightarrow V.S$ Remains same.

14. Two gases A and B having same initial state (P, V, n, T). Now gas 'A' is compressed to $\frac{V}{8}$ by isothermal process and other gas B is compressed to $\frac{V}{8}$ by adiabatic process. Find ratio of Final pressure of gas A and B (Both gases are monoatomic)

- (1) 1/4
- (2) 1/8
- (3) 1/12
- (4) 1/64

Ans. (1)

Sol. Isothermal process equation

$$PV = P_A (V/8)$$

$$8P = P_A$$

Adiabatic process equation

$$PV^{5/3} = P_B (V/8)^{5/3}$$

$$32P = 8^{5/3} P = P_B$$

$$\frac{P_A}{P_B} = \frac{8P}{32P} = \frac{1}{4}$$

15. Mirror is moved towards the object by 4 cm, then find how much distance image will shift

- (1) 8 cm (2) 4 cm (3) 12 cm (4) 16 cm

Ans. (1)

Sol. Image distance shift = $2 \times 4 = 8$ cm

16. The magnetic field intensity inside current carrying solenoid is $H = 2.4 \times 10^3$ A/m. If Length and no. of turns of solenoid is 15 cm and 60 turns. Find current flowing in solenoid.

- (1) 4 A (2) 6 A (3) 0.6 A (4) 60 A

Ans. (2)

Sol. $B = \mu_0 \frac{N}{L} i$

$$\frac{B}{\mu_0} = \frac{N}{L} i$$

$$H = \frac{N}{L} i$$

$$2.4 \times 10^3 = \frac{60}{15 \times 10^{-2}} i$$

$$6 \text{ A} = i$$

17. **Statement 1 :** Maximum power is dissipated when resonance occurs.

Statement 2 : Maximum power is dissipated containing pure resistance due to zero phase difference.

(1) Statement I and II both are correct and II is the correct explanation of I.

(2) Statement I and II both are correct and II is not the correct explanation of I.

(3) Both statement I and II are wrong.

(4) Statement I is true, II is false.

Ans. (1)

18. Base band signal of amplitude 3V is modulate with carrier wave of amplitude 15 V Ratio of maximum to minimum, amplitude in amplitude modulate wave

(1) $\frac{3}{4}$

(2) $\frac{4}{5}$

(3) $\frac{3}{2}$

(4) $\frac{3}{7}$


Ans. (3)

Sol. $A_{\max} = A_m + A_c = 18$

$$A_{\min} = A_c - A_m = 12$$

$$\frac{A_{\max}}{A_{\min}} = \frac{3}{2}$$

19. Radius of both wires is 0.2 cm, elongation in steel wire is $x \times 10^{-6}$ m and Young's modulus of steel is 2×10^{11} N/m². Find x.




Ans. 20

Sol. Tension in steel wire $T_2 = 2g + T_1$

$$T_2 = 20 + 11.4$$

$$= 31.4 \text{ N}$$



Elongation in steel wire $\Delta L = \frac{T_2 L}{A_y}$

$$\Delta L = \frac{31.4 \times 1.6}{\pi (0.2 \times 10^{-2})^2 \times 2 \times 10^{11}}$$

$$\Delta L = \frac{16}{2 \times 4 \times 10^{-6} \times 10^{11}}$$


$$= 2 \times 10^{-5} \text{ m}$$

$$= 20 \times 10^{-6} \text{ m}$$

20. A light of intensity 32 W/m^2 enters in a system of 3 polaroid's. Angle between 3rd and 1st polaroid is 90° . Light ray passes the system with intensity 3 W/m^2 . So angle between 1st and 2nd polaroid is.

Ans. 30°

Sol. $I_0 = 32 \text{ W/m}^2$



$$I_{\text{net}} = 3 = \frac{32}{2} \cos^2 \theta \cdot \sin^2 \theta$$

$$\frac{3}{4} = 4 \sin^2 \theta \cdot \cos^2 \theta = (\sin 2\theta)^2$$

$$\frac{\sqrt{3}}{2} = \sin(2\theta)$$

Hence, $\theta = 30^\circ$

- 21.** For an object radiating heat at 300 K, the wavelength corresponding to maximum intensity is λ . If the temperature of body is increased by 300 K, the new wavelength corresponding to maximum intensity will be

- (1) $\frac{\lambda}{2}$ (2) 2λ (3) $\frac{\lambda}{4}$ (4) 4λ

Ans. (1)

Sol. $\lambda = \frac{b}{T}$

$$T' \rightarrow 2T$$

$$\lambda' \rightarrow \frac{\lambda}{2}$$

- 22.** A quantity ℓ is given as $\ell = \frac{a^2 b^3}{c \sqrt{d}}$. Given error in the calculation of a, b, c and d are 1%, 2%, 3% and 4% respectively find the maximum percentage error in quantity ℓ .

Ans. 13

Sol.
$$\frac{\Delta L}{L} = 2 \left| \frac{\Delta a}{a} \right| + 3 \left| \frac{\Delta b}{b} \right| + \left| \frac{\Delta c}{c} \right| + \frac{1}{2} \left| \frac{\Delta d}{d} \right|$$


$$= \left(2 \times 1 + 3 \times 2 + 3 + \frac{1}{2} \times 4 \right) \%$$

$$= 13\%$$

- 23.** Three concentric spheres have charge densities σ , $-\sigma$, σ respectively. Radius of inner two spheres are 2 cm and 3 cm. If potential of inner and outer spherical shell are same. Then radius of outer sphere is _____ cm :

Ans. 5

Sol.



$$\frac{kq_1}{2} + \frac{kq_2}{3} + \frac{kq_3}{r}$$

$$= \frac{k(q_1 + q_2 + q_3)}{r}$$

$$\sigma \times 4\pi \times 2 - \sigma \times 4\pi \times 3$$

$$= \frac{\sigma[4\pi \times 2^2 - 4\pi \times 3^2]}{r}$$

$$\therefore r = 5 \text{ cm}$$

- 24.** The angular momentum of e^- in H-atom in first orbit is L . Find the change in angular momentum if e^- is in second orbit of H-atom.

- (1) $2L$ (2) L (3) $\frac{L}{2}$ (4) $4L$

Ans. (2)

Sol. $mvr = \frac{nh}{2\pi}$

$$L \propto n$$


$$\text{for } n = R, L' = 2L$$

$$\Delta L = L' - L = 2L - L = L$$

- 25.** A radioactive sample of nuclei X decays simultaneously into two different nuclei Y and Z with half-life of the decays processes as 12 minutes and 3 minutes respectively. Find the time after which 50% of nuclei of the sample X has decayed.


Ans. 2.4 min

Sol.



$$t_{1/2} = \frac{t_1 t_2}{t_1 + t_2} = \frac{3 \times 12}{15} = 2.4 \text{ min}$$

26. Zener breakdown voltage is 8 volt. If power of Zener Diode is 1.6 watt find R_0 .



Ans. 10Ω

Sol. $P_z = V_z I_z$

$$1.6 = 8 \cdot I_z$$

$$I_z = 0.2 \text{ A}$$

$$10 - 0.2R - 8 = 0$$

$$0.2R = 2$$

$$R = 10 \Omega$$