## **DPP - Daily Practice Problems**

## **Chapter-wise Sheets**

**Electrostatic Potential & Capacitance** 

Max. Marks : 180 Marking Scheme : (+4) for correct & (-1) for incorrect answer Time : 60 min.

**INSTRUCTIONS** : This Daily Practice Problem Sheet contains 45 MCQs. For each question only one option is correct. Darken the correct circle/ bubble in the Response Grid provided on each page.

6.

3.


abcd

2. abcd

1. If n drops, each charged to a potential V, coalesce to form a single drop. The potential of the big drop will be

(a) 
$$\frac{V}{n^{2/3}}$$
 (b)  $\frac{V}{n^{1/3}}$  (c)  $Vn^{1/3}$  (d)  $Vn^{2/3}$ 

2. The capacitance of a parallel plate capacitor is  $C_a$  (Fig. a). A dielectric of dielectric constant K is inserted as shown in fig. (b) and (c). If  $C_b$  and  $C_c$  denote the capacitances in fig. (b) and (c), then



(c) both 
$$C_b, C_c < C_a$$
 (d)  $C_a = C_b = C_c$ 

(a)

Response

Grid

3. The electric potential V(x) in a region around the origin is given by  $V(x) = 4x^2$  volts. The electric charge enclosed in a cube of 1 m side with its centre at the origin is (in coulomb) (a)  $8\varepsilon_0$  (b)  $-4\varepsilon_0$  (c) 0 (d)  $-8\varepsilon_0$ 

(a)b)(c)(d)

6. (a)b)c)d)

1.

- 4. A parallel plate condenser is immersed in an oil of dielectric constant 2. The field between the plates is
  - (a) increased, proportional to 2
  - (b) decreased, proportional to  $\frac{1}{2}$
  - (c) increased, proportional to -2
  - (d) decreased, proportional to  $-\frac{1}{2}$
- 5. What is the effective capacitance between points *X* and *Y*?

$$\begin{array}{c} C_1=6\mu F\\ \hline\\ C_3=6\mu FC_5=20\mu F\\ C_2=6\mu F\\ \hline\\ C_4=6\mu F\\ \hline\\ C_4=6\mu F\\ \hline\\ H\\ \hline\\ D_4\\ \hline\\ D_$$

(a)  $24 \mu F$  (b)  $18 \mu F$  (c)  $12 \mu F$  (d)  $6 \mu F$ Two identical particles each of mass m and having charges -q and +q are revolving in a circle of radius r under the influence of electric attraction. Kinetic energy of each


particle is 
$$\left(k = \frac{1}{4\pi\epsilon_0}\right)$$
  
(a)  $kq^2/4r$  (b)  $kq^2/2r$  (c)  $kq^2/8r$  (d)  $kq^2/r$ 

5.

abcd

4. (a)b)©(d)

7. Four metallic plates each with a surface area of one side A, are placed at a distance d from each other. The two outer plates are connected to one point A and the two other inner plates to another point B as shown in the figure. Then the capacitance of the system is




- 8. A parallel plate condenser with a dielectric of dielectric constant K between the plates has a capacity C and is charged to a potential V volt. The dielectric slab is slowly removed from between the plates and then reinserted. The net work done by the system in this process is
  - (a) zero (b)  $\frac{1}{2}(K-1) CV^2$ (c)  $\frac{CV^2(K-1)}{K}$ (d)  $(K-1) CV^2$
- 9. If a slab of insulating material  $4 \times 10^{-5}$  m thick is introduced between the plates of a parallel plate capacitor, the distance between the plates has to be increased by  $3.5 \times 10^{-5}$  m to restore the capacity to original value. Then the dielectric constant of the material of slab is (a) 8 (b) 6 (c) 12 (d) 10
- **10.** A unit charge moves on an equipotential surface from a point A to point B, then

(a) 
$$V_A - V_B = +ve$$
 (b)  $V_A - V_B = 0$ 

(c) 
$$V_A - V_B = -ve$$
 (d) it is stationary

**11.** Identify the false statement.

- (a) Inside a charged or neutral conductor, electrostatic field is zero
- (b) The electrostatic field at the surface of the charged conductor must be tangential to the surface at any point
- (c) There is no net charge at any point inside the conductor
- (d) Electrostatic potential is constant throughout the volume of the conductor
- 12. In a hollow spherical shell, potential (V) changes with respect to distance (s) from centre as





- 13. The 1000 small droplets of water each of radius r and charge Q, make a big drop of spherical shape. The potential of big drop is how many times the potential of one small droplet ?
  (a) 1 (b) 10 (c) 100 (d) 1000
- 14. The work done in carrying a charge q once around a circle of radius r with a charge Q placed at the centre will be

(d)  $Qq^2/(4\pi\epsilon_0 r)$ 

(a) 
$$Qq(4\pi\epsilon_0 r^2)$$
 (b)  $Qq/(4\pi\epsilon_0 r)$ 

15. A parallel plate condenser is filled with two dielectrics as shown. Area of each plate is A  $m^2$  and the separation is t m. The dielectric constants are  $k_1$  and  $k_2$  respectively. Its capacitance in farad will be

 $\frac{k_1 \quad k_2}{k_1 \quad k_2}$ 

(a) 
$$\frac{\varepsilon_0 A}{t} (k_1 + k_2)$$
 (b)  $\frac{\varepsilon_0 A}{t} \cdot \frac{k_1 + k_2}{2}$   
(c)  $\frac{2\varepsilon_0 A}{t} (k_1 + k_2)$  (d)  $\frac{\varepsilon_0 A}{t} \cdot \frac{k_1 - k_2}{2}$ 

- 16. Two metal pieces having a potential difference of 800 V are 0.02 m apart horizontally. A particle of mass  $1.96 \times 10^{-15}$  kg is suspended in equilibrium between the plates. If e is the elementary charge, then charge on the particle is
- (a) 8 (b) 6 (c) 0.1 (d) 3
  17. A one microfarad capacitor of a TV is subjected to 4000 V potential difference. The energy stored in capacitor is

  (a) 8 J
  (b) 16 J
  - (c)  $4 \times 10^{-3}$  J (d)  $2 \times 10^{-3}$  J
- 18. An unchanged parallel plate capacitor filled with a dielectric constant K is connected to an air filled identical parallel capacitor charged to potential  $V_1$ . If the common potential is  $V_2$ , the value of K is

(a) 
$$\frac{V_1 - V_2}{V_1}$$
 (b)  $\frac{V_1}{V_1 - V_2}$   
(c)  $\frac{V_2}{V_1 - V_2}$  (d)  $\frac{V_1 - V_2}{V_2}$ 

19. In the circuit given below, the charge in  $\mu$ C, on the capacitor having 5  $\mu$ F is (a) 4.5 (b) 9 (c) 7

w, the bacitor	$\begin{array}{c} 3\mu F & 2\mu F \\ e & & & \\ f & & & \\ a & & 4\mu F_{+} \\ \end{array} \begin{array}{c} 2\mu F \\ \hline & & \\ & & $	

Response Grid	12. a b c d		9. abcd 14.abcd	~ ~ ~ ~ ~	
	17.00000	18.0000	19.0000		


(d) 15

**20.** Two concentric, thin metallic spheres of radii  $R_1$  and  $R_2$   $(R_1 > R_2)$  bear charges  $Q_1$  and  $Q_2$  respectively. Then the potential at distance r between  $R_1$  and  $R_2$  will be

(a) 
$$k\left(\frac{Q_1+Q_2}{r}\right)$$
 (b)  $k\left(\frac{Q_1}{r}+\frac{Q_2}{R_2}\right)$   
(c)  $k\left(\frac{Q_2}{r}+\frac{Q_1}{R_1}\right)$  (d)  $k\left(\frac{Q_1}{R_1}+\frac{Q_2}{R_2}\right)$ 

21. Charge Q on a capacitor varies with voltage V as shown in the figure, where Q is taken along the X-axis and V along the Y-axis. The area of triangle OAB represents

(a) capacitance
(b) capacitive reactance
(c) magnetic field between the plates O




- 22. An alpha particle is accelerated through a potential difference of  $10^6$  volt. Its kinetic energy will be
  - (a) 1 MeV (b) 2 MeV (c) 4 MeV (d) 8 MeV
- **23.** Four point charges -Q, -q, 2q and 2Q are placed, one at each corner of the square. The relation between Q and q for which the potential at the centre of the square is zero is :

(a) 
$$Q = -q$$
 (b)  $Q = -\frac{1}{q}$  (c)  $Q = q$  (d)  $Q = \frac{1}{q}$ 

24. A parallel plate capacitor having a separation between the plates d, plate area A and material with dielectric constant K has capacitance  $C_0$ . Now one-third of the material is replaced by another material with dielectric constant 2K, so that effectively there are two capacitors one with area 1/3A, dielectric constant 2K and another with area 2/3A and dielectric constant K. If the capacitance of this new capacitor

is C then 
$$\frac{C}{C_0}$$
 is

(a) 1 (b) 4/3 (c)
25. Two condensers, one of capacity C and other of capacity C/2 are connected to a V-volt battery, as shown. The work done in charging fully both the condensers is



(d)  $2CV^2$ .

• A

 $CV^2$ 

•C

(a) 
$$\frac{1}{4}CV^2$$
 (b)  $\frac{3}{4}CV^2$  (c)  
A B and C are three points

- **26.** A, B and C are three points in a uniform electric field. The electric potential is
  - (a) maximum at B
  - (b) maximum at C
  - (c) same at all the three points A, B and C
  - (d) maximum at A

- 27. Three capacitors are connected in the arms of a triangle ABC as shown in figure 5 V is applied between A and B. The voltage between B and C is
  - (a) 2V (b) 1V

(c) 3V (d) 1.5V

**28.** Two parallel metal plates having charges + Q and –Q face each other at a certain distance between them. If the plaves are now dipped in kerosene oil tank, the electric field between the plates will

2uF

3uF

2μF

- (a) remain same (b) become zero
- (c) increases (d) decrease
- **29.** An air capacitor C connected to a battery of e.m.f. V acquires a charge q and energy E. The capacitor is disconnected from the battery and a dielectric slab is placed between the plates. Which of the following statements is correct ?
  - (a) V and q decrease but C and E increase
  - (b) V remains unchange, but q, E and C increase
  - (c) q remains unchanged, C increases, V and E decrease
  - (d) q and C increase but V and E decrease.
- 30. Choose the wrong statement about equipotential surfaces.
  - (a) It is a surface over which the potential is constant
  - (b) The electric field is parallel to the equipotential surface
  - (c) The electric field is perpendicular to the equipotential surface
  - (d) The electric field is in the direction of steepest decrease of potential
- **31.** Two spherical conductors A and B of radii a and b (b>a) are placed concentrically in air. The two are connected by a copper wire as shown in figure. Then the equivalent capacitance of the system is

(a) 
$$4\pi\epsilon_0 \frac{ab}{b-a}$$
 (b)  $4\pi\epsilon_0(a+b)$ 

(c)

**32.** A capacitor is charged to store an energy U. The charging battery is disconnected. An identical capacitor is now connected to the first capacitor in parallel. The energy in each of the capacitors is

(d)  $4\pi\varepsilon_0 a$ 


(a) 
$$U/2$$
 (b)  $3U/2$  (c) U (d)  $U/4$ 

- **33.** Equipotentials at a great distance from a collection of charges whose total sum is not zero are approximately
  - (a) spheres (b) planes
  - (c) paraboloids (d) ellipsoids

Response				23. abcd	
GRID	25.@b©d	26. abcd	27.@b©d	28. abcd	29. abcd
GRID	<b>30.</b> @bcd	<b>31.</b> @b©d	32.@b©d	<b>33.</b> @bcd	

$$\longrightarrow$$

34. Which of the following figure shows the correct equipotential surfaces of a system of two positive charges?



35. Two identical metal plates are given positive charges  $Q_1$ and  $Q_2$  (  $\leq Q_1$ ) respectively. If they are now brought close together to form a parallel plate capacitor with capacitance C, the potential difference between them is

(a) 
$$\frac{Q_1 + Q_2}{2C}$$
 (b)  $\frac{Q_1 + Q_2}{C}$  (c)  $\frac{Q_1 - Q_2}{C}$  (d)  $\frac{Q_1 - Q_2}{2C}$ 


**36.** The capacitance of the capacitor of plate areas  $A_1$  and  $A_2$  ( $A_1 < A_2$ ) at a distance d, as shown in figure is

(a) 
$$\frac{\epsilon_0 (A_1 + A_2)}{2d}$$
 (b)  $\frac{\epsilon_0 A_2}{d}$   
(c)  $\frac{\epsilon_0 \sqrt{A_1 A_2}}{d}$  (d)  $\frac{\epsilon_0 A_1}{d}$ 

- 37. In a given network the equivalent capacitance between A and B is  $[C_1 = C_4 =$  $1 \,\mu\text{F}, C_2 = C_3 = 2\mu\text{F}$ (a)  $3 \mu F$ (b) 6 µF
  - (d)  $2.5 \,\mu F$ (c)  $4.5 \,\mu F$
- 38. A parallel plate air capacitor is charged to a potential difference of V volts. After disconnecting the charging battery the distance between the plates of the capacitor is increased using an insulating handle. As a result the potential difference between the plates

34 00000

- (a) does not change (b) becomes zero
- (c) increases (d) decreases
- **39.** Figure shows three circular arcs, each of radius R and total charge as indicated. The net electric potential at the centre of curvature is



35 0000

 $A_1$ 

(a) 
$$\frac{Q}{2\pi\epsilon_0 R}$$
 (b)  $\frac{Q}{4\pi\epsilon_0 R}$   
(c)  $\frac{2Q}{\pi\epsilon_0 R}$  (d)  $\frac{Q}{\pi\epsilon_0 R}$ 

**40.** An electric field  $\vec{E} = (25\hat{i} + 30\hat{j})NC^{-1}$  exists in a region of space. If the potential at the origin is taken to be zero then the potential at x = 2 m, y = 2 m is :

(a) -110 V (b) -140 V (c) -120 V (d) -130 V

41. If a unit positive charge is taken from one point to another over an equipotential surface, then

-3C

40

B

38 @ 6 6 6 1

50

- (a) work is done on the charge
- (b) work is done by the charge
- (c) work done is constant
- (d) no work is done
- 42. Three large plates A, B and C are placed parallel to each other and charges are given as shown. The charge that appears on the left surface of plate B is

(a) 5C (b) 6C

- (c) 3C (d) -3C 43. Three charges  $2q_{1} - q_{2}$  and  $-q_{3}$  are located at the vertices of an equilateral triangle. At the centre of the triangle
  - (a) the field is zero but potential is non-zero
  - (b) the field is non-zero, but potential is zero
  - (c) both field and potential are zero
  - (d) both field and potential are non-zero
- 44. If a charge – 150 nC is given to a concentric spherical shell and a charge +50 nC is placed at its centre then the charge on inner and outer surface of the shell is
  - (a) -50 nC, -100 nC(b) +50 nC, -200 nC
  - (c) -50 nC, -200 nC(d) 50 nC, 100 nC
- 45. Two capacitors of capacitances  $C_1$  and  $C_2$  are connected in parallel across a battery. If Q1 and Q2 respectively be the

charges on the capacitors, then  $\frac{Q_1}{Q_2}$  will be equal to

(a) 
$$\frac{C_2}{C_1}$$
 (b)  $\frac{C_1}{C_2}$  (c)  $\frac{C_1^2}{C_2^2}$  (d)  $\frac{C_2^2}{C_1^2}$ 

37 00000

GRID 39		40.@bCd 45.@bCd	~ ~ ~ ~	42. a b c d	43. abcd	
DAILY PRACTICE PROBLEM DPP CHAPTERWISE CP16 - PHYSICS						
Total Questions		45	Total Marks		180	
Attempted			Correct			
Incorrect			Net Score			
Cut-off Score		50	Qualifying Score		70	
Success Gap = Net Score – Qualifying Score						
Net Score = (Correct × 4) – (Incorrect × 1)						

36 @@@@@