### CHAPTER-27 RAY OPTICS AND OPTICAL INSTRUMENTS

#### REFLECTION

#### LAWS OF REFLECTION

- The incident ray the reflected ray and normal to the surface of reflection at the point of incidence lie in the same plane, this plane is called the plane of incidence (also plane of reflection).
- The angle of incidence and the angle of reflection are equal  $\angle i = \angle r$



In vector form  $\hat{\mathbf{r}} = \hat{\mathbf{e}} - 2(\hat{\mathbf{e}} \cdot \hat{\mathbf{n}})\hat{\mathbf{n}}$ 

#### Object

<u>Real:</u> Point from which rays actually diverge.

Virtual: Point towards which rays appear to converge

#### Image:

Image is decided by reflected or refracted rays only. The point image for a mirror is that point towards which the rays reflected from the mirror, actually converge (real image).

#### OR

From which the reflected rays appear to diverge (virtual image).

# CHARACTERISTICS OF REFLECTION BY A PLANE MIRROR

The size of the image is the same as that of the object.

For a real object the image is virtual and for a virtual object the image is real. For a fixed incident light ray, if the mirror be rotated through an angle  $\theta$  the reflected ray turns through an angle  $2\theta$  in the same sense.

## Number of images (n) in inclined mirror Find $\frac{360}{4} = m$

- If m is even, then n = m 1, for all positions of object.
- If m is odd, then n = m, if object is not on bisector and n = m - 1, If object at bisector
- If m is fraction then n = nearest even number

# Spherical Mirrors

convex mirror

**Mirror Formula:**  $\frac{1}{f} = \frac{1}{v} + \frac{1}{u}$ . f = x-coordinate of focus u = x-coordinate of object v = x-coordinate of image **Note:** Valid only for paraxial rays. **Transverse Magnification:**  $m = \frac{h_2}{h_1} = -\frac{v}{u}$ 

 $h_2 = y$  co-ordinate of image  $h_1 = y$  co-ordinate of the object (both perpendicular to the principal axis of mirror) Longitudinal magnification:  $m_2 = \frac{\text{Length of image}}{\text{Length of object}}$ 

for small object  $m_2 = -m_t^2 m_t$  = transverse magnification.

Velocity of image of Moving Object (Spherical Mirror)

Velocity component along axis (Longitudinal velocity)



When an object is coming from infinite towards the focus of concave mirror

$$:: \frac{1}{v} + \frac{1}{u} = \frac{1}{f} \qquad :: -\frac{1}{v^2} \frac{dv}{dt} - \frac{1}{u^2} \frac{du}{dt} = 0 \Longrightarrow \vec{v}_{IM} = -\frac{v^2}{u^2} \vec{v}_{OM} = -m^2 \vec{v}_{OM}$$

- $V_{IM} = \frac{dv}{dt}$  = velocity of image with respect to mirror
- $V_{OM} = \frac{du}{dt}$  = velocity of object with respect to mirror.

#### Newton's Formula:

Applicable to a pair of real object and real image position only. They are called conjugate positions or foci,  $X_1$ ,  $X_2$  are the distance along the principal axis of the real object and real image respectively from the principal focus

$$X_1X_2 = f^2$$

**Optical Power:** Optical power of a mirror (in Diopters) =  $-\frac{1}{f}$  where f = focal length (in meters) with sign.

#### **REFRACTION-PLANE SURFACE**

#### Laws of Refraction (At any Refracting Surface)

(i) Incident ray, refracted ray and normal always lie in the same plane.



In vector form  $(\hat{e} \times \hat{n}) \cdot \hat{r} = 0$ 

(ii) The product of refractive index and sine of angle of incidence at a point in a medium is constant.

 $\mu_1 \sin i = \mu_2 \sin r$  (Snell's law)

#### Snell's Law

$$\frac{\sin i}{\sin r} = {}_1n_2 = \frac{n_2}{n_1} = \frac{v_1}{v_2} = \frac{\lambda_1}{\lambda_2}$$

In vector form  $\mu_1 |\hat{\mathbf{e}} \times \hat{\mathbf{n}}| = \mu_2 |\hat{\mathbf{r}} \times \hat{\mathbf{n}}|$ 

**Note:** Frequency of light does not change during refraction.

#### Deviation of a Ray due to refraction



angle of deviation,  $\delta = i - r$  (clockwise)

#### **Refraction Through a Parallel Slab**

Emerged ray is parallel to the incident ray, if medium is same on both sides.



Lateral shift  $x = \frac{t \sin(i-r)}{\cos r}$ ; t = thickness of slab **Note:** Emergent ray will not be parallel to the incidence of the state of

**Note:** Emergent ray will not be parallel to the incident ray if the medium on both the sides are different.

#### Apparent Depth of Submerged Object: (h' < h)



 $\Delta x$  = Apparent shift = t $\left(1 - \frac{1}{\mu}\right)$  always in direction of incident ray. Note: h and h' are always measured from surface. Critical Angle & Total Internal Reflection (TIR)



#### **Conditions of TIR**

- Ray is going from denser to rarer medium
- Angle of incidence should be greater than the critical angle (i > C).

Critical angle C =  $\sin^{-1}\frac{\mu_R}{\mu_D} = \sin^{-1}\frac{V_D}{V_R} = \sin^{-1}\frac{\lambda_D}{\lambda_R}$ 

#### **Refraction Through Prism:**



- 1 1 A
- $\delta = i + i' A$

Variation of  $\delta$  versus i



There is one and only one angle of incidence for which the angle of deviation is minimum. When δ = δ<sub>min</sub> then i = i' and r = r', the ray passes symmetrically about the prism, & then n = sin[A+δm<sup>-</sup>/2]/sin[A/2], where n = absolute R.I. of glass.

**Note:** When the prism is dipped in a medium then n = R.I. of glass w.r.t. medium.

- For a thin prism (A  $\leq 10^{\circ}$ );  $\delta = (n 1)A$
- **Dispersion of Light:** The angular splitting of a ray of white light into a number of components when it is refracted in a medium other than air is called Dispersion of Light.
- Angle of Dispersion: Angle between the rays of the extreme colours in the refracted (dispersed) light is called Angle of Dispersion.  $\theta = \delta_v \delta_r$
- Dispersive power ( $\omega$ ) of the medium of the material of prism.

 $\omega = \frac{\text{angular dispersion}}{\text{deviation of mean ray (yellow)}}$ 

٠



For small angled prism (A  $\leq 10^{\circ}$ );  $\boldsymbol{\omega} = \frac{\delta_v - \delta_R}{\delta_y} = \frac{n_v - n_R}{n-1}$ ;  $\mathbf{n} = \frac{n_v + n_R}{2}$ 

 $n_{v},\ n_{R}$  & n are R.I. of material for violet, red & yellow colours respectively.

#### **Refraction At Spherical Surface**

(A)  $\frac{\mu_2}{v} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R}$ v, u & R are to be kept with sign as v = PI u = -PO R = PC

(B)  $m = \frac{\mu_1 v}{\mu_2 u}$ 

#### Lens Formula

(A)



(B) 
$$\frac{1}{f} = (\mu - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right)$$

(C) 
$$m = \frac{v}{u}$$

#### **Power of Lenses**

Reciprocal of focal length in meter is known as power of lens. *SI unit:* dioptre (D)

Power of lens:  $P = \frac{1}{f(m)} = \frac{100}{f(m)} = \frac{100}{f(cm)}$  dioptre

#### **Combination of Lenses**

<u>Two thin lenses are placed in contact to each other</u> Power of combination.  $P = P_1 + P_2 \implies \frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2}$ Use sign convention while solving numerical.

#### Newton's Formula



$$x_1$$
 = distance of object from focus;

 $x_2$  = distance of image from focus.

#### Silvering of one surface of lens (use $P_{eq} = 2P_l + P_m$ )

• When plane surface is silvered

$$f = \frac{R}{2(\mu - 1)}$$

NI

• When convex surface is slivered



#### **OPTICAL INSTRUMENTS**

#### For Simple microscope

- Magnifying power when image is formed at D: M = 1 + D/f
- When image is formed at infinity M = D/f

#### For Compound microscope

- When image is formed at infinity  $M = -\frac{v_0}{u_0} \left( \frac{D}{u_e} \right)$
- Magnifying power when final image is formed at D,

$$\mathbf{M} = -\frac{\mathbf{v}_0}{\mathbf{u}_0} \left( 1 + \frac{\mathbf{D}}{\mathbf{f}_e} \right)$$
$$\mathbf{L} = \mathbf{v}_0 + |\mathbf{u}_0|$$

- Tube length  $L = v_0 + |u_e|$
- When final images is formed at infinity  $M = -\frac{v_0}{u_0} \times \frac{D}{f_e}$ And  $L = v_0 \times f_e$

# Astronomical Telescope: $M = -\frac{f_0}{u_a}$

• Magnifying power when final image is formed at D:

$$M = \frac{f_0}{f_e} \left( 1 + \frac{D}{f_e} \right)$$

- Tube length:  $L = f_0 + |u_e|$
- When final image is formed at infinity:  $M = \frac{f_0}{f_0} \& L = f_0 + f_e$

# **Limit of resolution for microscope:** $\frac{1.22 \lambda}{2a \sin \theta} = \frac{1}{\text{Resolving power}}$

**Limit of resolution for telescope:**  $\frac{1.22 \lambda}{a} = \frac{1}{\text{Resolving power}}$