IIT CHEMISTRY PHYSICAL CHEMISTRY

MOLE CONCEPT

BASICS OF CHEMISTRY

1. CLASSIFICATION OF MATTER

1.1 Physical classification: It is based on physical state under ordinary conditions of temperature and pressure, matter is classified into the following three types:
 (a) Solid (b) Liquid (c) Gas

(a) Solid: A substance is said to be solid if it possesses a definite volume and a definite shapeE.g. sugar, iron, gold, wood etc.

(b) Liquid: A substance is said to be liquid if it possesses a definite volume but not definite shape. They take up the shape of the vessel in which they are put.

E.g. water, milk, oil, mercury, alcohol etc.

(c) Gas: A substance is said to be gas if it neither possesses a definite volume nor a definite shape. This is because they fill up the whole vessel in which they are put.

E.g. hydrogen(H₂), oxygen(O₂), carbon dioxide(CO₂), etc.

1.2 Chemical classification:



On the basis of chemical nature matter is classified into the following two types :

(a) Pure Substance (b) Mixture

- (a) **Pure Substance:** When all constituent particles of a substance are same in chemical nature, it is said to be a pure substance. Two type of pure substances:
 - (i) Element (ii) Compound

(i) **Element:** an element consist of only one type of atoms. These particles may exist as Atoms or molecules. E.g. O_2 , P_4 , S_8 , etc.

- (ii) **Compound:** When two or more atoms of different elements combine together in a definite ratio. E.g. water, ammonia, carbon dioxide, sugar, etc.
- (b) Mixture: A mixture contains many types of particles. A mixture contains particles of two or more pure substances which may be present in it in any ratio. Hence, their composition is variable. Pure substances forming mixture are called its components. Many of the substances present around you are mixtures. For example, sugar solution in water, air, tea, etc., are all mixtures. Two types of mixtures:
 - (i) Homogeneous mixture (ii) Heterogeneous mixture

- (i) **Homogeneous mixture:** The components completely mix with each other. This means particles of components of the mixture are uniformly distributed throughout the bulk of the mixture and its composition is uniform throughout. E.g.: Sugar solution, air
- (ii) Heterogeneous mixture: In a heterogeneous mixture, the composition is not uniform throughout and sometimes different components are visible. E.g.: mixtures of salt and sugar, grains and pulses along with some dirt (often stone pieces)

2. INTRODUCTION:

There are a large number of objects around us which we can see and feel. It was John Dalton who firstly developed a theory on the structure of matter, later on which is known as Dalton's atomic theory.

2.1. DALTON'S ATOMIC THEORY:

- (i) Each element is composed of extremely small particles called atoms which can take part in chemical combination.
- (ii) All atoms of a given element are identical i.e., atoms of a particular element are all alike but differ from atoms of other elements.
- (iii) Atoms of different elements possess different properties (including different masses).
- (iv) Atoms are indestructible i.e., atoms are neither created nor destroyed in chemical reactions.
- (v) Atoms of elements combine to form molecules and compounds are formed when atoms of more than one element combine.
- (vi) In a given compound, the relative number and kind of atoms is constant.
- **2.2.** Atomic mass: It is the average relative mass of atom of element as compared with times the mass of an atom of carbon-12 isotope.

Atomic mass = $\frac{\text{Average mass of an atom}}{1/12 \times \text{Mass of an atom of C}^{12}}$

2.3. Gram atomic mass (GAM): Atomic mass of an element expressed in grams is called Gram atomic mass or gram atom or mole atom.

(i) Number of gram atoms =
$$\frac{\text{Mass of an element}}{\text{GAM}}$$

- (ii) Mass of an element in g = No. of gram atoms \times GAM
- (iii) Number of atoms in 1 GAM = 6.02×10^{23} Number of atoms in a given substance = No. of gram atoms $\times 6.02 \times 10^{23}$

(iv) Number of atoms in 1 g of element =
$$\frac{Mass}{GAM}$$

(v) Mass of one atom of the element (in g) = $\frac{\text{GAM}}{6.02 \times 10^{23}}$

2.4. Molecular mass: Molecular mass of a molecule, of an element or a compound may be defined as a number which indicates how many times heavier is a molecule of that element or compound as compared with $\frac{1}{12}$ of the mass of an atom of carbon-12. Molecular mass is also

expressed in a.m.u.

Molecular mass = $\frac{\text{Mass of one molecule of the substance}}{1/12 \times \text{Mass of one atom of C-12}}$

Actual mass of one molecule = Mol. mass (in amu) $\times 1.66 \times 10^{-24}$ g Molecular mass of a substance is the additive property and can be calculated by adding the atomic masses of atoms present in one molecule.

2.5. Gram molecular mass (GMM) : Molecular mass of an element or compound when expressed in g is called its gram molecular mass, gram molecule or mole molecule.

Number of gram molecules = $\frac{\text{Mass of substance}}{\text{GMM}}$

		011111		
Mass of substance	in $g = No$.	of gram	molecules >	< GMM

Element	R.A.M. (Relative Atomic Mass)	Mass of one atom	Gram Atomic mass /weight
Ν	14	14 amu	14 gm
He	4	4 amu	4 gm
С	12	12 amu	12 gm

2.6. Mole

One mole of any substance contains a fixed number (6.023×10^{23}) of any type of particles (atoms or molecules or ions) and has a mass equal to the atomic or molecular weight, in grams. Thus it is correct to refer to a mole of helium, a mole of electrons or a mole of any ion, meaning respectively Avogadro's number of atoms, electrons or ions.

Methods of Calculations of mole:

(1) If no. of some species is given, then no. of moles = $\frac{\text{Given no}}{N_{A}}$

(2) If weight of a given species is given, then no. of moles = $\frac{\text{Given wt.}}{\text{Atomic wt.}}$ (for atoms),

or =
$$\frac{\text{Given wt.}}{\text{Molecular wt.}}$$
 (for molecules)

(3) If volume of a gas is given along with its temperature (T) and pressure (P).

PV = nRT (n : Number of moles of gas)

P (Pressure of gas): Pressure of the gas is the force exerted by the gas per unit area of the walls of the container in all directions.

Thus, Pressure (P) = $\frac{\text{Force}(F)}{\text{Area}(A)} = \frac{\text{Mass}(m) \times \text{Acceleration}(a)}{\text{Area}(A)}$

Name	Symbol	Value
Bar	bar	$1 \text{ bar} = 10^5 \text{ Pa}$
Atmosphere	atm	$1 \text{ atm} = 1.01325 \times 10^5 \text{ Pa}$
Torr	Torr	1 Torr = $\frac{101325}{760}$ Pa × 133.322 Pa
millimeter of mercury	mm Hg	1 mm Hg = 133.322 Pa

V (Volume of gas): Volume is expressed in liters (L), milliliters (mL) or cubic centimeters (cm^3) , cubic meters (m^3) .

 $1 \text{ m}^3 = 10^3 \text{ dm}^3 = 10^6 \text{ cm}^3 = 10^6 \text{ mL} = 10^3 \text{ L}$

T (Temperature of gas): S.I. unit of temperature is Kelvin (K).

 $T(K) = t(^{\circ}C) + 273.15$ R (Universal gas constant) : Values of $R = 0.082 \text{ Latm} \text{K}^{-1} \text{mol}^{-1}$ =8.314 JK⁻¹mol⁻¹ $=1.987 \text{ CalK}^{-1} \text{mol}^{-1}$

- Note: 1 mole of atom is also termed as 1 g-atom 1 mole of ions is also termed as 1 g-ion 1 mole of molecule is also termed as 1 g-molecule
- How much time (in years) would it take to distribute one Avogadro number of wheat grains if *Ex.1* 10¹⁰ grains are distributed each second?
- Ans.
- 1.9×10^6 years (approx) 10^{10} grains are distributed in 1 second Sol.

 6.02×10^{23} grains are distributed in $\frac{6.02 \times 10^{23}}{10^{10}}$ sec $=\frac{6.02 \times 10^{23}}{10^{10} \times 60 \times 60 \times 24 \times 365}$ years $= 1.9 \times 10^6$ years (approx.)

Ex.2How many atoms are there in 100 amu of He?

Ans. 25

We know that, 1 amu = $\frac{1}{12}$ × weight of one ¹²C atom Sol. or weight of one ¹²C atom = 12 amu (at. wt. of C = 12 amu). Similarly, as the atomic weight of He is 4 weight of one He atom = 4 amu.

Thus, the number of atoms in 100 amu of He = $\frac{100}{\Lambda}$ = 25.

- Ex.3The weight of one atom of Uranium is 238 amu. Its actual weight in gm is _____:
- 396.74×10^{-4} Ans.
- Sol. Weight of aone atom = 238 amu

$$= 238 \times 1.667 \times 10^{-24}$$
$$= 396.74 \times 10^{-4}$$

Ex.4 Calculate the number of molecules in a drop of water weighing 0.09 g.

- Ans. 3.01×10^{21} molecules of H₂O Sol. number of mole = $\frac{0.09}{18}$ so number of molecules = $\frac{0.09}{18} \times N_A = = 3.01 \times 10^{21}$.
- *Ex.5* A sample of ethane has the same mass as 10.0 million molecules of methane. How many C_2H_6 molecules does the sample contain?

Ans. 5.34×10^6

Sol. Let the number of C_2H_6 molecules in the sample be n. As given, mass of C_2H_6 = mass of 10^7 molecules of CH_4

$$\frac{n}{N_A} \times \text{mol. wt. of } C_2 H_6 = \frac{10'}{N_A} \times \text{mol. wt. of } CH_4$$
$$\frac{n}{N_A} \times 30 = \frac{10^7}{N_A} \times 16$$
$$n = 5.34 \times 10^6.$$

Ex.6 If, from 10 moles NH_3 and 5 moles of H_2SO_4 , all the H-atoms are removed in order to form H_2 gas, then find the number of H_2 molecules formed.

Ans. 20 N_A

Sol. 10 mole NH₃ have mole of 'H' atom = 10×3 5 mole of H₂SO₄ have mole of 'H' atom = 10 Total mole of 'H' atom = 40 mole of H₂ = 20

DI

- *Ex.7* The weight of 350 mL of a diatomic gas at 0°C and 2 atm pressure is 1 g. The weight of one atom is :
- *Ans.* 16 amu

Sol.
$$PV = nRT;$$
 $n = \frac{PV}{RT}$
 $n = \frac{2 \times 0.350}{0.0821 \times 273} =$
 $n = \frac{Weight}{Atomic mass}$
Atomic mass = 16 amu

- *Ex.8* Oxygen is present in a 1-litre flask at a pressure of 7.6×10^{-10} mm of Hg at 0°C. Calculate the number of oxygen molecules in the flask.
- Ans. 0.44×10^{-13} . Sol. Pressure = 7.6×10^{-10} mm = 0.76×10^{-10} cm = atm (1 atom = 76 cm) = 10^{-12} atm. Volume = 1 litre, R = 0.0821 lit. atm/K/mole, temperature = 273 K. We know that PV = nRT or n = $\frac{pV}{RT}$ $n = \frac{10^{-12} \times 1}{0.082 \times 273} = 0.44 \times 10^{-13}$ Ex.9 Equal volumes of oxygen gas and a second gas weigh 1.00 and 2.375 grams respectively under the same experimental conditions. Which of the following is the unknown gas?
- Ex.9 Equal volumes of oxygen gas and a second gas weigh 1.00 and 2.375 grams respectively under the same experimental conditions. Which of the following is the unknown gas?
 (A) NO
 (B) SO₂
 (C) CS₂
 (D) CO
- Sol. Moles of O_2 = Moles of X(unknown gas) $\frac{1}{32} = \frac{2.375}{M_X}$ $M_X = 76$

Do yourself-1:

- **1.** The number of molecules in 16 g of methane is :
- 2. A sample of aluminium has a mass of 54.0 g. What is the mass of the same number of magnesium atoms? (At. wt. Al = 27, Mg = 24) (A) 12 g (B) 24 g (C) 48 g (D) 96 g
- **4.** The volume of a gas at 0° C and 700 mm pressure is 760 cc. The number of molecules present in this volume is:

5. Four 1-1 liter flasks are separately filled with the gases H_2 , He, O_2 and O_3 at the same temperature and pressure. The ratio of total number of atoms of these gases present in different flask would be:

- (A) 1:1:1:1 (B) 1:2:2:3 (C) 2:1:2:3 (D) 3:2:2:1
- 6. The weight of 2.01×10^{23} molecules of CO is-(A) 9.3 g (B) 7.2 g (C) 1.2 g (D) 3 g
- 7. How many moles of e⁻ weight one Kg : (A) 6.023×10^{23} (B) $\frac{1}{9.108} \times 10^{31}$ (C) $\frac{6.023}{9.108} \times 10^{54}$ (D) $\frac{1}{9.108 \times 6.023} \times 10^{8}$

3. LAWS OF CHEMICAL COMBINATION:

Antoine Lavoisier, John Dalton and other scientists formulate certain laws concerning the composition of matter and chemical reactions. These laws are known as the law of chemical combination.

3.1 THE LAW OF CONSERVATION OF MASS:

It is given by Antoine Lavoisier. In a chemical change total mass remains conserved that is mass before reaction is always equal to mass after reaction. Example:

 $H_{2}(g) + \frac{1}{2}O_{2}(g) \rightarrow H_{2}O(\ell)$ Before reaction initially $1 \mod \frac{1}{2} \mod R$ After the reaction $0 \quad 0 \quad 1 \mod R$ mass before reaction = mass of 1 mole $H_{2}(g) + \max \inf \frac{1}{2} \mod O_{2}(g)$ = 2 + 16 = 18 gmass after reaction = mass of 1 mole water = 18 g

3.2 LAW OF CONSTANT OR DEFINITE PROPORTION:

It is given by Proust. All chemical compounds are found to have constant composition irrespective of their method of preparation or sources.

- **Example:** In water (H_2O), Hydrogen and Oxygen combine in 2 : 1 molar ratio, the ratio remains constant whether it is tap water, river water or sea water or produced by any chemical reaction.
- *Ex.10* 1.80 g of a certain metal burnt in oxygen gave 3.0 g of its oxide. 1.50 g of the same metal heated insteam gave 2.50 g of its oxide. Show that these results illustrate the law of constant proportion.
- *Sol.* In the first sample of the oxide,

wt. of metal = 1.80 g,
wt. of oxygen =
$$(3.0 - 1.80)$$
 g = 1.2 g
 $\frac{\text{wt. of metal}}{\text{wt. of oxygen}} = \frac{1.80\text{g}}{1.2\text{g}} = 1.5$

In the second sample of the oxide,

 $\frac{\text{wt. of metal}}{\text{wt. of oxygen}} = \frac{1.50\text{g}}{1\text{g}} = 1.5$

Thus, in both samples of the oxide the proportions of the weights of the metal and oxygen are fixed. Hence, the results follow the law of constant proportion.

3.3 THE LAW OF MULTIPLE PROPORTIONS:

It is given by Dalton.

When one element combines with the other element to form two or more different compounds, the mass of one element, which combines with a constant mass of the other, bear a simple ratio to one another.

Note: Simple ratio here means the ratio between small natural numbers, such as 1: 1, 1: 2, 1: 3, Later on this simple ratio becomes the valency and then oxidation state of the element.

Example: Carbon and Oxygen when combine, can form two oxides, CO (carbon monoxide), CO₂(Carbon dioxides)

In CO, 12 g carbon combined with 16 g of oxygen.

In CO₂, 12 g carbon combined with 32 g of oxygen.

Thus, we can see the mass of oxygen which combine with a constant mass of carbon (12 g) bear simple ratio of 16:32 or 1:2.

3.4 LAW OF RECIPROCAL PROPORTION:

It is given by Richter.

The ratio of the weights of two elements A and B which combine separately with a fixed weight of the third element C is either the same or simple ratio of the weights in which A and B combine directly with each other.

Example:



- Ex.11 Ammonia contains 82.35% of nitrogen and 17.65% of hydrogen. Water contains 88.90% of oxygen and11.10% of hydrogen. Nitrogen trioxide contains 63.15% of oxygen and 36.85% of nitrogen. Show that these data illustrate the law of reciprocal proportions.
- Sol. In NH₃, 17.65g of H combine with N = 82.35g

1 g of H combine with N =
$$\frac{82.35}{17.65}$$
g = 4.67 g
In H₂O, 11.10 g of H combine with O = 88.90 g
1 g of H combine with O = $\frac{88.90}{11.10}$ g = 8.01g
Ratio of the weights of N and O which combine with fixed weight (=1g) of H
= 4.67 : 8.01 = 1 : 1.7
In N₂O₃, ratio of weights of N and O which combine with each other = 36.85 : 63.15
= 1 : 1.7
Thus the two ratios are the same. Hence it illustrates the law of reciprocal proportions.

3.5 GAY-LUSSAC'S LAW OF COMBINING VOLUME:

According to him elements combine in a simple ratio of atoms, gases combine in a simple ratio of their volumes provided all measurements should be done in the same temperature and pressure

$$\begin{array}{l} H_2(g) + \operatorname{Cl}_2(g) \longrightarrow 2H\operatorname{Cl}\\ 1 \text{ vol} \quad 1 \text{ vol} \quad 2 \text{ vol} \end{array}$$

3.6 AVOGADRO'S H YPOTHESIS:

Equal volumes of polyatomic all gases have equal number of molecules (not atoms) at same temperature and pressures conditions.

S.T.P. (Standard Temperature and Pressure)

At S.T.P. condition:

Temperature = 0° C or 273 K

Pressure = 1 bar

and volume of one mole of gas at STP is found to be equal to 22.7litres which is known as molar volume.

4. STOICHIOMETRY AND STOICHIOMETRIC CALCULATIONS

The word 'stoichiometry' is derived from two Greek words — *stoicheion*(meaning, *element*) and *metron*(meaning, *measure*).Stoichiometry, thus, deals with the calculation of masses (sometimes volumes also) of the reactants and the products involved in a chemical reaction

Example: When potassium chlorate (KClO₃) is heated it gives potassium chloride (KCl) and oxygen (O_2) .

 $\text{KClO}_3 \xrightarrow{\Delta} \text{KCl} + \text{O}_2$ (unbalanced chemical equation)

 $2\text{KClO}_3 \xrightarrow{\Delta} 2 \text{KCl} + 3\text{O}_2$ (balanced chemical equation)

• Remember a balanced chemical equation is one which contains an equal number atoms of each element on both sides of equation.

4.1 Interpretation of balanced chemical equations:

Once we get a balanced chemical equation then we can interpret a chemical equation by following ways

- (a) Mass mass analysis
- (b) Mass volume analysis
- (c) Mole mole analysis

(a) Mole - mole analysis :

This analysis is very much important for quantitative analysis point of view.

Now consider again the decomposition of KClO₃.

 $2 \text{ KClO}_3 \longrightarrow 2 \text{ KCl} + 3\text{O}_2$

In very first step of mole-mole analysis you should read the balanced chemical equation like 2 moles $KClO_3$ on decomposition gives you 2 moles KCl and 3 moles O_2 and from the stoichiometry of reaction we can write

Moles of $KClO_3$	Moles of KCl	Moles of O_2			
2	2	3			
Now for any general	balance chemi	cal equation like			
$a A + b B \longrightarrow c C$	+ d D				
you can write.					
Moles of A reacted	Moles of Br	eacted _ Moles of	of C produced	_ Moles of	D produced
a	b		с	_	d
Now for any general a A + b B \longrightarrow c C you can write. Moles of A reacted a	balance chemic + d D $\frac{1}{d} = \frac{\text{Moles of B r}}{b}$	$\frac{\text{eacted}}{\text{eacted}} = \frac{\text{Moles of }}{\text{Moles of }}$	$\frac{1}{c}$	_ Moles of	D produced

(b) Mass - mass analysis :

Consider the reaction 2 KClO₃ \longrightarrow 2KCl + 3O₂ According to stoichiometry of the reaction

or $\frac{\text{Mass of KClO}_3}{\text{Mass of KCl}} = \frac{2 \times 122.5}{2 \times 74.5} \qquad \frac{\text{Mass of KClO}_3}{\text{Mass of O}_2} = \frac{2 \times 122.5}{3 \times 32}$

Ex.12 Consider the balanced reaction

 $2\text{Cl}_2\text{O}_7 \longrightarrow 4\text{ClO}_2 + 3\text{O}_2 \qquad (\text{Cl} = 35.5)$

What can be concluded from the coefficients of species in this balanced equation?

(A) For this reaction, exactly 2 g of Cl_2O_7 must be taken to start the reaction

(B) For this reaction, exactly 2 mol of Cl₂O₇ must be taken to start the reaction

(C) Mole ratio of Cl_2O_7 , ClO_2 and O_2 during a chemical reaction at any instant are 2, 4 and 3 respectively

(D) The ratio of change in number of moles of Cl_2O_7 , ClO_2 and O_2 is 2:4:3

- *Sol.* It follows directly from definition of stoichiometry.
- *Ex.13* Calculate the weight of iron which will be converted into its oxide by the action of 36 g of steam.

 $(Given: 3Fe + 4H_2O \longrightarrow Fe_3O_4 + H_2)$

- *Ans.* 84 g
- Sol. Mole ratio of reaction suggests,

Mole of Fe =
$$\frac{3}{4}$$
 mol of H₂O
= $\frac{3}{4} \times \frac{36}{18} = \frac{3}{2}$
wt. of Fe = $\frac{3}{2} \times 56 = 84$ g

Ex.14 When Dinitrogen pentaoxide (N_2O_5 , a white solid) is heated, it decomposes into nitrogen dioxide and oxygen.

If a sample of N_2O_5 produces 1.6 g O_2 , then how many grams of NO_2 are formed ?

 $N_2O_5(s) \longrightarrow NO_2(g) + O_2(g)$ (not balanced) (A) 9.2 g (B) 4.6 g (C) 2.3 g (D) 18.4 g (A) Ans. $N_2O_5(s) \longrightarrow 2NO_2(s) + \frac{1}{2}O_2$ (Balanced reaction) Sol. $\frac{\text{Mole of O}_2}{1/2} = \frac{\text{Mole of NO}_2}{2}$ $\frac{1.6}{32} \times 2 \times 2 =$ Mole of NO₂ = 0.2 wt of $NO_2 = 0.2 \times 46 = 9.2$ g.

Mass - volume analysis : (c)

Sol.

Now again consider decomposition of KClO₃

$$2 \text{ KClO}_3 \longrightarrow 2 \text{ KCl} + 3 \text{O}_2$$

mass volume ratio : 2×122.5 g : 2×74.5 g : 3×22.4 L at STP we can use two relation for volume of oxygen

$$\frac{\text{Mass of KCIO}_3}{\text{volume of O}_2 \text{at STP}} = \frac{2 \times 122.5\text{g}}{3 \times 22.4\text{L}} \qquad \dots (i)$$

and
$$\frac{\text{Mass of KCl}}{\text{volume of O}_2 \text{ at STP}} = \frac{2 \times 74.5\text{ g}}{3 \times 22.4\text{L}} \qquad \dots (ii)$$

Ex.15 When oxygen gas is passed through Siemen's ozoniser, it completely gets converted into ozone gas. The volume of ozone gas produced at 1 atm and 273K, if initially 96 g of oxygen gas was taken, is :

(A) 44.8 L (C) 67.2 L (B) 89.6 L (D) 22.4 L Ans. (A) $3O_2 \longrightarrow 2O_3$ Mole = $\frac{96}{32} = 3$ mole = 2

Volume of O₃ gas at 1 atm and $273K = 2 \times 22.4 = 44.8 L$

Do yourself-2:

Assuming 100% yield of the reaction, how many moles of NaHCO₃ will produce 448 mL of 1. CO₂ gas at STP according to the reaction : NaHCO₃ \longrightarrow Na₂CO₃ + CO₂ + H₂O (unbalanced) (A) 0.04 (B) 0.4 (C) 4 (D) 40 2. Calculate the residue obtained on strongly heating 2.76 g Ag₂CO₃. $Ag_2CO_3 \xrightarrow{\Delta} 2Ag + CO_2 + \frac{1}{2}O_2$ 3. For the reaction $2P + Q \longrightarrow R$, 4 mol of P and excess of Q will produce : (A) 8 mol of R (B) $5 \mod \text{of } R$ (C) $2 \mod \text{of } R$ (D) $1 \mod \text{of } R$ 4. If 1.5 moles of oxygen combine with Al to form Al₂O₃, the weight of Al used in the reaction is: (A) 27 g (B) 40.5 g (C) 54g (D) 81 g 5. How many liters of CO₂ at STP will be formed when 0.01 mol of H₂SO₄ reacts with excess of Na_2CO_3 . $Na_2CO_3 + H_2SO_4 \longrightarrow Na_2SO_4 + CO_2 + H_2O$ (A) 22.7 L (B) 2.27 L (C) 0.227 L (D) 1.135 L

4.2 LIMITING REAGENT:

Many a time, reactions are carried out with the amounts of reactants that are different than the amounts as required by a balanced chemical reaction. In such situations, one reactant is in more amount than the amount required by balanced chemical reaction. The reactant which is present in the least amount gets consumed after sometime and after that further reaction does not take place whatever be the amount of the other reactant. Hence, the reactant, which gets consumed first, limits the amount of product formed and is, therefore, called the limiting reagent.

The reactant which consumed first into the reaction when we are dealing with balance chemical equation then if number of moles of reactants are not in the ratio of stoichiometric coefficient of balanced chemical equation, then there should be one reactant which should be limiting reactant.

Ex.16 Six mole of Na_2CO_3 is reacted with 4 moles of HCl solution. Find the volume of CO_2 gas produced at STP. The reaction is

$$Na_2CO_3 + 2 HCl \longrightarrow 2 NaCl + CO_2 + H_2O$$

Sol.From the reaction : $Na_2CO_3 + 2 HCl \longrightarrow 2 NaCl + CO_2 + H_2O$ gives moles3 mol6 molgiven mole ratio1:Stoichiometric coefficient ratio1:

See here given number of moles of reactants are not in stoichiometric coefficient ratio. Therefore there should be one reactant which consumed first and becomes limiting reagent. But the question is how to find which reactant is limiting, it is not very difficult you can easily

find it. According to the following method.

How t	o find limiting reagent:						
	Step : I						
	Divide the given moles of reactant by the respective stoich	iometric coefficient of that reactant.					
	Step : II	The resident having minimum					
	See for which reactant this division comes out to be minimum. The reactant having minimum						
	Ston • III						
	Now once you find limiting reagent then your focus should	be on limiting reagent					
	From Step I & II Na ₂ CO ₃	HCl					
	6	4					
	$\frac{-}{1} = 0$	$\frac{-}{2} = 2$ (Division in minimum)					
	∴ HCl is limiting reagent From Step III						
	Mole of HCl Mole of CO ₂ produced						
	From $\frac{2}{2} = \frac{1}{1}$						
	\therefore Mole of CO ₂ produced = 2 moles						
	\therefore Volume of CO ₂ produced at S.T.P. = 2 × 22.7 = 45.	.4 L					
	-						
Ex.17 Ans.	In the reaction $4A + 2B + 3C \longrightarrow A_4 B_2 C_3$ what will formed, starting from 2 moles of A, 1.2 moles of B & 1.44 (A) 0.5 (B) 0.6 (C) 0.48 (C)	be the number of moles of product moles of C : (D) 4.64					
Sol.	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$						
Ex.18	A 5 g mixture of SO ₂ and O ₂ gases is reacted to form SO ₃ g SO ₂ and O ₂ gases in mixture to obtain maximum amount of	gas. What should be the mass ratio of f SO3gas :					
	(A) 4 : 1 (B) 3 : 2 (C) 2 : 3	(D) 1 : 4					
Ans.	(A)						
Sol.	For maximum amount of product, the reactants should be p	present in their stoichiometric ratio.					
	$2SO_2(g) + O_2(g) \longrightarrow 2SO_3(g)$						
	mass x $5-x$						
	mole $\frac{x}{64} = \frac{5-x}{32}$						
	So, $\frac{\left(\frac{x}{64}\right)}{\left(\frac{5-x}{32}\right)} = 2:1$						
	Therefore, $x = 4$						
	$m_{SO2}: m_{O2} = 4: 1.$						

Ex.19 Calculate the weight of FeO from 4 g VO and 5.75 g of Fe₂O₃. Also report the limiting reactant. $VO + Fe_2O_3 \longrightarrow FeO + V_2O_5$ Weight of FeO formed = 5.17 g Ans. Sol. **Balanced** equation 2VO + $3Fe_2O_3 \longrightarrow 6FeO +$ V_2O_5 4 5.75 0 Moles before reaction 0 67 160 0.03590 0.5970 = $\left(\frac{6}{5} \times 0.0359\right) \left(\frac{1}{3} \times 0.0359\right)$ Moles after reaction (0.05970 - 0.0359)0 As 2 moles of VO react with 3 moles of Fe_2O_3 :. 0.05970 g moles of VO = $\frac{3}{2}$ 0.05970 = 0.08955 moles of Fe₂O₃ Moles of Fe_2O_3 available = 0.0359 only Hence, Fe_2O_3 is the limiting reagent. Moles of FeO formed = $\frac{6}{3}$ 0.0359 Weight of FeO formed = $0.0359 \times 2 \times 72 = 5.17$ g *.*.. $\left(\frac{n_{\text{FeO}}}{n_{\text{FeO}}} = \frac{6}{3}\right) \Longrightarrow n_{\text{FeO}} = \frac{6}{3} \times n_{\text{Fe}_2O_3}$

Do yourself-3:

1.	The reaction						
	$2C + O_2 \longrightarrow 2C$	$2C + O_2 \longrightarrow 2CO$					
	is carried out by taking 24 g of carbon and 128 g of O_2 .						
	Find out :						
	(i) Which reactan	t is left in excess?					
	(ii) How much of	it is left?					
	(iii) How many m	oles of CO are forme	d?				
	(iv) How many gra	ams of other reactant sh	hould be taken so that no	thing is left at the end of reaction?			
2.	How many mole	of Zn(FeS ₂) can be m	ade from 2 mole zinc,	3 mole iron and 5 mole sulphur.			
	(A) 2 mole	(B) 3 mole	(C) 4 mole	(D) 5 mole			
3.	Calculate the amo	ount of Ni needed in the	he Mond's process give	en below			
	Ni + 4CO	\longrightarrow Ni(CO) ₄					
	If CO used in this process is obtained through a process, in which 6 g of carbon is mixed wit						
	44 g CO ₂ . (Ni = 5	(9 [°] u)					
	(A) 14.675 g	(B) 29 g	(C) 58 g	(D) 28 g			
4.	0.5 mole of H ₂ SO	D_4 is mixed with 0.2	mole of Ca (OH) ₂ . Th	he maximum number of moles of			
	CaSO ₄ formed is	:					
	(A) 0.2	(B) 0.5	(C) 0.4	(D) 1.5			
5.	The mass of Na ₂ S	O ₄ produced from 19	6 gram of H_2SO_4 and 1	mole of NaOH.			
	(A) 49 g	(B) 98 g	(C) 61 g	(D) 34.3 g			

4.3 **PRINCIPLE OF ATOM CONSERVATION (POAC):**

Infect POAC is nothing but the conservation of mass, expressed before in the concepts of atomic theory. And if atoms are conserved, moles of atoms shall also be conserved. The principle is fruitful for the students when they don't get the idea of balanced chemical equation in the problem.

This principle can be under stand by the following example.

Consider the decomposition of KClO₃ (s) \rightarrow KCl (s) + O₂ (g) (unbalanced chemical react ion) Apply the principle of atom conservation (POAC) for K atoms. Moles of K atoms in reactant = moles of K atoms in products or moles of K atoms in KClO₃ = moles of K atoms in KCl Now, since 1 molecule of KClO₃ contains 1 atom of K or 1 mole of KClO₃ contains 1 mole of K, similarly 1 mole of KCl contains 1 mole of K Thus, moles of K atoms in KClO₃ = 1 × moles of KClO₃ and moles of K atoms in KCl = 1 × moles of KCl \therefore moles of KClO₃ = moles of KCl or $\frac{\text{wt. of KClO_3ing}}{\text{mol. wt. of KClO_3}} = \frac{\text{wt. of KCl in g}}{\text{mol. wt. of KClO_3}} = \frac{\text{wt. of KClO_3}}{\text{mol. wt. of KC$

The above equation gives the mass-mass relationship between KClO₃ and KCl which is important in stoichiometric calculations.

Again, applying the principle of atom conservation for O atoms,

moles of O in $KClO_3 = 3 \times moles$ of $KClO_3$

moles of O in $O_2 = 2 \times \text{moles of } O_2$

$$\therefore \qquad 3 \times \text{moles of KClO}_3 = 2 \times \text{moles of O}_2$$

or
$$\frac{\text{wt. of KClO}_3}{\text{mol. wt. of KClO}_3} = 2 \times \frac{\text{vol. of O}_2 \text{ at NTP}}{\text{standard molar vol.}(22.4\text{lt})}$$

* The above equations thus give the mass-volume relationship of reactants and products.

Ex.20 27.6g K_2CO_3 was treated by a series of reagents so as to convert all of its carbon to K_2Zn_3 [Fe(CN)₆]₂. Calculate the weight of the product.

[mol. wt. of $K_2CO_3 = 138$ and mol. wt. of K_2Zn_3 [Fe(CN)₆]₂ = 698]

Ans. 11.6 g

*

Sol. Here we have not knowledge about series of chemical reactions but we known about initial reactant and final product accordingly

$$K_2CO_3 \xrightarrow{\text{Several}} K_2Zn_3 [Fe(CN)_6]_2$$

Since C atoms are conserved, applying POAC for C atoms, Moles of C in $K_2CO_3 =$ moles of C in K_2Zn_3 [Fe(CN)₆]₂ $1 \times$ moles of $K_2CO_3 = 12 \times$ moles of K_2Zn_3 [Fe(CN)₆]₂

(:: 1 mole of K_2CO_3 contains 1 moles of C)

 $\frac{\text{wt. of } \text{K}_2\text{CO}_3}{\text{mol. wt. of } \text{K}_2\text{CO}_3} = 12 \frac{\text{wt. of the product}}{\text{mol. wt. of product}}$ wt. of K₂Zn₃ [Fe(CN)₆]₂ = $\frac{27.6}{138} \frac{698}{12} = 11.6 \text{ g}$ *Ex.21* In a gravimetric determination of P of an aqueous solution of dihydrogen phosphate in $H_2PO_4^-$ is treated with a mixture of ammonium and magnesium ions to precipitate magnesium ammonium phosphate, Mg(NH₄)PO₄.6H₂O. This is heated and decomposed to magnesium pyrophosphate, Mg₂P₂O₇. A solution of $H_2PO_4^-$ yielded 2.054 g of (Mg₂P₂O₇). What weight of NaH₂PO₄ was present originally?

Ans. 2.22 g

Sol.
$$NaH_2PO_4 + Mg^{2+} + NH_4^+ + \longrightarrow Mg(NH_4)PO_4.6H_2O \xrightarrow{\Delta} Mg_2P_2O_7$$

As P atoms are conserved, applying POAC for P atoms, moles of P in $NaH_2PO_4 = Moles$ of P in $Mg_2P_2O_7$

$$\Rightarrow$$
 1 × Moles of NaH₂PO₄ = 2 × Moles of Mg₂P₂O₇

$$\therefore \qquad \frac{W_{NaH_2PO_4}}{M_{NaH_2PO_4}} = 2 \times \frac{W_{Mg_2P_2O_7}}{M_{Mg_2P_2O_7}} \Longrightarrow \frac{W_{NaH_2PO_4}}{120} = 2 \times \frac{2.054}{222}$$
$$\therefore \qquad W_{NaH_2PO_4} = 2.22 \text{ g}$$

4.4 **PERCENTAGE YIELD:**

In general, in any chemical reaction, the amount of product formed is always less than the calculated amount due to reversibility in the chemical reaction. Therefore, yield of a chemical reaction (Y) comes into picture and is given by :

The percentage yield of product = $\frac{\text{actual yield}}{\text{the theoretical maximum yield}} \times 100$

Ex.22In a certain operation 358 g of TiCl₄ is reacted with 96 g of Mg. Calculate % yield of Ti if 32 g
of Ti is actually obtained [At. wt. Ti = 48, Mg = 24]
(A) 35.38 % (B) 66.6 % (C) 100 % (D) 60 %Ans.(A)Sol.TiCl₄ + 2Mg \longrightarrow Ti + 2MgCl₂

Initial mole $\frac{358}{190} = 1.88$ $\frac{96}{24} = 4$ final mole 0 $4 - 2 \times 1.88$ 1.88 2×1.88 wt of Ti obtained $= \frac{358}{190} \times 48$ % yield $= \frac{32 \times 100}{\frac{358 \times 48}{190}} = 35.38$ %

- *Ex.23*0.05 mole of LiAlH4 in ether solution was placed in a flask containing 74g (1 mole) of t-butyl alcohol. The product LiAlHC12H27O3 weighed 12.7 g. If Li atoms are conserved, the percentage yield is : (Li = 7, Al = 27, H = 1, C = 12, O = 16).
(A) 25%(B) 75%(C) 100%(D) 15%
- Ans. (C)
- Sol. Li AlH₄ + t-butyl alcohol $\xrightarrow{\text{Ether}}$ LiAlH C₁₂H₂₇O₃(M.W. = 254) 0.05 mole 12.7 gram

$$=\frac{12.7}{254}=0.05$$
 mole

Li atom remain conserved so

No. of mole of LiAlH₄ = No. of mole of LiAlHC₁₂H₂₇O₃ So No. of mole of LiAlHC₁₂H₂₇O₃ = 0.05 % yield = $\frac{0.05}{0.05} \times 100 = 100\%$

4.5 **PERCENTAGE PURITY:**

- *Ex.24* How much marble of 90.5 % purity would be required to prepare 10 litres of CO_2 at 1 atm ,0 °Cwhen the marble is acted upon by dilute HCl ?
- *Ans.* 49.326 g
- Sol. $CaCO_3 + 2HCI \longrightarrow CaCl_2 + H_2O + CO_2$ 100 g 22.4 litre 22.4 L of CO₂ at STP will be obtained from 100 g of CaCO₃ 100

 \therefore 10 L of CO₂ will be obtained from pure CaCO₃ = $\frac{100}{22.4} \times 10 = 44.64$ g

 $\therefore \qquad \text{Impure marble required} = \frac{100}{90.5} \times 44.64 = 49.326 \text{ g}$

4.6 SEQUENTIAL REACTIONS:

This reaction is defined as that reaction which proceeds from reactants to final products through one or more intermediate stages. The overall reaction is a result of several successive or consecutive steps.

Example: $A \rightarrow B \rightarrow C$

Ex.25 Minimum amount of Ag_2CO_3 (s) required to produce sufficient oxygen for the complete combustion of C_2H_2 which produces 11.2 L of CO_2 at 1 atm and 273K after combustion is:

$$[Ag = 108]$$

$$Ag_2CO_3 (s) \longrightarrow 2Ag (s) + CO_2 (g) + 1/2 O_2 (g)$$

$$C_2H_2 + 5/2 O_2 \longrightarrow 2CO_2 + H_2O$$

Ans. 345 g

Sol. Ag₂CO₃ (s) \longrightarrow 2Ag (s) + CO₂ (g) + 1/2 O₂ (g) C₂H₂ + 5/2 O₂ \longrightarrow 2CO₂ + H₂O By Stoichiometry of reaction Moles of CO₂formed $=\frac{11.2}{22.4}=\frac{1}{2}$ Moles of O₂ required $=\frac{5}{4}\times\frac{1}{2}=\frac{5}{8}$ Moles of Ag₂CO₃required $=2\times\frac{5}{8}=\frac{5}{4}$ Mass of Ag₂CO₃ required $=\frac{5}{4}\times276=345$ g

Ex.26 NX is produced by the following step of reactions

 \longrightarrow M X₂ $M + X_2$ $3MX_2 + X_2 \longrightarrow M_3X_8$ $M_3 X_8 + N_2 CO_3 \longrightarrow$ $NX + CO_2 + M_3O_4$ How much M (metal) is consumed to produce 206 g of NX. (Take at wt of M = 56, N=23, X = 80) (A) 42 g (B) 56 g (C) 52g (D) 64 g (A) $M + X_2 \longrightarrow$ $MX_2 + X_2 \longrightarrow$ $N_{22}CO_3 \longrightarrow$ Ans. (A) Sol. MX_2 M_3X_8 $NX + CO_2 + M_3O_4$ mole of NX = $\frac{206}{103}$ = 2 POAC for X Atom : = No. of X Atom in NX No. of X atom in M_3X_8 8 [No. of mole of M_3X_8] = 1 [No. of mole of NX] No. of mole of $M_3X_8 = \left\lceil \frac{2}{8} \right\rceil = \frac{1}{4}$ mole Now POAC for M Atom 3 [No. of mole of M_3X_8] = 1 × [No. of Mole of M] $3 \times \frac{1}{4}$ = No. of mole of M ... weight of M atom $=\frac{3}{4} \times 56 = 42$ gram

Ex.27 The following process has been used to obtain iodine from oil-field brines in California.

 $\begin{array}{rcl} Nal + AgNO_{3} \longrightarrow & Agl + NaNO_{3} \ ; \\ 2AgI + Fe \longrightarrow & Fel_{2} + 2Ag \\ 2FeI_{2} + 3Cl_{2} \longrightarrow & 2FeCl_{3} + 2I_{2} \end{array}$

How many grams of $AgNO_3$ are required in the first step for every 254 kg I₂ produced in the third step.

(A) 340 kg (B) 85 kg (C) 68 kg (D) 380 kg

Ans. (A)

Sol. Balanced equation :

$$Nal + AgNO_{3} \longrightarrow Agl + NaNO_{3} \quad (1)$$

$$2AgI + Fe \longrightarrow Fel_{2} + 2Ag \quad (2)$$

$$2FeI_{2} + 3Cl_{2} \longrightarrow 2FeCl_{3} + 2I_{2} \quad (3)$$

From (3)

$$\frac{\text{mole of } I_2}{2} = \frac{\text{mole of } \text{FeI}_2}{2}$$

$$\frac{\text{mole of } \text{FeI}_2}{1} = \frac{\text{mole of } \text{AgI}}{2}$$

$$\frac{\text{mole of } \text{AgI}}{1} = \frac{\text{mole of } \text{AgNO}_3}{1}$$

$$\therefore \text{ mole of } I_2 = (\text{mole of } \text{FeI}_2)$$

$$= \left(\frac{\text{mole of } \text{AgI}}{2}\right) = \left(\frac{\text{mole of } \text{AgNO}_3}{2}\right)$$

$$\frac{254 \times 10^3}{254} = \frac{\text{mole of } \text{AgNO}_3}{2}$$

$$2 \times 10^3 = \text{mole of } \text{AgNO}_3 = \frac{\text{mass of } \text{AgNO}_3}{\text{molar mass of } \text{AgNO}_3}$$

$$\text{mass of } \text{AgNO}_3 = 170 \times (2 \times 10^3) \text{ g} = 340 \times 10^3 \text{ g} = 340 \text{ kg.}$$

4.7 PARALLEL REACTIONS:

The reactions in which a substance reacts or decomposes in more than one way are called parallel or side reactions.

Example: $A \longrightarrow B$

 $A \longrightarrow C$

Ex.28 Find out moles of CO_2 & CO produced by combustion of 2 mol carbon with 1.25 O_2 leaving number residue:

Ans. $CO_2 = 0.5 \text{ mol}$, $CO_2 = 1.5 \text{ mol}$ Sol. $C + O2 \longrightarrow CO_2$ $x \quad x \qquad x$ $C + \frac{1}{2} O_2 \longrightarrow CO$ $2-x \qquad 2-x \qquad 2-x$ $x + 1 - \frac{x}{2} = 1.25$ $\frac{x}{2} = \frac{125}{100}$ x = 0.5 mol, $CO_2 = 0.5 \text{ mol}$, $CO_2 = 1.5 \text{ mol}$

4.8 MIXTURE ANALYSIS:

The analysis of a chemical reaction is generally carried out in the form of mass of reacting species taking part in a given reaction (**gravimetric analysis**) or in terms of concentrations of reacting species taking part in a given reaction (**volumetric analysis**). In Gravimetric Analysis, we generally analyse reactions such as : decomposition of compounds under heat to produce a residue and a gas, or displacement reactions, action of acids on metals, or simple balanced chemical equations involving Weight (solid) – Volume (gas) relationships. In Volumetric Analysis, we generally analyse Neutralisation and Redox Titrations involving aqueous solutions in general.

(i) Except Li carbonates of all the alkali metals are thermally stable and does not decompose on heating. $\text{Li}_2\text{CO}_3 \xrightarrow{\Delta} \text{Li}_2\text{O} + \text{CO}_2$ $M_2\text{CO}_3 \xrightarrow{\Delta} \text{No reaction}$

(M = Na, K, Rb, Cs)

(ii) All the carbonates of alkaline earth metals are thermally unstable and decompose on heating as follow.

$$MCO_3 \xrightarrow{\Delta} MO+ CO_2$$

(iii) Bicarbonates of both alkali metals and alkaline earth metals are decomposed at relatively low temperature as follow.

$$2MHCO_3 \xrightarrow{\Lambda} M_2CO_3 + H_2O + CO_2$$
$$M(HCO_3)_2 \xrightarrow{\Lambda} MCO_3 + CO_2 + H_2O$$

- **Ex.29** A sample of 3 g containing Na₂CO₃ and NaHCO₃ loses 0.248 g when heated to 300°C, the temperature at which NaHCO₃ decomposes to Na₂CO₃, CO₂ and H₂O. What is the percentage of Na₂CO₃ in the given mixture?
- Ans. 77.48%

Sol. The loss in weight is due to removal of CO_2 and H_2O which escape out on heating. wt. of Na₂CO₃ in the product = 3.00 - 0.248 = 2.752 g Let wt. of Na₂CO₃ in the mixture be x g \therefore wt. of NaHCO₃ = (3.00 - x) g Since Na₂CO₃ in the products contains x g of unchanged reactant Na₂CO₃ and rest produced from NaHCO₃. The wt. of Na₂CO₃ produced by NaHCO₃ = (2.752 - x)g $NaHCO_3 \longrightarrow Na_2CO_3 + (H_2O + CO_2)$ (3.0 - x)(2.752 - x)Applying POAC for Na atom 1 × moles of NaHCO₃ = 2 × moles of Na₂CO₃ $\Rightarrow \frac{(3-x)}{84} = 2x \frac{(2.752-x)}{106}$ \therefore x = 2.3244 g :. % of Na₂CP₃ = $\frac{2.3244}{3} \times 100 = 77.48\%$

Ex.30 10 g of a sample of a mixture of $CaCl_2$ and NaCl is treated to precipitate all the calcium as CaCO₃. This Ca CO₃ is heated to convert all the Ca to CaO and the final mass of CaO is 1.62 g. The percent by mass of $CaCl_2$ in the original mixture is. (\mathbf{A}) 20 1 \mathbf{a} (D) 1 (0) 0/(0) 01 0 0/

(D) 11 0 0

(A)
$$32.1 \%$$
 (B) 16.2% (C) 21.8% (D) 11.0%
Ans. (A)
Sol. $CaCl_2 + NaCl = 10 g$
Let weight of $CaCl_2 = x g$
 $CaCl \rightarrow CaCO_3 \rightarrow CaO$
1 mol 1 mol 1 mol
 $\frac{x}{111} \text{ mol } \frac{x}{111} \text{ mol } \frac{x}{111} \text{ mol } \frac{x}{111}$
Mole of $CaO = \frac{1.62}{56}$
 $\therefore \frac{x}{111} = \frac{1.62}{56}$
 $x = 3.21 g$
% of $CaCl_2 = \frac{3.21}{10} \times 100 = 32.1 \%$

Do yourself-4:

1.	3.0 g an impure sample of sodium sulphate dissolved in water was treated with excess of barium chloride solution when 1.74 g of $BaSO_4$ was obtained as dry precipitate. Calculate the percentage purity of sample.				
2.	If the percentage yie produced, if 8 moles NaNO ₃ (s) \longrightarrow Na ₂ (A) 4.2 mole	eld of given reaction i of NaNO ₃ are taken in $O(s) + N_2(g) + O_2(g)$ (B) 2.4 mole	is 30%, how many nitially : (unbalanced) (C) 4.8 mole	y total moles of the gases will be (D) 2.1 mole	
3.	A 5 g mixture of SO SO ₂ and O ₂ gases in (A) 4 : 1	$_2$ and O_2 gases is reacted mixture to obtain maximum (B) 3 : 2	ed to form SO_3 gas imum amount of S (C) 2 : 3	s. What should be the mass ratio of O ₃ gas : (D) 1 : 4	
4.	 25.4 g of iodine and 14.2g of chlorine are made to react completely to yield a mixture of IC and ICl₃. Calculate the number of moles of ICl and ICl₃ formed. (A) 0.1 mole, 0.1 mole (B) 0.1 mole, 0.2 mole (C) 0.5 mole, 0.5 mole (D) 0.2 mole, 0.2 mole 				
5.	When 1 mole of A reacts with $\frac{1}{2}$ mole of B ₂ (A + $\frac{1}{2}$ B ₂ \longrightarrow AB), 100 Kcal heat is liberated and when 1 mole of A reacted with 2 mole of B ₂ (A + 2B ₂ \longrightarrow AB ₄), 200 Kcal heat is liberated. When 1 mole of A is completely reacted with excess, of B ₂ to form AB as well as AB ₄ , 140 Kcal heat is liberated calculate the mole of B ₂ used. [Write your answer as number of mole of B ₂ used × 10]				
6.	A solid mixture weighing 5.00 g containing lead nitrate and sodium nitrate was heated below 600° C until the mass of the residue was constant. If the loss of mass is 30 %, find the mass of lead nitrate and sodium nitrate in mixture. $2Pb(NO_3)_2 \xrightarrow{\Delta} 2PbO + 4NO_2\uparrow + O_2\uparrow$ $2NaNO_3 \xrightarrow{\Delta} 2NaNO_2 + O_2\uparrow$ (At. wt. of Pb = 207, Na = 23, N = 14, O = 16)				

5. **DENSITY:**

It is of two type.

1. Absolute density 2. Relative density

(a) For liquid and solids

Absolute density = $\frac{\text{mass}}{\text{volume}}$

Relative density or specific gravity = $\frac{\text{density of the substance}}{\text{density of water at 4°C}}$

(b) For gasses:

Absolute density (mass / volume) = $\frac{PM}{RT}$

where P is pressure of gas, M = mol. wt. of gas, R is the gas constant, T is the temperature.

Relative density andVapour density:

Vapour density is defined as the density of the gas with respect to hydrogen gas at the same temperature and pressure.

Vapour density =
$$\frac{d_{gas}}{d_{H_2}} = \frac{PM_{gas} / RT}{PM_{H_2} / RT}$$

V.D. = $\frac{M_{gas}}{M_{H_2}} = \frac{M_{gas}}{2} \Rightarrow \boxed{M_{gas} = 2V.D.}$

Ex.31 Find the relative density of SO₃ gas with respect to methane :

(A) 8 (B) 3.5 (C) 2.5 (D) 5
Ans. (D)
Sol. R.D.
$$= \frac{M_{SO_3}}{M_{CH_4}} = \frac{80}{16} = 5.$$

Ex.32 The atomic mass of a metal is 27 u. If its valency is 3, the vapour density of the volatile metal chloride will be:

(A) 66.75 (B) 321 (C) 267 (D) 80.25

Ans. (A)

Sol. Element must be Al

Hence, volatile chloride will be AlCl₃ so V.D. = $\frac{M_{AlCl_3}}{2} = \frac{133.5}{2} = 66.75$

Ex.33 The density of water at 4°C is 1×10^3 kg m⁻³. Assuming no empty space to be present between water molecules, the volume occupied by one molecule of water is approximately : (A) 3×10^{-23} mL (B) 6×10^{-23} mL (C) 3×10^{-22} mL (D) 6×10^{-22} mL **Ans.** (A) **Sol.** 1×10^3 kg/m³ = 1 g/mL. [Since, $1m^3 = 10^6$ cm³ = 10^6 mL]. = 1 gm/cc 6.022×10^{23} H₂O molecule weigh ...18 g 1 H₂O molecule weigh ... $\frac{18}{6.022 \times 10^{23}}$ g = 3×10^{-23} g d = $\frac{\text{mass}}{\text{volume}}$, So, volume = $\frac{3 \times 10^{-23} \text{ g}}{1(\text{g}/\text{mL})}$ = 3×10^{-23} mL.

6. **AVERAGE/ MEAN ATOMIC MASS :**

The weighted average of the isotopic masses of the element's naturally occuring isotopes.

Mathematically, average atomic mass of X (A_x) = $\frac{a_1 x_1 + a_2 x_2 + \dots + a_n x_n}{100}$

Where :

a₁, a₂, a₃ atomic mass of isotopes.

 x_1, x_2, x_3 mole % of isotopes. and

Ex.34 Naturally occurring chlorine is 75.53% Cl³⁵ which has an atomic mass of 34.969 amu and 24.47% Cl³⁷ which has a mass of 36.966 amu. Calculate the average atomic mass of chlorine -(A) 35.5 amu (B) 36.5 amu (C) 71 amu (D) 72 amu

Sol. (A) Average atomic mass =

(% of I isotope \times Its atoms mass)+(% II isotope \times its atomic mass)/100

<u>75.53×34.969+24.47×36.96</u> 100

= 35.5 amu.

6.1 AVERAGE MOLAR MASS OR AVERAGE GRAM MOLECULAR MASS:

The average molar mass of the different substance present in the container $=\frac{n_{1}M_{1}+n_{2}M_{2}+....n_{n}M_{n}}{n_{1}+n_{2}+...n_{n}}$

Where :

M₁, M₂, M₃ are molar masses.

 n_1, n_2, n_3 moles of substances.

Average molecule wt. = $\frac{\sum n_i M_i}{\sum n_i}$ where n_i = no. of moles of compound, m_i = molecular mass of

compound

Ex.35 The molar composition of polluted air is as follows :

Gas	mole percentage composition		
Oxygen	16%		
Nitrogen	80%		
Carbon dioxide	03%		
Sulphurdioxide	01%		
What is the average molecular wai	man male evilor weight of the siver rellected our 9		

What is the average molecular weight of the given polluted air?

Ans. 29.48

Sol.
$$M_{avg} = \frac{16 \times 32 + 80 \times 28 + 44 \times 3 + 64 \times 1}{100} = \frac{512 + 2240 + 132 + 64}{100} = \frac{2948}{100} = 29.48$$

6.2 **DEGREE OF DISSOCIATION** (α):

Degree of dissociation represents the fraction of one mole dissociated into the products. (Defined for one mole of substance)

- So, $\alpha = no.$ of moles dissociated / initial no. of moles taken
 - = fraction of moles dissociated out of 1 mole.

Note : % dissociation = $\alpha \times 100$

Suppose 5 moles of PCl₅ is taken and if 2 moles of PCl₅ dissociated then $\alpha = \frac{2}{5} = 0.4$

6.3 RELATIONSHIP BETWEEN AVERAGE MOLAR MASS& DEGREE OF DISSOCIATION (α):

Let a gas A_n dissociates to give n moles of A as follows-

$$\begin{array}{ccc} A_n\left(g\right) & \overleftarrow{\qquad} & n \mathrel{A}\left(g\right) \\ t = 0 & a & 0 \\ t = t_{eq} & a - x & nx & \alpha = \frac{x}{a} \implies & x = a\alpha. \end{array}$$

 $a - a \alpha = a(1-\alpha) n a \alpha$

Total no. of moles = $a - a \alpha + n a \alpha$

 $= [1 + (n - 1) \alpha] a$

Average molecular weight of $mixture(g) = \frac{molecular weight of A_n(g)}{total no. of moles at equilibrium}$

$$= \frac{a.M_{th}}{a(1+(n-1)\alpha)}$$
$$M_{avg} = \frac{M_{th}}{[1+(n-1)\alpha]}$$

where M_{th} = theoritical molecular weight (n = atomicity)

$$M_{\text{mixture}} = \frac{M_{A_n}}{[1 + (n-1)\alpha]}, M_{A_n} = \text{Molar mass of gas } A_n$$

Vapour density (V.D). : Density of the gas divided by density of hydrogen under same temperature & pressure is called vapour density.

D = vapour density without dissociation =
$$\frac{M_{A_n}}{2}$$

d = vapour density of mixture = averagevapour density
 $\frac{D}{d} = 1 + (n - 1) \alpha$
 $\alpha = \frac{D - d}{(n - 1) \times d} = \frac{M_T - M_o}{(n - 1)M_0}$

Ex.36 NH₃ decomposes into N₂ & H₂. If average molar mass of reaction mixture is 10 then, find α ? Ans. 0.7

Sol. $NH_{3} \longrightarrow \frac{1}{2}N_{2} + \frac{3}{2}H_{2}$ $n_{1} \quad 1 \qquad 0 \qquad 0$ $1 - \alpha \qquad \frac{\alpha}{2} \qquad \frac{3\alpha}{2}$ $10 = \frac{17}{1 - \alpha + \frac{\alpha}{2} + \frac{3\alpha}{2}}$ $10 = \frac{17}{1 + \alpha}$ $1 + \alpha = 1.7$ $\alpha = 0.7$

7. PERCENTAGE COMPOSITION AND MOLECULAR FORMULA :

Here we are going to find out the percentage of each element in the compound by knowing the molecular formula of compound.

We known that according to law of definite proportion any sample of a pure compound always possess constant ratio with their combining elements.

• Example:

Every molecule of ammonia always has formula NH_3 irrespective of method of preparation or sources. i.e. 1 mole of ammonia always contains 1 mol of N and 3 mole of H. In other wards 17 g of NH_3 always contains 14 g of N and 3 g of H. Now find out % of each element in the compound.

Mass % of N in NH₃ =
$$\frac{\text{Mass of N in 1 mol NH}_3}{\text{Mass of 1 mol of NH}_3}$$
 100 = $\frac{14g}{17}$ 100 = 82.35 %
Mass % of H in NH₃ = $\frac{\text{Mass of H in 1 mol NH}_3}{\text{Mass of 1 mol of NH}_3}$ 100 = $\frac{3}{17}$ 100 = 17.65%

7.1. EMPIRICAL AND MOLECULAR FORMULA:

We have just seen that knowing the molecular formula of the compound we can calculate percentage composition of the elements. Conversely if we know the percentage composition of the elements initially, we can calculate the relative number of atoms of each element in the molecules of the compound. This gives as the empirical formula of the compound. Further if the molecular mass is known then the molecular formula can be easily determined.

Thus, the empirical formula of a compound is a chemical formula showing the relative number of atoms in the simplest ratio, the molecular formula gives the actual number of atoms of eachelement in a molecule. **Empirical formula:** An empirical formula represents the simple whole number ratio of various atoms present in a compound,

Molecular formula: whereas, the molecular formula shows the exact number of different types of atoms present in a molecule of a compound

The molecular formula is generally an integral multiple of the empirical formula.

That is :

 $molecular formula = empirical formula \times n$ where n = $\frac{molecular formula mass}{emprirical formula mass}$

Ex.37 An organic substance containing carbon, hydrogen and oxygen gave the following percentage composition.

C=40.687 % ; H=5.085 % and O=54.228 %

The molecular weight of the compound is 118. Calculate the molecular formula of the compound.

- Ans. $C_4H_6O_4$
- Sol. Step -1

To calculate the empirical formula of the compound.

Element	Symbol	Percentage of element	At. mass of element	Relative no. of atoms = <u>Percentage</u> At.mass	Simplest Atomic ratio	Simplest whole no. atomic ratio
Carbon	С	40.687	12	$\frac{40.687}{12} = 3.390$	$\frac{3.390}{3.389} = 1$	2
Hydrogen	Н	5.085	1	$\frac{5.085}{1} = 5.035$	$\frac{5.085}{3.389}$ =1.5	3
Oxygen	0	54.228	16	$\frac{54.228}{16} = 3.389$	$\frac{3.389}{3.389} = 1$	2

• Step - 2

To calculate the empirical formula mass.

The empirical formula of the compound is $C_2H_3O_2$.

.: Empirical formula mass

 $=(2 \times 12) + (3 \times 1) + (2 \times 16) = 59.$

• Step - 3

To calculate the value of 'n'

 $n = \frac{\text{molecular mass}}{\text{Empirical formula nass}} = \frac{118}{59} = 2$

Step - 4 To calculate the molecular formula of the salt Molecular formula $= n \times (\text{Empirical formula})$ $= 2 \times C_2 H_3 O_2 = C_4 H_6 O_4$ Thus the molecular formula is $C_4 H_6 O_4$.

- *Ex.38* Chlorophyll the green colouring material of plants contains 3.68 % of magnesium by mass. Calculate the number of magnesium atom in 5.00 g of the complex.
- Sol. Mass of magnesium in5.0 g of complex $=\frac{3.68}{100}$ 5.00 = 0.184 g Atomic mass of magnesium = 24 24 g of magnesium contain = 6.023×10^{23} atoms 0.184 g of magnesium would contain $=\frac{6.023 \times 10^{23}}{24}$ 0.184 = 4.617 × 1021 atom Therefore, 5.00 g of the given complex would contain 4.617 × 10²¹ atoms of magnesium.

Ex.39 A sample of CaCO₃ has Ca = 40%, C = 12% and O = 48% by mass. If the law of constant proportions is true, then the mass of Ca in 5 g of CaCO₃ obtained from another source will be: (A) 0.2 g (B) 2 g (C) 0.6 g (D) Cannot be determined *Ans.* (B)

Sol. Mass of $Ca = 5 \times \frac{40}{100} = 2g$.

8. EXPERIMENTAL METHODS TO DETERMINE ATOMIC & MOLECULAR MASSES

8.1 For determination of atomic mass :

(a) Dulong's & Pettit' slaw :

In case of metals, it is observed that product of atomic weight and specific heat capacity is constant.

The equation connecting the two parameters was given by Dulong's and Petit's Law.

Atomic weight of metal \times specific heat capacity (cal/gm[•]C) = 6.4.

In should be remembered that this law is an empirical observation and this gives an approximate value of atomic weight. Also this law can be applied only to metals at high temperature conditions only.

8.2 Experimental methods for molecular mass determination.

- (a) Victor Meyer's Method
- (b) Silver Salt Method
- (c) Chloroplatinate Salt Method

(a) Victor Meyer's Method: (Applicable for volatile substance)

A known mass of the volatile substance taken in the Hoffmann's bottle and is vapourised by throwing the Hoffmann's bottle into the Vector Meyer's tube. The vapour displace an equal volume of the moist air. Which vapours is measured at the room temperature and atmospheric pressure. The barometric pressure and the room temperature is recorded. Following diagram gives the experimental set-up for the Victor-Meyer's process.

Calculation involved

Let the mass of the substance taken by = Wg

Volume of moist vapours collected = Vcm^3

Room temperature = TK

Barometric pressure = P mm Aqueous tension at TK = p mmPressure of dry vapour = (P - p) mmCalculation of molecular mass (M)

$$\frac{(P-p)}{760} \times \frac{V}{1000} = \frac{W}{M} \times RT$$
$$\implies M = \frac{W \times RT \times 760 \times 1000}{(P-p) \times V}$$



(b) Silver salt Method: (A used for organic acids)

A known mass of the acid is dissolved in water followed by the subsequent addition of silver nitrate solution till the precipitation of silver salt is complete. The precipitate is separated, dried, weighed

Organic acid $\xrightarrow{AgNO_3}$ Silver salt \xrightarrow{Ignite} Ag

and ignited till decomposition is complete. The residue of pure silver left behind is weighed. Calculations involved

Let the mass of the silver salt formed = W g

The mass of Ag formed = x g

Let us understand to calculations by considering the monobasic acid MX.

$$\underset{\text{Organic acid}}{\text{HA}} \xrightarrow{\text{AgNO}_3} \underset{\text{Silver salt(Wg)}}{\text{AgA}} \xrightarrow{\text{Tgnite}} \underset{\text{Silver(xg)}}{\text{Ag}} Ag$$

Mass of the salt that gives x gm of Ag = W g

Mass of the salt gives 108g (1g-atom) of $Ag = \frac{108W}{x}g$

Molar mass of salt = $\frac{108W}{x}g$

Molar mass of acid = (Molar mass of salt) – (Atomic mass of Ag) + (Atomic mass of H) = $\frac{108W}{x} - 108 + 1 = \left(\frac{108W}{x} - 107\right) g \text{ mol}^{-1}$

For polybasic acid of the type H_nX (n is basicity)

 $H_{n}A \xrightarrow{AgNO_{3}} Ag_{n}A \xrightarrow{Tgnite} nAg_{Silver(xg)}$ Mass of the silver that gives x g of Ag = W g Mass of the silver that gives (108n g) of Ag = $\frac{108nW}{x}g$ Molar mass of salt = $\frac{108 \times nW}{x}g$ Molar mass of acid = (molar mass of salt) = $\frac{108 \times nW}{x} - n \times 108 - n \times 108$ = $n\left(\frac{108W}{x} - 107\right)g mol^{-1}$

(c) Platinic chloride Method : (Applicable for finding the molecular masses of organic bases). A known mass of organic base is allowed to react with chloroplatinic acid (H₂PtCl₆) in conc. HCl to form insoluble platinic chloride. The precipitate of platinic chloride is separated, dried, weighed and subsequently ignited till decomposition is complete. The residue left is platinum which is again weighed. The molecular mass is then calculated by knowing the mass of the platinic chloride salt and that of platinum left.

If B represents the molecule of monoacidic organic base, then the formula of platinic chloride salt is

$$\begin{array}{c} B_{2}H_{2}PtCl_{6} \\ B_{Organic \ base} \xrightarrow{H_{2}PtCl_{6}} B_{2}H_{2}PtCl_{6} \\ \xrightarrow{H_{2}PtCl_{6}} B_{2}H_{2}PtCl_{6} \\ \xrightarrow{\text{lg nite}} Pt \\ P_{\text{latinic chloridesalt(Wg)}} \xrightarrow{\text{lg nite}} Pt \\ \xrightarrow{(xg)} \end{array}$$

Let the mass of platinic chloride salt = W g, The mass of the platinum residue left = x g It may be noted that salt formed with diacidic base would be B_2 (H₂PtCl₆)₂

With triacidic base it would be B_2 (H_2PtCl_6)₃ and with polyacidic base would be $B_2(H_2PtCl_6)_n$.

Mass of salt which gives 195 g (1 g-atom) of Pt $\frac{W \times 195}{x}$ g

Molar mass of salt = $\frac{W \times 195}{x}$ g mol⁻¹

Now from the formula $B_2(H_2PtCl_6)$

Molar mass of salt = $(2 \times \text{molar mass of base}) + (\text{Molar mass of H}_2\text{PtCl}_6)$

Molar mass of base = $\frac{1}{2}$ (molar mass of salt – Molar mass of H₂PtCl₆)

$$= \frac{1}{2} \left(\frac{W \times 195 \times n}{x} - n \times 410 \right) = \frac{n}{2} \left(\frac{W \times 195}{x} - 410 \right) \operatorname{g mol}^{-1}$$

Do y	vourself-5:				
1.	120 g Mg is burn HCl to form MgC formed:	t in air to give a mix 12 and NH ₄ Cl. If 10 ^o	kture of MgO and Mg ₃ 7 g NH ₄ Cl is produced	N_2 . The mixture is now diss then determine the moles of	olved in f MgCl ₂
	(A) 2.5	(B) 4	(C) 2	(D) 5	
2.	Penicillin V was A 9.6 mg sampl V is x %. If ther amu. Report your ansy	s treated chemically e of penicillin V ga e is one sulphur ato wer as y/x.	to convert the sulphur ve 4.66 mg BaSO ₄ . The m in the molecule, the	present to barium sulphate, e percentage of sulphur in P molecular weight of Pencilli	BaSO ₄ . renicillin n V is y
3.	From the follo $Cl_2 + 2I$ 3KCIO $4KCIO_3$ Calculate the r (A) 142 g	wing reaction seq $KOH \longrightarrow KCl +$ $\longrightarrow 2KCl$ $\longrightarrow 3KC$ mass of chlorine r (B) 284 g	puence : $- \text{KClO} + \text{H}_2\text{O}$ $+ \text{KClO}_3$ $1\text{O}_4 + \text{KCl}$ needed to produce 13 (C) 432 g	88.5 g of KClO4 : (D) None of these	,
4.	The density of a (A) 0.015	ir at STP is 0.0013 g (B) 15	mL^{-1} . Its vapour densi (C) 1.5	ty is : (D) Data insufficien	t
5.	$SO_3(g) \rightleftharpoons S$	$O_2(g) + \frac{1}{2} O_2(g)$			
	If observed vapo (A) 0.28	our density of mixtur (B) 0.38	re at equilibrium is 35 th (C) 0.48	hen find out value of α: (D) 0.58	
6.	A sample of a c What is empirica	ompound contains 9 al formula of compo	9.75 g Zn, 9×10^{22} ato und? (Atomic Mass Zn	ms of Cr and 0.6 gram-ator $= 65$)	ns of O.
	(A)ZnCrO ₄	$(B)ZnCr_2O_4$	$(C)Zn_2CrO_4$	(D) None of these	
7.	An organic con The molecular n (A) 100000 u	mpound on analysi nass of the compoun (B) 10000 u	is was found to cont d, if it's one molecule o (C) 20000 u	ain 0.032% of sulphur b contains two sulphur atoms, (D) 200000 u	y mass. is :
8.	In an organic co by weight. Mole	mpound of molar m cular formula can be	$aass108 \text{ g mol}^{-1} \text{ C, H and }$	nd N atoms are present in 9	: 1 : 3.5
	$(A) C_6 H_8 N_2$	(B) $C_7 H_{10} N$	$(C) C_5 H_6 N_3$	(D) $C_4H_{18}N_3$	
9.	At 100° C and 1 0.0006 g cm ^{-3} temperature is :	atmp, if the densit, then the volume	ty of liquid water is 1.0 occupied by water n) g cm ^{-3} and that of water v nolecules in 1 L of steam	apour is at that
	$(A) 6 cm^3$	(B) 60 cm^3	(C) 0.6 cm^3	(D) 0.06 cm^3	

ANSWER KEY

DO YOURSELF

Do yourself-1:

is, $S = 2$ atoms, $O = 8$ atoms.
ns, $S = 6$ atoms, $O = 18$ atoms.
(\mathbf{C})
g of carbon is needed.
5. 11 6. 0.95 g
(Δ) 6 (Δ)
1

EXERCISE (S- I)

PROBLEMS RELATED WITH DIFFERENT TYPES OF ATOMIC MASSES & BASIC CONCEPT OF MOLE

- 1. How much time (in seconds) would it take to distribute one Avogadro number of wheat grains if 10^{10} grains are distributed each second?
- 2. What is the mass of one ${}^{12}C$ atom in gram ?
- 3. Calculate the weight of 12.046×10^{23} atoms of carbon.
- **4.** Find :
 - (i) No. of moles of Cu atom in 10^{20} atoms of Cu.
 - (ii) Mass of 200 atoms of ${}^{16}_{8}$ O in amu
 - (iii) Mass of 100 atoms of ${}^{14}_{7}$ N in gm.
 - (iv) No. of molecules & atoms in $54 \text{ gm H}_2\text{O}$.
 - (v) No. of atoms in 88 gm CO_2 .
- 5. Calculate mass of O atoms in $6 \text{ gm CH}_3\text{COOH}$?
- 6. Calculate mass of water present in 499 gm $CuSO_4.5H_2O$? (Atomic mass – Cu = 63.5, S = 32, O = 16, H = 1)
- 7. What mass of Na₂SO₄.7H₂O contains exactly 6.023×10^{22} atoms of oxygen ?
- 8. Find the total number of nucleons present in $12 \text{ g of } {}^{12}\text{C}$ atoms.
- 9. Calculate the number of electrons, protons and neutrons in 1 mole of ${}^{16}O^{-2}$ ions.
- 10. The density of liquid mercury is 13.6 g/cm^3 . How many moles of mercury are there in 1 litre of the metal? (Atomic mass of Hg = 200.)
- 11. A sample of ethane has the same mass as 10.0 million molecules of methane. How many C_2H_6 molecules does the sample contain?
- 12. If, from 10 moles NH_3 and 5 moles of H_2SO_4 , all the H-atoms are removed in order to form H_2 gas, then find the number of H_2 molecules formed.

STOICHIOMETRY

- 13. Chlorine can be prepared by reacting HCl with MnO_2 . The reaction is represented by the equation: $MnO_2(s) + 4HCl \longrightarrow Cl_2(g) + MnCl_{2(aq)} + 2H_2O(l)$ Assuming the reaction goes to completion, what mass of HCl solution is needed to produce 142 g of Cl₂
- 14. Calculate the volume of O_2 needed for combustion of 1.2 kg of carbon at STP. Reaction : $C + O_2 \xrightarrow{\Delta} CO_2$.

- 15. Methyl-t-butyl ether, C₅H₁₂O, is added to gasoline to promote cleaner burning. How many moles of oxygen gas, O₂ are required to burn 1.0 mol of this compound completely to form carbon dioxide and water?
- 16. Aluminum carbide (Al_4C_3) liberates methane on treatment with water : $Al_4C_4 + 12H_2O \longrightarrow 3CH_4 +$ 4Al(OH), Find mass of aluminum carbide required to produce 11.35 L of methane under STP conditions.
- 17. Calculate mass of phosphoric acid required to obtain 53.4g pyrophosphoric acid. $2H_3PO_4 \rightarrow H_4P_2O_7 + H_2O_7$
- 18. Nitric acid is manufactured by the Ostwald process, in which nitrogen dioxide reacts with water. $3 \operatorname{NO}_{2}(g) + \operatorname{H}_{2}O(l) \rightarrow 2 \operatorname{HNO}_{3}(aq) + \operatorname{NO}(g)$ How many grams of nitrogen dioxide are required in this reaction to produce 25.2 gm HNO₃?
- 19. Flourine reacts with uranium to produce uranium hexafluoride, UF₆, as represented by this equation $U(s) + 3F_2(g) \rightarrow UF_2(g)$ How many fluorine molecules are required to produce 2.0 mg of uranium hexafluoride, UF₆, from an excess of uranium? The molar mass of UF_6 is 352 gm/mol.
- 20. XeF_6 fluorinates I₂ to IF₇ and liberates Xenon(g). 3.5 mmol of XeF₆ can yield a maximum of_ mmol of IF_7 .
- What total volume, in litre at 600°C and 1 atm, could be formed by the decomposition of 16 gm of 21. $\begin{array}{c} \mathrm{NH_4NO_3} ?\\ 2 \ \mathrm{NH_4NO_3} \rightarrow 2\mathrm{N_2} + \mathrm{O_2} + 4\mathrm{H_2O_{(g)}} \end{array}$

LIMITING REACTANT

- 50 g of CaCO₃ is allowed to react with 73.5 g of H_3PO_4 . Calculate : 22. (i) Amount of $Ca_3(PO_4)_2$ formed (in moles) (ii) Amount of unreacted reagent (in moles)
- Reaction $4A + 2B + 3C \longrightarrow A_4B_2C_3$, is started from 2 moles of A, 1.2 moles of B & 1.44 moles 23. of C. find number of moles of product formed.
- 24. Potassium superoxide, KO₂, is used in rebreathing gas masks to generate oxygen : $KO_2(s) + H_2O(l) \rightarrow KOH(s) + O_2(g)$ If a reaction vessel contains 0.158 mol KO₂ and 0.10 mol H₂O, how many moles of O₂ can be produced?
- 25. A chemist wants to prepare diborane by the reaction $6 \operatorname{LiH} + 8\operatorname{BF}_3 \longrightarrow 6\operatorname{Li}\operatorname{BF}_4 + \operatorname{B}_2\operatorname{H}_6$ If he starts with 2.0 moles each of LiH & BF₃. How many moles of B_2H_6 can be prepared.
- Titanium, which is used to make air plane engines and frames, can be obtained from titanium tetrachloride, 26. which in turn is obtained from titanium oxide by the following process : $3 \operatorname{TiO}_{2}(s) + 4C(s) + 6Cl_{2}(g) \longrightarrow 3\operatorname{TiCl}_{4}(g) + 2CO_{2}(g) + 2CO(g)$ A vessel contains 4.32 g TiO₂, 5.76 g C and; 7.1 g Cl₂, suppose the reaction goes to completion as written, how many gram of TiCl₄ can be produced? (Ti = 48)

- 27. Carbon reacts with chlorine to form CCl_4 . 36 gm of carbon was mixed with 142 g of Cl_2 . Calculate mass of CCl_4 produced and the remaining mass of reactant.
- 28. Sulphuric acid is produced when sulphur dioxide reacts with oxygen and water in the presence of a catalyst : $2SO_2(g) + O_2(g) + 2H_2O(l) \rightarrow 2H_2SO_4$. If 5.6 mol of SO₂ reacts with 4.8 mol of O₂ and a large excess of water, what is the maximum number of moles of H₂SO₄ that can be obtained?

PROBLEMS RELATED WITH MIXTURE

- **29.** One gram of an alloy of aluminium and magnesium when heated with excess of dil. HCl forms magnesium chloride, aluminium chloride and hydrogen. The evolved hydrogen collected at 0°C has a volume of 1.12 litres at 1 atm pressure. Calculate the composition of (% by mass) of the alloy.
- **30.** A sample containing only $CaCO_3$ and $MgCO_3$ is ignited to CaO and MgO. The mixture of oxides produced weight exactly half as much as the original sample. Calculate the percentages of $CaCO_3$ and $MgCO_3$ (by mass) in the sample.
- 31. Determine the percentage composition (by mass) of a mixture of anhydrous sodium carbonate and sodium bicarbonate from the following data:
 wt. of the mixture taken = 2g
 Loss in weight on heating = 0.11 gm.
- **32.** 92 g mixture of $CaCO_3$, and MgCO₃ heated strongly in an open vessel. After complete decomposition of the carbonates it was found that the weight of residue left behind is 48 g. Find the mass of MgCO₃ in grams in the mixture.
- **33.** When 4 gm of a mixture of NaHCO₃ and NaCl is heated, 0.66 gm CO_2 gas is evolved. Determine the percentage composition (by mass) of the original mixture.

PERCENTAGE YIELD AND PERCENTAGE PURITY

- **34.** 200 g impure $CaCO_3$ on heating gives 11.35 L CO₂ gas at STP. Find the percentage of calcium in the lime stone sample.
- **35.** A power company burns approximately 474 tons of coal per day to produce electricity. If the sulphur content of the coal is 1.30% by weight, how many tons SO₂ are dumped into the atmosphere each day?
- 36. Calculate the percent loss in weight after complete decomposition of a pure sample of potassium chlorate. $KClO_3(s) \longrightarrow KCl(s) + O_2(g)$
- **37.** A sample of calcium carbonate is 80% pure, 25 gm of this sample is treated with excess of HCl.How much volume of CO₂ will be obtained at 1 atm & 273 K?
- **38.** Cyclohexanol is dehydrated to cyclohexene on heating with conc. H_2SO_4 . If the yield of this reaction is 75%, how much cyclohexene will be obtained from 100 g of cyclohexanol?

 $C_6H_{12}O \xrightarrow{\text{con.H}_2SO_4} C_6H_{10}$
- 39. If the yield of chloroform obtainable from acetone and bleaching powder is 75%. What is the weight of acetone required for producing 30 gm of chloroform ?
 2CH₃COCH₃ + 6CaOCl₂ → Ca(CH₃COO)₂ + 2CHCl₃ + 3CaCl₂ + 2Ca(OH)₂
- 40. The percent yield for the following reaction carried out in carbon tetrachloride (CCI_{4}) solution is 80%

 $Br_2 + CI_2 \longrightarrow 2BrCI$

- (a) What amount of BrCI would be formed from the reaction of 0.025 mol Br₂ and 0.025 mol CI₂?
- (b) What amount of Br_2 is left unchanged?

SEQUENTIAL & PARALLEL REACTIONS

- 41. Sulphur trioxide may be prepared by the following two reactions : $S_8 + 8O_2(g) \rightarrow 8SO_2(g)$ $2SO_2(g) + O_2(g) \rightarrow 2SO_3(g)$ How many grams of SO₃ will be produced from 1 mol of S₈?
- 42. $2PbS + 3O_2 \rightarrow 2PbO + 2SO_2$ $3SO_2 + 2HNO_3 + 2H_2O \rightarrow 3H_2SO_4 + 2NO$ According to the above sequence of reactions, how much H₂SO₄ will 1075.5 gm of PbS produce?
- **43.** Potassium superoxide, KO_2 , is utilised in closed system breathing apparatus. Exhaled air contains CO_2 and H_2O , both of which are removed and the removal of water generates oxygen for breathing by the reaction

 $4\text{KO}_2(s) + 2\text{H}_2\text{O}(1) \rightarrow 3\text{O}_2(g) + 4\text{KOH}(s)$

The potassium hydroxide removes CO₂ from the apparatus by the reaction :

 $\operatorname{KOH}(s) + \operatorname{CO}_{2}(g) \rightarrow \operatorname{KHCO}_{3}(s)$

- (a) What mass of KO₂ generates 20 gm of oxygen ?
- (b) What mass of CO_2 can be removed from the apparatus by 100 gm of KO_2 ?

MISCELLANEOUS PROBLEM

- 44. In a determination of P an aqueous solution of NaH_2PO_4 is treated with a mixture of ammonium and magnesium ions to precipitate magnesium ammonium phosphate $Mg(NH_4)PO_4$. $6H_2O$. This is heated and decomposed to magnesium pyrophosphate, $Mg_2P_2O_7$ which is weighed. A solution of NaH_2PO_4 yielded 1.054 g of $Mg_2P_2O_7$. What weight of NaH_2PO_4 was present originally?
- 45. Calculate the amount of H_2SO_4 produced (in gm) when 40 ml H_2O (d = 0.9 gm/ml) reacts with 50 litre SO_3 at 1 atm. and 300 K, according to the following reaction ? $H_2O + SO_3 \rightarrow H_2SO_4$
- **46.** 0.80g of the chloroplatinate of a mono acid base on ignition gave 0.262g of ppt. Calculate the molecular weight of the base.

47. Calculate the atomic mass (average) of chlorine using the following data :

	% Natural Abundance	Molar Mass		
³⁵ Cl	75.77	34.9689		
³⁷ Cl	24.23	36.9659		

- **48.** Average atomic mass of Magnesium is 24.31 amu. This magnesium is composed of 79 mole % of ²⁴Mg and remaining 21 mole % of ²⁵Mg and ²⁶Mg. Calculate mole % of ²⁶Mg.
- **49.** A moth repellent has the composition 49% C, 2.7% H and 48.3% Cl. Its molecular weight is 147 gm. Determine its molecular formula
- **50.** Haemoglobin contains 0.25% iron by mass. The molecular mass of of Haemoglobin is 89600 then the number of iron atoms per molecule of Haemoglobin (Atomic mass of Fe = 56)

- 1. Sodium chlorate, NaClO₃, can be prepared by the following series of reactions: $2KMnO_4 + 16 HCl \rightarrow 2 KCl + 2 MnCl_2 + 8H_2O + 5 Cl_2$ $6Cl_2 + 6 Ca(OH)_2 \rightarrow Ca(ClO_3)_2 + 5 CaCl_2 + 6H_2O$ $Ca(ClO_3)_2 + Na_2SO_4 \rightarrow CaSO_4 + 2 NaClO_3$ What mass of NaClO₃ can be prepared from 100 ml of concentrated HCl (density 1.18 gm/ml and 36% by mass)? Assume all other substances are present in excess amounts.
- 2. Two substance $P_4 & O_2$ are allowed to react completely to form mixture of $P_4 O_6 & P_4 O_{10}$ leaving none of the reactants. Using this information calculate the composition of final mixture when mentioned amount of $P_4 & O_2$ are taken.

$$P_4 + 3O_2 \longrightarrow P_4O_6$$

$$P_4 + 5O_2 \longrightarrow P_4O_{10}$$

- (i) If 1 mole $P_4 \& 4$ mole of O_2
- (ii) If 3 mole $P_4 \& 11$ mole of O_2
- (iii) If 3 mole $P_4 \& 13$ mole of O_2
- 3. By the reaction of carbon and oxygen, a mixture of CO and CO_2 is obtained. What is the composition (% by mass) of the mixture obtained when 20 grams of O_2 reacts with 12 grams of carbon?
- 4. Nitrogen (N), phosporus (P), and potassium (K) are the main nutrients in plant fertilizers. According to an industry convention, the numbers on the label refer to the mass % of N, P_2O_5 , and K_2O , in that order. Calculate the N : P : K ratio of a 30 : 10 : 10 fertilizer in terms of moles of each elements, and express it as x : y : 1.0. Find y.
- 5. A 10 g sample of a mixture of calcium chloride and sodium chloride is treated with Na_2CO_3 to precipitate calcium as calcium carbonate. This $CaCO_3$ is heated to convert all the calcium to CaO and the final mass of CaO is 1.12gm. Calculate % by mass of NaCl in the original mixture.
- 6. A mixture of Ferric oxide (Fe_2O_3) and Al is used as a solid rocket fuel which reacts to give Al_2O_3 and Fe. No other reactants and products are involved. On complete reaction of 1 mole of Fe_2O_3 , 200 units of energy is released.

(a) Write a balance reaction representing the above change.

(b) What should be the ratio of masses of Fe_2O_3 and Al taken so that maximum energy per unit mass of fuel is released.

(c) What would be energy released if 16 kg of Fe₂O₃ reacts with 2.7 kg of Al.

7. 1 gm sample of $KClO_3$ was heated under such conditions that a part of it decomposed according to the equation

(1) $2\text{KClO}_3 \longrightarrow 2\text{KCl} + 3\text{O}_2$

and remaining underwent change according to the equation.

(2) $4\text{KClO}_3 \longrightarrow 3 \text{KClO}_4 + \text{KCl}$

If the amount of O_2 evolved was 112 ml at 1 atm and 273 K., calculate the % by weight of KClO₄ in the residue.

- 8. 5.33 mg of salt $[Cr(H_2O)_5Cl].Cl_2$. H_2O is treated with excess of $AgNO_3(aq.)$ then mass of AgCl ppt.obtained will be: Given: [Cr = 52, Cl = 35.5]
- 9. If mass % of oxygen in monovalent metal carbonate is 48%. Then find the number of atoms of metal present in 5mg of this metal carbonate sample is $(N_A = 6.0 \times 10^{23})$
- 10. To find formula of compound composed of A & B which is given by $A_x B_y$, it is strongly heated in oxygen as per reaction- $A_x B_y + O_2 \rightarrow AO + Oxide \text{ of } B$

If 2.5gm of $A_x B_y$ on oxidation gives 3gm oxide of A, Find empirical formula of $A_x B_y$, [Take atomic mass of A = 24 & B = 14]

- 11. Calculate maximum mass of CaCl₂ produced when 2.4×10^{24} atoms of calcium is taken with 96 litre of Cl₂ gas at 380 mm pressure and at 27°C. [R : 0.08 atm L/mole-K & N_A = 6 × 10²³]
- 12. $P_4S_3 + 8O_2 \longrightarrow P_4O_{10} + 3SO_2$ Calculate minimum mass of P_4S_3 is required to produce at least 1 gm of each product.
- 13. Consider the given reaction $H_4P_2O_7 + 2NaOH \rightarrow Na_2H_2P_2O_7 + 2H_2O$ If 534 gm of $H_4P_2O_7$ is reacted with 30×10^{23} molecules of NaOH then total number of molecules produced in the product is

EXERCISE (O-I)

Single	Correct :			
	PROBLEMS REL	ATED WITH DIFFE	RENT TYPES OF AT	OMIC MASSES &
		BASIC CONCI	EPT OF MOLE	
1.	Which of the following	g has the Maximum mass	?	
	(A) 1 g-atom of C		(B) $\frac{1}{2}$ mole of CH ₄	
	(C) 10 mL of water		(D) 3.011×10^{23} atoms	s of oxygen
2.	The number of molecu	ales of CO ₂ present in 44	ag of CO ₂ is :	
	(A) 6.0×10^{23}	(B) 3×10 ²³	(C) 12×10^{23}	(D) 3×10 ¹⁰
3.	The number of mole of	fammonia in 4.25 g of an	nmonia is :	
	(A) 0.425	(B) 0.25	(C) 0.236	(D) 0.2125
4.	The charge on 1 gram	ions of Al^{3+} is : ($N_A = Av$	ogadro number, e = cha	rge on one electron)
	(A) $\frac{1}{27}$ N _A e coulomb	(B) $\frac{1}{3} \times N_A e$ coulomb	(C) $\frac{1}{9} \times N_{A}$ ecoulomb	(D) $3 \times N_A e$ coulomb
5.	The atomic weights of how many atoms are p	two elements A and B ar present in 2x g of B?	e 40u and 80u respective	ely. If x g of A contains y atoms,
	(A) $\frac{y}{2}$	(B) $\frac{y}{4}$	(C) y	(D) 2y
6.	A sample of aluminium $(At. wt. Al = 27, Mg =$	n has a mass of 54.0 g. Wh = 24)	nat is the mass of the same	e number of magnesium atoms?
	(A) 12 g	(B) 24 g	(C) 48 g	(D) 96 g.
7.	The weight of a molect (A) 1.09×10^{-21} g	ule of the compound C_{60} (B) 1.24×10^{-21} g	H ₂₂ is: (C) 5.025×10^{-23} g	(D) 16.023×10^{-23} g
8.	The number of electro	on in 3.1 mg NO_3^- is -		
	(A) 32	(B) 1.6×10^{-3}	(C) 9.6×10^{20}	(D) 9.6×10^{23}
9.	A gaseous mixture co molecules of $CO_2(g)$	ntains CO ₂ (g) and N ₂ O and N ₂ O (g) is	(g) in a 2 : 5 ratio by ma	ass. The ratio of the number of
	(A) 5 : 2	(B) 2 : 5	(C) 1 : 2	(D) 5:4
10.	Which of the following	g contain largest number	of carbon atoms?	
	(A) 15 gm ethane, $C_2 H$	H_6	(B) 40.2 gm sodium oz	$xalate, Na_2C_2O_4$
	(C) 72 gm glucose, C_{e}	$H_{12}O_6$	(D) 35 gm pentene, C_5	H ₁₀
11.	The number of hydrog	gen atoms in 0.9 gm gluc	ose, $C_6 H_{12} O_6$, is same as	3
	(A) 0.048 gm hydrazin	he, N_2H_4	(B) 0.17 gm ammonia,	NH ₃
	(C) 0.30 gm ethane, C	C_2H_6	(D) 0.03 gm hydrogen	$, H_2$

12.	The weight of 1×10^{22} (A) 41.59 g	molecules of CuSO ₄ . 5 (B) 415.9 g	² H ₂ O is : (C) 4.159 g	(D) 2.38 g
13.	The number of carbor 1.2×10^{-3} g is	atoms present in a sign	nature, if a signature wr	itten by carbon pencil weights
	(A) 12.04×10^{20}	(B) 6.02×10^{19}	(C) 3.01×10^{19}	(D) 6.02×10^{20}
14.	Ethanol, C_2H_5OH , is the 293 K. If 1.2 mole of end measured out?	e substance commonly c thanol are needed for a p	called alcohol. The densit articular experiment, wh	ry of liquid alcohol is 0.8 g/ml at aat volume of ethanol should be
	(A) 55 ml	(B) 58 ml	(C) 69 ml	(D) 79 ml
15.	112.0 ml of NO_2 at 1 at the volume of and the r	m & 273 K was liquefie number of molecules in t	d, the density of the liqu he liquid NO ₂ .	id being 1.15 gm/ml. Calculate
	(A) 0.10 ml and 3.01	× 10 ²²	(B) 0.20 ml and 3.01	× 10 ²¹
	(C) 0.20 ml and 6.02	$\times 10^{23}$	(D) 0.40 ml and 6.02	× 10 ²¹
16.	X gm A atoms on comb the molecular weight o	oining with Y atoms of B of compound formed. (A	form 5 molecules of a contract tomic weight of $B = M$)	mpound containing A & B. Find
	(A) $\frac{(XN_A + MY)}{5}$	(B) $\frac{X+M}{5}$	(C) $\frac{X+MY}{5}$	$(D)\left(\frac{X + MYN_A}{5}\right)$
17.	At same temperature a	nd pressure, two gases h	nave the same number of	molecules. They must
	(A) have same mass(C) have a volume of 2	22.7 dm ³ each	(B) have equal volumes (D) have an equal num	s ber of atoms
18.	An iodized salt contain ions going into his bod	ns 0.5 % of NaI. A person y everyday is	a consumes 3 gm of salt e	everyday. The number of iodide
	(A) 10 ⁻⁴	(B) 6.02 ×10 ⁻⁴	(C) 6.02×10^{19}	(D) 6.02×10^{23}
19.	Equal volumes of oxy same experimental con	gen gas and a second g ditions. Which of the fol	as weigh 1.00 and 2.375 lowing is the unknown g	5 grams respectively under the as?
	(A) NO	(B) SO ₂	(C) CS_2	(D) CO
20.	Four 1-1 litre flasks are pressure. The ratio of t	e separately filled with th otal number of atoms of	he gases H_2 , He, O_2 and C_2 these gases present in di	O ₃ at the same temperature and afferent flask would be :
	(A) 1 : 2 :3 : 4	(B) 2 : 1 : 2 : 4	(C) 2 : 1 : 2 : 3	(D) 2 : 1 :2 : 3
		STOICHI	OMETRY	
21.	For the reaction $2P + 0$	$Q \rightarrow R, 8 \text{ mol of } P \text{ and } c$	excess of Q will produc	e :
	$(A) 8 \mod of R$	(B) 5 mol of R	$(C) 4 \mod of R$	(D) 13 mol of R
22.	If 1.5 moles of oxygen (A) 27 g	combine with Al to form (B) 40.5 g	$n Al_2O_3$, the weight of Al (C) 54g	used in the reaction is : (D) 81 g

- 23. 74 gm of a sample on complete combustion gives 132 gm CO_2 and 54 gm of H_2O . The molecular formula of the compound may be (A) C_5H_{12} (B) $C_4H_{10}O$ (C) $C_3H_6O_2$ (D) $C_3H_7O_2$
- 24. The mass of CO_2 produced from 620 gm mixture of $C_2H_4O_2$ & O_2 , prepared to produce maximum energy is (Combustion reaction is exothermic) (A) 413.33 gm (B) 593.04 gm (C) 440 gm (D) 320 gm
- 25. The minimum mass of mixture of A_2 and B_4 required to produce at least 1 kg of each product is : (Given At. mass of 'A' = 10; At. mass of 'B' = 120)

$$5A_2 + 2B_4 \longrightarrow 2AB_2 + 4A_2B$$

(A) 2120 gm (B) 1060 gm (C) 560 gm (D) 1660 gm

LIMITING REAGENT

26. The mass of Mg₃N₂ produced if 48 gm of Mg metal is reacted with 34 gm NH₃ gas is Mg + NH₃ \longrightarrow Mg₃N₂ + H₂

(A)
$$\frac{200}{3}$$
 gm (B) $\frac{100}{3}$ gm (C) $\frac{400}{3}$ gm (D) $\frac{150}{3}$ gm

- 27. The mass of P_4O_{10} produced if 440 gm of P_4S_3 is mixed with 384 gm of O_2 is $P_4S_3 + O_2 \longrightarrow P_4O_{10} + SO_2$ (A) 568 gm (B) 426 gm (C) 284 gm (D) 396 gm
- 28. Mass of sucrose $C_{12}H_{22}O_{11}$ produced by mixing 84 gm of carbon, 12 gm of hydrogen and 56 lit. O_2 at 1 atm & 273 K according to given reaction, is $C(s) + H_2(g) + O_2(g) \longrightarrow C_{12}H_{22}O_{11}(s)$ (A) 138.5 (B) 155.5 (C) 172.5 (D) 199.5
- 29. $0.5 \text{ mole of } H_2SO_4 \text{ is mixed with } 0.2 \text{ mole of } Ca (OH)_2$. The maximum number of moles of $CaSO_4$ formed is (A) 0.2 (B) 0.5 (C) 0.4 (D) 1.5

SEQUENTIAL & PARALLEL REACTIONS

- 30. 25.4 g of iodine and 14.2g of chlorine are made to react completely to yield a mixture of ICl and ICl₃. Calculate he number of moles of ICl and ICl₃ formed.
 (A) 0.1 mole, 0.1 mole (B) 0.1 mole, 0.2 mole (C) 0.5 mole, 0.5 mole (D) 0.2 mole, 0.2 mole
- **31.** What weights of P_4O_6 and P_4O_{10} will be produced by the combustion of 31g of P_4 in 32g of oxygen leaving no P_4 and O_2 .

(A) 2.75 g, 219.5 g (B) 27.5 g, 35.5 g (C) 55 g, 71 g (D) 17.5 g, 190.5 g

32. What weight of $CaCO_3$ must be decomposed to produce the sufficient quantity of carbon dioxide to convert 21.2 kg of Na_2CO_3 completely in to $NaHCO_3$. [Atomic mass Na = 23, Ca = 40]

 $\begin{array}{ccc} CaCO_{3} \longrightarrow CaO + CO_{2} \\ Na_{2} CO_{3} + CO_{2} + H_{2}O \longrightarrow 2NaHCO_{3} \\ (A) 100 \text{ Kg} \qquad (B) 20 \text{ Kg} \qquad (C) 120 \text{ Kg} \end{array} \tag{D} 30 \text{ Kg} \end{array}$

33. The following process has been used to obtain iodine from oil-field brines in California. NaI + AgNO₃ \longrightarrow AgI + NaNO₃ ; 2AgI + Fe \longrightarrow Fel₂ + 2Ag 2Fel₂ + 3Cl₂ \longrightarrow 2FeCl₃ + 2I₂ How many grams of AgNO₃ are required in the first step for every 254 kg I₂ produced in the third step. (A) 340 kg (B) 85 kg (C) 68 kg (D) 380 kg

34. 10 g of a sample of a mixture of CaCl₂ and NaCl is treated to precipitate all the calcium as CaCO₃. This Ca CO₃ is heated to convert all the Ca to CaO and the final mass of CaO is 1.62 g. The percent by mass of CaCl₂ in the original mixture is. (A) 32.1 % (B) 16.2 % (C) 21.8 % (D) 11.0 %

MISCELLANEOUS PROBLEM

- **35.** 40 gm of a carbonate of an **alkali metal** or **alkaline earth metal** containing some inert impurities was made to react with excess HCl solution. The liberated CO_2 occupied 12.315 lit. at 1 atm & 300 K. The correct option is
 - (A) Mass of impurity is 1 gm and metal is Be
 - (B) Mass of impurity is 3 gm and metal is Li
 - (C) Mass of impurity is 5 gm and metal is Be
 - (D) Mass of impurity is 2 gm and metal is Mg
- 36. In chemical scale, the relative mass of the isotopic mixture of X atoms (X^{20}, X^{21}, X^{22}) is approximately equal to : $(X^{20} has 99 percent abundance)$ (A) 20.002 (B) 21.00 (C) 22.00 (D) 20.00
- 37. Calculate percentage change in M_{avg} of the mixture, if PCl₅ undergo 50% decomposition in a closed vessel. PCl₅ \longrightarrow PCl₃ + Cl₂

(A) 50% (B) 66.66% (C) 33.33% (D) Zero

- **38.** A compound possess 8% sulphur by mass. The least molecular mass is :(A) 200(B) 400(C) 155(D) 355
- 39. The empirical formula of a compound of molecular mass 120 is CH₂O. The molecular formula of the compound is :
 (A) C₂H₄O₂
 (B) C₄H₈O₄
 (C) C₃H₆O₃
 (D) all of these
- 40. Calculate the molecular formula of compound which contains 20% Ca and 80% Br (by wt.) if molecular weight of compound is 200. (Atomic wt. Ca = 40, Br = 80) (A) Ca_{1/2}Br (B) CaBr₂ (C) CaBr (D) Ca₂Br

41.Cortisone is a molecular substance containing 21 atoms of carbon per molecule. The mass percentage of
carbon in cortisone is 69.98%. Its molar mass is :
(A) 176.5(B) 252.2(C) 287.6(D) 360.1

42. One gram of the silver salt of an organic dibasic acid yields, on strong heating, 0.5934 g of silver. If the weight percentage of carbon in it 8 times the weight percentage of hydrogen and one-half the weight percentage of oxygen, determine the molecular formula of the acid. [Atomic weight of Ag = 108] (A) $C_4H_6O_4$ (B) $C_4H_6O_6$ (C) $C_2H_6O_2$ (D) $C_5H_{10}O_5$

43. The percentage by mole of NO_2 in a mixture of $NO_2(g)$ and NO(g) having average molecular mass 34 is :

(A) 25% (B) 20% (C) 40% (D) 75%

EXERCISE (O-II)

ONE OR MORE THAN ONE MAY BE CORRECT :

- 1. Select the correct statement(s) for $(NH_4)_3PO_4$.
 - (A) Ratio of number of oxygen atom to number of hydrogen atom is 1:3
 - (B) Ratio of number of cation to number of anion is 3 : 1
 - (C) Ratio of number of gm-atom of nitrogen to gm-atoms of oxygen is 3:2
 - (D) Total number of atoms in one mole of $(NH_4)_3PO_4$ is 20.
- 2. 12 g of Mg was burnt in a closed vessel containing 32 g oxygen. Which of the following is not correct.(A) 2 gm of Mg will be left unburnt.
 - (B) 0.75 gm-molecule of O_2 will be left unreacted.
 - (C) 20 gm of MgO will be formed.
 - (D) The mixture at the end will weight 44 g.
- 3. 50 gm of CaCO₃ is allowed to react with 68.6 gm of H₃PO₄ then select the correct option(s)- $3CaCO_3 + 2H_3PO_4 \rightarrow Ca_3(PO_4)_2 + 3H_2O + 3CO_2$ (A) 51.67 gm salt is formed (B) Amount of unreacted reagent = 35.93 gm (C) n_{co2} = 0.5 moles (D) 0.7 mole CO₂ is evolved
- 4. Industrially TNT ($C_7H_5N_3O_6$, explosive material) is synthesized by reacting toluene (C_7H_8) with nitric acid in presence of sulphuric acid. Calculate the maximum weight of $C_7H_5N_3O_6$ which can be produced by 140.5 gm of a mixture of C_7H_8 and HNO₃. $C_7H_8 + 3HNO_3 \longrightarrow C_7H_5N_3O_6 + 3H_2O$
 - (A) 140.5 (B) 113.5 (C) $\frac{140.5}{2}$ (D) 140.5 (3 × 18)
- 5. 'A' reacts by following two parallel reaction to give B & C If half of 'A' goes into reaction I and other half goes to reaction-II. Then, select the correct statement(s)

$$A + N \xrightarrow{I} B + L$$

$$A + N \xrightarrow{\Pi} \frac{1}{2} B + \frac{1}{2}(C) + L$$

(A) B will be always greater than C

- (B) If 2 mole of C are formed then total 2 mole of B are also formed
- (C) If 2 mole of C are formed then total 4 mole of B are also formed
- (D) If 2 mole of C are formed then total 6 mole of B are also formed

6.

Silver metal in ore is dissolved by potassium cyanide solution in the presence of air by the reaction

 $4 \operatorname{Ag} + 8 \operatorname{KCN} + \operatorname{O}_2 + 2\operatorname{H}_2\operatorname{O} \longrightarrow 4 \operatorname{K}[\operatorname{Ag}(\operatorname{CN})_2] + 4 \operatorname{KOH}$

- (A) The amount of KCN required to dissolve 100 g of pure Ag is 120 g.
- (B) The amount of oxygen used in this process is 0.742 g (for 100 g pure Ag)
- (C) The amount of oxygen used in this process is 7.40 g (for 100 g pure Ag)
- (D) The volume of oxygen used at STP is 5.20 litres.

7. Crude calcium carbide, CaC_2 , is made in an electric furnace by the following reaction,

 $CaO + 3C \longrightarrow CaC_2 + CO$

The product contain 85% CaC₂ and 15% unreacted CaO.

- (A) 1051.47 kg of CaO is to be added to the furnace charge for each 1000 kg of CaC_2 .
- (B) 893.8 kg of CaO is to be added to the furnace charge for each 1000 kg of crude product.
- (C) 708.2 kg of CaO is to be added to the furnace charge for each 1000 kg of CaC_2 .
- (D) 910.3 kg of CaO is to be added to the furnace charge for each 1000 kg of crude product.

8. Given following series of reactions:

- (I) $NH_3 + O_2 \longrightarrow NO + H_2O$
- (II) $NO + O_2 \longrightarrow NO_2$
- (III) $NO_2 + H_2O \longrightarrow HNO_3 + HNO_2$
- (IV) $HNO_2 \longrightarrow HNO_3 + NO + H_2O$

Select the correct option(s):

(A) Moles of HNO_3 obtained is half of moles of Ammonia used if HNO_2 is not used to produce HNO_3 by reation (IV)

(B) $\frac{100}{6}$ % more HNO₃ will be produced if HNO₂ is used to produce HNO₃ by reaction (IV) than if

 HNO_2 is not used to produce HNO_3 by reaction (IV)

(C) If HNO₂ is used to produce HNO₃ then $\frac{1}{4}$ th of total HNO₃ is produced by reaction (IV)

(D) Moles of NO produced in reaction (IV) is 50% of moles of total HNO₃ produced.

9. In the quantitative determination of nitrogen, N₂ gas liberated from 0.42 gm of a sample of organic compound was collected over water. If the volume of N₂ gas collected was $\frac{100}{11}$ ml at total pressure 860 mm Hg at 250 K, % by mass of nitrogen in the organic compound is [Aq. tension at 250 K is 24 mm Hg and R = 0.08 L atm mol⁻¹ K⁻¹]

(A)
$$\frac{10}{3}\%$$
 (B) $\frac{5}{3}\%$ (C) $\frac{20}{3}\%$ (D) $\frac{100}{3}\%$

Assertion Reason:

10. Statement $-1: 2A + 3B \longrightarrow C$

4/3 moles of 'C' are always produced when 3 moles of 'A' & 4 moles of 'B' are added.

Statement -2 : 'B' is the limiting reactant for the given data.

(A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.

- (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
- (C) Statement-1 is false, statement-2 is true.
- (D) Statement-1 is true, statement-2 is false.

Match the column :

One type of artifical diamond (commonly called YAG for yttrium aluminium garnet) can be represented 11. by the formula $Y_3Al_5O_{12}$. [Y = 89, Al = 27]

	Column I		Column II
	Element		Weight percentage
(P)	Y	(1)	22.73%
(Q)	Al	(2)	32.32%
(R)	0	(3)	44.95%

The recommended daily dose is 17.6 milligrams of vitamin C (ascorbic acid) having formula $C_6H_8O_6$. 12. Match the following. Given : $\mathrm{N}_{\mathrm{A}}^{}=6\times10^{23}$

	Column I		Column II
(A)	O-atoms present	(P)	10 ⁻⁴ mole
(B)	Moles of vitamin C in 1 gm of vitamin C	(Q)	5.68×10^{-3}
(C)	Moles of vitamin C that should be consumed daily	(R)	3.6×10^{20}

Matching list type : 13.

(C)

(D)

4

2

3

3

1

1

2

4

Column-I

	Colı	ımn-I				Colu	mn-II
	(ma	ss of pro	oduct)				
(P)	2H ₂	$+ O_2 \rightarrow$	· 2H ₂ O			(1)	1.028 g
	1g	1g					
(Q)	$3H_2$	$+ N_2 \rightarrow$	· 2NH ₃			(2)	1.333 g
	1g	1g					
(R)	$H_2 +$	$-\operatorname{Cl}_2 \to 2$	2HC1			(3)	1.125 g
	1g	1g					
(S)	2H ₂	$+ C \rightarrow$	CH ₄			(4)	1.214 g
	1g	1g					
Code	:						
	Р	Q	R	S			
(A)	3	4	1	2			
(B)	2	4	1	3			

		COMPREHI	ENSION:	
14.	NaBr, used to produc	e AgBr for use in photog	raphy can be sel	f prepared as follows :
	$Fe + Br_2$	\rightarrow FeBr ₂	(i)	
	$\mathrm{FeBr}_{2} + \mathrm{Br}_{2}$ -	$\longrightarrow \mathrm{Fe}_{3}\mathrm{Br}_{8}$	(ii)	(not balanced)
	$\mathrm{Fe}_{3}\mathrm{Br}_{8} + \mathrm{Na}_{2}\mathrm{CO}_{3}$ —	\rightarrow NaBr + CO ₂ + Fe ₃ O	4(iii)	(not balanced)
(a)	Mass of iron require	d to produce 2.06×10^3 k	kg NaBr	
	(A) 420 gm	(B) 420 kg	(C) 4.2×10^{5}]	kg (D) 4.2×10^8 gm
(b)	If the yield of (ii) is 6 NaBr	0% & (iii) reaction is 70	% then mass of	iron required to produce 2.06×10^3 kg
	(A) 10^{5} kg	(B) 10 ⁵ gm	(C) 10 ³ kg	(D) None
(c)	If yield of (iii) reaction	on is 90% then mole of C	O ₂ formed when	2.06×10^3 gm NaBr is formed
	(A) 20	(B) 10	(C) 40	(D) None
15.	Preparation of cobalt	Metaborate involves the	following steps c	of reactions:
	(i) $Ca_{2}B_{6}O_{11} + Na_{2}C$	$O_3(aq) \xrightarrow{Boiled} CaCO$, (insoluble) + N	$Na_{2}B_{4}O_{7} + 2NaBO_{2}$
	(ii) Na ₂ B ₄ O7 $\xrightarrow{\Delta}$	$NaBO_2 + B_2O_3$	5	2 7 7 2
	(iii) $CoO + B_2O_2$	\sim Co(BO ₂) ₂ .		
	(Atomic weight : B =	= 11, Co = 59)		
(a)	Mass of $Ca_2B_6O_{11}$ in reaction is	kg required to produce	14.5 kg of Co(BO_2 , assuming 100% yield of each
	(A) 32.2	(B) 40	(C) 28.2	(D) 30
(b)	If the yield of reaction	$(i), (ii) \& (iii) is 60\%, \frac{20}{3}$	$\frac{90}{5}$ % & 32.2 % re	spectively, then mass of $Ca_2B_6O_{11}$ in kg
	required to produce	14.5 kg of Co(BO ₂) ₂ is		
	(A) 250	(B) 200	(C) 190	(D) 150
16.	Water is added to 3.52 and 0.8 gram of a gas fluorine.	2 grams of UF ₆ . The produ s only. The gas [containir	acts are 3.08 grar ng fluorine and h	ns of a solid [containing only U, O & F] ydrogen only], contains 95 % by mass
	[Assume that the en	npirical formula is sam	e as molecular	formula.]
(a)	The empirical formula	a of the gas is	(- 1)	
	(A) HF_2	$(\mathbf{B})\mathbf{H}_{2}\mathbf{F}$	(C) HF	(D) HF_3
(b)	The empirical formula $(A) UF_2O_2$	a of the solid product is (B) UFO ₂	(C) UF ₂ O	(D) UFO
(c)	The percentage of flue (A) 66.66 %	orine of the original comp (B) 33.33 %	oound which is co (C) 50 %	onverted into gaseous compound is (D) 89.9 %

		EXERCI	SE - JEE MAIN		
1.	The weight of 2.01	$\times 10^{23}$ molecules of C	CO is–		[AIEEE 2002]
	(1) 9.3 g	(2) 7.2 g	(3) 1.2 g	(4) 3 g	
2.	In an organic compo Molecular formula ((1) C.H.N.	ound of molar mass108 can be : (2) C-H. N	8 g mol ⁻¹ C, H and N aton (3) C.H.N.	ns are present in 9 (4) C.H.,N	: 1 : 3.5 by weight. [AIEEE 2002]
3.	If we consider that unit, the mass of one (1) be a function of t (2) remain unchange (3) increase two fold (4) decrease twice	1/6, in place of 1/12, 1 e mole of the substanc the molecular mass of ed d	mass of carbon atom is ta e will :- Thesubstance	aken to be the rela	ative atomic mass [AIEEE-2005]
4.	How many moles of (1) 3.125×10^{-2}	f magnesium phospha (2) 1.25×10^{-2}	te, Mg ₃ (PO ₄) ₂ will contai (3) 2.5×10^{-2}	n 0.25 mole of ox (4) 0.02	ygen atoms? [AIEEE 2006]
5.	A transition metal M of chlorine the form $(1) \text{MCl}_2$	A forms a volatile chlo ula of the metal chloric (2) MCl ₄	ride which has a vapour c le will be (3) MCl ₅	lensity of 94.8. If i [AIEE] (4) MCl ₃	t contains 74.75% E 2012 (Online)]
6.	The ratio of number $16.0 \text{ g oxygen } (\text{O}_2) \text{ i}$ (Atomic mass : C =	er of oxygen atoms (C is :- = 12, O = 16 and Avog	D) in 16.0g ozone (O_3), 2 gadro's constant $N_A = 6.0$	28.0 g carbon mc) × 1023 mol ⁻¹) [AIEE	onoxide (CO) and E 2012 (Online)]
	(1) 3 : 1 : 1	(2) 1 : 1 : 2	(3) 3 : 1 : 2	(4) 1 : 1 : 1	
7.	A gaseous hydroca formula of the hydro	rbon gives upon com ocarbon is	ubustion 0.72 g of water	and 3.08 g of Co [J]	D ₂ . The empirical EE(Main)-2013]
	$(1) C_2 H_4$	(2) $C_{3}H_{4}$	(3) $C_6 H_5$	(4) $C_7 H_8$	
8.	The ratio of masses of their molecule is : (1) 1 : 8	of oxygen and nitroge (2) 3 : 16	n in a particular gaseous (3) 1 : 4	mixture is 1 : 4. T [J (4) 7 : 32	he ratio of number EE(Main)-2014]
9.	The molecular form (Mol. Wt. 206) What per gram resin?	ula of a commercial re at would be the maxim	sin used for exchanging ionum uptake of Ca ²⁺ ions	ons in water softer by the resin when J	ning is C ₈ H ₇ SO ₃ Na expressed in mole EE(Main)-2015]
	$(1) \frac{1}{103}$	(2) $\frac{1}{206}$	$(3) \frac{2}{309}$	$(4) \frac{1}{412}$	

10. In Carius method of estimation of halogens, 250 mg of an organic compound give 141 mg of AgBr. The percentage of bromine in the compound is : (at. mass Ag = 108; Br = 80)

[JEE(Main)-2015]

(1) 24 (2) 36 (3) 48 (4) 60

11.	The most abundant elements by mass in the body of a healthy human adult are : Oxygen (61.4%) ; Carbon (22.9%), Hydrogen (10.0%) ; and Nitrogen (2.6%). The weight which a 75 kg person would gain if all ¹ H atoms are replaced by ² H atoms is					
	(1) 15 kg	(2) 37.5 kg	(3) 7.5 kg	(4) 10 kg		
12.	1 gram of a carbonate molar mass of M_2CO_2 (1) 1186	e (M_2CO_3) on treatmen $_3$ in g mol ⁻¹ is : (2) 84.3	t with excess HCl prod (3) 118.6	uces 0.01186 mole of CO ₂ . the [JEE(Main)-2017] (4) 11.86		
13.	The reaction of mass percent of C and H of an organic compound $(C_X H_Y O_Z)$ is 6 : 1, If one molecule of the above compound $(C_X H_Y O_Z)$ contains half as much oxygen as required to burn on molecule of compound $C_X H_Y$ completely to CO_2 and H_2O . The empirical formula of compound $C_X H_Y O_Z$ is : [JEE(Main)-2018]					
	(1) $C_2 H_4 O_3$	(2) $C_{3}H_{6}O_{3}$	$(3)C_2H_4O$	(4) $C_{3}H_{4}O_{2}$		
14.	For the following read	ction, the mass of water	produced from 445 g of	⁵ C ₅₇ H ₁₁₀ O ₆ is:		
	$2C_{57}H_{110}O_6(s) + 163$	$O_2(g) \longrightarrow 114 CO_2(g)$) + 110 H ₂ O(l)	[JEE(Main)-2019(Jan)]		
	(1) 490 g	(2) 890 g	(3) 445 g	(4) 495 g		
15.	A 10 mg effervescent T = 298.15 K and p = percentage of sodium	tablet containing sodiur 1 bar. If molar volume bicarbonate in each tab	n bicarbonate and oxalio of CO ₂ is 25.0 L under s olet ? [Molar mass of Na	c acid releases 0.25 ml of CO_2 at such condition, what is the $0.000 \text{ HCO}_3 = 84 \text{ g mol}^{-1}$]		
				[JEE(Main)-2019(Jan)]		
	(1) 0.84	(2) 8.4	(3) 16.8	(4) 33.6		
16.	An organic compound 4 moles of H_2O and 1	d is estimated through D mole of nitrogen gas. Th	umus method and was f ne formula of the compo	found to evolve 6 moles of CO_2 , bund is : LIEE(Main)-2019(Jan)]		
	(1) $C_6 H_8 N_2$	(2) $C_6 H_8 N$	(3) $C_{12}H_8N_2$	(4) $C_{12}H_8N$		
17.	The percentage comp	osition of carbon by mol	e in methane is :	[JEE(Main)-2019(April)]		
	(1) 25%	(2) 75%	(3) 20%	(4) 80%		
18.	For a reaction,					
	$N(\sigma) + 3H(\sigma) \rightarrow 2N$	H (g): identify dihydro	gen (H) as a limiting re	eagent in the following reaction		

 $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$; identify dihydrogen (H₂) as a limiting reagent in the following reaction mixtures. [JEE(Main)-2019(April)]

- (1) 28 g of N_2 + 6 g of H_2 (2) 56 g of N_2 + 10 g of H_2
- (3) 14 g of N_2 + 4 g of H_2 (4) 35 g of N_2 + 8 g of H_2

19. The minimum amount of $O_2(g)$ consumed per gram of reactant is for the reaction : (Given atomic mass : Fe = 56, O = 16, Mg = 24, P = 31, C = 12, H = 1)

[JEE(Main)-2019(April)]

(1) $4Fe(s)+3O_2(g) \rightarrow 2Fe_2O_3(s)$ (2) $C_3H_8(g)+5O_2(g) \rightarrow 3CO_2(g)+4H_2O(l)$ (3) $2Mg(s)+O_2(g) \rightarrow 2MgO(s)$ (4) $P_4(s)+5O_2(g) \rightarrow P_4O_{10}(s)$

20. 5 moles of AB₂ weigh 125×10^{-3} kg and 10 moles of A₂B₂ weigh 300×10^{-3} kg. The molar mass of A(MA) and molar mass of B (MB) in kg mol⁻¹ are:

[JEE(Main)-2019(April)]

(1) $M_A = 5 \times 10^{-3}$ and $M_B = 10 \times 10^{-3}$ (2) $M_A = 50 \times 10^{-3}$ and $M_B = 25 \times 10^{-3}$ (3) $M_A = 25 \times 10^{-3}$ and $M_B = 50 \times 10^{-3}$ (4) $M_A = 10 \times 10^{-3}$ and $M_B = 5 \times 10^{-3}$

- **21.**25 g of an unknown hydrocarbon upon burning produces 88 g of CO2 and 9 g of H2O. This unknown
hydrocarbon contains:[JEE(Main)-2019(April)]
 - $(1)\,18\,g\,of\,carbon$ and $7\,g\,of\,hydrogen$
 - (2) 22 g of carbon and 3 g of hydrogen
 - (3) 24 g of carbon and 1 g of hydrogen
 - (4) 20 g of carbon and 5 g of hydrogen
- **22.** At 300 K and 1 atmospheric pressure, 10 mL of a hydrocarbon required 55 mL of O_2 for complete combustion, and 40 mL of CO_2 is formed. The formula of the hydrocarbon is:

[JEE(Main)-2019(April)] (4) C_4H_6

(1) $C_4 H_7 Cl$ (2) $C_4 H_{10}$ (3) $C_4 H_8$

EXECISE - JEE ADVANCED

1.	How many moles of e-	[JEE '2002 (Scr), 1]		
	(A) 6.023×10^{23}	(B) $\frac{1}{9.108} \times 10^{31}$	(C) $\frac{6.023}{9.108} \times 10^{54}$	(D) $\frac{1}{9.108 \times 6.023} \times 10^8$

- 2. Calculate the amount of Calcium oxide required when it reacts with 852 g of P_4O_{10} . [JEE 2005] 6CaO + $P_4O_{10} \longrightarrow 2 Ca_3 (PO_4)_2$
- Given that the abundances of isotopes ⁵⁴Fe, ⁵⁶Fe and ⁵⁷Fe are 5%, 90% and 5%, respectively, the atomic mass of Fe is : [JEE 2009]

(A) 55.85 (B) 55.95 (C) 55.75 (D) 56.05

	AN	NSW	ER KEY
	E	XERC	CISE (S-1)
1.	6.023×10^{13}	2.	$1.99 \times 10^{-23} g$
3.	24g	4.	(i) $\frac{10^{20}}{N_A}$ moles ,(ii) 3200 amu , (iii) $14 \times 1.66 \times 10^{-24}$ g
			$(iv) 3N_A, 9N_A, (v) 6N_A$
5.	3.2 g	6.	180 g
7.	2.5 g	8.	$7.227 imes 10^{24}$.
9.	$10 \times 6.023 \times 10^{23}, 8 \times 6.023 \times 10^{23},$	8 × 6.0	0.023×10^{23} .
10.	68 mole	11.	$5.34 imes 10^{6}$.
12.	20 N _A	13.	292 g
14.	2270 L	15.	7.5 moles
16.	24	17.	58.8 g
18.	27.6 gm	19.	1.0×10^{19}
20.	3	21.	50.14 L
22.	(i) 1/6 mole (ii) 5/12 mole	23.	0.48
24.	0.1185	25.	0.25 mole
26.	9.5	27.	$w_c = 24 \text{ gm}; W_{CCl_4} = 154 \text{ gm}$
28.	5.6	29.	Al = 60%; Mg = 40%
30.	$CaCO_3 = 28.4\%; MgCO_3 = 71.6\%$	31.	NaHCO ₃ = 14.9 %; Na ₂ CO ₃ = 85.1 %
32.	42 g	33.	63 % , 37%
34.	10 %	35.	12.3
36.	39.18	37.	4.48 litre
38.	61.5 gm	39.	19.4 gm
40.	(a) 0.050 mol , (b) 0.050 mol	41.	640.0
42.	441 gm	43.	(a)59.17 gm (b) 61.97 gm
44.	1.14 gm	45.	0196
46.	92.7 gm/mole	47.	35.4527
48.	10	49.	$C_{6}H_{4}C_{12}$ 50. 4

			E	XERC	CISE (S-	·II)			
1.	12.9 gm			2.	(i) 0.5,	0.5; (ii) 2, 1	(iii) 1, 2	2	
3.	% CO = 65	5.625;%	$CO_2 = 34.375$	4.	10:0.6	66 : 1			
5.	%NaCl = 7	7.8%	_						
6.	(i) $Fe_{2}O_{3} + $	$2 \text{Al} \longrightarrow$	$Al_2O_3 + 2Fe;$ (i	i) 80 : 2 [°]	7; (iii)10,0	000 units			
7.	59.72%								
8.	Ans. 5.74 g	m							
	$[Cr(H_2O)_5 O_{10}]$ 0.02 mo	C1].C1 ₂ . H ol	$_{2}$ O + 2AgNO ₃ (a	aq.)— 0.041	→2AgCl、 mol	\downarrow + [Cr(H ₂ O) ₅	,Cl](NC	$(D_3)_2$	
9.	Ans. $6 \times 10^{\circ}$)20							
	$M_2CO_3 \Rightarrow$	$48 = \left(\frac{1}{2N}\right)$	$\left(\frac{48}{1+60}\right) \times 100 \Rightarrow$	> 2M +	60 = 100	, M 200			
	$n = \frac{5 \times 10^{-3}}{100}$	$= 5 \times 10^{\circ}$	⁻⁵ mol						
	$n_{M} = 10 \times 1$ $n0_{M} = 6.00$	$0^{-5} \text{ mol} =$ × 10 ¹⁹ ato	= 10 ⁻⁴ mol						
10.	Ans.(A B)								
100	2 5	3							
	$\frac{2.5}{24x+14y}$ ×	$x = \frac{3}{40}$	× 1						
11.	222 gm		12. 1.145	58		13. 7.5 ×	Na		
			E	XERC	CISE (O	9-I)			
1.	А	2.	А	3.	B	4.	D	5.	С
6.	C	 7.	В	8.	C	9.	B	10.	D
11.	С	12.	С	13.	В	14.	С	15.	В
16.	А	17.	В	18.	С	19.	С	20.	С
21.	С	22.	С	23.	С	24.	С	25.	А
26.	А	27.	В	28.	В	29.	А	30.	А
31.	В	32.	В	33.	A	34.	А	35.	В
36.	А	37.	С	38.	В	39.	В	40.	В
41.	D	42.	В	43.	А				
			E	XERC	ISE (O	-II)			
1.	A,B	2.	А	3.	A,B,C	4.	В	5.	A,D
6.	A,C,D	7.	A,B	8.	A,C,D	9.	А	10.	С
11.	(A)→ R ; (B	P;(C)	→Q	12.	(A)→F	R;(B)→Q;(C)→P		
13.	А	14.	(a) B (b) C (c)B	15.	(a)A(b)A	16.	(a) C (b) A (c	c)A

EXERCISE - JEE MAIN									
1.	(1)	2.	(1)	3.	(2)	4.	(1)	5.	(2)
6.	(4)	7.	(4)	8.	(4)	9.	(4)	10.	(1)
11.	(3)	12.	(2)	13.	(1)	14.	(4)	15.	(2)
16.	(1)	17.	(3)	18.	(2)	19.	(1)	20.	(1)
21.	(3)	22.	(4)						

EXERCISE - JEE ADVANCED

1.	(D)	2.	1008 g	3.	(B)
----	-----	----	--------	----	-----

PHYSICAL CHEMISTRY

HINT & SOLUTIONS : MOLE CONCEPT

EXERCISE # S-I

- 1. 10^{10} grains are distributed in 1 sec $\Rightarrow 6.023 \times 10^{23}$ grains are distributed in $\frac{6.023 \times 10^{23}}{10^{10}}$ sec $= 6.023 \times 10^{13}$ seconds.
- 2. Mass of 6.023×10^{23} atom = 12 gm \Rightarrow Mass of 1 atom = $\frac{12}{6.023 \times 10^{23}}$ gm = 1.99×10^{-23} gm
- 3. Weight of 6.023×10^{23} atoms = 12 gm \Rightarrow Weight of 12.046×10^{23} atom = $\frac{12 \times 12.046 \times 10^{23}}{6.023 \times 10^{23}}$ = 24 gm

4. (i) Number of moles of Cu atom in 10²⁰ atoms of Cu =
$$\left(\frac{10^{20}}{N_A}\right)$$
 moles

- (ii) Mass of 200 atoms of ${}^{16}_{8}$ O in amu \rightarrow Mass of 1 atom = 16 amu \Rightarrow Mass of 200 atoms = 16 × 200 amu = 3200 amu
- (iii) Mass of 100 atoms of ${}^{14}_{7}$ N in gm \rightarrow Mass of 1 atom of ${}^{14}_{7}$ N in gm = 14×1.66×10⁻²⁴ gm \Rightarrow Mass of 100 atom of ${}^{14}_{7}$ N = 14 × 100 × 1.66 × 10⁻²⁴ gm i.e. 1400 × 1.66 × 10⁻²⁴ gm
- (iv) Number of molecules in 54 gm H₂O = $\left(\frac{54}{18} \times N_A\right) = 3N_A$ Number of atoms in 54 gm H₂O = $(3N_A) \times 3 = 9N_A$

(v) Number of atoms in 88 gm
$$CO_2 = \left(\frac{88}{44}\right) \times 3N_A$$
 i.e. $6N_A$

5. Mass of O atoms in 6 gm CH₃COOH = $n_{CH_3COOH} = \frac{6}{60}$ i.e. $\frac{1}{10}$ In 1 mole of CH₃COOH, mass of O atom = 32 gm \Rightarrow Mass of O atom in $\frac{1}{10}$ mole CH₃COOH = $\frac{32}{10}$ i.e. 3.2 gm

6.
$$n_{CuSO_4.5H_2O} = \frac{499}{249.5} = 2 \text{ mole}$$

1 mole of CuSO_4.5H_2O contains 90 gm H_2O
 $\Rightarrow 2 \text{ mole of CuSO}_4.5H_2O \text{ contains } (90 \times 2) \text{ i.e. } 180 ; gm H_2O.$

- 7. 1 mole of $Na_2SO_4.7H_2O$ contains $11N_A'O'$ atoms
 - $\Rightarrow 6.023 \times 10^{22} \text{ atom of 'O' are present in } \frac{6.023 \times 10^{22}}{11 \times N_A} = \frac{6.023 \times 10^{22}}{11 \times 6.022 \times 10^{23}} = \frac{1}{110} \text{ mole}$ i.e. 2.5 gm
- 8. Number of Nucleon present in 12 gm of ${}^{12}C$ atoms = 12 N_A = $12 \times 6.023 \times 10^{23}$ = 7.227×10^{24}
- $\begin{array}{lll} \textbf{9.} & \text{In 1 mole of } {}^{16}\text{O}^{-2} \text{ ions} \\ \text{Number of Electrons} = 10 N_{\text{A}} \text{ i.e. } 10 \times 6.023 \times 10^{23} \\ \text{Number of Protons} = 8 N_{\text{A}} & \text{i.e. } 8 \times 6.023 \times 10^{23} \\ \text{Number of Neutrons} = 8 N_{\text{A}} & \text{i.e. } 8 \times 6.023 \times 10^{23} \\ \end{array}$
- **10.** Mass of liquid mercury = 13.6 gm

$$\Rightarrow$$
 Moles of liquid mercury = $\frac{13.6}{200}$ i.e. 0.068

 \Rightarrow Moles of liquid mercury in 1 lit of the metal = $0.068 \times 1000 = 68$ mole

11. Mass of C₂H₆ sample =
$$\left(\frac{10^7}{N_A}\right)$$
 moles of CH₄ i.e. $\left(\frac{16 \times 10^7}{N_A}\right)$ g
 \Rightarrow Mole of C₂H₆ sample = $\left(\frac{16 \times 10^7}{N_A \times 30}\right)$
 \Rightarrow Number of C₂H₆ molecules in sample = $\left(\frac{16 \times 10^7}{N_A \times 30}\right) \times N_A$ i.e. 5.34 × 10⁶

12. Number of H-atom removed =
$$(30N_A + 10N_A) = 40 N_A$$

 \Rightarrow Number of H₂ molecules formed = $(20 N_A)$

- 13. $\begin{array}{ll} \text{MnO}_{2(\text{s})} + 4\text{HCl} \longrightarrow \text{Cl}_{2(\text{g})} + \text{MnCl}_{2(\text{aq})} + 2\text{H}_2\text{O}(l) \\ 1 \text{ mole } \text{Cl}_2 \text{ is produced from 4 mole HCl} \\ \Rightarrow & 142 \text{ gm Cl}_2 \text{ or 2 mole } \text{Cl}_2 \text{ is produced from 8 mole HCl} & \text{i.e. } (8 \times 36.5) = 292 \text{ gm HCl} \\ \end{array}$
- 14. $C + O_2 \longrightarrow CO_2$ $n_C \rightarrow \frac{1.2 \times 10^3}{12}$ i.e. 100 mole.

Mole of O_2 needed for 1 mole C = 1 mole

- \Rightarrow Mole of O₂ needed for 100 mole C = 100 mole
- \Rightarrow Volume of O₂ needed = $100 \times 22.7 = 2270$ lits.

$$15. \quad C_5H_{12}O + \frac{15}{2}O_2 \longrightarrow 5CO_2 + 6H_2O$$

 \Rightarrow Moles of O₂ required to burn 1 mole of this compound completely is 7.5 moles.

16. Al₄C₃ + 12H₂O → 3CH₄ + 4Al(OH)₃
n_{CH₄} =
$$\frac{11.35}{2.70} = \frac{1}{2}$$
 mole
3 mole CH₄ is produced from 1 mole Al₄C₃.
⇒ $\frac{1}{2}$ mole CH₄ is produced from 1 mole $(\frac{1}{3} \times \frac{1}{2})$ mole Al₄C₃
i.e. $\frac{1}{6}$ mole Al₄C₃ or $\frac{1}{6} \times 144$
i.e. 24 gm Al₄C₃
17. 2H₃PO₄ → H₄P₂O₇ + H₂O
n<sub>H₄P₅O₇ → $\frac{534}{178} = 0.3$ mole
1 mole H₄P₂O₇ is obtained from 2 mole H₃PO₄
⇒ 0.3 mole H₃PO₄ → 106 mole H₃PO₄
i.e. (0.6 × 98) = 58.5 g H₃PO₄
18. 3NO₂ + H₂O → 2HNO₃ + NO
n_{NNO₄} = $\frac{25.2}{63} = 0.4$
2 mole HNO₃ is produced from 3 mole NO₂
⇒ 0.4 mole HNO₃ is produced from $(\frac{3}{2} \times 0.4) = 0.6$ mole NO₂
or (0.6 × 46)g NO₂
i.e. 27.6g NO₂
19. U + 3F₂ → UF₆
(Excess)
n_{UF₆} = $\frac{2 \times 10^{-3}}{352}$ ⇒ n_{UF₆} = 5.6×10^{-6}
1 mole UF₆ is obtained from 3 mole F₂
⇒ 5.6×10^{-6} mole UF₆ is obtained from $-\rightarrow 5.6 \times 10^{-6} \times 3 \times 6.023 \times 10^{23}$
= 101.1×10^{17}
= 1×10^{19}
20. 7XeF₆ + 3I₂ → 6IF₇ + 7Xe
7 mole XeF₆ produces 6 mole IF₇
⇒ 3.5×10^{-3} mole XeF₆ produces $(\frac{6}{7} \times 3.5 \times 10^{-3})$ i.e. 3 m mol IF₇</sub>

21. 2NH₄NO₃ → 2N₂ + O₂ + 4H₂O.
mole initial
$$\frac{1}{5}$$
 0 0 0 0
moles final 0 $\frac{1}{5}$ $\frac{1}{10}$ $\frac{2}{5}$
 \Rightarrow n_T = $\left(\frac{1}{5} + \frac{1}{10} + \frac{2}{5}\right) = \left(\frac{7}{10}\right)$
Apply = Pv = nRT \Rightarrow v = $\frac{nRT}{P}$ = $\left(\frac{7}{10} \times 0.0821 \times 873\right)$ = 50.14 litre
22. 3CaCO₃ + 2H₃PO₄ → Ca₃(PO₄)₂ + 3H₂O + 3CO₂
Moles $\rightarrow \frac{50}{100}$ $\frac{73.5}{98}$
 $= \frac{1}{2}$ $\frac{3}{4}$
 $\frac{moles}{SC} \rightarrow \frac{1}{6}$ $\frac{3}{8}$
 \Rightarrow CaCO₃ is L. R
(i) Amount of Ca₃(PO₄)₂ formed = $\frac{1}{6}$ mole
(ii) Amount of unreacted reagent = $\left(\frac{3}{4} - \frac{1}{3}\right) = \frac{9-4}{12} = \left(\frac{5}{12}\right)$ moles.
23. $\frac{4A + 2B + 3C}{SC} \rightarrow A_4 B_2 C_3$
 $\frac{Moles}{SC} \rightarrow \frac{2}{4}$ $\frac{1.2}{1}$ $\frac{1.44}{3}$
 \Rightarrow C is L.R
 \Rightarrow moles of product formed = $\frac{1.44}{3} = 0.48$ moles.
24. $\frac{4KO_2 + 2H_2O \longrightarrow 4 \text{ KOH + 3O_2}}{Moles \rightarrow 0.158}$.10
 $\frac{moles}{SC} \rightarrow \frac{0.158}{4}$.10
 $\frac{moles}{SC} \rightarrow \frac{0.158}{4}$.10
 $\frac{moles}{SC} \rightarrow 0.158$ L.R.
 \Rightarrow Moles of O₂ produced is $\frac{3\times0.158}{4}$ i.e. 0.1185 mole

 $6 \text{ LiH} + 8\text{BF}_3 \longrightarrow 6 \text{ LiBF}_4 + B_2H_6$ 25. Moles : 2 2 $\frac{2}{8}$ $\frac{2}{6}$ moles \rightarrow S.C BF₃ is L.R. \Rightarrow Moles of B₂H₆ prepared $=\frac{1}{8} \times 2 = \frac{1}{4}$ moles i.e. 0.25 mol \Rightarrow $3\text{TiO}_2 + 4\text{C} + 6\text{Cl}_2 \longrightarrow 3\text{TiCl}_4 + 2\text{CO}_2 + 2\text{CO}_2$ 26. 4.32 5.76 7.1 moles 80 12 71 i.e. 0.054 0.48 0.1 $\frac{0.054}{3} \frac{0.48}{4}$ $\frac{0.1}{6}$ moles \rightarrow S.C \Rightarrow Cl_2 is L.R. Amount of TiCl₄ obtained = $\left(\frac{3}{6} \times 0.1\right) \times 190 = 9.5$ g \Rightarrow 27. $C + 2Cl_2 \longrightarrow CCl_4$ $\frac{1\overline{42}}{71}$ moles $\rightarrow \frac{36}{12}$ 3 2 i.e. moles $\rightarrow 3$ 1 S.C Cl₂ is L.R. \Rightarrow Mass of CCl₄ produced = 1×154 i.e. 154 gm (i) Remaining mass of reactants = $(3 - 1) \times 12 = 24$ gm (ii)

28.

 $\begin{array}{rcl} 2SO_2 + O_2 + 2H_2O \longrightarrow 2H_2SO_4 \\ moles \rightarrow & 5.6 & 4.8 \ (excess) \\ \hline \frac{moles}{S.C} \rightarrow & \frac{5.6}{2} & \frac{4.8}{1} \\ \Rightarrow & SO_2 \ is \ L.R. \\ \Rightarrow & maximum \ number \ of \ moles \ of \ H_2SO_4 \ that \ can \ be \ obtained = 5.6 \ mole. \end{array}$

29.
$$2Al + 6HCl \longrightarrow 2AlCl_3 + 3H_2$$

Moles $\frac{x}{27}$ (excess)
 \Rightarrow moles of H₂ obtained = $\left(\frac{3}{2} \times \frac{x}{27}\right)$ (1)
mg + 2HCl \longrightarrow MgCl₂ + H₂

Moles $\rightarrow \frac{(1-x)}{24}$ (excess) moles of H₂ obtained = $\left(\frac{1-x}{24}\right)$ (2) \Rightarrow Total moles of H₂ obtained = $\left(\frac{3x}{54} + \frac{1-x}{24}\right)$ (3) \Rightarrow Now, $n_{\rm H_2} = \frac{1.12}{22.4}$ i.e. 0.05(4) \Rightarrow Now, equation (3) & (4) $\frac{3x}{54} + \frac{1-x}{24} = 0.05$ $\frac{12x+9-9x}{216} = 0.05$ \Rightarrow $m = 10.8 - 9 = \frac{1.8}{3} = 60\%$ 3x + 9 = 10.8 $3x = 1.8 \implies x = 0.6$ % by mass Al $\longrightarrow 60\%$ & % by mass mg $\longrightarrow 40\%$ \Rightarrow \Rightarrow

$$CaCO3 \longrightarrow CaO + CO_2$$

moles $\rightarrow a$ 0 0
- a a
MgCO_3 \longrightarrow MgO + CO_2
moles $\rightarrow b$ 0 0
- b b

Now, According to question,

Mass of oxide produced is exactly half as much as the original sample.

$$\Rightarrow \qquad \mathbf{a} \times \mathbf{56} + \mathbf{b} \times \mathbf{80} = \frac{1}{2} (100 \ \mathbf{a} + \mathbf{84} \ \mathbf{b})$$

 \Rightarrow b = 3a

30.

$$\Rightarrow \qquad \% \text{ weight of } CaCO_3 = \left(\frac{100a}{100a + 84b}\right) \times 100$$
$$= \left(\frac{100a}{352a} \times 100\right) = 28.4 \%$$

% weight of $MgCO_3 = (100 - 28.4)$ i.e 71.6 %

31. Na₂CO₃ → x Let mass of Na₂CO₃ be x gm 2 mass of NaHCO₃ be (2-x) gm
2Na(HCO₃) → Na₂CO₃ + CO₂ + H₂O
mole →
$$\frac{2-x}{84}$$
 $\frac{1.89 - x}{406}$
Now, $\frac{\text{moles of NaHCO_3}}{2} = \frac{\text{moles of Na_2CO_3}}{1}$
 $\frac{(2-x)}{84 \times 2} = \frac{1.89 - x}{106}$
⇒ 212 - 106x = 84 × 2 (1.89 - x)
⇒ 212 - 106x = 317.52 - 168 x
⇒ 168x - 106x = 317.52 - 212
⇒ 62x = 105.52
⇒ x = $\frac{105.52}{62} = 1.70 \text{ g}$
% by mass of Na₂CO₃ = $\frac{1.70}{2} \times 100 = 85.1 \%$

% by mass of $NaHCO_3 = 14.9\%$

32. let CaCO₃ be x gm
2 MgCo₃ be (92 - x) gm
CaCO₃
$$\longrightarrow$$
 CaO + CO₂
Mole: $\frac{x}{100}$ 0 0
 $-\frac{x}{100} \frac{x}{100}$
MgCO₃ \longrightarrow MgO + CO₂
Mole $\rightarrow \left(\frac{92 - x}{84}\right)$ - -
 $-\left(\frac{92 - x}{84}\right)$ $\left(\frac{92 - x}{84}\right)$
Now, weight of residue = 48
 $\Rightarrow \frac{x}{100} \times 56 + \left(\frac{92 - x}{84}\right) \times 40 = 48$
 \Rightarrow on solving we get x = 50
 \Rightarrow weight of MgCO₃ = 42 gm.

33. NaCl \longrightarrow x Let NaCl be x gm & NaHCO₃ be (4 - x) gm 2NaHCO₃ \longrightarrow Na₂CO₃ + CO₂ + H₂O Mole $\rightarrow \frac{4-x}{84}$ 0 0 0 $-\frac{(4-x)}{84\times 2}$

Now, According to question

$$44 \times \frac{(4-x)}{84 \times 2} = 0.66$$

$$\Rightarrow \quad (4-x) = \frac{0.66 \times 84}{22 \times 100}$$

$$\Rightarrow \quad x = 1.48 \text{ gm} \quad \& \quad \text{Weight of NaHCO}_3 = 2.52 \text{ gm}$$

$$\Rightarrow \quad \% \text{ by mass of NaCl} \rightarrow \frac{1.48}{4} \times 100 = 37 \%$$

$$\& \qquad \% \text{ by mass of NaHCO}_3 \text{ is } 63\%$$

34.
$$CaCO_3 \longrightarrow CaO + CO_2$$

 $n_{CO_2} = \frac{11.35}{22.70} \rightarrow \frac{1}{2}$ mole
 $n_{CaCO_3} \longrightarrow 2$ mole
 \Rightarrow 1 mole CaCO_3 produces 1 mole CO_2
 \Rightarrow 2 mole CaCO_3 produces 2 mole CO_2
But produced mole is $\frac{1}{2}$ mole
 \Rightarrow % of Ca in lime stone sample = $\left(\frac{20}{200} \times 100\right)$ i.e. 10%

35. Sulphur present in 1.30 gm per 100g of coal. $S + O_2 \longrightarrow SO_2$

$$Moles \rightarrow \frac{1.30}{32} \text{ (excess)}$$

$$- \left(\frac{1.30}{32}\right)$$
⇒ weight of SO₂ produced = $\left(\frac{1.30}{32} \times 64\right) = 2.60 \text{ gm}$

100 g coal sample produced 2.60 gm SO_2

$$\Rightarrow$$
 474 tons will produced $\frac{2.60}{100} \times 474$ i.e. (12.3 tons)

36. $2KClO_3 \longrightarrow 2KCl + 3O_2$ The loss in weight of sample is because of O_2 gas produced. 2 mole KClO₃ is producing 3 mole O_2 *.*.. i.e. for 1 mole loss is $\left(\frac{3}{2} \times 32\right)$ i.e 48g O₂. \Rightarrow % loss in weight = $\left(\frac{48}{122.5}\right) \times 100 = 39.18$ % 37. $CaCO_3 + 2HCl \longrightarrow CaCl_2 + CO_2 + H_2O$ 1 mole CaCO₃ will produce 1 mole CO₂ i.e. $\left(.8 \times \frac{1}{4}\right)$ mole will produce $\left(.8 \times \frac{1}{4}\right)$ mole CO₂ Volume of CO_2 produced = $.2 \times 22.4$ litres = 4.48 litres. \Rightarrow $C_6H_{12}O \xrightarrow{\text{ cons.H}_2SO_4} C_6H_{10} + H_2O$ 38. 1 .25 .75 .75 1 mole $C_6H_{12}O$ produces 0.75 mole C_6H_{10} . \Rightarrow 100 gm C₆H₁₂O produces $(.75 \times 82)$ g C₆H₁₀ \Rightarrow \Rightarrow $61.5 \text{ g C}_6\text{H}_{10}$ is produced. $2CH_{3}COCH_{3} + 6CaOCl_{2} \longrightarrow Ca(CH_{3}COO)_{2} + 2CHCl_{3} + 3CaCl_{2} + 2Ca(OH)_{2}$ 39. x(0.75)Х $\mathbf{x} \times .75 = \frac{30}{119.5}$ \Rightarrow $x = \frac{30}{119.5 \times 0.75} \qquad \Rightarrow x = 0.334 \text{ mole}$ \Rightarrow \Rightarrow Mass of $CH_3COCH_3 = 0.334 \times 58$ i.e. 19.4 gm 40. $Br_2 + Cl_2 \longrightarrow 2BrCl$ mole. 0.025 0.025 $(0.025 \times 2) \times .8 = 0.04$ (i) amount of BrCl formed = 0.04(ii) Br_2 left unchanged = 0.025 - 0.02 = 0.00541. $S_8 + 8O_2 \longrightarrow 8SO_2$ $2SO_2 + O_2 \longrightarrow 2SO_3$ 1 mole $S_8 = 8$ mole SO_2(1) 2 mole $SO_2 = 2$ mole SO_3 \Rightarrow 8 mole $SO_2 = 8$ mole SO_3 (2) i.e. From (1) & (2) SO₃ obtained from 1 mol of $S_8 = (8 \times 80)g$ SO₃ i.e. 640g SO₃

42. 2Pbs + 3O₂ → 2PbO + 2SO₂
3SO₂ + 2HNO₃ + 2H₂O → 3H₂SO₄ + 2NO
n_{Pbs} =
$$\frac{1075.5}{239.2}$$
 = 4.49
2 mole Pbs = 2 mole SO₂(1)
3 mole SO₂ = 3 mole H₂SO₄
⇒ 2 mole SO₂ = 2 mole H₂SO₄(3)
From (1) & (3)
1 mole Pbs = 1 mole H₂SO₄
⇒ 4.49 mole Pbs = 4.49 mole H₂SO₄ or 4.50 mole H₂SO₄
⇒ mass of H₂SO₄ = (4.50 × 98) = 441 g.m
43. 4KO₂ + 2H₂O → 3O₂ + 4KOH
KOH + CO₂ → KHCO₃
(a) 3 mole O₂ is produced by $\frac{4}{3}$ mole KO₂
⇒ 1 mole O₂ is produced by $\frac{4}{3}$ mole KO₂
⇒ $\left(\frac{20}{32}\right)$ mole O₂ is produced by $\left(\frac{4}{3} \times \frac{20}{22} \times 71\right)$ g KO₂ = 59.17 gm.
(b) 1 mole KO₂ = 1 mole KOH
= 1 mole CO₂
⇒ $\left(\frac{100}{71}\right)$ mole KO₂ = $\left(\frac{100}{71} \times 44\right)$ g CO₂ = 61.97 g CO₂
44. NaH₂PO₄ + NH₄⁺ + Mg⁺² → Mg(NH₄) PO₄.6H₂O
 \downarrow Λ
Mg₂P₂O₇

Applying P.O.A.C on 'P' atom $1 \times n_{NaH_2PO_4} = 2 \times n_{Mg_2P_2O_7}$ weight 1.054

$$1 \times \frac{\text{weight}}{120} = 2 \times \frac{1.034}{222}$$

Weight of NaH₂PO₄ = $\frac{2 \times 1.054 \times 120}{222} = 1.14$ gm

$$45. H_2O + SO_3 \longrightarrow H_2SO_4$$

$$n_{H_{2}O} = \left(\frac{40 \times .9}{18}\right) = 2$$

 $n_{SO_3} = \left(\frac{1 \times 50}{0.0821 \times 300}\right) = 2.03 = 2$

 \Rightarrow Weight of H₂SO₄ produced = 2 × 98 = 196 gm.

46.
$$A_{2}[H_{2}PtCl_{6}] \xrightarrow{\Lambda} Pt$$

$$n_{Pt} = \frac{0.262}{195}$$

$$\Rightarrow \text{ moles of salt} = \frac{0.80}{2A + 410}$$

$$\Rightarrow \frac{0.80}{2A + 410} = \frac{0.262}{195}$$

$$\Rightarrow A = \left(\frac{.80 \times 195}{0.262} - 410\right) \times \frac{1}{2}$$

$$\Rightarrow A = 92.70 \text{ gm/mole.}$$

47. Atomic mass of chlorine =
$$\frac{75.77 \times 34.9689 + 24.23 \times 36.9659}{100}$$
$$= \frac{2649.59355 + 895.683757}{100} = 35.4527$$

48.
$$24.31 = \frac{79 \times 24 + (21 - x) \times 25 + x \times 26}{100}$$
$$\Rightarrow 2431 = 1896 + 525 - 25x + 26x$$
$$\Rightarrow x = 10$$

49.

%	Atomic	Relative number of	Simplest	Simplest whole
	mass	atom = $\frac{\%}{$	atomic ratio	number ratio
		At mass		
$C \rightarrow 49\%$	12	$\frac{49}{12} = 4.08$	3	3
$H \rightarrow 2.7\%$	1	$\frac{2.7}{1} = 2.70$	1.98 = 2	2
$Cl \rightarrow 48.3\%$	35.5	$\frac{48.3}{35.5} = 1.36$	1	1

 $\Rightarrow \qquad \text{Emperica formula} = C_3H_2Cl$

$$\Rightarrow \qquad n = \frac{147}{73.5} = 2$$

 \Rightarrow molecular formula = (C₆H₄Cl₂)

50.
$$.25 = \frac{x \times 56}{89600} \times 100$$
$$\Rightarrow \quad x = \frac{.25 \times 89600}{56 \times 100} \qquad \Rightarrow x = 4$$

EXERCISE # S-II

1.
$$\frac{\text{mole of HCl}}{16} = \frac{\text{mole of Cl}_2}{5}$$
$$\frac{\text{mole of Cl}_2}{6} = \frac{\text{mole of Ca(ClO}_3)_2}{1}$$
$$\frac{\text{mole of Ca(ClO}_3)_2}{1} = \frac{\text{mole of NaClO}_3}{2}$$
Also, moles of HCl in 100 ml = 1.164
$$\Rightarrow \text{ moles of NaClO}_3 = \left(\frac{5 \times 1.164 \times 2}{16 \times 6}\right)$$
$$\Rightarrow \text{ mass of NaClO}_3 \text{ produced} = \left(\frac{5 \times 1.164 \times 2}{16 \times 6}\right) \times 106.5 = 12.9 \text{ g}$$
2. (i) $P_4 + O_2 \longrightarrow P_4O_6 + P_4O_{10}$
$$\text{Let } n_{P_4O_6} \longrightarrow x \quad \& \quad n_{P_4O_6} \longrightarrow y$$
$$\frac{P.O.A.C \text{ on } P:}{4 \times n_{P_4} = 4 \times n_{P_4O_6} + 4 \times n_{P_4O_6}} \longrightarrow y$$
$$\frac{P.O.A.C \text{ on } P:}{4 \times n_{P_4} = 4 \times n_{P_4O_6} + 4 \times n_{P_4O_6}} \longrightarrow y$$
$$\frac{P.O.A.C \text{ on } P:}{4 \times n_{P_4} = 4 \times n_{P_4O_6} + 4 \times n_{P_4O_6}} \longrightarrow y$$
$$\frac{P.O.A.C \text{ on } P:}{4 \times n_{P_4} = 4 \times n_{P_4O_6} + 4 \times n_{P_4O_6}} \longrightarrow y$$
$$\frac{P.O.A.C \text{ on } P:}{4 \times n_{P_4} = 4 \times n_{P_4O_6} + 4 \times n_{P_4O_6}} \longrightarrow y$$
$$\frac{P.O.A.C \text{ on } O:}{2 \times n_{O_2} = 6x + 10y} \longrightarrow 3x + 5y = 4 \qquad \dots (1)$$
$$\frac{P.O.A.C \text{ on } O:}{2 \times 16 \times 5} \implies \text{ moles of } P_4O_6 \text{ obtained } = 0.5 \quad \& \text{ moles of } P_4O_{10} \text{ obtained } = 0.5$$
(ii) $P_4 + O_2 \longrightarrow P_4O_6 + P_4O_{10}$
$$\frac{P.O.A.C \text{ on } P:}{3 \times 4 = 4x + 4y} \implies \dots (1)$$
$$\frac{P.O.A.C \text{ on } P:}{3 \times 4 = 4x + 4y} \implies \dots (1)$$
$$\frac{P.O.A.C \text{ on } P:}{3 \times 4 = 4x + 4y} \implies \dots (1)$$
$$\frac{P.O.A.C \text{ on } P:}{3 \times 4 = 4x + 4y} \implies \dots (1)$$
$$\frac{P.O.A.C \text{ on } P:}{3 \times 4 = 4x + 4y} \implies \dots (1)$$

3x + 3y = 93x + 5y = 11 $\frac{-}{-2y = -2}$ y = 1 & $\mathbf{x} = 2$ \Rightarrow moles of $P_4O_6 = 2$ & moles of $P_4O_{10} = 1$ \Rightarrow $P_4 + O_2 \longrightarrow P_4O_6 + P_4O_{10}$ (iii) P.O.A.C on P: $4 \times 3 = 4x + 4y$ \Rightarrow x + y = 3.....(1) P.O.A.C on O: $2 \times 13 = 6x + 10y$ 3x + 5y = 13 \Rightarrow(2) Solving equation (1) & (2)3x + 3y = 93x + 5y = 13 $\frac{-}{-2y = -4}$ y = 2 \Rightarrow \Rightarrow moles of $P_4O_6 = 1$ & moles of $P_4O_{10} = 1$ $C + O_2 \longrightarrow CO_2 + CO$ 3. Let moles of CO be x & moles of CO_2 ben y. <u>P.O.A.C on C :</u> 1 = x + y.....(1) P.O.A.C on O: $2 \times \frac{20}{22} = x + 2y$(2) Solving equation (1) & (2), we get & y = 0.25 $x \rightarrow 0.75$ mass % of CO $\rightarrow \frac{.75 \times 28}{0.75 \times 28 + 0.25 \times 44} \times 100 = \frac{21}{32} \times 100 = 65.625\%$ mass % of CO₂ = 34.375 % 4. Ν : P_2O_5 K_2O : 30 10 10 : mole: : 14 94 142 For : Р Κ Ν : : mole: $\frac{30}{14}$ 10 10 : 71 47 Simplest: 10 : 0.66 : 1 ratio

5. Let CaCl₂ be x gm & NaCl be
$$(10 - x)$$
 gm
CaCl₂ $\xrightarrow{Na_2CO_3}$ CaCO₃ \longrightarrow CaO
moles: $\frac{x}{111}$ $\frac{x}{111}$ $\frac{x}{111}$
As, $\frac{x}{111} = \frac{1.12}{56}$
 \Rightarrow x = 2.22 gm
% NaCl = $\frac{7.78}{10} \times 100 = 77.8$ %

(a)
$$Fe_2O_3 + 2Al \longrightarrow Al_2O_3 + 2Fe$$

(b) Fe_2O_3 & Al reacts in mole ratio $1:2$
 \Rightarrow ratio of mass $= \frac{100}{54}$ or $80:27$

(c)
$$nFe_2O_3 \longrightarrow 100$$

moles of Fe_2O_3 reacted = 50 moles
 \Rightarrow energy released = 50 × 200 = 10,000 units

 $\left(\frac{3x}{2 \times 122.5}\right)$

7.
$$2\text{KClO}_{3} \longrightarrow 2\text{KCl} + 3\text{O}_{2}$$
$$\text{mole:} \quad \left(\frac{x}{122.5}\right) \qquad \frac{x}{122.5} \quad \left(\frac{x}{2\times 122.5}\right)$$
$$n_{\text{O}_{2}} = \frac{112}{22400} = \frac{3x}{2\times 122.5}$$
$$\implies \qquad x = 0.41 \text{ gm}$$

$$\Rightarrow \qquad \text{mass of KCl obtained from 0.41 gm KClO}_3$$

$$= \left(\frac{2 \times 74.5}{2 \times 122.5} \times 0.41\right) = 0.25 \text{ gm KCl}$$

$$4\text{KClO3} \longrightarrow 3\text{KClO}_4 + \text{KCl}$$

$$\begin{pmatrix} 0.59/_{122.5} \end{pmatrix}$$

& mass of KCl obtained from $0.59g \text{ KClO}_3$

$$= \frac{74.5 \times 0.59}{4 \times 122.5} = 0.089 \text{g KCl}$$

& mass of KClO₄ obtained from 0.59g KClO₃

$$= \frac{3 \times 138.5}{4 \times 122.5} \times 0.59 = 0.500 \text{ gm}$$

 \Rightarrow % by weight of KClO₄ in residue

$$= \left(\frac{0.500}{.84}\right) \times 100 = 59.72 \%$$

8.	$\left[\operatorname{Cr}(\operatorname{H}_{2}\operatorname{O})_{5}\operatorname{Cl}\right].\operatorname{Cl}_{2}.\operatorname{H}_{2}\operatorname{O} + \operatorname{AgNO}_{3} \longrightarrow 2\operatorname{AgCl} + \left[\operatorname{cr}(\operatorname{H}_{2}\operatorname{O})_{5}\operatorname{Cl}\right].(\operatorname{NO}_{3})_{2}\right]$							
	moles	of $[Cr(H_2O)_5 Cl] \cdot Cl_2 \cdot H_2O$ is $\frac{5.33}{266.5}$ i.e. 0.02 moles						
	\Rightarrow	moles of AgCl obtained = $2 \times 0.02 = 0.04$ mole mass of AgCl obtained = $0.04 \times 143.5 = 5.74$ gm						
9.	Let the	e metal carbonate be M_2CO_3						
	As, ma	ass % of O = 48 = $\frac{48}{2x+60} \times 100$						
	\Rightarrow	x = 20 i.e. molar mass of metal = 20gm/mole						
	\Rightarrow	moles of M ₂ CO ₃ = $\frac{5 \times 10^{-3}}{(20 \times 2 + 60)} = 5 \times 10^{-5}$ mole.						
	$\begin{array}{c} \Rightarrow \\ \Rightarrow \end{array}$	moles of metal is 10^{-4} mole Number of atoms of metal present = $10^{-4} \times 6 \times 10^{23}$ i.e. 6×10^{19} atoms						
10.	A _x B _y - Apply	$+ O_2 \longrightarrow AO + \text{oxide of B.}$ P. O. A. C on A,						
	$x \times \frac{1}{24}$	$\frac{2.5}{4x+14y} = 1 \times \frac{5}{40}$						
	$\begin{array}{c} \Rightarrow \\ \Rightarrow \end{array}$	x : y = 3 : 2 Emperical formula of compound is $3 : 2$						
11.	moles-	$Ca + Cl_2 \longrightarrow CaCl_2$ $\rightarrow \left(\frac{2.4 \times 10^{24}}{6 \times 10^{23}}\right) \left(\frac{380 \times 96}{760 \times 0.08 \times 300}\right)$						
	\rightarrow	$\simeq 4 \simeq 2$						
	\Rightarrow	mass of $CaCl_2 = 2 \times 111 = 222$ gm						
12.	mole:	$P_4 S_3 + 8O_2 \longrightarrow P_2O_{10} + 3SO_2$ $\left(\frac{1}{64} \times \frac{1}{2}\right) \qquad \qquad \frac{1}{284} - \frac{1}{64}$						
	⇒	mass of P ₄ S ₃ required = $\frac{1}{64 \times 3} \times 220 = 1.1458$ gm						
13.	$\begin{array}{l} \text{moles:} \\ \Rightarrow \\ \Rightarrow \\ \end{array}$	$\begin{array}{l} H_4P_2O_7 + 2NaOH \longrightarrow Na_2H_2P_2O_7 + 2H_2O \\ 3 & 5 \end{array}$ NaOH is L.R. number of molecules Na_2H_2P_2O_7 formed = (2.5) N _A & number of molecules H_2O formed = (5) N _A Total number of molecules formed in product is (7.5)N _A .						

(A) 1 of - atom of c = 12 g1. (B) $\frac{1}{2}$ mole CH₄ = 8 g (C) 10 ml of $H_2O = 10 \text{ g}$ (D) 3.011×10^{23} atoms of oxygen = 8g option (A) is \Rightarrow $n_{CO_2} = \frac{44}{44} = 1$ mole 2. \Rightarrow the molecules of CO₂ = N_A i.e. 6×10^{23} . \Rightarrow correct option is (A) $n_{\rm NH_3} = \frac{4.25}{17}$ i.e. mole 3. \Rightarrow option (B) is correct. Charge on 1 gram ions of Al⁺³ is 3N_Ae Coulomb 4. \Rightarrow option (D) is correct. 5. Atomic weight of A = 40 u& Atomic weight of B = 80 u $\frac{2x}{80} N_A \rightarrow ? \qquad \dots \dots (2)$ Comparing (1) and (2), we get 2xg of B = y \Rightarrow option (C) is correct $n_{AI} \longrightarrow \frac{54}{27} = 2$ mole. 6. mass of same number of magnesium atoms = 48 gm \Rightarrow \Rightarrow Correct option is (C). 7. Weight of molecule of compounds C₆₀H₂₂ $= (60 \times 12 + 22 \times 1) \text{ amu} = (720 + 22) \text{ amo}$ = 742 × 1.66 × 10⁻²⁴ g = 1.24 × 10⁻²¹ gm Option (B) is correct. \Rightarrow $n_{_{NO_{3}^{-}}} = \frac{3.1 \times 10^{^{-3}}}{62 \times 10}$ 8. $= 0.5 \times 10^{-4}$ $= 5 \times 10^{-5}$ Number of electron in 3.1 mg $NO_3^{\Theta} = 5 \times 10^{-5} \times 32 \times 6.022 \times 10^{23} = 9.6 \times 10^{20}$. \Rightarrow Correct option is (C)
9. Ratio of number of molecules of CO₂ & N₂O $=\frac{2x}{44} \times N_A \times \frac{44}{5x \times N_A} = 2:5$ \Rightarrow correct option is (B) (A) $n_{C_2H_6} = \frac{15}{30} = \frac{1}{2}$ mole 10. \Rightarrow number of carbon atoms = $\frac{N_A}{2} \times 2$ i.e. N_A $n_{Na_2C_2O_4} = \frac{40.2}{134} = 0.3$ mole (B) \Rightarrow number of carbon atoms = $0.3 \times 2N_A = 0.6 N_A$ $n_{glucose} = \frac{72}{180} = 0.4$ (C) \Rightarrow number of carbon atoms = $0.4 \times 6N_A = 2.4 N_A$ $n_{C_5H_{10}} = \frac{35}{70} = 0.5$ (D) \Rightarrow number of carbon atoms = $0.5 \times 5N_A = 2.5 N_A$ \Rightarrow Correct option is (D) number of H-atom in 0.9 gm glucose = $\frac{0.9}{180} \times 12N_A = 0.06 N_A$ 11. $nN_2H_4 = \frac{0.048}{32}$ (A) number of H-atoms = $\frac{0.048}{32} \times 4 \text{ N}_{\text{A}} = 0.006 \text{ N}_{\text{A}}$ \Rightarrow $n_{\rm NH_3} = \frac{0.17}{17}$ (B) number of H-atom = $0.01 \times 3N_A = 0.03 N_A$ \Rightarrow $n_{C_2H_6} = \frac{0.30}{30} = 0.01$ (C) number of H-atoms = $0.06 N_A$. \Rightarrow $n_{H_2} \longrightarrow \frac{0.03}{2} = \frac{0.03}{2}$ mole (D) number of H-atoms = $\frac{0.03}{2} \times 2 \text{ N}_{\text{A}} = 0.03 \text{ N}_{\text{A}}$ \Rightarrow Correct option is (C) \Rightarrow

12. $n_{CuSO_4.5H_2O} = \left(\frac{1 \times 10^{22}}{6.022 \times 10^{23}}\right) = 1.66 \times 10^{-2} \text{ mole}$ $\Rightarrow \qquad \text{weight} = 1.66 \times 10^{-2} \times 249.5 = 4.159 \text{ gm}$ $\Rightarrow \qquad \text{Correct option is (C)}$

13.
$$n_{C} \longrightarrow \frac{1.2 \times 10^{-3}}{12}$$
 mole
 \Rightarrow number of carbon atoms $= \frac{1.2 \times 10^{-3}}{12} \times N_{A}$
 $= 6.02 \times 10^{19}$ atoms
 \Rightarrow Correct option is (B).
14. mass of 1.2 mole ethanol $= (1.2 \times 46)$ g
 \Rightarrow Volume $= \frac{mass}{density} = \frac{1.2 \times 46}{0.8} = 69$ ml
 \Rightarrow Correct option is (C).
15. $n_{NO_{2}} = \frac{112}{22.4 \times 10^{3}}$
 \Rightarrow number of molecules $= \frac{112}{22.4 \times 10^{3}} \times 6.02 \times 10^{23} = 3.1 \times 10^{21}$
Now, volume $= \frac{mass}{density} = \frac{5 \times 10^{-3} \times 46}{1.15}$
 $= 200 \times 10^{-3} = 0.20$ ml
 \Rightarrow Correct option is (B).
16. Let the molecule be $A_{x}B_{y}$
Molecular weight of compounds formed $= \left(\frac{XN_{x} + MY}{5}\right)$
 \Rightarrow Correct option is (A).
17. By Avogadro's Hypothesis,
for gas A, $P.v_{A} = n_{A}$, RT
for gas B, $P.v_{B} = n_{B}$. RT
If $n_{A} = n_{B}$
 $\Rightarrow v_{A} = v_{B}$
 \Rightarrow Correction option is (B).
18. Nal \longrightarrow Na' $+ \Gamma$
As, 100 g salt contains 0.5 g Nal
 $= 0.015$ g Nal.
 $n_{Nal} \longrightarrow \frac{0.015}{150}$ mole
 \Rightarrow number of Γ ions $= \frac{0.015}{150} \times 6.02 \times 10^{23}$
 $= 6.02 \times 10^{19}$.
 \Rightarrow Correct option is (C).

19. By Avogadro's hypothesis $n_{O_2} = n_{unknown gas}$ $\frac{1}{32} = \frac{2.375}{M}$ (M \rightarrow molar mass of unknown gas) \Rightarrow \Rightarrow $M = 2.375 \times 32$ M = 76 \Rightarrow \Rightarrow correct option is (C) 20. Ratio of number of atoms = 2:1:2:3 \Rightarrow Correct option is (C). 21. $2P + Q \longrightarrow R$ mole : 8 (Excess) 0 4 0 \Rightarrow Correct option is (C). 22. $4 \text{ Al} + 3\text{O}_2 \longrightarrow 2\text{Al}_2\text{O}_3$ (Excess) 1.5 0 0 1 weight of Al 54 gm. \Rightarrow $C_XH_YO_Z + \left(X + \frac{Y}{4} - \frac{Z}{2}\right)O_2 \longrightarrow X CO_2 + \frac{Y}{2}H_2O$ 23. $n_{CO_2} = \frac{132}{44} = 3$ X = 3 \Rightarrow $n_{H_{2}O} \longrightarrow \frac{54}{18} = 3$ Y = 6. \Rightarrow Correct option is (C). \Rightarrow 24. $C_2H_4O_2 + 2O_2 \longrightarrow 2CO_2 + 2H_2O_2$ 1 mole C₂H₄O₂ & 2 mole O₂ produces 2 mole CO₂ i.e. 124 g mixture produces 88 gm CO₂ \Rightarrow 620g mixture will produces $\frac{88}{124} \times 620 = 440$ gm \Rightarrow Correct option is (C) 25. $5A_2 + 2B_4 \longrightarrow 2AB_2 + 4A_2B.$ 5 mole A_2 produce 2 mole AB_2 \Rightarrow (2 × 250)g AB₂ is produced from 100g A₂ \Rightarrow 1000g AB₂ is produced from $\left(\frac{100}{2 \times 250}\right) \times 1000$ g of A₂ $= 200 \text{ g of } A_2$

Also, 2 mole AB_2 is produced from 2 mole B_4 \Rightarrow (2 × 250)g AB₂ is produced from (2 × 120 × 4) g of B₄ \Rightarrow 1000 g AB₂ is produced from $\left(\frac{2 \times 120 \times 4}{2 \times 250}\right) \times 1000 = 1920$ gm. \Rightarrow Minimum mass of mixture of A₂ & B₂ is (1920 + 200) i.e. 2120 gm \Rightarrow Correct option is (A) 26. $3Mg + 2NH_3 \longrightarrow Mg_3N_2 + 3H_2$ Mole : $\frac{48}{24}$ $\frac{34}{17}$ = 2 = 2 \Rightarrow Mg is L. R. \Rightarrow mass of Mg₃N₂ produced is $\frac{2}{3} \times (100)g = \frac{200}{3}g$ \Rightarrow Correct option is (A) 27. $P_4S_3 + 8O_2 \longrightarrow P_4O_{10} + 3SO_2$ Mole : $\frac{440}{220} = \frac{384}{32}$ = 2= 12 \Rightarrow O₂ is L.R \Rightarrow mass of P₄O₁₀ produced = $\frac{(12 \times 284)}{8}$ g = 426 gm. \Rightarrow Correct option is (B). $12C + 11 H_2 + \frac{11}{2}O_2 \longrightarrow C_{12} H_{22} O_{11}$ 28. Mole: $\frac{84}{12}$ $\frac{12}{2}$ $\frac{56}{22.4}$ 2.5 6 7 \Rightarrow O₂ is L.R. \Rightarrow Mass of sucrose produced = $\left(\frac{2}{11} \times 2.5\right) \times 342 \text{ g} = 155.5 \text{ g}$ \Rightarrow Correct option is (B). $H_2SO_4 + Ca(OH)_2 \longrightarrow CaSO_4 + 2H_2O.$ 29. Mole .5 0.2 $Ca(OH)_2$ is L.R. \Rightarrow number of moles of CaSO₄ formed = 0.2 \Rightarrow Correct option is (A)

30. $I_2 + 2Cl_2 \longrightarrow ICl + ICl_3$ 14.2 25.4 Mole: 254 71 0.1 0.2 = No. L. R \Rightarrow moles of ICl produced = 0.1 & moles of ICl_3 produced = 0.1 \Rightarrow Correct option is (A) $2P_4 + 8O_2 \longrightarrow P_4O_6 + P_4O_{10}$ 31. Mole : $\frac{31}{124} = \frac{32}{32}$ $\frac{1}{4}$ 1 = No. L. R Weights of P₄O₆ produced = $\frac{1}{2} \times \frac{1}{4} \times 220 = 27.5$ g \Rightarrow & weight of P₄O₁₀ produced = $\frac{1}{8} \times 284 = 35.5$ g Correct option is (B) \Rightarrow 32. $CaCO_3 \longrightarrow CaO + CO_2$ $Na_2CO_3 + CO_2 + H_2O \longrightarrow 2 NaHCO_3$ $n_{Na_2CO_3} = \frac{21.2 \times 10^3}{106} = 2 \times 10^2$ moles. Moles of $CaCO_3 = mole of CO_2$(1) Moles of CO_2 = mole of Na_2CO_3(2) From (1) & (2)Mole of $CaCO_3 = 2 \times 10^2$ Mass o f CaCO₃ = $2 \times 10^2 \times 100 = 20$ kg \Rightarrow Correct option is (B). \Rightarrow $NaI + AgNO_3 \longrightarrow AgI + NaNO_3$. 33. $2AgI + Fe \longrightarrow FeI_2 + 2Ag$ $2\text{FeI}_2 + 3\text{Cl}_2 \longrightarrow 2\text{FeCl}_3 + 2\text{I}_2.$ $n_{I_2} = (1 \times 10^3)$ $\frac{\text{mole of AgNO}_3}{1} = \frac{\text{mole of AgI}}{1}$(1) $\frac{\text{mole of AgI}}{2} = \frac{\text{mole of FeI}_2}{1}$(2) $\frac{\text{mole of FeI}_2}{2} = \frac{\text{mole of I}_2}{2}$(3)

From (1), (2) & (3) Mole of I_2 = mole of $FeI_2 = \frac{mole of Ag I}{2}$ *.*.. $=\frac{\text{mole of AgNO}_3}{2}$ $10^3 = \frac{\text{wt of AgNO}_3}{170 \times 2}$ \Rightarrow wt. of $AgNO_3 = 340 \text{ kg}$ \Rightarrow Correct option is (A) \Rightarrow 34. Let $CaCl_2$ be x g & NaCl be (10 - x) g $CaCl_2 \longrightarrow CaCO_3 \longrightarrow CaO$ Moles $\frac{x}{111}$ $\frac{x}{111}$ $\frac{x}{111}$ moles of CaO $\longrightarrow \frac{1.62}{56}$ \Rightarrow moles of CaCl₂ = $\frac{1.62}{56}$ $\Rightarrow \frac{1.62}{56} = \frac{x}{111}$ \Rightarrow x = $\frac{111 \times 1.62}{56}$ $\Rightarrow x = 3.21$ \Rightarrow % by mass of CaCl₂ = $\frac{3.21}{10} \times 100 = 32.1$ % \Rightarrow Correct option is (A) 35. for Alkali metal carbonate $M_2CO_3 + HCl$ \longrightarrow MCl + CO₂ + H₂O (Excess) For Alkaline Earth metal carbonate $MCO_3 + HCl \longrightarrow MCl_2 + CO_2 + H_2O$ $n_{CO_2} = \frac{1 \times 12.315}{0.0821 \times 300} = 0.5$ mole $Li_2CO_3 + 2HCl \longrightarrow 2LiCl + CO_2 + H_2O$ 1 mole CO_2 is produced from 1 mole Li_2CO_3 0.5 mole CO₂ is produced from .5 mole Li₂CO₃ OR $= .5 \times (74)$ \Rightarrow $= 37 \text{ gm. of } \text{Li}_2\text{CO}_3 \&$ mass of impurity = 3 gmCorrect option is (B) \Rightarrow

36.
$$M = \frac{99 \times 20 + \frac{1}{2} \times (21 + 22)}{100} = 20.002$$

$$\Rightarrow \text{ Correct option is (A)}$$
37.
$$PCl_5 \longrightarrow PCl_3 + Cl_2$$

$$M_{Avg.} = \frac{M_{Theo.}}{1 + (n - 1)\alpha} = \frac{208.5}{1 + 0.5} = 139$$

$$\Rightarrow \qquad \% \text{ change in } M_{Avg.} \text{ of the mixture} = \left(\frac{208.5 - 139}{208.5}\right) \times 100 = 33.33 \%$$

$$\Rightarrow \qquad \text{Correct option is (C)}$$
38.
$$8 = \frac{1 \times 32}{M} \times 100$$

$$\Rightarrow \qquad M = 400$$

Correct option is (B). \Rightarrow

molecular formula mass 39. n = Emperical formula mass 120 4 Ï

$$\Rightarrow$$
 n = $\frac{120}{30}$ = 4

 \Rightarrow molecular formula = (CH₂O) × 4 i.e. (C₄H₈O₄)

 \Rightarrow Correct option is (B)

40.

Element	%	Relative number	Simplest atomic	Simplest whole
		of atoms	Ratio	Number ratio
Ca	20	$\frac{20}{40} = \frac{1}{2}$	1	1
Br	80	$\frac{80}{80} = 1$	2	2

Empirical formula is CaBr₂ \Rightarrow

As,
$$n = \frac{200}{200} = 1$$

Molecular formula is CaBr₂ \Rightarrow

correct option is (B). \Rightarrow

41.
$$69.98 = \frac{21 \times 12}{M} \times 100$$
$$\Rightarrow M = \frac{21 \times 12}{69.98} \times 100 \qquad \Rightarrow M = 360.1$$
$$\Rightarrow \text{Correct option is (D)}$$

42. Method-1

Let the compound be $C_x H_y O_z$ Now, weight % of $C = 8 \times$ (weight % of H) $\Rightarrow \frac{x}{y} = \frac{2}{3}$ Also, weight % of C = $\frac{1}{2}$ × (weight % of O) $\Rightarrow \frac{x}{z} = \frac{2}{3}$ The correct option is (B) \Rightarrow Method-2 $Ag_2A \longrightarrow Ag$ P.O. A.C. on Ag $2 \times \frac{1}{216 + M_{\rm A}} = \frac{0.5934}{108} \times 1$ \Rightarrow M_A \rightarrow 148 \Rightarrow molar mass of Acid = 150 С Η 0 8 weight % 1 16 2 mole % 3 3 \Rightarrow Empirical formula = C₂H₃O₃ \Rightarrow molecular formula is C₄H₆O₆ \Rightarrow n = 2 \Rightarrow Correct option is (B)

43. Let % of NO₂ be x NO % of number be (100 - x) $\Rightarrow 34 = \frac{x \times 46 + (100 - x) \times 30}{100} \Rightarrow x = 25 \%$ \Rightarrow Correct option is (A) 1. For $(NH_4)_3PO_4$:

2.

3.

4.

5.

Ratio of number of O atoms to number of H atoms = $\frac{4}{12}$ i.e. (1:3) Ratio of number of cations to number of anions = 3:1Number of gm-atoms of nitrogen to atoms of oxygen = $\frac{3}{4}$ Total number of atoms in 1 mole of $(NH_4)_3 PO_4 = 20N_A$ Correct options are (A), (B) \Rightarrow $2 \text{ Mg} + \text{O}_2 \longrightarrow 2 \text{ MgO}$ moles: $\frac{12}{24} = \frac{32}{32}$ Mg is L.R, So is 100% consumed \Rightarrow i.e. number MgO is left unburnt Amount of O_2 left unreacted = 0.75 gm molecule Amount of MgO formed = 0.5×40 i.e. 20 gm The mixture at the end will weight 44 gm. \Rightarrow Correct option (A) $3 \operatorname{CaCO}_3 + 2H_3\operatorname{PO}_4 \longrightarrow \operatorname{Ca}_3(\operatorname{PO}_4)_2 + 3H_2O + 3\operatorname{CO}_2$ moles : $\frac{50}{100} = 0.5$ $\frac{68.6}{98} = 0.7$ \Rightarrow CaCO₃ is L.R. Amount of salt formed = $\left(\frac{1}{3} \times 0.5 \times 310\right)$ = 35.93 gm $n_{CO_2} = 0.5$ Correct option are (A), (B), (C) \Rightarrow $C_7H_8 + 3HNO_3 \longrightarrow C_7H_5N_3O_6 + 3H_2O$ As, C_7H_8 & HNO_3 reacts in 1 : 3 ratio $x \times 92 + 3x \times 62 = 140.5$ \Rightarrow \Rightarrow x = 0.5 \Rightarrow Maximum weight of $C_7H_5N_3O_6$ which can be produced is 0.5×227 Correct option is (B). $A + N \xrightarrow{I} B + L$ $A + N \xrightarrow{\Pi} \frac{1}{2}B + \frac{1}{2}(C) + L$ 4 2 2 \Rightarrow B will always be greater than C & If 2 moles of C are formed the total 6 mole of B are also formed Correct option are (A), (D) \Rightarrow

i.e 113.5 gm

6. $4Ag + 8KCN + O_2 + 2H_2O \longrightarrow 4K [Ag (CN)_2] + 4 KOH$

 $\Rightarrow 4 \times 108g \text{ of Ag reacts with } 8 \times 65g \text{ of KCN } 100g \text{ of Ag reacts with } \frac{8 \times 65}{4 \times 108} \times 100$

Hence statement A is correct

- $\Rightarrow \qquad 4\times 108 \text{g of Ag require 32 gm of } O_2$
- $\Rightarrow 100g \text{ of Ag require} = \frac{32}{4 \times 108} \times 100 = 7.40 \text{ gm}$

Hence option (C) is correct.

Volume of O₂ required =
$$\frac{7.4}{32} \times 22.4 = 5.20$$
 liters

- 7. CaO $+3C \longrightarrow CaC_2 + CO$
 - (A) Find product contains 85% CaC₂ & 15% CaO. Let mass of product = 100 gm
 - $\therefore \qquad \text{mass of } CaC_2 = 85 \text{ gm} \\ \text{mass of } CaO = 15 \text{ gm} \end{cases}$

used mole of CaO = mole of CaC₂ produced = $\left(\frac{85}{64}\right)$

$$\therefore$$
 mass of CaO producing 85 gm Cal₂ = $\frac{85}{64} \times 56$ =74.375 gm

:. Initial mass of CaO = (74.375 + 15) = 89.37585 gm CaC₂ obtained from 89.38 gm CaO 1000 kg CaC₂ obtained from $\frac{89.38}{85} \times 10^3$ kg CaO. = 1051.47 kg CaO.

(B) 100 gm produced requires CaO = 89.38g

$$\Rightarrow$$
 10³ kg product requires CaO = $\frac{89.38}{100} \times 10^3 = 893.8$ kg CaO.

8.
$$2NH_3 + \frac{5}{2}O_2 \longrightarrow 2NO + 3H_2O.$$

 $2NO + O_2 \longrightarrow 2NO_2$
 $2NO_2 + H_2O \longrightarrow HNO_3 + HNO_2$
 $3HNO_2 \longrightarrow HNO_3 + NO + H_2O$

- (A) Moles of HNO₃ produced is help of moles of Ammonia used if HNO₂ is not used to produce HNO₃ by reaction (IV).
- (B) Incorrect
- (C) $\frac{1}{4}$ th of total HNO₃ is produced by reaction (IV) if HNO₂ is used to produce HNO₃.
- (D) Moles of number produced in reaction (IV) is 50% of moles of total HNO₃ produced.

9. mass of substance = 0.42 gm. Volume of N₂ = $\frac{100}{11}$ ml Temperature = 250 KPressure = 860 - 24 = 836 mm Hg**Step (1)** Volume of N₂ at S.T.P. i.e. $V_2 = \frac{P_1 V_1}{T_1} \times \frac{T_2}{P_2}$ $V_2 = \frac{836 \times 100 \times 273}{760 \times 11 \times 250} \implies V_2 = 10.92 \text{ ml}$ \Rightarrow **Step (2)** % of N₂ in organic compound $= \frac{28 \times 10.92}{22700 \times 0.42} \times 100 = \frac{10}{3}$ % Correct option is (A) \Rightarrow 10. $2A + 3B \longrightarrow C$ moles: 3 4 B is L.R. & C formed is $\frac{4}{3}$ moles only if the yeild is 100%. \Rightarrow \Rightarrow Correct option (C). 11. Y₃Al₅O₁₂ weight % of Y $\longrightarrow \frac{89 \times 3}{594} \times 100 = 44.95$ % weight % of Al $\longrightarrow \frac{27 \times 5}{594} \times 100 = 22.73 \%$ weight % of O $\longrightarrow \frac{12 \times 6}{594} \times 100 = 32.32 \%$ $n_{C_6H_8O_6} = \frac{17.6 \times 10^{-3}}{176} = 10^{-4}$ moles 12. O - atoms present = $6 \times 6 \times 10^{23} \times 10^{-4} = 3.6 \times 10^{20}$ moles of vitamin C in 1 gm of vitamin C = 5.68×10^{-3} . moles of vitamin C that should be consumed daily = 10^{-4} 13. $2H_2 + O_2 \longrightarrow 2H_2O$ moles: $\frac{1}{2}$ $\frac{1}{32}$ \Rightarrow O₂ is L.R. \Rightarrow mass of H₂O produced = $\frac{1}{32} \times 2 \times 18 = 1.125$ g $3H_2 + N_2 \longrightarrow 2NH_3$

moles: $\frac{1}{2}$ $\frac{1}{28}$ \Rightarrow H₂ is L.R. mass of NH₃ produced = $\frac{1}{28} \times 2 \times 17 = 1.214$ g \Rightarrow $H2 + Cl_2 \longrightarrow 2HCl$ moles: $\frac{1}{2}$ $\frac{1}{71}$ \Rightarrow Cl₂ is L.R mass of HCl produced = $2 \times 36.5 \times \frac{1}{71} = 10028$ g \Rightarrow $2H_2 + C \longrightarrow CH_4$ moles: $\frac{1}{2}$ $\frac{1}{12}$ \Rightarrow C is L. R. Mass of CH₄ produced = $\frac{1}{12} \times 16 = 1.333$ g Correct option is (A) \Rightarrow 14. $Fe + Br_2 \longrightarrow FeBr_2$ $Br_2 + 3FeBr_2 \longrightarrow Fe_3Br_8$ $Fe_3Br_8 + 4Na_2CO_3 \longrightarrow 8NaBr + 4CO_2 + Fe_3O_4.$ $n_{\text{NaBr}} = \frac{2.06 \times 10^3 \times 10^3}{103} = 2 \times 10^4$ (a) $\frac{\text{moles of Fe}}{1} = \frac{\text{moles of FeBr}_2}{1}$ $\frac{\text{moles of FeBr}_2}{2} = \frac{\text{moles of Fe}_3\text{Br}_8}{1}$ $\frac{\text{moles of Fe}_{3}\text{Br}_{8}}{1} = \frac{\text{mole of NaBr}}{2}$ & moles of Fe = $\frac{\text{moles of NaBr}}{8} \times 3 = \frac{2 \times 10^4}{8} \times 3$ \Rightarrow \Rightarrow mass of Fe required = $\frac{6 \times 10^4 \times 56}{8} = 420 \text{ kg} \Rightarrow$ Correct option is (B) $mole: \frac{3FeBr_2}{8} + Br_2 \longrightarrow Fe_3Br_8$ $\frac{3}{8} \times 2 \times 10^4 \times \frac{100}{70} \times \frac{100}{60} \qquad \frac{1}{8} \times 2 \times 10^4 \times \frac{100}{70}$ (b) $Fe_{3}Br_{8} + 4Na_{3}CO_{3} \longrightarrow 8 NaBr + 4CO_{2} + Fe_{3}O_{4}$ mole : $\frac{1}{8} \times 2 \times 10^4 \times \frac{100}{70}$ 2×10^{4} $\Rightarrow \qquad \frac{10^6}{8 \times 7} = 0.01786 \times 10^6$ $= 1.786 \times 10^4$ moles $\Rightarrow \qquad \text{mass of Fe required} = \frac{1.786 \times 10^4 \times 56}{10^2} = 17.86 \times 56$

$$= 1000 \text{ kg} \text{ or } 10^3 \text{ kg}$$

$$\Rightarrow \text{ Correct option is (C)}$$
(c) moles of CO₂ formed = $\frac{1}{2} \times 2 \times 10 = 10 \text{ moles}$

$$\Rightarrow \text{ Correct option is (B).}$$
15. (a) Ca₂B₆O₁₁ + 2Na₂CO₃ \longrightarrow 2CaCO₃ + Na₂B₆O7 + 2NaBO₂

$$\left(\frac{3}{200} \times \frac{10^2}{32.2} \times \frac{100}{60}\right) \times 100 \times 100 \qquad \left(\frac{3}{200} \times \frac{10^2}{32.2} \times 100\right) \times 100$$
Na₂B₆O₇ \longrightarrow 2NaBO₂ + B₂O₃

$$\left(\frac{10^2}{32.2} \times \frac{10^3}{32.2} \times 100\right) \qquad \left(\frac{10^2}{32.2}\right) \times 100$$
COO + B₂O₃ \longrightarrow CO(BO₂)₂

$$\left(\frac{10^2}{32.2}\right) \times 100 \qquad 10^2$$
n_{cooreo}_{2)₂} $\left(\frac{10^2}{1450}\right) \times 100 \times 10^2$
n_{cooreo}_{2)₂} $\left(\frac{14.5 \times 10^3}{1450} = 10^2 \text{ moles}\right)$
 $\Rightarrow \text{ mass of Ca2B6O11 required = $10^2 \times (80 + 66 + 176) = 322 \times 10^2 \text{g} = 32.2 \text{ kg}$
 $\Rightarrow \text{ correct option is (A)}$
(b) mass of Ca₂B₆O₁₁ obtained = $\frac{3 \times 10^6}{200 \times 32.2 \times 60} \times 3220 = \frac{10^6}{400} \text{g}$
 $= \frac{10^6}{4 \times 10^3} \text{kg} = \frac{1000}{4} \text{kg} = 250 \text{ kg}$
 $\Rightarrow \text{ correct option is (A)}$
16. UF₆ + H₂O \longrightarrow U_xO_y F_z + $\left(\frac{H_{\frac{5}{3}} \text{ F}_{\frac{5}{3}}}{10}\right) \text{ or } (\text{HF})$
(a) The empirical formula of gas is HF
 \Rightarrow Correct option is (C)
(b) Mass of H₂O = 0.2
 \Rightarrow UF₆ + 2H₂O \longrightarrow U₁O₂F₂ + 4HF
 \Rightarrow Empirical formula of solid is UF₂O₂
 \Rightarrow Correct option is (A)
(c) 1 mole UF₆ gives 4 more HF
 \Rightarrow % of fluorine converted in gaseous product
 $= 100 - \left(\frac{114 - 76}{114}\right) \times 100 = 66.66 \%$$

EXERCISE # JEE-MAINS

- 1. Moles pg CO = $\frac{2.01 \times 10^{23}}{6.02 \times 10^{23}} = 0.33$ Mass of CO = 0.33×28 = 9.3 gm 2. C H N
 - 9 WT 1 3.5 $\frac{9}{12}$ $\frac{1}{1}$ 3.5 Moles 14 $\frac{3}{4}$ 1 1 4 3 4 1 $EF = C_3H_4N_1 \implies$ Mass (54) MF = n (EF)108 = n (54)n = 2 $MF = C_6H_8N_2$
- **3.** Remain unchanged

The mass of 1 mole of the substance will remain unchanged.

 \Rightarrow Correct option is (2)

4. 8 mole oxygen atom is present in 1 mole
$$Mg_3(PO_4)_2$$

$$\Rightarrow$$
 0.25 mole oxygen atom is present in $\left(\frac{1}{8} \times .25\right) = 3.125 \times 10^{-2}$ mole.

 \Rightarrow Correct option is (1)

5.
$$V.D = 94.8 \implies molar mass = 2 \times 94.8 = 189.6 \text{ gm}$$

$$\Rightarrow$$
 mass of chlorine = 74.75% of 189.6 = 141.726 gm

4

$$\Rightarrow$$
 mole of Cl = $\frac{141.726}{35.5}$ =

- \Rightarrow Formed of metal chloride will be MCl₄.
- \Rightarrow Correct option is (2)

6.
$$n_{O_3} \longrightarrow \frac{16}{48} = \frac{1}{3}$$
 mole
 $n_{CO} \longrightarrow \frac{28}{28} \sim 1$ mole
 $n_{O_2} \longrightarrow \frac{16}{32} = \frac{1}{2}$ mole
 \Rightarrow Ratio of oxygen atoms = 1 : 1 : 1
 \Rightarrow Correct option is (D)

7.
$$C_{x}H_{y} + \left(x + \frac{y}{4}\right)O_{2} \longrightarrow xCO_{2} + \frac{y}{2}H_{2}O$$

$$n_{CO_{2}} \rightarrow \frac{3.08}{44} = 0.07$$

$$n_{H_{2}O} \rightarrow \frac{.72}{18} = 0.04$$

$$\therefore \quad C : H = 0.07 : 0.04 = 7 : 8$$

$$\Rightarrow \quad \text{The empirical formula of compounds is (C_{7}H_{8})}$$

$$\Rightarrow \quad \text{Correct option is (4)}$$

8.

 $\begin{array}{cccc} 0 & \mathrm{N} \\ \text{ratio of masses} \longrightarrow 1 & : & 4 \\ \text{ratio of mole} & \longrightarrow \frac{1}{16} & : & \frac{4}{14} \\ \text{ratio of molecules} \longrightarrow \frac{\mathrm{N}_{\mathrm{A}}}{16} & : & \frac{4\mathrm{N}_{\mathrm{A}}}{14} \\ \Rightarrow & \text{Ratio of number of molecules} = 7 : 32 \\ \Rightarrow & \text{Correct option is (D)} \end{array}$

9.
$$2C_8H_7SO_3Na + Ca^{+2} \longrightarrow Ca (C_8H_7SO_3)_2 + 2Na^+$$

mole : $\frac{1}{206}$

$$\Rightarrow$$
 maximum uptake of Ca⁺² ions = $\frac{1}{412}$

 \Rightarrow Correct option is (4).

10. By carius method,

% Br =
$$\frac{80 \times \text{weight of AgBr}}{188 \times \text{Weight of organic Halide}} \times 100$$

= $\frac{80}{188} = \frac{141 \times 10^{-3}}{250 \times 10^{-3}} \times 100 = 24$
 \Rightarrow Correct option is (1)

11.
$$100 \text{ kg} \longrightarrow (10 \text{ kg}^1 \text{ H})$$

 \downarrow
 $(20 \text{ kg}^2 \text{ H})$
 $\Delta W = 10 \text{ kg}$
 \Rightarrow weight gain is 10% of 75 kg i.e. 7.5 kg
 \Rightarrow Correct option is (3)

 $M_2CO_3 + 2HCl \longrightarrow 2MCl + CO_2 + H_2$ 12. $n_{CO_2} \longrightarrow 0.01186$ 1 mole CO₂ is produced by 1 mole M₂CO₃ 0.01186 mole CO₂ is produced by $\left(\frac{1 \times 0.01186}{1}\right)$ mole M₂CO₃ \Rightarrow $\frac{1}{M_{(M,C0_{1})}} = 0.01186$ \Rightarrow $M_{M_2CO_3} = \frac{1}{0.01186} = 84.3 \text{ gm}$ \Rightarrow Correct option is (2) \Rightarrow С 13. Element Η Mass ratio 6 1 2 Mole ratio 1 So, empirical formula : CH₂ For buring CH₂ unit ; oxygen required is $\frac{3}{2}$ mole Empirical formula is $(CH_2O_{3/2})$ i.e. $C_2H_4O_3$ \Rightarrow Correct option is (1) \Rightarrow Moles of $C_{57}H_{110}O_6 = \frac{445}{890} = 0.5$ moles 14. From 2 moles of $C_{57}H_{110}O_6 \longrightarrow 110$ moles of H_2O is produced 0.5 moles of $C_{57}H_{110}O_6 \longrightarrow 110 \times \frac{0.5}{2}$ moles H_2O \Rightarrow 27.5 moles Mass of $H_2O = 27.5 \times 18$ gram = 495 gram 15. $2 \text{ NaHCO}_3 + \text{H}_2\text{C}_2\text{O}_4 \longrightarrow \text{Na}_2\text{C}_2\text{O}_4 + 2\text{CO}_2 + 2\text{H}_2\text{O}$ $=\frac{0.25\times10^{-3}}{25}$ Moles of CO₂ $= 10^{-5}$ moles Moles of NaHCO₃ = $\frac{2}{2} \times 10^{-5}$ = 10^{-5}

wt. of NaHCO₃ = $10^{-5} \times 84$

% Mass
$$= \frac{84 \times 10^{-5}}{10 \times 10^{-3}} \times 100 = 8.4\%$$

16.		$C_x H_y N_z - \frac{I}{M}$	$\xrightarrow{\text{Duma}} 6\text{CO}_2 + 4\text{H}_2\text{O} + \text{N}_2$
	Clearl	y, x = 6 y = 8 z = 2	
	Hence	$C_6H_8N_2$	
17.	Metha % mol	ne CH_4 1 mole \sim le of C = $\frac{1}{1+4}$	1 mole C 4 mole H $-\frac{4}{4} \times 100$
18.	Optior	$N_2 + 3H_2$ n B 56 gm 10 gm Clearl	
19.	(1)	4 Fe + 4×56 gm 1 gm	$3O_2 \longrightarrow 2 \operatorname{Fe}_2O_3$ $3 \times 32 \operatorname{gm}$ $\frac{3 \times 32}{4 \times 56} = \frac{3}{7} \operatorname{gram}$
	(2)	C ₃ H ₈ + 44 g 1 gram	$SO_2 \longrightarrow 3CO_2 + 4H_2O$ 80 gm $\frac{80}{44} = \frac{20}{11} \text{ gram}$
	(3)	2Mg + 2×24 1 gram	$O_2 \longrightarrow 2MgO_2$ 32 $\frac{32}{2 \times 24} = \frac{2}{3} \text{ gram}$
	(4)	P ₄ + 4×31 1 gram Ans. (1)	$SO_2 \longrightarrow P_4O_{10}$ 5×16 $\frac{80}{124} = \frac{20}{31} \text{ gram}$

 $\label{eq:mass} \begin{array}{ll} \mbox{20.} & \mbox{Mass of 1 mol of } AB_2 = M_A + 2M_B = 25 \times 10^{-3} \mbox{ kg} \\ & \mbox{Mass of 1 mol of } A_2B_2 = 2M_A + 2M_B = 30 \times 10^{-3} \mbox{ kg} \\ & \mbox{M}_A = 5 \times 10^{-3} \mbox{ kg} \mbox{ / mol} \\ & \mbox{M}_B = 10 \times 10^{-3} \mbox{ kg} \mbox{ / mol} \end{array}$

22.
$$C_xH_y + (x + \frac{y}{4})O_2 \longrightarrow x CO_2 + \frac{y}{2}H_2O$$

10 ml 55 ml

40 ml

For gases, volume is proportional to moles

$$\frac{10}{55} = \frac{1}{x + \frac{4}{4}} \qquad \dots \dots (1)$$
$$\frac{1}{x} = \frac{10}{40} \qquad \dots \dots (2)$$
$$x = 4$$
$$y = 6$$
$$C_4 H_6$$

EXERCISE # JEE-ADVANCED

1. Mass of 1 $e^- = 9.1 \times 10^{-31} \text{ kg}$

$$\Rightarrow \qquad \text{moles of } e^- \text{ weighing 1 } \text{kg} = \frac{1}{9.108 \times 10^{-31}} \times \frac{1}{\text{N}_{\text{A}}} = \frac{1}{9.108 \times 6.023} \times 10^8$$

 \Rightarrow Correct option is (D)

2.
$$6CaO + P_4O_{10} \longrightarrow 2Ca_3(PO_4)_2$$

 $nP_4O_{10} \longrightarrow \frac{852}{284} = 3 \text{ mole}$

1 mole P_4O_{10} reacts with 6 mole CaO

 $\Rightarrow 3 \text{ mole } P_4O_{10} \text{ reacts with } 18 \text{ mole } CaO \text{ or } 18 \times 56 \text{ CaO}$ i.e. 1008g CaO.

3. Atomic mass of Fe =
$$\frac{54 \times 5 + 56 \times 90 + 57 \times 5}{100} = 55.95$$

 \Rightarrow Correct option is (B)