



Telangana State Board of  
**INTERMEDIATE** Education  
**FIRST YEAR**



# **MATHEMATICS**

## **I B**

**BASIC LEARNING MATERIAL**  
For The Academic Year : 2021-2022





**TELANGANA STATE BOARD OF  
INTERMEDIATE EDUCATION**

**MATHEMATICS - IB**

**FIRST YEAR  
(English Medium)**

**BASIC LEARNING MATERIAL**

**ACADEMIC YEAR  
2021-2022**

## **Coordinating Committee**

**Sri Syed Omer Jaleel, IAS**  
Commissioner, Intermediate Education &  
Secretary, Telangana State Board of Intermediate Education  
Hyderabad

**Dr. Md. Abdul Khaliq**  
Controller of Examinations  
Telangana State Board of Intermediate Education

### **Educational Research and Training Wing**

**Ramana Rao Vudithyala**  
Reader

**Mahendar Kumar Taduri**  
Assistant Professor

**Vasundhara Devi Kanjarla**  
Assistant Professor

### **Learning Material Contributors**

**S.S.R.S RAGHUVeer, M.Sc., PGDCA**  
Junior Lecturer in Mathematics,  
Chitanya Kalasala,  
B.N. Reddy Nagar,  
**HYDERBAD**

**B. RAJASRI, M.Sc.**  
Junior Lecturer in Mathematics,  
Maharshi Veda Vignan Mahavidyalaya  
Junior College,  
Begumpet, **HYDERBAD**

**B. MRUTYUNJAYA, M.Sc.**  
Junior Lecturer in Mathematics,  
Government Junior College,  
PATANCHERU  
**SANGAREDDY DIST.**

**M. VIJITH KUMAR, M.Sc. B.Ed.**  
Junior Lecturer in Mathematics,  
Government Junior College,  
**LUXETTIPET**

**J. RAVI KIRAN, M.Sc., B.Ed.**  
Junior Lecturer in Mathematics,  
Government Junior College Boys,  
**NIRMAL**

**T. SREEDHAR, M.Sc., PGDCA**  
Junior Lecturer in Mathematics,  
Government Junior College (A&C),  
**KARIMNAGAR**

## **PREFACE**

The ongoing Global Pandemic Covid-19 that has engulfed the entire world has changed every sphere of our life. Education, of course is not an exception. In the absence of Physical Classroom Teaching, Department of Intermediate Education Telangana has successfully engaged the students and imparted education through TV lessons. In the back drop of the unprecedented situation due to the pandemic TSBIE has reduced the burden of curriculum load by considering only 70% syllabus for class room instruction as well as for the forthcoming Intermediate Examinations. It has also increased the choice of questions in the examination pattern for the convenience of the students.

To cope up with exam fear and stress and to prepare the students for annual exams in such a short span of time , TSBIE has prepared “Basic Learning Material” that serves as a primer for the students to face the examinations confidently. It must be noted here that, the Learning Material is not comprehensive and can never substitute the Textbook. At most it gives guidance as to how the students should include the essential steps in their answers and build upon them. I wish you to utilize the Basic Learning Material after you have thoroughly gone through the Text Book so that it may enable you to reinforce the concepts that you have learnt from the Textbook and Teachers. I appreciate ERTW Team, Subject Experts, who have involved day in and out to come out with the, Basic Learning Material in such a short span of time.

I would appreciate the feedback from all the stake holders for enriching the learning material and making it cent percent error free in all aspects.

The material can also be accessed through our website [www.tsbie.cgg.gov.in](http://www.tsbie.cgg.gov.in).

**Commissioner & Secretary**  
Intermediate Education, Telangana.

# CONTENTS

<b>Unit 1</b>	<b>Locus</b>	<b>2</b>
<b>Unit 2</b>	<b>Transformation of Axes</b>	<b>8</b>
<b>Unit 3</b>	<b>The Straight Line</b>	<b>14</b>
<b>Unit 4</b>	<b>Pair of Straight Lines</b>	<b>46</b>
<b>Unit 5</b>	<b>Three Dimensional Coordinates</b>	<b>58</b>
<b>Unit 6</b>	<b>Direction Cosines and Direction Ratios</b>	<b>61</b>
<b>Unit 7</b>	<b>The Plane</b>	<b>67</b>
<b>Unit 8</b>	<b>Limits and Continuity</b>	<b>69</b>
<b>Unit 9</b>	<b>Differentiation</b>	<b>76</b>
<b>Unit 10</b>	<b>Applications of Derivatives</b>	<b>87</b>

## Locus

**Def :** The path of a point moving subject to the given condition is called locus.

The equation representing the locus is called equation of locus.

### Steps to find locus

**Step 1 :** Let the point P be  $P(h, k)$

**Step 2 :** Use the formula related to given condition

**Step 3 :** Simplify

**Step 4 :** The simplified algebraic equation represents locus

### Short Answer Questions (4 Marks)

1. Find the equation of locus of a point which is at a distance 5 from  $A(4, -3)$ .

**Sol:** Let  $P(h, k)$  be any point

Given point  $A(4, -3)$

$|PA| = 5$  given condition

$$\Rightarrow |PA|^2 = 25$$

$$(h - 4)^2 + (k + 3)^2 = 25 \quad (\text{distance between two points} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2})$$

$$\text{equation of locus } (x - 4)^2 + (y + 3)^2 = 25$$

$$x^2 + y^2 - 8x + 6y + 25 = 25 \Rightarrow x^2 + y^2 - 8x + 6y = 0$$

2. **Find the equation of locus of a point which is equidistant from the points  $A(-3, 2)$ , and  $B(0, 4)$**

**Sol:** Let  $P(h, k)$  be any point

$|PA| = |PB|$  given condition

$$\Rightarrow |PA|^2 = |PB|^2$$

$$(h + 3)^2 + (k - 2)^2 \Rightarrow h^2 + (k - 4)^2$$

$$h^2 + 6h + 9 + k^2 - 4k + 4 = h^2 + k^2 - 8k + 16$$

$$6h - 4k + 13 = -8x + 16$$

$$6h + 4k = 3$$

$$\therefore \text{equation of locus } 6x + 4y - 3 = 0$$

3. Find the equation of locus of a point P such that the distance of P from the origin is twice the distance of P and A(1,2).

Sol: Let P(h, k) be any point

O(0, 0) be the origin

A(1, 2) given point

|PO| = 2|PA| given condition

$$\Rightarrow |PO|^2 = 4 |PA|^2$$

$$h^2 + k^2 = 4[(h-1)^2 + (k-2)^2]$$

$$\text{simplifying } 3h^2 + 3k^2 - 8h - 16k + 20 = 0$$

$$\text{equation of locus } = 3x^2 + 3y^2 - 8x - 16y + 20 = 0$$

4. Find the equation of locus of a point which is equidistant from the coordinate axes.

Sol: Let P(h, k) be any point


distance from X-axis = |PM|

distance from Y-axis = |PN|

$$|PM| = |PN| \Rightarrow |PM|^2 = |PN|^2$$

$$k^2 = h^2.$$

$$\text{equation of locus } y^2 = x^2 \Rightarrow x^2 = y^2$$



5. Find the equation of locus of a point equidistant from A(2,0) and the Y-axis.


Sol: Let P(h, k) be any point

given point = A(2, 0)

given condition |PA| = h

$$|PA|^2 = h^2 \Rightarrow (h-2)^2 + k^2 = h^2$$

$$\text{equation of locus } y^2 - 4x^2 + 4 = 0.$$



6. Find the equation of locus of a point P such that  $|PA|^2 + |PB|^2 = 2c^2$  where A(a, 0),

B(-a, 0) and  $0 < |a| < |c|$

Sol: Let P(h, k) be any point

given point = A(a, 0), B(-a, 0)

given condition  $|PA|^2 + |PB|^2 = 2c^2$

$$(h-a)^2 + k^2 + (h+a)^2 + k^2 = 2c^2$$

$$\text{simplifying } h^2 + k^2 = c^2 - a^2$$

$$\therefore \text{equation of locus } x^2 + y^2 = c^2 - a^2.$$

7. Find the equation of locus of P, if the line segment joining (2, 3) and (-1, 5) subtends a right angle at P.

**Sol:** Let P( $h$ ,  $k$ ) be any point

given point = A(2, 3), B(-1, 5)

$$\text{slope of PA } m_1 = \frac{y_2 - y_1}{x_2 - x_1} = \frac{k - 3}{h - 2}$$

$$\text{slope of PB } m_2 = \frac{k - 5}{h + 1}$$


PA  $\perp$  PB

$$m_1 \times m_2 = -1$$

$$\Rightarrow \frac{k - 3}{h - 2} \times \frac{k - 5}{h + 1} = -1$$

simplifying  $h^2 + k^2 - h - 8k + 13 =$

$$\therefore \text{equation of locus } x^2 + y^2 - x - 8y + 13 = 0.$$



8. The ends of the hypotenuse of a right angle triangle are (0,6) and (6,0). Find the equation of the locus of its third vertex.

**Sol:** Let P( $h$ ,  $k$ ) be any point

given point = (6, 0), (0, 6)

$$\text{slope of PA } m_1 = \frac{y_2 - y_1}{x_2 - x_1} = \frac{k - 0}{h - 0}$$

$$\text{slope of PB } m_2 = \frac{k - 6}{h - 0}$$

given condition PA  $\perp$  PB


$$m_1 \times m_2 = -1$$

$$\Rightarrow \frac{k}{h - 6} \times \frac{k - 6}{h} = -1$$

$$\Rightarrow k^2 - 6k = [h^2 - 6h]$$

simplifying  $h^2 + k^2 - 6h - 6k = 0.$

$$\therefore \text{equation of locus } x^2 + y^2 - 6x - 6y = 0.$$



9. Find the equation of the locus of a point, the difference of whose distance from (-5, 0) and (3, 0) is 8.

**Sol:** Let P( $h$ ,  $k$ ) be any point

given point = A(-5, 0), B(3, 0)



given condition  $|PA - PB| = 8$

$$|PA| = 8 + |PB|$$

squaring both sides

$$|PA|^2 = [8 + |PB|]^2$$

simplifying  $|PA|^2 - |PB|^2 - 64 = 16 |PB|$

$$[(h+5)^2 + k^2] - [(h-5)^2 + k^2] - 64 = 16 |PB|$$

$$\Rightarrow 20h - 64 = 16 |PB|$$

$$\Rightarrow 5h - 16 = 4 |PB|$$

$$\Rightarrow (5h - 16)^2 = 16 |PB|^2$$

$$\Rightarrow (5h - 16)^2 = 16 [(h-5) + k^2]$$

simplifying  $\frac{h^2}{16} - \frac{k^2}{9} = 1$

$\therefore$  equation of locus  $\frac{x^2}{16} - \frac{y^2}{9} = 1$ .

**10. Find the equation of the locus of P, if A(4, 0), B(-4, 0) and  $|PA - PB| = 4$**

**Sol:** Let P(h, k) be any point

given points = A(4, 0), B(-4, 0)

given condition  $|PA - PB| = 4$

$$\Rightarrow |PA| = 4 + |PB|$$

$$[|PA|^2 - |PB|^2] - 16 = 8 |PB|$$

$$[(h-4)^2 + k^2] - [(h+4)^2 + k^2] - 16 = 8 |PB|$$

$$-16h - 16 = 8 |PB|$$

$$4(h+1)^2 = (h+4)^2 + k^2$$

simplifying  $3h^2 - k^2 = 12$

$\therefore$  equation of locus  $\frac{x^2}{4} - \frac{y^2}{12} = 1$ .

**11. Find the equation of the locus of a point, the sum of whose distances from (0,2) and (0,-2) is 6.**

**Sol:** Let P(h, k) be any point

given points = A(2, 3), B(2, -3)

given condition  $|PA| + |PB| = 8$

$$\Rightarrow |PA| = 8 - |PB|$$

$$[|PA|^2 - |PB|^2] - 64 = -16 |PB|$$

$$[(h-2)^2 + (k-3)^2] - [(h-2)^2 + (k+3)^2] - 64 = -16 |PB|$$

$$-12k - 64 = -16 |PB|$$

$$(3k + 16)^2 = 16[(h - 2)^2 + (k + 3)^2]$$

$$\text{simplifying } 16h^2 + 7k^2 - 64h - 48 = 0$$

$$\therefore \text{equation of locus } 16x^2 + 7y^2 - 64x - 48 = 0.$$


12. **A(5, 3), B(3, -2) are two fixed points. Find the equation of the locus of P, so that the area of triangle PAB is 9.**

**Sol:** Let 3rd vertex be P(h, k)

given points = A(5, 3), B(3, -2)

given condition |PA| + |PB| = 8

$$\text{area of } \Delta PAB = \frac{1}{2} \begin{vmatrix} h & k \\ 5 & 3 \\ 3 & -2 \\ h & k \end{vmatrix} = 9$$



$$|3h - 5k - 10 - 9 - 36k + 2h| = 18$$

$$|5h - 2k - 19| = 18$$

$$(5h - 2k - 1)(5h - 2k - 37) = 0$$

$$\therefore \text{equation of locus } (5h - 2k - 1)(5h - 2k - 37) = 0.$$

### Practice Problem

**A(2,3) and B(-3,4) are two given points. Find the equation of locus of P so that the area of the triangle PAB is 8.5**

13. **If the distance from P to the points (2, 3), (2, -3) are in the ratio 2 : 3, then find the equation of the locus of P.**

**Sol:** Let (2, 3), (2, -3) (h, k) are the vertices of the triangle

Let P(h, k) be any point

Given points = A(2, 3), B(2, -3)

$$\text{given condition } \frac{|PA|}{|PB|} = \frac{2}{3}$$

$$\Rightarrow 3|PA| = 2|PB|$$

$$9|PA|^2 = 4|PB|^2$$

$$9[(h - 2)^2 + (k - 3)^2] = 4[(h - 2)^2 + (k + 3)^2]$$

$$\text{simplifying } 5h^2 + 5k^2 - 20h - 78k + 65 = 0$$

$$\therefore \text{equation of locus } 5x^2 + 5y^2 - 20x - 78y + 65 = 0.$$

14. **A(1, 2), B(2, -3) and C(-2, 3) are three points. A point P moves such that  $|PA|^2 + |PB|^2 = 2|PC|^2$ . Show that the equation to the locus of P is  $7x - 7y + 4 = 0$ .**

**Sol:** Let P( $h, k$ ) be any point

$$|PA|^2 + |PB|^2 = (h-1)^2 + (k-2)^2 + (h-2)^2 + (k+3)^2$$

$$2|PC|^2 = 2[(h+2)^2 + (k-3)^2]$$

substituting in  $|PA|^2 + |PB|^2 = 2|PC|^2$

simplifying  $7h - 7k + 4 = 0$

$\therefore$  equation of locus  $7x - 7y + 4 = 0$ .

\*\*\*\*\*

TSBIE

## Transformation of Axis

**Def :** Without changing the direction of co-ordinate axes of the origin is shifted to a given point then the change occurred is called translation of axes.

Let  $P(x, y)$  original coordinates

and  $P(X, Y)$  transformed coordinates

$$x = PQ = ON = OL + LN = OL + O'M = h + X = X + h$$

$$y = PN = PM + MN = Y + O'L = Y + k$$

$$\therefore x = X + h, y = Y + k \Rightarrow X = x - h, Y = y - k$$

$$P(x, y) = (X + h, Y + k)$$

$$P(X, Y) = (x - h, y - k)$$

$$(h, k) = (x - X, y - Y)$$

original equation of the curve  $f(x, y)$

transformed equation of the curve  $f(X, Y)$

**Rotation of axes :** without the changing the position of origin the axes are rotated through an angle then it is called rotation of axes.

$$x = OL = OQ - LQ = X\cos\theta - NM$$

$$= X\cos\theta - Y\sin\theta.$$

$$y = PL = PN + NL = PN + MQ$$

$$= PM\cos\theta + OM\sin\theta$$

$$= Y\cos\theta + X\sin\theta = X\sin\theta + Y\cos\theta.$$

$$P(x, y) = (X\cos\theta - Y\sin\theta, X\sin\theta + Y\cos\theta)$$

$\theta$	X	Y
x	$\cos\theta$	$-\sin\theta$
y	$\sin\theta$	$\cos\theta$

## Short Answer Questions (4 Marks)

1. When the origin is shifted to  $(4, -5)$  by the translation of axes, find the coordinates of the following points with reference to new axes.

**Sol :** (i)  $(0, 3)$

$$\begin{array}{llll} x = 0, & y = 3 & h = 4 & k = -5 \\ X = x - h & Y = y - k & & \\ X = -4 & Y = 8 & \text{Ans: } (4, -8) & \end{array}$$

(ii)  $(-2, 4)$

$$\begin{array}{llll} x = -2 & y = 4 & h = 4 & k = -5 \\ X = x - h & Y = y - k & & \\ X = -4 - 2 = -6 & Y = 4 + 5 = 9 & \text{Ans: } (-6, 9) & \end{array}$$

(iii)  $(4, -5)$

$$\begin{array}{llll} x = 4 & y = -5 & h = 4 & k = -5 \\ (X, Y) = (x - h, y - k) = (4 - 4, -5 - (-5)) = (0, 0) & & & \end{array}$$

2. The origin is shifted to  $(2, 3)$  by the translation of axes. If the coordinates of a point P change as follows, find the coordinates of P in the original system.

**Sol :** (i)  $(4, 5)$

$$\begin{array}{llll} X = 4 & Y = 5 & h = 2 & k = 3 \\ x = X + h & y = Y + k & & \\ x = 6 & Y = 8 & \text{Ans: } (6, 8) & \end{array}$$

(ii)  $(-4, 3)$

$$\begin{array}{llll} X = -4 & Y = 3 & h = 2 & k = 3 \\ x = X + h & y = Y + k & & \\ x = -2 & Y = 6 & \text{Ans: } (-2, 6) & \end{array}$$

(iii)  $(0, 0)$

$$\begin{array}{llll} X = 0 & Y = 0 & h = 2 & k = 3 \\ x = X + h & y = Y + k & & \\ x = 2 & Y = 3 & \text{Ans: } (2, 3) & \end{array}$$

3. Find the point to which the origin is to be shifted so that the point  $(3, 0)$  may change to  $(2, -3)$ .

**Sol :**  $(x, y) = (3, 0)$  given point  
 $(X, Y) = (2, -3)$  transformed point  
 $(h, k) = (x - X, y - Y) = (3 - 2, 0 - (-3)) = (1, 3)$

4. When the origin is shifted to  $(-1, 2)$  by the translation of axes, find the transformed equations of the following.

**Sol :** (1)  $x^2 + y^2 + 2x - 4y + 1 = 0$

$$(h, k) = (-1, 2)$$

$$x = X + h = X - 1 \qquad y = Y + k = Y + 2$$

$$\text{substituting } (X-1)^2 + (Y+2)^2 + 2(X-1) - 4(Y+2) + 1 = 0$$

simplifying  $X^2 + Y^2 - 4 = 0$  required equation

$$(2) \quad 2x^2 + y^2 - 4x + 4y = 0$$

$$x = X + h = X - 1$$

$$y = Y + k = Y + 2$$

$$\text{substituting } 2(X-1)^2 + (Y+2)^2 - 4(X-1) + 4(Y+2) = 0$$

$$\text{simplifying } 2[X^2 - 2X + 1] + [Y^2 + 4Y + 4] - 4X + 4 + 4Y + 8 = 0$$

$$2X^2 + Y^2 - 8X + 8Y + 18 = 0 \text{ required equation}$$

- 5. The point to which the origin is shifted and the transformed equation are given below. Find the original equation.**

**Sol :** (1)  $(3, -4), x^2 + y^2 = 4$

Let the transformed equation indicated by  $X^2 + Y^2 = 4$  ..... (1)

simplifying  $(h, k) = (3, -4)$

$$X = x - h \quad Y = y - k$$

$$X = x - 3 \quad Y = y + 4$$

original equation after substitution and simplification

$$x^2 + y^2 - 6x + 8y + 21 = 0$$

$$(2) \quad (-1, 2), x^2 + y^2 + 16 = 0$$

changed origin is  $(h, k) = (-1, 2)$

$$\text{given transformed equation } x^2 + 2y^2 + 16 = 0$$

Let this denoted by  $X^2 + 2Y^2 + 16 = 0$  .....(1)

$$X = x - h = x + 1 \quad Y = y - k = y - 2$$

original equation after substitution and simplification

$$x^2 + 2y^2 + 2x - 8y + 25 = 0$$

- 6. Find the point to which the origin is to be shifted so as to remove the first degree terms from the equation.**

**Sol :** given equation  $f(x, y) = 4x^2 + 9y^2 - 8x + 36y + 4 = 0$

transformed equation

$$f(x+h, y+k) = f(x, y) = 4(X+h)^2 + 9(Y+k)^2 - 8(X+h) + 36(Y+k) + 4 = 0$$

X, Y to eliminate X, Y terms equate coefficient of X and coefficient of Y to 0

$$8h - 8 = 0 \Rightarrow h = 1$$

$$18k + 36 = 0 \Rightarrow k = -2$$

the point to which the origin to be shifted is  $(h, k) = (1, -2)$  or

use the formula  $\left( \frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right)$

7. Find the angle through which the axes are to be rotated so as to remove the  $xy$  term in the equation  $x^2 + 4xy + y^2 - 2x + 2y - 6 = 0$

**Sol :** transformed equation  $x^2 + 3xy - 2y^2 + 17x - 7y - 11 = 0$

for convenient let the equation  $X^2 + 3XY - 2Y^2 + 17X - 7Y - 11 = 0$  ..... (1)

$(h, k) = (2, 3)$

$X = x - h = x - 2$                        $Y = y - k = y - 3$

substituting in (1)

$$(x - 2)^2 + 3(x - 2)(y - 3) - 2(y - 3)^2 + 17(x - 2) - 7(y - 3) - 11 = 0$$

original equation  $x^2 + 3xy - 2y^2 + 4x - y - 20 = 0$

**Very Short Answer Questions (2 Marks)**

1. When the axes are rotated through an angle  $30^\circ$ , find the new coordinates of the following points.

**Sol :** (i)  $(0, 5)$                        $\theta = 30^\circ$

$$X = x \cos \theta + y \sin \theta = 5 \sin 30^\circ = \frac{5}{2}$$

$$Y = -x \cos \theta + y \cos \theta = 5 \cos 30^\circ = \frac{5\sqrt{3}}{2}$$

$$(X, Y) = \left( \frac{5}{2}, \frac{5\sqrt{3}}{2} \right)$$

(ii)  $(-2, 4)$                        $\theta = 30^\circ$

$$X = -2 \cos 30^\circ + 4 \sin 30^\circ = \frac{-2\sqrt{3}}{2} + \frac{4}{2} = 2 - \sqrt{3}$$

$$Y = -2 \sin 30^\circ + 4 \cos 30^\circ = \frac{-2}{2} + \frac{4\sqrt{3}}{2} = 2\sqrt{3} - 1$$

$$(2 - \sqrt{3}, 2\sqrt{3} - 1)$$

(iii)  $(0, 0)$                        $\theta = 30^\circ$                       (Home work)

2. When the axes are rotated through an angle  $60^\circ$ , the new coordinates of three points are the following.

**Sol :** (i)  $(X, Y) = (3, 4)$                        $\theta = 60^\circ$

$$x = X \cos \theta - Y \sin \theta = 3 \cos 60^\circ - 4 \sin 60^\circ = \frac{3 - 4\sqrt{3}}{2}$$

$$y = X\sin\theta + Y\cos\theta = 3\sin 60^\circ + 4\cos 60^\circ = \frac{3\sqrt{3} + 4}{2}$$

$$(x, y) = \left( \frac{3 - 4\sqrt{3}}{2}, \frac{3\sqrt{3} + 4}{2} \right)$$

(ii)  $(-7, 2) = (X, Y) \quad \theta = 60^\circ$

$$x = X\cos\theta - Y\sin\theta = -7\cos 60^\circ - 2\sin 60^\circ = \frac{-7 - 2\sqrt{3}}{2}$$

$$y = X\sin\theta + Y\cos\theta = -7\sin 60^\circ + 2\cos 60^\circ = \frac{2 - 7\sqrt{3}}{2}$$

$$\text{Ans: } \left( \frac{-7 - 2\sqrt{3}}{2}, \frac{2 - 7\sqrt{3}}{2} \right)$$

(iii)  $(2, 0) = (X, Y) \quad \theta = 60^\circ$

$$x = X\cos\theta - Y\sin\theta = 2\cos 60^\circ - 0\sin 60^\circ = 1$$

$$y = X\sin\theta + Y\cos\theta = 2\sin 60^\circ + 0\cos 60^\circ = \frac{2\sqrt{3}}{2} = \sqrt{3}$$

$$(x, y) = (1, \sqrt{3})$$

3. When the axes are rotated through an angle  $\frac{\pi}{6}$ , find the transformed equation of

$$x^2 + 2\sqrt{3}xy - y^2 = 2a^2$$

Sol :  $\theta = \pi/6$

$$f(x, y) = x^2 + 2\sqrt{3}xy - y^2 = 2a^2$$

$$x = X\cos\theta - Y\sin\theta = X\cos\frac{\pi}{6} - Y\sin\frac{\pi}{6} = \frac{X\sqrt{3}}{2} - \frac{Y}{2}$$

$$y = X\sin\theta + Y\cos\theta = X\sin\frac{\pi}{6} + Y\cos\frac{\pi}{6} = \frac{X}{2} + \frac{Y\sqrt{3}}{2}$$

transformed equation =  $f(X, Y)$

$$= \left( \frac{\sqrt{3}X - Y}{2} \right)^2 + 2\sqrt{3} \left( \frac{\sqrt{3}X - Y}{2} \right) \left( \frac{X + Y\sqrt{3}}{2} \right) - \left( \frac{X + Y\sqrt{3}}{2} \right)^2 = 2a^2$$

simplifying  $X^2 - Y^2 = a^2$ .



4. When the axes are rotated through an angle  $\pi/4$ , find the transformed equation of

$$3x^2 + 10xy + 3y^2 = 9$$

Sol :  $\theta = \pi/4$

$$f(x, y) = 3x^2 + 10xy + 3y^2 = 9$$

$$x = X\cos\theta - Y\sin\theta = \frac{X - Y}{\sqrt{2}}$$

$$y = X\sin\theta + Y\cos\theta = \frac{X + Y}{\sqrt{2}}$$

by substituting and simplifying the transformed equation  $8X^2 - 2Y^2 = 9$

5. When the axes are rotated through an angle  $\alpha$ , find the transformed equation of  $x\cos\alpha + y\sin\alpha = p$

Sol :  $x = X\cos\alpha - Y\sin\alpha$

$$y = X\sin\alpha + Y\cos\alpha$$

by substituting

$$(X\cos\alpha - Y\sin\alpha)\cos\alpha + (X\sin\alpha + Y\cos\alpha)\sin\alpha = p$$

$$X(\cos^2\alpha + \sin^2\alpha) = p$$

$$X = p.$$

6. When the axes are rotated through an angle  $45^\circ$ , the transformed equation of a curve is  $17x^2 - 16xy + 17y^2 = 225$ . Find the original equation of the curve.

Sol : angle of rotation  $= \theta = 45^\circ$

let the transformed equation be

$$17X^2 - 16XY + 17Y^2 = 225$$

$$X = x\cos\theta + y\sin\theta \quad Y = -x\sin\theta + y\cos\theta$$

$$= \frac{x+y}{\sqrt{2}} \quad = \frac{-x+y}{\sqrt{2}}$$

simplifying

$$17\left(\frac{x+y}{\sqrt{2}}\right)^2 - 16\left(\frac{x+y}{\sqrt{2}}\right)\left(\frac{-x+y}{\sqrt{2}}\right) + 17\left(\frac{-x+y}{\sqrt{2}}\right)^2 = 225$$

$$\text{required equation } 25x^2 + 9y^2 = 225$$

\*\*\*\*\*

## The Straight Line

### Chapter 3(a)

#### Note :

1. If a non - vertical straight line makes an angle ' $\theta$ ' with the X- axis measured counter - clock wise from the positive direction of X - axis, then  $\tan\theta$  is called the slope of the line L.

$$m = \tan\theta$$

2. Slope of X- axis & its parallel line is zero
3. Slope of Y- axis & its parallel line is not define
4. A line which is passing through  $A(x_1, y_1)$ ,  $B(x_2, y_2)$  then its slope  $(m) = \frac{y_2 - y_1}{x_2 - x_1}$ .
5. The equation of the S.L. which cut off non - zero intercepts ' $a$ ' and ' $b$ ' on the X-axis and the Y-axis respectively is  $\frac{x}{a} + \frac{y}{b} = 1$

#### Intercept Form

6. The equation of the S.L. with slope ' $m$ ' and cutting off y - intercept ' $C$ ' is  $y = mx + c$  (slope - intercept form)
7. If it passes through origin then the equation is  $y = mx$
8. **Point - slope form :**  
The equation of the straight line with slope ' $m$ ' and passing through the point  $(x_1, y_1)$  is  $y - y_1 = m(x - x_1)$ .
9. The equation of the S.L. passing through the points  $A(x_1, y_1)$  and  $B(x_2, y_2)$  is  $y - y_1 = \left( \frac{y_2 - y_1}{x_2 - x_1} \right) (x - x_1)$  (Two point form).

$$10. \quad A(x_1, y_1), B(x_2, y_2), C(x_3, y_3) \text{ are collinear} \Leftrightarrow \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = 0 \quad (\text{OR})$$

$$\text{Slope of AC} = \text{Slope of AB} \quad (\text{OR}) \quad \text{Area of } \triangle ABC = 0.$$

### Questions

1. Find the equation of the S.L. passing through the points  $(at_1^2, 2at_1)$  and  $(at_2^2, 2at_2)$

Sol:  $A(at_1^2, 2at_1), B(at_2^2, 2at_2)$

$$y - y_1 = \left( \frac{y_2 - y_1}{x_2 - x_1} \right) (x - x_1)$$

$$y - 2at_1 = \left( \frac{2at_2 - 2at_1}{at_2^2 - at_1^2} \right) (x - at_1^2)$$

$$y - 2at_1 = \frac{2a(t_2 - t_1)}{a(t_2^2 - t_1^2)} (x - at_1^2)$$

$$y - 2at_1 = \frac{2(t_2 - t_1)}{(t_2 - t_1)(t_2 + t_1)} (x - at_1^2)$$

$$(t_2 + t_1)(y - 2at_1) = 2(x - at_1^2)$$

$$(t_1 + t_2)y - (t_1 + t_2)2at_1 = 2x - 2at_1^2$$

$$x - 2at_1^2 - (t_1 + t_2)y + (t_1 + t_2)2at_1 = 0$$

$$x - 2at_1 - (t_1 + t_2)y + 2at_1^2 + 2at_1t_2 = 0$$

$$2x - (t_1 + t_2)y + 2at_1t_2 = 0$$

2. Find the value of x, if the slope of the line passing through (2, 5) and (x, 3) is 2.

Sol:  $A(2, 5), B(x, 3), m = 2$

$$\text{Slope of AB} = \frac{y_2 - y_1}{x_2 - x_1}$$

$$= \frac{3 - 5}{x - 2} = 2$$

$$-2 = 2(x - 2)$$

$$2x - 4 = -2$$

$$2x = -2 + 4 \Rightarrow 2x = 2 \Rightarrow x = 1.$$

3. Find the value of  $y$  if the line joining points  $(3, 4)$  and  $(2, 7)$  is parallel to the line the points  $(-1, 4)$  and  $(0, 6)$

Sol:  $A(3, 4), B(2, 7), C(-1, 4), D(0, 6)$

Slope of AB = Slope of CD

$$\frac{7-y}{2-3} = \frac{6-4}{0-(-1)} \Rightarrow \frac{7-y}{-1} = \frac{2}{1}$$

$$7-y = -2 \Rightarrow 7+2 = y \Rightarrow y = 9.$$

4. Find the condition for the points  $(a, 0)$  and  $(0, b)$ , where  $ab \neq 6$ , to be collinear

Sol:  $A(a, 0), B(h, k), C(0, b)$

points A, B & C are collinear  $\Leftrightarrow$  Slope of AC = Slope of AB

$$\frac{b-0}{0-a} = \frac{k-0}{h-a} \Rightarrow \frac{-b}{a} = \frac{k}{h-a}$$

$$\Rightarrow -b(h-a) = ak$$

$$\Rightarrow -bh + ab = ak \Rightarrow ak + bh = ab$$

$$\frac{ak}{ab} + \frac{bh}{ab} = \frac{ab}{ab} \Rightarrow \frac{h}{a} + \frac{k}{b} = 1$$

5. Find the equation of the S.L. which makes the  $150^\circ$  with the position X-axis in the positive direction and which passes through the point  $(-2, -1)$

Sol:  $\theta = 150^\circ, A(-2, -1)$

$$m = \tan 150^\circ = \tan(90 + 60) = -\cot 60^\circ = -\frac{1}{\sqrt{3}}$$

The equation of the S.L. with slope  $-\frac{1}{\sqrt{3}}$  and passing through the point  $(-2, -1)$  is

$$y - y_1 = m(x - x_1)$$

$$y + 1 = \frac{-1}{\sqrt{3}}(x + 2)$$

$$\sqrt{3}(y + 1) = -(x + 2)$$

$$\sqrt{3}y + \sqrt{3} = -(x + 2)$$

$$\sqrt{3}(y + 1) = -x - 2$$

$$x + \sqrt{3}y + 2 + 2 + \sqrt{3} = 0$$

**Exercise:**

Find the slopes of the lines (i) parallel to and (ii) perpendicular to the line passing through (6, 3) and (-4, 5)

6. The angle made by a S.L. with the positive X-axis in the positive direction is  $60^\circ$  and the X- intercept cut off by it is 3. Find the equation of the S.L

Sol:  $\theta = 60^\circ, C = 3$

$\therefore$  Required equation  $y = mx + c$ . ( $m = \tan 60^\circ = \sqrt{3}$ )

$$y = \sqrt{3}x + 3$$

$$\Rightarrow \sqrt{3}x - y + 3 = 0$$

7. Find the equation of the S.L. passing through the origin and making equal angles with the coordinate axes.

Sol: O(0, 0)

Since making equal angles with the coordinate axes

$$\theta = 45^\circ, 135^\circ.$$

$$m = \tan 45^\circ \text{ or } \tan 135^\circ$$

$$m = 1 \text{ or } \tan 135^\circ = \tan(180^\circ - 45^\circ) = -\tan 45^\circ = -1$$

$$m = \pm 1$$

$\therefore$  Equation of S.L. with slope  $\pm 1$  and passing through origin is

$$(y - 0) = \pm 1(x - 0)$$

$$y = \pm x$$

$$x + y = 0 \text{ or } x - y = 0$$

8. Find the equation of the S.L. passing through the point (2, 3) and making non - zero intercepts on the axes of coordinatas whose sum is zero

Sol: A(2, 3)

sum of the intercepts = zero

$$a + b = 0 \Rightarrow b = -a$$

Equation of a S.L. in intercept form

$$\frac{x}{a} + \frac{y}{b} = 1 \Rightarrow \frac{x}{a} + \frac{y}{-a} = 1$$

$$\Rightarrow \frac{x - y}{a} = 1 \Rightarrow x - y = a \quad \dots(1)$$

But it is passing through A(2, 3)

$$2 - 3 = a \Rightarrow a = -1$$

$\therefore$  From (1)

$$x - y = -1$$

$$x - y + 1 = 0$$

9. Find the equation of the S.L. passing through  $(-4, 5)$  and cutting off equal and non zero intercepts on the coordinate axes.

Sol:  $A = (-4, 5)$

According to the problem  $a = b$   
equation of a S.L. in intercept form

$$\boxed{\frac{x}{a} + \frac{y}{b} = 1}$$

$$\Rightarrow \frac{x}{a} + \frac{y}{a} = 1$$

$$\Rightarrow \frac{x+y}{a} = 1 \Rightarrow x+y = a \quad \dots(1)$$

But it is passing through  $A(-4, 5)$

$$-4 + 5 = a \Rightarrow a = 1$$

$\therefore$  Equation of the required S.L. is

$$x + y = 1$$

$$x + y - 1 = 0$$

10. Find the equation of the S.L. passing through  $A(-1, 3)$  and (i) parallel (ii) perpendicular to the S.L. passing through  $B(2, -5)$  and  $C(4, 6)$

Sol:  $A(-1, 3), B(2, -5), C(4, 6)$

$$\text{Slope of BC} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6 - (-5)}{4 - 2} = \frac{6 + 5}{2} = \frac{11}{2}$$

(i) Slope of a parallel line to BC also  $= \frac{11}{2}$

equation of S.L. with slope  $\frac{11}{2}$  and passing through  $A(-1, 3)$  is

$$y - 3 = \frac{11}{2}(x - (-1))$$

$$2(y - 3) = 11(x + 1)$$

$$2y - 6 = 11x + 11$$

$$11x - 2y + 17 = 0$$

(ii) Slope of a line which is perpendicular to BC is  $= \frac{-1}{m}$

$$= -\frac{1}{\left(\frac{11}{2}\right)} = \frac{-2}{11}$$

equation of a S.L. with slope  $\frac{-2}{11}$  and passing through A(-1, 3) is

$$y - 3 = \frac{-2}{11}(x - (-1))$$

$$11(y - 3) = -2(x + 1)$$

$$11y - 33 = -2x - 2$$

$$2x + 11y - 31 = 0$$

## The Straight Line

### Exercise 2b

#### Note :

- The equation of the straight line, whose distance from the origin is P and the normal ray of which drawn from the origin makes an angle  $\alpha$  with the positive direction of the X-axis measured counter clock - wise is

$$x \cos \alpha + y \sin \alpha = P$$

It is called Normal Form

- The equation of the straight line passing through  $(x_1, y_1)$  and making an angle  $\theta$  with the positive direction of the X-axis measured counter - clock wise is

$$(x - x_1) : \cos \theta = (y - y_1) : \sin \theta$$

- $\frac{x - x_1}{\cos \theta} = \frac{y - y_1}{\sin \theta} = r$  say

$x = x_1 + r \cos \theta$ ,  $y = y_1 + r \sin \theta$  is called parametric form of the point P.

$|r|$  denotes the distance of the point  $(x_1, y_1)$  from the point  $(x, y)$  on the straight line.

- General form of a straight line is  $ax + by + c = 0$

$$\text{Its slope } (m) = \frac{-a}{b}$$

### Questions

- Transform the equation  $x + y + 1 = 0$  into normal form**

**Sol:**  $x + y + 1 = 0$

Normal form of  $ax + by + c = 0$  is

$$\frac{-ax}{\sqrt{a^2 + b^2}} + \frac{-(b)y}{\sqrt{a^2 + b^2}} = \frac{c}{\sqrt{a^2 + b^2}}, \quad \text{where } c > 0$$

$$\text{Divide with } \sqrt{a^2 + b^2} = \sqrt{(1)^2 + (1)^2} = \sqrt{1+1} = \sqrt{2}$$

$$\frac{x}{\sqrt{2}} + \frac{y}{\sqrt{2}} + \frac{1}{\sqrt{2}} = 0$$

$$\left(\frac{-1}{\sqrt{2}}\right)x + \left(\frac{-1}{\sqrt{2}}\right)y = \frac{1}{\sqrt{2}} \quad \dots\dots(1)$$

Compare with  $x \cos \alpha + y \sin \alpha = P$

$$\cos \alpha = \frac{-1}{\sqrt{2}}, \sin \alpha = \frac{-1}{\sqrt{2}}, P = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \alpha = \frac{5\pi}{4}$$

$\therefore$  Equation of the required straight line

$$x \cos\left(\frac{5\pi}{4}\right) + y \sin\left(\frac{5\pi}{4}\right) = \frac{1}{\sqrt{2}}$$

- 2. Transform the equation  $4x - 3y + 12 = 0$  into (a) slope - intercept form (b) intercept form and (c) normal form**

**Sol. :**  $L = 4x - 3y + 12 = 0$

- (a) Slope - intercept form ( $y = mx + c = 0$ )

$$\Rightarrow 4x + 12 = 3y$$

$$y = \frac{4x + 12}{3}$$

$$y = \left(\frac{4}{3}\right)x + 4$$

$$\Rightarrow m = \frac{4}{3}, c = 4$$

- (b) Intercept form  $\frac{x}{a} + \frac{y}{b} = 1$

$$\Rightarrow 4x - 3y + 12 = 0$$

$$-4x + 3y = 12$$

$$\frac{-4x}{12} + \frac{3y}{12} = \frac{12}{12}$$

$$\frac{x}{(-3)} + \frac{y}{(4)} = 1$$

$$\Rightarrow a = -3, b = 4$$

- (c) Normal form ( $x \cos \alpha + y \sin \alpha = P$ )

$$4x - 3y + 12 = 0$$



$$-4x + 3y = 12$$

Divide with  $\sqrt{a^2 + b^2} = \sqrt{(-4)^2 + (3)^2} = \sqrt{16 + 9} = 5$

$$\frac{-4x}{5} + \frac{3y}{5} = \frac{12}{5}$$

$$\cos \alpha = \frac{-4}{5}, \sin \alpha = \frac{3}{5}, P = \frac{12}{5}$$

3. Transform the equation  $\sqrt{3}x + y = 4$  into (a) slope intercept form (b) intercept form and (c) normal form.

Sol. :  $L = \sqrt{3}x + y = 4$

- (a) Slope - intercept form  $y = mx + c = 0$

$$\sqrt{3}x + y = 4$$

$$\Rightarrow y = -\sqrt{3}x + 4$$

$$\Rightarrow m = -\sqrt{3}, c = 4$$

- (b) Intercept form  $\frac{x}{a} + \frac{y}{b} = 1$

$$\sqrt{3}x + y = 4$$

$$\Rightarrow \frac{\sqrt{3}}{4}x + \frac{y}{4} = \frac{4}{4}$$

$$\frac{x}{\left(\frac{4}{\sqrt{3}}\right)} + \frac{y}{(4)} = 1$$

$$\Rightarrow a = \frac{4}{\sqrt{3}}, b = 4$$

- (c) Normal form ( $x \cos \alpha + y \sin \alpha = P$ )

$$\sqrt{3}x + y = 4$$

Divide with  $\sqrt{a^2 + b^2} = \sqrt{(\sqrt{3})^2 + (1)^2} = \sqrt{3 + 1} = 2$

$$\frac{\sqrt{3}x}{2} + \frac{y}{2} = \frac{4}{2}$$

$$2\left(\frac{\sqrt{3}}{2}\right) + y\left(\frac{1}{2}\right) = 2$$

$$\cos \alpha = \frac{\sqrt{3}}{2}, \sin \alpha = \frac{1}{2}, P = 2$$

$$\Rightarrow \alpha = 30^\circ = \frac{\pi}{6}$$

$$\therefore x \cos \frac{\pi}{6} + y \sin \frac{\pi}{6} = 2$$

**Home Work**

II 3i, iv, v, vi

4. A straight line parallel to the line  $y = \sqrt{3}x$  passes through  $Q(2, 3)$  and cuts the line  $2x + 4y - 27 = 0$  at  $P$ . Find the length  $PQ$

**Sol:**  $y = \sqrt{3}x$  .....(1)       $2x + 4y - 27 = 0$  .....(2)

Slope of a equation which is parallel to (1) also  $\sqrt{3}$ .

$$m = \sqrt{3} = \tan 60^\circ \Rightarrow \theta = 60^\circ$$

$$\theta = 60^\circ \text{ and } Q(2, 3)$$

$$P = (x, y) = (x_1 + r \cos \theta, y_1 + r \sin \theta)$$

$$P = (2 + r \cos \theta, 3 + r \sin \theta)$$

$$= \left( 2 + r \left( \frac{1}{2} \right), 3 + r \left( \frac{\sqrt{3}}{2} \right) \right)$$

$$= \left( 2 + \frac{r}{2}, 3 + \frac{\sqrt{3}r}{2} \right)$$

But  $P$  is on equation (2)

$$2 \left( 2 + \frac{r}{2} \right) + 4 \left( 3 + \frac{\sqrt{3}r}{2} \right) - 27 = 0$$

$$2 \left( \frac{4+r}{2} \right) + 4 \left( \frac{6+\sqrt{3}r}{2} \right) = 27$$

$$4 + r + 12 + 2\sqrt{3}r = 27$$

$$(2\sqrt{3}r + 1)r = 11$$

$$\Rightarrow r = \frac{11}{2\sqrt{3}+1} = \frac{11}{2\sqrt{3}+1} \times \frac{2\sqrt{3}-1}{2\sqrt{3}-1} = \frac{11(2\sqrt{3}-1)}{12-1}$$

$$\Rightarrow r = \frac{11(2\sqrt{3}-1)}{11} = 2\sqrt{3}-1$$

$$\therefore PQ = |r| = 2\sqrt{3}-1$$

5. If the area of the triangle formed by the straight lines  $x = 0$ ,  $y = 0$  and  $3x + 4y = a$  ( $a > 0$ ) is 6, find the value of  $a$ .

Sol:  $3x + 4y = a$

$$\frac{3x}{a} + \frac{4y}{a} = \frac{a}{a} \Rightarrow \frac{x}{\left(\frac{a}{3}\right)} + \frac{y}{\left(\frac{a}{4}\right)} = 1 \quad \dots\dots\dots (1)$$

Area of the triangle with coordinates axes and  $\frac{x}{a} + \frac{y}{b} = 1$  is  $\Delta = \frac{1}{2}|ab|$

$$\therefore \Delta = \frac{1}{2} \left| \frac{a}{3} \times \frac{a}{4} \right| = 6 \Rightarrow \frac{a^2}{12} = 12$$

$$\Rightarrow a^2 = 144 \Rightarrow a = \sqrt{144}$$

$$\therefore a = 12.$$

6. Transform the equation  $\frac{x}{a} + \frac{y}{b} = 1$  into the normal form when  $a > 0$  and  $b > 0$ . If the perpendicular distance of the straight line from the origin is P, deduce that

$$\frac{1}{p^2} = \frac{1}{a^2} + \frac{1}{b^2}.$$

Sol:  $\frac{x}{a} + \frac{y}{b} = 1$

Divide with  $\sqrt{a^2 + b^2} = \sqrt{\left(\frac{1}{a}\right)^2 + \left(\frac{1}{b}\right)^2}$

$$\frac{\left(\frac{x}{a}\right)}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}} + \frac{\left(\frac{y}{b}\right)}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}} = \frac{1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}}$$

$$x \left( \frac{1}{a\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}} \right) + y \left( \frac{1}{b\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}} \right) = \frac{1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}}$$

According to the problem  $P = \frac{1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}} \Rightarrow \frac{1}{P} = \sqrt{\frac{1}{a^2} + \frac{1}{b^2}}$

Squaring on Both Sides

$$\frac{1}{p^2} = \frac{1}{a^2} + \frac{1}{b^2}$$

7. Line L has intercepts  $a$  and  $b$  on the axes of coordinates. When the axes are rotated through a given angle, keeping the origin fixed, the same line L has intercepts  $p$  and

$q$  on the transformed axes. Prove that  $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2} + \frac{1}{q^2}$

**Sol:** Equation of S.L. is in intercept form  $\frac{x}{a} + \frac{y}{b} = 1$  .....(1)

Let ' $\theta$ ' be the angle, where axes are notabled

$$x = x' \cos \theta - y' \sin \theta, \quad y = x' \sin \theta + y' \cos \theta$$

Transformed equation of (1) is

$$\frac{x' \cos \theta - y' \sin \theta}{a} + \frac{x' \sin \theta + y' \cos \theta}{b} = 1$$

$$\frac{b(x' \cos \theta - y' \sin \theta) + a(x' \sin \theta + y' \cos \theta)}{ab} = 1$$

$$\frac{bx' \cos \theta - by' \sin \theta + ax' \sin \theta + ay' \cos \theta}{ab} = 1$$

$$\frac{(a \sin \theta + b \cos \theta)x'}{ab} + \frac{(a \cos \theta - b \sin \theta)y'}{ab} = 1$$

$$\frac{\left(\frac{x'}{ab}\right)}{(a \sin \theta + b \cos \theta)} + \frac{\left(\frac{y'}{ab}\right)}{(a \cos \theta - b \sin \theta)} = 1 \quad \dots\dots(2)$$

According to the problem

$$p = \frac{ab}{a \sin \theta + b \cos \theta}, \quad q = \frac{ab}{a \cos \theta - b \sin \theta}$$

$$\frac{1}{p^2} + \frac{1}{q^2} = \left(\frac{1}{p}\right)^2 + \left(\frac{1}{q}\right)^2$$

$$= \left(\frac{a \sin \theta + b \cos \theta}{ab}\right)^2 + \left(\frac{a \cos \theta - b \sin \theta}{ab}\right)^2$$

$$= \frac{a^2 \sin^2 \theta + b^2 \cos^2 \theta + 2ab \sin \theta \cos \theta + a^2 \cos^2 \theta + b^2 \sin^2 \theta - 2ab \cos \theta \sin \theta}{(ab)^2}$$

$$= \frac{a^2(\sin^2 \theta + \cos^2 \theta) + b^2(\sin^2 \theta + \cos^2 \theta)}{a^2 b^2} = \frac{a^2(1) + b^2(1)}{a^2 b^2}$$

$$= \frac{a^2}{a^2b^2} + \frac{b^2}{a^2b^2} = \frac{1}{b^2} + \frac{1}{a^2}$$

$$\therefore \frac{1}{p^2} + \frac{1}{q^2} = \frac{1}{a^2} + \frac{1}{b^2}$$

## The Straight Line

### Exercise 3c

**Note :**

1  $L_1 = a_1x + b_1y + c_1 = 0$ ,  $L_2 = a_2x + b_2y + c_2 = 0$  then intersecting point of  $L_1, L_2$  is

$$\left( \frac{b_1c_2 - b_2c_1}{a_1b_2 - a_2b_1}, \frac{c_1a_2 - c_2a_1}{a_1b_2 - a_2b_1} \right)$$

2.  $L = ax + by + c = 0$  & two points  $A(x_1, y_1), B(x_2, y_2)$  then notation of

$$L_{11} = ax_1 + by_1 + c_1 = 0, \text{ \& } L_{22} = ax_2 + by_2 + c_2 = 0$$

(a) If A & B are same side to  $L = 0$

$\Rightarrow L_1$  &  $L_2$  has same sign

(b) If A & B are opposite side to  $L = 0$

$\Rightarrow L_{11}$  &  $L_{22}$  has opposite signs.

3. (a) X-axis divides line segment  $\overline{AB}$  in the ratio  $= -y_1:y_2$

(a) Y-axis divides the line segment  $\overline{AB}$  in the ratio  $= -x_1:x_2$

4.  $L_1 = 0, L_2 = 0, L_3 = 0$  are concurrent lines

$$\Leftrightarrow \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = 0 \quad \text{OR}$$

$$a_1(b_2c_3 - b_3c_2) + b_1(c_2a_3 - c_3a_2) + c_1(a_2b_3 - a_3b_2) = 0$$

## Questions

1. Find the ratio in which the straight line  $2x + 3y = 5$  divides the line segment joining the  $(0, 0)$  and  $(-2, 1)$ . State whether the points lie on the same side or neither side of the S.L.

**Sol:**  $L \equiv 2x + 3y = 5$ ,  $A(0, 0), B(-2, 1)$

$$L_{11} \equiv 2(0) + 3(0) = 5$$

$$L_{11} = -5 < 0$$

$$L_{22} \equiv 2(-2) + 3(1) - 5 = -4 + 3 - 5 = -6$$

$$L_{22} = -6 < 0$$

$L = 0$  divides the line segment  $\overline{AB}$  in the ratio  $= -L_{11} : L_{22}$

$$\text{Ratio} = -(-5) : (-6) = -5:6$$

$$L_{11} < 0, L_{22} < 0 \ \& \ L_{11}, L_{22} < 0$$

$\therefore$  A, B are lie on the same side of the line  $L = 0$

- 2. Find the value of  $k$ , if the lines  $2x - 3y + k = 0$ ,  $3x - 4y - 13 = 0$  and  $8x - 11y - 33 = 0$  are concurrent.**

**Sol:**  $L_1 \equiv 2x - 3y + k = 0$ ,  $L_2 \equiv 3x - 4y - 13 = 0$ ,  $L_3 \equiv 8x - 11y - 33 = 0$

$L_1, L_2, L_3$  are concurrent

$$\Rightarrow \begin{vmatrix} 2 & -3 & k \\ 3 & -4 & -13 \\ 8 & -11 & -33 \end{vmatrix} = 0$$

$$2(132 - 143) + 3(-99 + 104) + k(-33 + 32) = 0$$

$$2(-11) + 3(5) + k(-1) = 0$$

$$\therefore k = -7.$$

Home work  $I_5$  &  $II_7$

- 3. A triangle of area 24 sq. units is formed by a S.L. and the coordinate axes in the first quadrant. Find the equation of the S.L. if it passes through (3, 4).**

**Sol:**  $x = 0, y = 0, A(3, 4)$

intercept form of a S.L.  $\frac{x}{a} + \frac{y}{b} = 1$  .....(1)

Area of the triangle with coordinate axes & eqn (1)  $\Delta = 24$  sq. units.

$$\frac{1}{2}|ab|24 \Rightarrow ab = 48 \Rightarrow b = \frac{48}{a} \quad \text{.....(2)}$$

But equation (1) is passing through A

$$\frac{3}{a} + \frac{4}{b} = 1 \Rightarrow \frac{3}{a} + \frac{4}{\left(\frac{48}{a}\right)} = 1$$

From equation (2)

$$\Rightarrow \frac{3}{a} + \frac{4a}{48} = 1 \Rightarrow \frac{36 + a^2}{12a} = 1$$

$$\Rightarrow 36 + a^2 = 129$$

$$\Rightarrow a^2 - 129 + 36 = 0$$

$$\Rightarrow (a - 6)^2 = 0$$

$$\therefore a = 6.$$

From equation (2)  $b = \frac{48}{6} = 8$

From (1) : Equation of required S.L.

$$\frac{x}{6} + \frac{y}{8} = 1 \Rightarrow \frac{4x + 3y}{24} = 1$$

$$4x + 3y = 24$$

$$4x + 3y - 24 = 0.$$

- 4. If  $3a + 2b + 4c = 0$ , then show that the equation  $ax + by + c = 0$  represents a family of concurrent straight lines and find the point of concurrency**

**Sol:**  $3a + 2b + 4c = 0$

Divide with 4

$$\frac{3a}{4} + \frac{2b}{4} + \frac{4c}{4} = 0$$

$$a\left(\frac{3}{4}\right) + b\left(\frac{1}{2}\right) + c = 0 \quad \dots\dots(1)$$

$\therefore$  Each member of the family of S.Lines given by  $ax + by + c = 0$  passes through the fixed point  $\left(\frac{3}{4}, \frac{1}{2}\right)$ . Hence the set of lines  $ax + by + c = 0$  for parametric values of  $a$ ,  $b$  and  $c$  is

a family of concurrent lines and the point of concurrency is  $\left(\frac{3}{4}, \frac{1}{2}\right)$ .

H.W page No. 55 Example 4

- 5. Find the point on the straight line  $3x + y + 4 = 0$  which is equidistant from the points  $(-5, 6)$  and  $(3, 2)$ .**

**Sol:**  $A(-5, 6)$ ,  $B(3, 2)$ ,  $L \equiv 3x + y + 4 = 0$

Let  $P(a, b)$  be the required print

$$AP = BP$$

$$\sqrt{(a+5)^2 + (b-6)^2} = \sqrt{(a-3)^2 + (b-2)^2}$$

Squaring on B.S.

$$a^2 + 25 + 10a + b^2 + 36 - 12b = a^2 + 9 - 6a + b^2 + 4 - 4b$$

$$10a - 12b + 61 + 6a + 4b - 13 = 0$$

$$16a - 8b + 48 = 0$$

$$\text{Divide with } 2a - b + 6 = 0 \quad \dots\dots(1)$$

But P is on L = 0

$$3a + b + 4 = 0 \quad \dots\dots(2)$$

Now (1) + (2)

$$2a - b + 4 = 0$$

$$\underline{3a + b + 4 = 0}$$

$$5a + 10 = 0 \Rightarrow 5a = -10$$

$$\therefore a = -2$$

$$\text{From (1) } 2(-2) - b + 6 = 0 \Rightarrow -4 + 6 = b$$

$$\therefore b = 2$$

$\therefore$  Required point P = (-2, 2).

6. A straight line through P(3, 4) makes an angle of  $60^\circ$  with the positive direction of the X-axis. Find the coordinates of the points on the line which are 5 units away from P.

Sol: P(3, 4),  $r = 5$ ,  $\theta = 60^\circ$

Parametric form of a point

$$(x, y) = (x_1 \pm r \cos\theta, y_1 \pm r \sin\theta)$$

The point on the line which is at a distance of 5 units from P

$$(x, y) = (3 \pm 5 \cos 60^\circ, 4 \pm 5 \sin 60^\circ)$$

$$= \left[ 3 \pm 5 \left( \frac{1}{2} \right), 4 \pm 5 \left( \frac{\sqrt{3}}{2} \right) \right]$$

$$= \left[ \frac{6 \pm 5}{2}, \frac{8 \pm 5\sqrt{3}}{2} \right]$$

$$= \left( \frac{6+5}{2}, \frac{8+5\sqrt{3}}{2} \right), \left( \frac{6-5}{2}, \frac{8-5\sqrt{3}}{2} \right)$$

$\therefore$  Required points =  $\left( \frac{11}{2}, \frac{8+5\sqrt{3}}{2} \right), \left( \frac{1}{2}, \frac{8-5\sqrt{3}}{2} \right)$



7. A S.L through  $Q(\sqrt{3}, 2)$  makes an angle  $\pi/6$  with the positive direction of the X-axis.  
If the S.L. intersects the line  $\sqrt{3}x - 4y + 8 = 0$  at P, find the distance PQ

**Sol:**  $\sqrt{3}x - 4y + 8 = 0$  .....(1)

$$Q(\sqrt{3}, 2), \quad \theta = \frac{\pi}{6} = \frac{180}{6} = 30 \Rightarrow m = \tan 30^\circ = \frac{1}{\sqrt{3}}$$

Equation of a S.L. with slope  $\frac{1}{\sqrt{3}}$  and passing through  $Q(\sqrt{3}, 2)$

$$y - 2 = \frac{1}{\sqrt{3}}(x - \sqrt{3})$$

$$\sqrt{3}(y - 2) = 1(x - \sqrt{3})$$

$$\sqrt{3}y - 2\sqrt{3} = x - \sqrt{3}$$

$$x - \sqrt{3}y + \sqrt{3} = 0 \quad \text{.....(2)}$$

For P: (1)  $-\sqrt{3}$ (2)

$$\sqrt{3}x - 4y + 8 = 0$$

$$\sqrt{3}x - 3y + 3 = 0$$

$$-y + 5 = 0 \Rightarrow y = 5.$$

from (1)  $\sqrt{3}x - 4(5) + 8 = 0 \Rightarrow \sqrt{3}x = 12 \Rightarrow x = \frac{12}{\sqrt{3}}$

$$x = \frac{4 \times 3}{\sqrt{3}} = \frac{4 \times \sqrt{3} \times \sqrt{3}}{\sqrt{3}} = 4\sqrt{3}$$

$$P = (4\sqrt{3}, 5)$$

$$\therefore PQ = \sqrt{(\sqrt{3} - 4\sqrt{3})^2 + (2 - 5)^2}$$

$$= \sqrt{(-3\sqrt{3})^2 + (-3)^2} = \sqrt{9(3) + 9} = \sqrt{36}$$

$$\therefore PQ = 6.$$

H.W. Page 54 : Example 3

## The Straight Line

### Exercise 3d

**Note :**

1. Let  $\theta$  be the angle between the line  $L_1$  &  $L_2$  then  $\theta = \cos^{-1} \left( \left| \frac{a_1 a_2 + b_1 b_2}{\sqrt{(a_1^2 + b_1^2)(a_2^2 + b_2^2)}} \right| \right)$

2.  $L_1 \perp L_2 \Leftrightarrow a_1 a_2 + b_1 b_2 = 0$  OR  $m_1, m_2 = -1$

3. Perpendicular distance from  $P(x_1, y_1)$  to  $L \equiv ax + by + c = 0$  is  $(d) = \left| \frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}} \right|$

4. Distance between the parallel lines  $L_1 \equiv ax + by + c_1 = 0$  &  $L_2 \equiv ax + by + c_2 = 0$

is  $= \left| \frac{c_1 - c_2}{\sqrt{a^2 + b^2}} \right|$

### Questions

1. Find the value of  $k$ , if the angle between the straight lines  $4x - y + 7 = 0$  and  $kx - 5y - 9 = 0$  is  $45^\circ$

**Sol:**  $L_1 \equiv 4x - y + 7 = 0$ ,  $L_2 \equiv kx - 5y - 9 = 0$ ,  $\theta = 45^\circ$

$$\cos \theta = \frac{a_1 a_2 + b_1 b_2}{\sqrt{(a_1^2 + b_1^2)(a_2^2 + b_2^2)}}$$

$$\cos 45^\circ = \left| \frac{4k + (-)(-5)}{\sqrt{(4^2 + (-1)^2)(k^2 + (-5)^2)}} \right|$$

$$\frac{1}{\sqrt{2}} = \frac{4k + 5}{\sqrt{(16 + 1)(k^2 + 25)}}$$

$$\frac{1}{2} = \frac{16k^2 + 25 + 40k}{17(k^2 + 25)}$$

$$17k^2 + 425 = 32k^2 + 50 + 80k$$

$$15k^2 + 80k - 375 = 0$$

$$3k^2 + 16k - 75 = 0$$

$$3k^2 + 25k - 9k - 75 = 0$$

$$k(3k + 25) - 3(3k + 25) = 0$$

$$(3k + 25)(k - 3) = 0$$

$$3k + 25 = 0 \text{ OR } k - 3 = 0$$

$$3k = -25 \text{ OR } k = 3$$

$$\therefore k = 3 \text{ OR } \frac{-25}{3}$$

2. Find the value of  $k$ , if the straight lines  $y - 3kx + 4 = 0$  and  $(2k - 1)x - (8k - 1)y - 6 = 0$  are perpendicular

Sol:  $L_1 \equiv y - 3kx + 4 = 0 \Rightarrow L_1 \equiv 3kx - y - 4 = 0$  ..... (1)

$L_2 \equiv (2k - 1)x - (8k - 1)y - 6 = 0$  ..... (2)

$L_1 \perp L_2 \Leftrightarrow a_1 a_2 + b_1 b_2 = 0$

$3k(2k - 1) + (-1)[-(8k - 1)] = 0$

$6k^2 - 3k + 8k - 1 = 0$

$6k^2 + 5k - 1 = 0$


$6k^2 + 6k - k - 1 = 0$

$6k(k + 1) - 1(k + 1) = 0$

$(k + 1)(6k - 1) = 0$

$k + 1 = 0 \text{ OR } 6k - 1 = 0$

$k = -1 \text{ OR } k = \frac{1}{6}$



3. Find the length of the perpendicular distance from the point  $(-2, -3)$  to the line  $5x - 2y + 4 = 0$ .

Sol:  $L \equiv 5x - 2y + 4 = 0$ ,  $P(-2, -3)$

$$(d) = \left| \frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}} \right|$$

Distance between parallel line (1) & (2)  $(d) = \left| \frac{5(-2) - 2(-3) + 4}{\sqrt{(5)^2 + (-2)^2}} \right| = \frac{-10 + 6 + 4}{\sqrt{25 + 4}}$

$\therefore d = 0.$

Home Work I<sub>9,10</sub>

4. Find the distance between parallel lines  $3x + 4y - 3 = 0$ ,  $6x + 8y - 1 = 0$

Sol:  $3x + 4y - 3 = 0$

$6x + 8y - 1 = 0$  ..... (2)

$2 \times (1) \Rightarrow 6x + 8y - 6 = 0$  ..... (1)

$$\begin{aligned} \text{distance between parallel lines} &= \left| \frac{-6 - (-1)}{\sqrt{(6)^2 + (8)^2}} \right| \\ &= \frac{-6 + 1}{\sqrt{36 + 64}} = \left| \frac{-5}{\sqrt{100}} \right| = \left| \frac{-5}{10} \right| = \left| \frac{-1}{2} \right| \end{aligned}$$

$$\therefore d = \frac{1}{2}$$

5. If  $Q(h, k)$  is the foot of the perpendicular from  $P(x_1, y_1)$  on the straight line  $ax + by + c = 0$ , then

$$(h - x_1) : a = (k - y_1) : b = -(ax_1 + by_1 + c) : (a^2 + b^2)$$

Sol:  $L \equiv ax + by + c = 0$ ,  $P(x_1, y_1)$ ,  $Q(h, k)$

$$\text{Slope of } L (m_1) = \frac{-a}{b}$$

$$\text{Slope of } PQ (m_2) = \frac{k - y_1}{h - x_1}$$

$$L \perp PQ \Leftrightarrow m_1 m_2 = -1$$

$$\left( \frac{-a}{b} \right) \left( \frac{k - y_1}{h - x_1} \right) = -1$$

$$\frac{k - y_1}{h - x_1} = \frac{b}{a} \Rightarrow \frac{k - y_1}{b} = \frac{h - x_1}{a}$$

$$\frac{h - x_1}{a} = \frac{k - y_1}{b} = \lambda \text{ say}$$

$$\frac{h - x_1}{a} = \lambda \qquad \frac{k - y_1}{b} = \lambda$$

$$h - x_1 = a\lambda \qquad k - y_1 = b\lambda$$

$$h = x_1 + a\lambda \qquad k = y_1 + b\lambda$$


But  $Q$  is on  $L = 0$

$$a(x_1 + a\lambda) + b(y_1 + b\lambda) + c = 0$$

$$ax_1 + a^2\lambda + by_1 + b^2\lambda + c = 0$$

$$(a^2 + b^2)\lambda = -ax_1 - by_1 - c$$

$$\therefore \frac{h - x_1}{a} = \frac{k - y_1}{b} = \frac{-(ax_1 + by_1 + c)}{a^2 + b^2}$$



6. Find the foot of the perpendicular from  $(-1, 3)$  on the straight line  $5x - y - 18 = 0$

Sol: Let  $Q(h, k)$  be the foot of the perpendicular from  $(-1, 3)$  to  $5x - y - 18 = 0$

$$\frac{h - x_1}{a} = \frac{k - y_1}{b} = \frac{-(ax_1 + by_1 - c)}{a^2 + b^2}$$

$$\frac{h - (-1)}{5} = \frac{k - 3}{1} = \frac{-(-5 - 3 - 18)}{(5)^2 + (1)^2}$$

$$\frac{h + 1}{5} = \frac{k - 3}{-1} = \frac{26}{25 + 1}$$

$$\frac{h + 1}{5} = \frac{k - 3}{-1} = \frac{26}{26} = 1$$

$$\frac{h + 1}{5} = 1 \qquad \frac{k - 3}{-1} = 1$$

$$h + 1 = 5 \qquad k - 3 = -1$$

$$h = 5 - 1 \qquad k = -1 + 3$$

$$h = 4 \qquad k = 2$$

$$\therefore (h, k) = (4, 2)$$

Home Work II<sub>5</sub>

7. If  $Q(h, k)$  is the image of the point  $P(x_1, y_1)$  w.r.t. the straight line  $ax + by + c = 0$  then  $(h - x_1) : a = (k - y_1) : b = -2(ax_1 + by_1 + c) : (a^2 + b^2)$

Sol:  $L \equiv ax + by + c = 0$   $P(x_1, y_1)$   $Q(h, k)$

$$\text{Slope of } L = m_1 = \frac{-a}{b}$$

$$\text{Slope of } PQ (m_2) = \frac{k - y_1}{h - x_1}$$


$$PQ \perp L \Leftrightarrow m_1 m_2 = -1$$

$$\left(\frac{-a}{b}\right) \left(\frac{k - y_1}{h - x_1}\right) = -1$$

$$\frac{k - y_1}{h - x_1} = \frac{b}{a} \Rightarrow \frac{k - y_1}{b} = \frac{h - x_1}{a}$$

$$\frac{h - x_1}{a} = \frac{k - y_1}{b} = \lambda \text{ let } \dots\dots\dots(1)$$

$$\frac{h - x_1}{a} = \lambda \qquad \frac{k - y_1}{b} = \lambda$$



$$\begin{aligned} h - x_1 &= a\lambda & k - y_1 &= b\lambda \\ h &= x_1 + a\lambda & k &= y_1 + b\lambda \end{aligned} \quad \dots\dots(2)$$

Since M is the midpoint of P & Q

$$M = \left( \frac{x_1 + h}{2}, \frac{y_1 + k}{2} \right)$$

It is on L = 0

$$a \left( \frac{x_1 + h}{2} \right) + b \left( \frac{y_1 + k}{2} \right) + c = 0$$

$$\frac{ax_1 + ah + by_1 + bk + 2c}{2} = 0$$

$$ax_1 + a(x_1 + a\lambda) + by_1 + b(y_1 + b\lambda) + 2c = 0$$

$$ax_1 + ax_1 + a^2\lambda + by_1 + by_1 + b^2\lambda + 2c = 0$$

$$\lambda(a^2 + b^2) = -2ax_1 - 2by_1 - 2c$$

$$\therefore \lambda = \frac{-2(ax_1 + by_1 + c)}{a^2 + b^2}$$

(1) ಸಂಘಟಿಸಿ

$$\frac{h - x_1}{a} = \frac{k - y_1}{b} = \frac{-2(ax_1 + by_1 + c)}{a^2 + b^2}$$

**8. Find the image of (1, -2) w.r.t. the straight line  $2x - 3y + 5 = 0$ .**

**Sol:** (h, k) is the image of (1, -2) w.r.t the line  $2x - 3y + 5 = 0$

$$\frac{h-1}{2} = \frac{k+2}{-3} = \frac{-2(2(1) - 3(-2) + 5)}{(2)^2 + (-3)^2}$$

$$\frac{h-1}{2} = \frac{k+2}{-3} = \frac{-2(2+6+5)}{4+9} = \frac{-2(13)}{13} = -2$$

$$\frac{h-1}{2} = -2 \qquad \frac{k+2}{-3} = -2$$

$$h-1 = -4 \qquad k+2 = 6$$

$$h = -3 \qquad k = 4$$

$$\therefore (h, k) = (-3, 4)$$

Home Work II<sub>8</sub>

9. Find the equations of the straight lines passing through the point  $(-3, 2)$  and making an angle of  $45^\circ$  with the straight line  $3x - y + 4 = 0$

Sol:  $P(-3, 2)$ ,  $L \equiv 3x - y + 4 = 0$ ,  $\theta = 45^\circ$

Let  $m$  be the slope of a line which is passing through P

$$y - 2 = m(x + 3) \quad \dots(*)$$

$$y - 2 = mx + 3m$$

$$mx - y + (2 + 3m) = 0 \quad \dots(1)$$

According to the problem,  $45^\circ$  is the angle between

$L = 0$  & (1)

$$\cos 45^\circ = \frac{3(m) + (-1)(-1)}{\sqrt{(3)^2 + (-1)^2(m^2 + (-1)^2)}} \Rightarrow \frac{1}{\sqrt{2}} = \frac{3m + 1}{\sqrt{(9 + 1)(m^2 + 1)}}$$

Squaring on B.S.

$$\frac{1}{2} = \frac{9m^2 + 1 + 6m}{10(m^2 + 1)}$$

$$10(m^2 + 1) = 2(9m^2 + 1 + 6m)$$

$$10m^2 + 10 = 18m^2 + 2 + 12m$$

$$8m^2 + 10m - 8 = 0$$

$$2m^2 + 3m - 2 = 0$$

$$2m^2 + 4m - 1m - 2 = 0$$

$$2m(m + 2) - 1(m + 2) = 0$$

$$(m + 2)((2m - 1) = 0$$

$$m + 2 = 0 \quad \text{OR} \quad 2m - 1 = 0$$

$$m = -2 \quad \text{OR} \quad m = \frac{1}{2}$$

Case (i):

$m = -2$  then, from equation (\*)

$$y - 2 = -2(x + 3)$$



$$y - 2 = -2x - 6$$

$$2x + y + 4 = 0$$

Case (ii):

$m = \frac{1}{2}$  then, from equation (\*)

$$y - 2 = \frac{1}{2}(x + 3)$$



$$2y - 4 = x + 3$$

$$x - 2y + 7 = 0$$

∴ Equation of required s. lines

$$2x + 4y + 4 = 0,$$

$$x - 2y + 7 = 0$$

**10. Find the angles of the triangle whose sides are  $x + y - 4 = 0$ ,  $2x + y - 6 = 0$  and  $5x + 3y - 15 = 0$ .**

**Sol:**  $L_1 \equiv x + y - 4 = 0$

$$L_2 \equiv 2x + y - 6 = 0$$

$$L_3 \equiv 5x + 3y - 15 = 0$$

$$\cos A = \frac{1(2) + 1(1)}{\sqrt{(1^2 + 1^2)(2^2 + 1^2)}} = \frac{2 + 1}{\sqrt{(1+1)(4+1)}} = \frac{3}{\sqrt{10}}$$

$$A = \cos^{-1}\left(\frac{3}{\sqrt{10}}\right)$$

$$\cos B = \frac{2(5) + 1(3)}{\sqrt{(2^2 + 1^2)(5^2 + 3^2)}} = \frac{10 + 3}{\sqrt{(4+1)(25+9)}} = \frac{13}{\sqrt{170}}$$

$$B = \cos^{-1}\left(\frac{13}{\sqrt{170}}\right)$$

$$\cos C = \frac{5(1) + 3(1)}{\sqrt{(5^2 + 3^2)(1^2 + 1^2)}} = \frac{5 + 3}{\sqrt{(25+9)(1+1)}} = \frac{8}{\sqrt{68}} = \frac{8}{2\sqrt{17}} = \frac{4}{\sqrt{17}}$$

$$C = \cos^{-1}\left(\frac{4}{\sqrt{17}}\right)$$

**11. Find the equations of the straight lines passing through the point of intersection of the lines  $3x + 2y + 4 = 0$ ,  $2x + y = 1$  and whose distance from  $(2, -1)$  is 2**

**Sol:**  $L_1 \equiv 3x + 2y + 4 = 0$ ,  $L_2 \equiv 2x + y = 1$ ,  $A = (2, -1)$ ,  $d = 2$

For the point of intersection of  $L_1$  &  $L_2$

$$2L_1 - 3L_2$$

$$6x + 4y + 8 = 0$$

$$6x + 15y - 3 = 0$$

$$-11y + 11 = 0 \Rightarrow 11y = 11 \Rightarrow y = 1$$

From  $L_1$   $3x + 2(1) + 4 = 0 \Rightarrow 3x = -6 \Rightarrow x = -2$

$P(2, -1)$



Let  $m$  be the slope of a line which is passing through P

$$y - 1 = m(x + 2) \quad \dots(*)$$

$$y - 1 = mx + 2m$$

$$mx - y + (1+2m) = 0 \quad \dots(1)$$

Given that perpendicular distance from

A to equation (1) = 2

$$\left| \frac{m(2) - (-1) + 1 + 2m}{\sqrt{m^2 + (-1)^2}} \right| = 2$$

$$\frac{2m + 1 + 1 + 2m}{\sqrt{m^2 + 1}} = 2 \Rightarrow \frac{4m + 2}{\sqrt{1 + m^2}} = 2$$

$$2(2m + 1) = 2\sqrt{1 + m^2} \Rightarrow 2m + 1 = \sqrt{1 + m^2}$$

Squaring on both sides

$$4m^2 + 1 + 4m = 1 + m^2$$

$$3m^2 + 4m = 0$$

$$m(3m + 4) = 0$$

$$m = 0 \text{ OR } m = \frac{-4}{3}$$

Case (i):  $m = 0$  then, from (\*)

$$y - 1 = 0(x + 3)$$

$$y = 1$$

Case (ii):  $m = \frac{-4}{3}$  then, from (\*)

$$y - 1 = -\frac{4}{3}(x + 2)$$

$$3y - 3 = -4x - 8$$

$$4x + 3y + 5 = 0$$

$\therefore$  Equation of required straight lines  $y = 1$ ,  $4x + 3y + 5 = 0$ .

**12. Find the equation of the line perpendicular to the line  $3x + 4y + 6 = 0$  and making an intercept -4 on the X-axis.**

**Sol:**  $L_1 \equiv 3x + 4y + 6 = 0$ , X-intercept = -4

Equation of any line which is perpendicular to  $L = 0$

$$4x - 3y + k = 0 \quad \dots(*)$$

$$4x - 3y - k = 0$$

$$\frac{4x}{(-k)} - \frac{3y}{(-k)} = \frac{-k}{(-k)}$$

$$\frac{x}{\left(\frac{-k}{4}\right)} - \frac{y}{\left(\frac{-k}{3}\right)} = 1 \quad \dots(1)$$

$$\text{X-intercept of (1)} = \frac{-k}{4}$$

$$\text{According to the problem } \frac{-k}{4} = -4 \Rightarrow k = 16$$

$$\therefore \text{Equation of required straight line } 4x - 3y + 16 = 0$$

### The Straight Line

#### Exercise 3e

#### Note :

1. The medians of a triangle are concurrent. The concurrent point of medians is the centroid.
2. The altitudes of a triangle are concurrent. The concurrent point of the altitudes is ortho-center.
3. The internal bisectors of the angles of a triangle are concurrent. The concurrent point is called incenter.
4. The perpendicular bisectors of the sides of a triangle are concurrent. The point of concurrency is the 'circumcenter' of the triangle.

#### Questions

1. Find the equation of the straight line passing through the origin and also through the point of intersection of the lines  $2x - y + 5 = 0$  and  $x + y + 1 = 0$ .

$$\text{Sol: } O(0, 0), L_1 \equiv 2x - y + 5 = 0, L_2 \equiv x + y + 1 = 0$$

$$\text{From } L_1 + L_2 \quad 2x - y + 5 = 0$$

$$\underline{x + y + 1 = 0}$$

$$3x + 6 = 0 \Rightarrow 3x = -6 \Rightarrow x = -2$$

$$\text{From } L_1: 2(-2) - y + 5 = 0 \Rightarrow 1 - y = 0 \Rightarrow y = 1$$

intersecting point of  $L_1$  &  $L_2$   $A(-2, 1)$

Equation of required straight line passing through  $O$  &  $A$

$$y - 0 = \left( \frac{1 - 0}{-2 - 0} \right) (x - 0)$$

$$-2(y) = 1(x)$$

$$x + 2y = 0$$

2. Find the equation of the straight line parallel to the line  $3x + 4y = 7$  and passing through the point of intersection of the lines  $x - 2y - 3 = 0$  and  $x + 3y - 6 = 0$

Sol:  $L_1 \equiv 3x + 4y = 7$ ,  $L_2 \equiv x - 2y - 3 = 0$ ,  $L_3 \equiv x + 3y - 6 = 0$

To find the point intersection of  $L_2, L_3$

$$\begin{array}{r} x - 2y - 3 = 0 \\ x + 3y - 6 = 0 \\ \hline - \quad - \quad + \\ -5y + 3 = 0 \\ \Rightarrow 5y = 3 \Rightarrow y = \frac{3}{5} \end{array}$$

From  $L_2$   $x - 2\left(\frac{3}{5}\right) - 3 = 0$

$$x = \frac{6}{5} + 3 = \frac{6+15}{5} = \frac{21}{5}$$

Equation of any line parallel to  $A = \left(\frac{21}{5}, \frac{3}{5}\right)$

$L_1 \equiv 3x + 4y = 7$  Equation of any line parallel to  $L_1 = 0$  is  $3x + 4y + k = 0$

But it is passing through A

$$3\left(\frac{21}{5}\right) + 4\left(\frac{3}{5}\right) + k = 0$$

$$k + \frac{63+12}{5} = 0 \Rightarrow k = \frac{-75}{5} = -15$$

$\therefore$  Equation of required straight line  $3x + 4y - 15 = 0$

3. Find the equation of the straight line making non - zero equal intercepts on the coordinate axes and passing through the point of intersection of the lines  $2x - 5y + 1 = 0$  and  $x - 3y - 4 = 0$

Sol:  $L_1 \equiv 2x - 5y + 1 = 0$ ,  $L_2 \equiv x - 3y - 4 = 0$

To find the point intersection of  $L_1, L_2$

$$\begin{array}{r} 2x - 5y + 1 = 0 \\ 2x - 6y - 8 = 0 \\ \hline - \quad + \quad + \\ y + 9 = 0 \Rightarrow y = -9 \end{array}$$

From  $L_1$   $2x - 5(-9) + 1 = 0$

$$2x - 45 + 1 = 0$$

$$2x = -46$$

$$x = -23$$

$A = (-23, -9)$

Since equal intercepts on the coordinate axes

$$\Rightarrow a = b$$

Equation of S.L. in intercept form

$$\frac{x}{a} + \frac{y}{b} = 1 \Rightarrow \frac{x}{a} + \frac{y}{a} = 1 \quad (\text{as } a = b)$$

But it is passing through A

$$-23 - 9 = a$$

$$-32 = a$$

Equation of a required straight line  $x + y = -32 \Rightarrow x + y + 32 = 0$ .

- 4. Find the length of the perpendicular drawn from the point of intersection of the lines  $3x + 2y + 4 = 0$  and  $2x + 5y - 1 = 0$  to the straight line  $7x + 24y - 15 = 0$**

**Sol:**  $L_1 \equiv 3x + 2y + 4 = 0$ ,  $L_2 \equiv 2x + 5y - 1 = 0$ ,  $L_3 \equiv 7x + 24y - 15 = 0$

Point of intersection of  $L_1$  &  $L_2$

$$\frac{x}{2(-) - 5(4)} = \frac{y}{4(2) - (-1)(3)} = \frac{1}{3(5) - 2(2)}$$

$$\frac{x}{-2 - 20} = \frac{y}{8 + 3} = \frac{1}{15 - 4}$$

$$\frac{x}{-22} = \frac{y}{11} = \frac{1}{11}$$

$$A = (-2, 1)$$

Perpendicular distance from A to  $L_3 = 0$

$$d = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$$

$$= \frac{|7(-2) + 24(1) - 15|}{\sqrt{(7)^2 + (24)^2}} = \frac{|-14 + 24 - 15|}{\sqrt{49 + 576}}$$

$$= \frac{|-5|}{\sqrt{625}} = \frac{5}{25}$$

$$\therefore d = \frac{1}{5}$$

- 5. The base of an equilateral triangle is  $x + y - 2 = 0$  and the opposite vertex is  $(2, -1)$ . Find the equations of the remaining sides.**

**Sol:**  $L \equiv x + y - 2 = 0$ ,  $A(2, -1)$

Let 'm' be the slope of a line which is passing through A.

Equation of a line with slope  $m$  and passing through A is

equation of the line passing through A and intersecting  $L = 0$  at B

$$y + 1 = m(x - 2) \quad \dots(*)$$

$$y + 1 = mx - 2m$$


$$mx - y - (1 + 2m) = 0 \quad \dots(1)$$

Since  $\Delta ABC$  is equilateral triangle

$$\Rightarrow \angle A = \angle B = \angle C$$

Angle between  $L$   $L = 0$ , (1)

$$\cos 60^\circ = \frac{1(m) + 1(-1)}{\sqrt{((1)^2 + (1)^2)(m^2 + (-1)^2)}}$$



Squaring on both sides

$$\frac{1}{2} = \frac{m - 1}{\sqrt{(1+1)(1+m^2)}}$$

$$\frac{x}{-2 - 20} = \frac{y}{8 + 3} = \frac{1}{15 - 4}$$

$$\frac{1}{4} = \frac{m^2 + 1 - 2m}{2(1 + m^2)}$$

$$(1 + m^2) = 2m^2 + 2 - 4m$$

$$2m^2 + 2 - 4m - 1 - m^2 = 0$$

$$m^2 - 4m + 1 = 0$$

$$m = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(1)}}{2(1)}$$

$$= \frac{4 \pm \sqrt{16 - 4}}{2} = \frac{4 \pm \sqrt{12}}{2}$$

$$= \frac{4 \pm 2\sqrt{3}}{2} = \frac{2(2 \pm \sqrt{3})}{2}$$

$$m = 2 \pm \sqrt{3}$$

From (\*) : Equation of required S.L.  $y + 1 = (2 \pm \sqrt{3})(x - 2)$ .

Home work II<sub>2</sub>

6. Find the incenter of the triangle whose vertices are  $(1, \sqrt{3})$ ,  $(2, 0)$  and  $(0, 0)$

Sol:  $A(0, 0)$ ,  $B(2, 0)$ ,  $C(1, \sqrt{3})$

$$c = AB = \sqrt{(2 - 0)^2 + (0 - 0)^2} = \sqrt{4} = 2$$

$$b = AC = \sqrt{(1-0)^2 + (\sqrt{3}-0)^2} = \sqrt{1+3} = 2$$

$$a = BC = \sqrt{(1-2)^2 + (\sqrt{3}-0)^2} = \sqrt{1+3} = 2$$


$$\text{Incentre } I = \left( \frac{ax_1 + bx_2 + cx_3}{a+b+c}, \frac{ay_1 + by_2 + cy_3}{a+b+c} \right)$$

$$I = \left( \frac{2(0) + 2(2) + 2(1)}{2+2+2}, \frac{2(0) + 2(0) + 2(\sqrt{3})}{2+2+2} \right)$$

$$= \left( \frac{0+4+2}{6}, \frac{0+0+2\sqrt{3}}{6} \right)$$

$$= \left( \frac{6}{6}, \frac{2\sqrt{3}}{6} \right) = \left( 1, \frac{\sqrt{3}}{3} \right)$$

$$\therefore I = \left( 1, \frac{1}{\sqrt{3}} \right)$$



7. Find the orthocenter of the triangle whose vertices are  $(-5, -7)$ ,  $(13, 2)$  and  $(-5, 6)$

Sol:  $A(-5, -7)$ ,  $B(13, 2)$ ,  $C(-5, 6)$

$$\text{Slope of BC} = \frac{6-2}{-5-13} = \frac{4}{-18} = \frac{-2}{9}$$

Let AD be the perpendicular drawn from A to BC

Altitude drawn from CF to AB

Let BE = be the perpendicular drawn from B To  $\overline{AC}$

$$AD \perp BC$$

$$\text{Slope of AD} = \frac{-1}{m} = \frac{-1}{\left(\frac{-2}{9}\right)} = \frac{9}{2}$$

Equation of a line with Slope  $\frac{9}{2}$  and passing through A is

$$y + 7 = \frac{9}{2}(x + 5)$$


$$2y + 14 = 9x + 45$$

$$9x - 2y + 31 = 0 \quad \dots(1)$$

$$\text{Slope of AC} = \frac{6+7}{-5-(-5)} = \frac{13}{0} = \infty$$

$BE \perp AC \Rightarrow \text{Slope of BE} = 0$

$$\text{Equation of BE} \quad y - 2 = 0(x - 13)$$



$$\begin{aligned} y - 2 &= 0 \\ y &= 2 \quad \dots(2) \end{aligned}$$

Intersecting point of (1), (2) is our required orthocenter

$$\begin{aligned} \text{From (1) \& (2)} \quad 9x - 2(2) + 31 &= 0 \\ 9x + 27 = 0 \Rightarrow 9x &= -27 \Rightarrow 9x = -3. \\ \therefore O &= (-3, 2) \end{aligned}$$

**8. If the equations of the sides of a triangle are  $7x + y - 10 = 0$ ,  $x - 2y + 5 = 0$  and  $x + y + 2 = 0$ , find the orthocenter of the triangle.**

**Sol:**  $L_1 \equiv 7x + y - 10 = 0$ ,  $L_2 \equiv x - 2y + 5 = 0$ ,  $L_3 \equiv x + y + 2 = 0$

For A  $2L_1 + L_2$

$$\begin{aligned} 14x + 2y - 20 &= 0 \\ \underline{x - 2y + 5} &= 0 \\ 15x - 15 &= 0 \Rightarrow x = 1. \end{aligned}$$

From  $L_1$   $7(1) + y - 10 = 0 \Rightarrow y - 3 = 0 \Rightarrow y = 3 \Rightarrow A = (1, 3)$

For B  $L_1 - L_3$

$$\begin{aligned} 7x + y - 10 &= 0 \\ \underline{x + y + 2} &= 0 \\ - - - & \\ 6x - 12 &= 0 \Rightarrow x = 2. \end{aligned}$$

from  $L_1$   $7(2) + y - 10 = 0 \Rightarrow y + 4 = 0 \Rightarrow y = -3 \Rightarrow B = (2, -4)$

For C  $L_2 - L_3$

$$\begin{aligned} x - 2y + 5 &= 0 \\ \underline{x + y + 2} &= 0 \\ -3y + 3 &= 0 \Rightarrow y = 1. \end{aligned}$$

$L_2$  from  $x - 2(1) + 5 = 0 \Rightarrow x + 3 = 0 \Rightarrow x = -3$   
 $\Rightarrow C = (-3, 1)$

Slope of BC =  $\frac{-1}{(1)} = -1$

$AD \perp BC \Rightarrow$  Slope of AD =  $\frac{-1}{(-1)} = 1$

Equation of AD  $y - 3 = 1(x - 1)$   
 $y - 3 = (x - 1)$   
 $x - y + 2 = 0 \quad \dots(1)$

Slope of AC =  $\frac{-1}{(-2)} = \frac{1}{2}$

$$BE \perp AC \Rightarrow \text{Slope of BE} = \frac{-1}{(1/2)} = -2$$

$$\begin{aligned} \text{Equation of BE} \quad y + 4 &= -2(x - 2) \\ y + 4 &= -2x - 4 \\ 2x + y &= 0 \quad \dots(2) \end{aligned}$$

Intersecting point of (1) & (2) is our required orthocenter (1) + (2)

$$\begin{aligned} x - y + 2 &= 0 \\ \underline{2x + y} &= 0 \\ 3x + 2 &= 0 \Rightarrow x = -2/3 \end{aligned}$$

$$\text{From (1)} \quad \frac{-2}{3} - y + 2 = 0 \Rightarrow \frac{-2+6}{3} = y \Rightarrow y = \frac{4}{3}$$

$$\therefore \text{required orthocentre } O = \left( \frac{-2}{3}, \frac{4}{3} \right)$$

Home Work I<sub>3</sub>, II<sub>4</sub> & III<sub>1</sub>

**9. Find the circumcenter of the triangle whose vertices are (-2, 3), (2, -1) and (4, 0)**

**Sol:** A(-2, 3), B(2, -1), C(4, 0)

Let S = (a, b) be the circumcenter of the  $\Delta ABC$

We know that SA = SB = SC

From : SA = SB

$$\begin{aligned} \sqrt{(-2-a)^2 + (3-b)^2} &= \sqrt{(2-a)^2 + (-1-b)^2} \\ 4 + a^2 + 4a + 9 + b^2 - 6b &= 4 + a^2 - 4a + 1 + b^2 + 2b \\ 4a - 6b + 13 + 4a - 2b - 5 &= 0 \\ 8a - 8b + 8 &= 0 \end{aligned}$$

$$\text{Divide with 8} \quad a - b + 1 = 0 \quad \dots\dots(1)$$

SA = SB

$$\sqrt{(-2-a)^2 + (3-b)^2} = \sqrt{(4-a)^2 + (0-b)^2}$$

Squaring on B.S

$$\begin{aligned} 4 + a^2 + 4a + 9 + b^2 - 6b &= 16 + a^2 - 8a + b^2 \\ 4a - 6b + 13 + 8a - 16 &= 0 \\ 12a - 6b - 3 &= 0 \\ 4a - 2b - 1 &= 0 \quad \dots\dots(2) \end{aligned}$$

Intersecting point of (1) & (2) is our required circumcenter

$$\begin{aligned} 4(1) - (2) \quad \quad \quad 4a - 4b + 4 &= 0 \\ \underline{4a - 2b - 1} &= 0 \\ - \quad + \quad + & \end{aligned}$$



$$-2b + 5 = 0 \Rightarrow b = 5/2$$

$$\text{From (1) } a - \frac{5}{2} + 1 = 0 \Rightarrow a = \frac{5}{2} - 1 \Rightarrow a = \frac{5-2}{2} = \frac{3}{2}$$

$$\therefore \text{ required circumcentre } S = \left( \frac{3}{2}, \frac{5}{2} \right)$$

Home Work (1) page 73 Example (2) 4 II<sub>iii</sub> (3) III<sub>2</sub> (4) I<sub>13</sub>

**10. If  $p$  and  $q$  are the lengths of the perpendiculars from the origin to the straight lines  $x \sec \alpha + y \operatorname{cosec} \alpha = a$  and  $x \cos \alpha - y \sin \alpha = a \cos 2\alpha$  Prove that  $4p^2 + q^2 = a^2$ .**

**Sol:**  $L_1 \equiv x \sec \alpha + y \operatorname{cosec} \alpha = a$ ,  $L_2 \equiv x \cos \alpha - y \sin \alpha - a \cos 2\alpha = 0$

$p$  = perpendicular distance from  $(0, 0)$  to  $L_1 = 0$

$$\begin{aligned} p &= \frac{|(0) \sec \alpha + (0) \operatorname{cosec} \alpha - a|}{\sqrt{(\sec \alpha)^2 + (\operatorname{cosec} \alpha)^2}} = \frac{|-a|}{\sqrt{\frac{1}{\cos^2 \alpha} + \frac{1}{\sin^2 \alpha}}} \\ &= \frac{a}{\sqrt{\frac{\sin^2 \alpha + \cos^2 \alpha}{\cos^2 \alpha \sin^2 \alpha}}} = \frac{a}{\sqrt{\frac{1}{(\sin \alpha \cos \alpha)^2}}} = \frac{a}{\left( \frac{1}{\sin \alpha \cos \alpha} \right)} \end{aligned}$$

$$p = a \sin \alpha \cos \alpha$$

Now  $q$  = perpendicular distance from  $(0, 0)$  to  $L_2 = 0$

$$\begin{aligned} q &= \frac{|(0) \cos \alpha - (0) \sin \alpha - a \cos 2\alpha|}{\sqrt{(\cos \alpha)^2 + (-\sin \alpha)^2}} = \frac{|0 - 0 - a \cos 2\alpha|}{\sqrt{\cos^2 \alpha + \sin^2 \alpha}} \\ &= \frac{|-a \cos 2\alpha|}{\sqrt{1}} \end{aligned}$$

$$\therefore q = a \cos 2\alpha$$

$$\begin{aligned} \text{Now } 4p^2 + q^2 &= 4(a \sin \alpha \cos \alpha)^2 + (a \cos 2\alpha)^2 \\ &= a^2 (2 \sin \alpha \cos \alpha)^2 + a^2 \cos^2 2\alpha \\ &= a^2 [\sin^2 2\alpha + \cos^2 2\alpha] \\ &= a^2 (1) \quad [\because \sin^2 \theta + \cos^2 \theta = 1] \\ \therefore 4p^2 + q^2 &= a^2 \end{aligned}$$

## Pair of Straight Lines

### Key Concepts

- If  $a, b, h$  are real numbers, not all zero, then  $H \equiv ax^2 + 2hxy + by^2 = 0$  is called a homogeneous equation of second degree in  $x$  and  $y$   
 $S \equiv ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  is called a general equation of second degree in  $x$  and  $y$ .
- If  $a, b, h$  are not all zero, then the equation  $H \equiv ax^2 + 2hxy + by^2 = 0$  represents a pair of straight lines  $\Leftrightarrow h^2 \geq ab$ .
- Let the equation  $ax^2 + 2hxy + by^2 = 0$  represent a pair of straight lines. Then the angle  $\theta$  between the lines is given by

$$\cos \theta = \frac{|a + b|}{\sqrt{(a - b)^2 + 4h^2}}$$

- (i) If  $h^2 = ab$  then the lines are coincide  
 (ii) If  $a + b = 0$  then the lines are perpendicular
- If the second degree equation  $S \equiv ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  in the two variables  $x$  and  $y$  represents a pair of straight lines
- $\Leftrightarrow$  (i)  $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$   
 (ii)  $h^2 \geq ab, g^2 \geq ac, f^2 \geq bc$

### Long Answer Questions (7 Marks)

1. If  $\theta$  is an angle between the lines represented by  $ax^2 + 2hxy + by^2 = 0$  then prove that

$$\cos \theta = \frac{|a + b|}{\sqrt{(a - b)^2 + 4h^2}}$$

**Sol:** Let  $ax^2 + 2hxy + by^2 = 0$  represents the lines

$$l_1x + m_1y = 0 \quad \dots(1)$$

$$l_2x + m_2y = 0 \quad \dots(2)$$

$$\therefore ax^2 + 2hxy + by^2 = (l_1x + m_1y)(l_2x + m_2y)$$

$$\Rightarrow ax^2 + 2hxy + by^2 = l_1l_2x^2 + (l_1m_2 + l_2m_1)xy + m_1m_2y^2$$

by comparing like term coefficients on both sides

$$l_1l_2 = a, l_1m_2 + l_2m_1 = 2h, m_1m_2 = b$$

If  $\theta$  is an angle between (1) and (2) then

$$\begin{aligned} \cos\theta &= \frac{|l_1l_2 + m_1m_2|}{\sqrt{l_1^2 + m_1^2} \sqrt{l_2^2 + m_2^2}} \\ &= \frac{|l_1l_2 + m_1m_2|}{\sqrt{(l_1^2 + m_1^2)(l_2^2 + m_2^2)}} \\ &= \frac{|l_1l_2 + m_1m_2|}{\sqrt{l_1^2l_2^2 + l_1^2m_2^2 + l_2^2m_1^2 + m_1^2m_2^2}} \\ &= \frac{|l_1l_2 + m_1m_2|}{\sqrt{(l_1l_2 - m_1m_2)^2 + 2l_1l_2m_1m_2 + (l_1m_2 + l_2m_1)^2 - 2l_1m_2m_2l_1}} \\ &= \frac{|a + b|}{\sqrt{(a - b)^2 + (2h)^2}} \quad [\because l_1l_2 = a, m_1m_2 = b, l_1m_2 + l_2m_1 = 2h] \\ &= \frac{|a + b|}{\sqrt{(a - b)^2 + 4h^2}} \end{aligned}$$

2. Prove that the product of perpendiculars from  $(\alpha, \beta)$  to the pair of straight lines

$$ax^2 + 2hxy + by^2 = 0 \text{ is}$$

$$\frac{|a\alpha^2 + 2h\alpha\beta + b\beta^2|}{\sqrt{(a - b)^2 + 4h^2}}$$

**Sol:**  $ax^2 + 2hxy + by^2 = 0$  Let the pair of straight lines represents the lines

$$l_1x + m_1y = 0 \quad \dots(1)$$

$$l_2x + m_2y = 0 \quad \dots(2)$$

$$\therefore ax^2 + 2hxy + by^2 = (l_1x + m_1y)(l_2x + m_2y)$$

$$= l_1l_2x^2 + (l_1m_2 + l_2m_1)xy + m_1m_2y^2$$

by comparing like term coefficients on both sides

$$l_1l_2 = a, l_1m_2 + l_2m_1 = 2h, m_1m_2 = b$$

Let  $d_1$  = perpendicular distance from a point  $(\alpha, \beta)$  to  $l_1x + m_1y = 0$

$$d_1 = \frac{|l_1\alpha + m_1\beta|}{\sqrt{l_1^2 + m_1^2}}$$

similarly

Let  $d_2$  = perpendicular distance from a point  $(\alpha, \beta)$  to  $l_2x + m_2y = 0$

$$d_2 = \frac{|l_2\alpha + m_2\beta|}{\sqrt{l_2^2 + m_2^2}}$$

$\therefore$  Product of perpendicular distances =  $d_1 \times d_2$

$$\begin{aligned} &= \frac{|l_1\alpha + m_1\beta|}{\sqrt{l_1^2 + m_1^2}} \times \frac{|l_2\alpha + m_2\beta|}{\sqrt{l_2^2 + m_2^2}} \\ &= \frac{|(l_1\alpha + m_1\beta)(l_2\alpha + m_2\beta)|}{\sqrt{(l_1^2 + m_1^2)(l_2^2 + m_2^2)}} \\ &= \frac{|l_1l_2\alpha^2 + (l_1m_1 + l_2m_1)\alpha\beta + m_1m_2\beta^2|}{\sqrt{l_1^2l_2^2 + l_1^2m_2^2 + l_2^2m_1^2 + m_1^2m_2^2}} \\ &= \frac{|l_1l_2\alpha^2 + (l_1m_1 + l_2m_1)\alpha\beta + m_1m_2\beta^2|}{\sqrt{(l_1l_2 - m_1m_2)^2 + 2l_1l_2m_1m_2 + (l_1m_2 + l_2m_1)^2 - 2l_1l_2m_1m_2}} \\ &= \frac{|l_1l_2\alpha^2 + (l_1m_1 + l_2m_1)\alpha\beta + m_1m_2\beta^2|}{\sqrt{(l_1l_2 - m_1m_2)^2 + (l_1m_2 + l_2m_1)^2}} \\ &= \frac{|a\alpha^2 + 2h\alpha\beta + b\beta^2|}{\sqrt{(a-b)^2 + (2h)^2}} \quad [\because l_1l_2 = a, m_1m_2 = b, l_1m_2 + l_2m_1 = 2h] \\ &= \frac{|a\alpha^2 + 2h\alpha\beta + b\beta^2|}{\sqrt{(a-b)^2 + 4h^2}} \end{aligned}$$

3. Show that the area of a triangle formed by the lines  $ax^2 + 2hxy + by^2 = 0$  and

$$lx + my + n = 0 \text{ is } \left| \frac{n^2 \sqrt{h^2 - ab}}{am^2 - 2hlm + bt^2} \right|$$

**Sol:** Let the equation  $ax^2 + 2hxy + by^2 = 0$  represents the lines

$$l_1x + m_1y = 0 \quad \dots(1)$$

$$l_2x + m_2y = 0 \quad \dots(2)$$

$$\begin{aligned} \therefore ax^2 + 2hxy + by^2 &= (l_1x + m_1y)(l_2x + m_2y) \\ &= l_1l_2x^2 + (l_1m_2 + l_2m_1)xy + m_1m_2y^2 \end{aligned}$$

Let the given line be  $lx + my + n = 0 \dots(3)$

$$l_1l_2 = a, l_1m_2 + l_2m_1 = 2h, m_1m_2 = b$$

Let A be the intersection point of (1) and (3)

$$\begin{aligned} \frac{x}{m_1n - o} &= \frac{y}{o - nl_1} = \frac{1}{l_1m - lm_1} \\ \Rightarrow x &= \frac{m_1n}{l_1m - lm_1}, y = \frac{-nl_1}{l_1m - lm_1} \end{aligned}$$

$$\therefore A = \left( \frac{m_1n}{l_1m - lm_1}, \frac{-nl_1}{l_1m - lm_1} \right)$$

Similarly, Let B be the intersection point of (2) and (3)

$$\begin{aligned} \frac{x}{m_2n - o} &= \frac{y}{o - nl_2} = \frac{1}{l_2m - lm_2} \\ \therefore B &= \left( \frac{m_2n}{l_2m - lm_2}, \frac{-nl_2}{l_2m - lm_2} \right) \end{aligned}$$

clearly  $O(0,0)$  be the point of intersection of (1) and (2)

[Area of a triangle whose vertices  $(0, 0), (x_1, y_1), (x_2, y_2)$  is  $= \frac{1}{2}|x_1y_2 - x_2y_1|$ ]

$$\begin{aligned} \therefore \text{Area of } \Delta OAB &= \frac{1}{2} \left| \frac{m_1n}{l_1m - lm_1} \times \frac{(-nl_2)}{l_2m - lm_2} - \frac{m_2n}{l_2m - lm_2} \times \frac{(-nl_1)}{l_1m - lm_1} \right| \\ &= \frac{1}{2} \left| \frac{l_1m_2n^2 - m_1l_2n^2}{(l_1m - lm_1)(l_2m - lm_2)} \right| \\ &= \frac{1}{2} \left| \frac{n^2(l_1m_2 - m_1l_2)}{l_1l_2m^2 - (l_1m_2 + l_2m_1)lm + m_1m_2l^2} \right| \\ &= \frac{1}{2} \left| \frac{n^2\sqrt{(l_1m_2 - l_2m_1)^2}}{l_1l_2m^2 - (l_1m_2 + l_2m_1)lm + m_1m_2l^2} \right| \\ &= \frac{1}{2} \left| \frac{n^2\sqrt{(l_1m_2 + l_2m_1)^2 - 4l_1m_2l_2m_1}}{l_1l_2m^2 - (l_1m_2 + l_2m_1)lm + m_1m_2l^2} \right| \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \left| \frac{n^2 \sqrt{(2h)^2 - 4ab}}{am^2 - 2hlm + bl^2} \right| \quad [\because l_1 l_2 = a, m_1 m_2 = b, l_1 m_2 + l_2 m_1 = 2h] \\
&= \frac{1}{2} \left| \frac{n^2 \sqrt{4h^2 - 4ab}}{am^2 - 2hlm + bl^2} \right| \\
&= \left| \frac{n^2 \sqrt{h^2 - ab}}{am^2 - 2hlm + bl^2} \right|
\end{aligned}$$

4. Show that the straight lines represented by  $(x^2 + 2a)^2 - 3y^2 = 0$  and  $x = 0$  form an equilateral triangle.

**Sol:** Given equation of pair of straight lines is  $(x^2 + 2a)^2 - (\sqrt{3}y)^2 = 0$

$$\Rightarrow (x + 2a - \sqrt{3}y)(x + 2a + \sqrt{3}y) = 0$$

$\therefore$  The lines represented by the given equation of pair of straight lines are

$$x + \sqrt{3}y + 2a = 0 \quad \dots(1)$$

$$x - \sqrt{3}y + 2a = 0 \quad \dots(2)$$

$$\text{Given line be } x - a = 0 \quad \dots(3)$$

Let A be the angle between (1) and (3) then

$$\cos A = \frac{|1 \cdot 1 + \sqrt{3} \cdot 0|}{\sqrt{1+3}\sqrt{1+0}} = \frac{1}{2} \Rightarrow A = 60^\circ$$

Again B be the angle between (2) and (3) then

$$\cos B = \frac{|1 \cdot 1 + (-\sqrt{3}) \cdot 0|}{\sqrt{1+3}\sqrt{1+0}} = \frac{1}{2} \Rightarrow B = 60^\circ$$

$\therefore$  Since (1) and (2) are different lines and two angles are  $60^\circ, 60^\circ$

$\therefore$  Third angle is also  $60^\circ$

$\therefore$  Given lines form an equilateral triangle.

5. Find the centroid and area of a triangle formed by the lines  $2y^2 - xy - 6x^2 = 0$  and  $x + y + 4 = 0$

**Sol:** Given equation of straight lines  $2y^2 - xy - 6x^2 = 0 \quad \dots(1)$

$$\Rightarrow 2y^2 - 4xy + 3xy - 6x^2 = 0$$

$$\Rightarrow 2y(y - 2x) + 3x(y - 2x) = 0$$

$$\Rightarrow (y - 2x)(2y + 3x) = 0$$

∴ The lines represented by the given equation  $2y^2 - xy - 6x^2 = 0$  are given by

$$\Rightarrow 2x - y = 0 \quad \dots(2)$$

$$\text{and } 3x + 2y = 0 \quad \dots(3)$$

$$\text{and given straight line } x + y + 4 = 0 \quad \dots(4)$$

Clearly O(0, 0) is point of intersection of (2) and (3)

Let A be the point of intersection of (2) and (4)

$$\frac{x}{-4-0} = \frac{y}{0-8} = \frac{1}{2+1}$$

$$\therefore A = \left( \frac{-4}{3}, \frac{-8}{3} \right)$$

B be the point of intersection of (3) and (4)

$$\frac{x}{8-0} = \frac{y}{0-12} = \frac{1}{3-2}$$

$$\therefore B = (8, -12)$$

$$\text{Centroid of } \triangle ABC = \left( \frac{0 - \frac{4}{3} + 8}{3}, \frac{0 - \frac{8}{3} - 12}{3} \right) = \left( \frac{20}{9}, \frac{-44}{9} \right)$$

[Area of a triangle whose vertices are O(0, 0), A( $x_1, y_1$ ), B( $x_2, y_2$ ) is  $= \frac{1}{2} |x_1 y_2 - x_2 y_1|$ ]

$$\begin{aligned} \therefore \text{ Required area of triangle} &= \frac{1}{2} \left| \left( \frac{-4}{3} \right) (-12) - 8 \left( \frac{-8}{3} \right) \right| \\ &= \frac{1}{2} \left| \frac{48}{3} + \frac{64}{3} \right| = \frac{112}{2 \times 3} = \frac{56}{3} \text{ Sq.units} \end{aligned}$$

6. Show that the lines represented by  $(lx + my)^2 - 3(mx - ly) = 0$  and  $lx + my = 0$  form

an equilateral triangle with area  $\frac{n^2}{\sqrt{3}(l^2 + m^2)}$

Sol: Given straight line  $lx + my = 0 \quad \dots(1)$

$$\text{Slope of (1) } m_1 = \frac{-l}{m}$$

Let the equation of line  $y = m_2 x$  which makes an angle of  $60^\circ$  with (1)

$$\therefore \tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

$$\tan 60^\circ = \left| \frac{-l/m - y/x}{1 + (-l/m)y/x} \right| \quad \left[ \because m_2 = y/x \right]$$

$$\Rightarrow \sqrt{3} = \left| \frac{-lx - my}{mx - ly} \right| = \left| \frac{-(lx + my)}{mx - ly} \right|$$

squaring both sides

$$3(mx - ly)^2 = (lx + my)^2$$

$$\Rightarrow (lx + my)^2 - 3(mx - ly)^2 = 0 \quad \dots(2)$$

which is given equation of pair of straight in the data

$\therefore (lx + my)^2 - 3(mx - ly)^2 = 0$  and  $lx + my + n = 0$  form a equilateral triangle

Also O (0, 0) is point of intersection of lines in (2)

$\therefore$  Perpendicular distance to (1) from O(0,0)  $P = \frac{|n|}{\sqrt{l^2 + m^2}}$

Area of that equilateral triangle

$$\frac{P^2}{\sqrt{3}} = \frac{n^2}{\sqrt{3}(l^2 + m^2)}$$

7. If  $(\alpha, \beta)$  is the centroid of the triangle formed by the lines  $ax^2 + 2hxy + by^2 = 0$  and

$lx + my = 0$  then prove that  $\frac{\alpha}{bl - hm} = \frac{\beta}{am - hl} = \frac{2}{3(bl^2 - 2hlm + am^2)}$

**Sol:** Let  $ax^2 + 2hxy + by^2 = 0$  represent the lines

$$l_1x + m_1y = 0 \quad \dots(1)$$

$$l_2x + m_2y = 0 \quad \dots(2)$$

అనుకొనుము.

$$\therefore ax^2 + 2hxy + by^2 = (l_1x + m_1y)(l_2x + m_2y)$$

$$= l_1l_2x^2 + (l_1m_2 + l_2m_1)xy + m_1m_2y^2$$

comparing both sides like terms

$$l_1l_2 = a, l_1m_2 + l_2m_1 = 2h, m_1m_2 = b$$

$$\text{Given equation } lx + my = 0 \quad \dots(3)$$

Clearly O(0, 0) is point of intersection of (1) and (2)

Let A be the point of intersection of (1) and (3)



$$\frac{x}{-m_1 - 0} = \frac{y}{0 + l_1} = \frac{1}{ml_1 - lm_1}$$

$$\therefore A = \left( \frac{-m_1}{l_1m - lm_1}, \frac{l_1}{l_1m - lm_1} \right)$$

Again let B be the point of intersection of (2) and (3)

$$\frac{x}{-m_2 - 0} = \frac{y}{0 + l_2} = \frac{1}{ml_2 - lm_2}$$

$$\therefore B = \left( \frac{-m_2}{l_2m - lm_2}, \frac{l_2}{l_2m - lm_2} \right)$$

centroid of  $\Delta OAB = (\alpha, \beta)$

$$\therefore (\alpha, \beta) = \left( \frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right)$$

$$\Rightarrow (\alpha, \beta) = \frac{1}{3} \left[ \frac{-m_1}{l_1m - lm_1} - \frac{m_2}{l_2m - lm_2}, \frac{l_1}{l_1m - lm_1} + \frac{l_2}{l_2m - lm_2} \right]$$

$$\therefore \alpha = \frac{1}{3} \left[ \frac{-m_1}{l_1m - lm_1} - \frac{m_2}{l_2m - lm_2} \right], \beta = \left[ \frac{l_1}{l_1m - lm_1} + \frac{l_2}{l_2m - lm_2} \right]$$

Now

$$\alpha = \frac{1}{3} \left[ \frac{-m_1(l_2m - lm_2) - m_2(l_1m - lm_1)}{(l_1m - lm_1)(l_2m - lm_2)} \right]$$

$$= \frac{1}{3} \left[ \frac{-m_1l_2m + lm_1m_2 - l_1mm_2 + lm_1m_2}{l_1l_2m^2 - (l_1m_2 + l_2m_1)lm + m_1m_2l^2} \right]$$

$$= \frac{1}{3} \left[ \frac{2m_1ml - m(l_1m_2 + l_2m_1)}{l_1l_2m^2 - (l_1m_2 + l_2m_1)lm + m_1m_2l^2} \right]$$

$$= \frac{1}{3} \left[ \frac{2bl - m(2h)}{am^2 - 2hlm + bl^2} \right]$$

$$\Rightarrow \frac{\alpha}{bl - hm} = \frac{2}{3(bl^2 - 2hlm + am^2)}$$

similarly we can prove  $\frac{\beta}{am - hl} = \frac{2}{3(bl^2 - 2hlm + am^2)}$

$$\therefore \frac{\alpha}{bl - hm} = \frac{\beta}{am - hl} = \frac{2}{3(bl^2 - 2hlm + am^2)}$$

8. Prove that the equation  $2x^2 + 3xy - 2y^2 + 3x + y + 1 = 0$  represents a pair of perpendicular lines.

**Sol:** compare the given equation  $2x^2 + 3xy - 2y^2 + 3x + y + 1 = 0$  with

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$$

$$a = 2, h = \frac{3}{2}, b = -2, g = \frac{3}{2}, f = \frac{1}{2}, c = 1$$

$$\text{Now } abc + 2fgh - af^2 - bg^2 - ch^2$$

$$= 2(-2)(1) + 2\left(\frac{1}{2}\right)\left(\frac{3}{2}\right)\left(\frac{3}{2}\right) - 2\left(\frac{1}{2}\right)^2 - (-2)\left(\frac{3}{2}\right)^2 - 1\left(\frac{3}{2}\right)^2$$

$$= -4 + \frac{9}{4} - \frac{1}{2} + \frac{9}{2} - \frac{9}{4}$$

$$= -4 + \frac{8}{2} = 0$$

$$\text{also } h^2 - ab = \frac{9}{4} - 2(-2) = \frac{9}{4} + 4 = \frac{9+16}{4} = \frac{25}{4} > 0$$

$$g^2 - ac = \frac{9}{4} - 2(1) = \frac{9}{4} - 2 = \frac{9-8}{4} = \frac{1}{4} > 0$$

$$f^2 - bc = \frac{1}{4} - (-2) = \frac{1}{4} + 2 = \frac{1+8}{4} = \frac{9}{4} > 0$$

$\therefore$  given equation  $2x^2 + 3xy - 2y^2 + 3x + y + 1 = 0$  represents a pair of straight line also  
coeff.  $x^2$  + coeff.  $y^2 = 2 - 2 = 0$

$\therefore$  The lines in the pair of straight lines are perpendicular.

9. Show that the product of the perpendicular distance from the origin to the pair of straight lines represented by  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$  is

$$\frac{|c|}{\sqrt{(a-b)^2 + 4h^2}}$$

**Sol:** Let the equation  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

represents the lines  $l_1x + m_1y = 0$  ....(1)

$$l_2x + m_2y = 0 \quad \dots(2)$$

$$\therefore ax^2 + 2hxy + by^2 + 2gx + 2fy + c = (l_1x + m_1y + n_1)(l_2x + m_2y + n_2)$$

$$= l_1l_2x^2 + (l_1m_2 + l_2m_1)xy + m_1m_2y^2 + (l_1n_2 + l_2n_1)x + (m_1n_2 + m_2n_1)y + n_1n_2$$

comparing both sides the coefficients of like terms

$$l_1l_2 = a, l_1m_2 + l_2m_1 = 2h, m_1m_2 = b, l_1n_2 + l_2n_1 = 2g, m_1n_2 + m_2n_1 = 2f, n_1n_2 = c$$

Now perpendicular distance from O(0, 0) to (1) =  $\frac{|n_1|}{\sqrt{l_1^2 + m_1^2}}$

similarly perpendicular distance from O(0, 0) to (2) =  $\frac{|n_2|}{\sqrt{l_2^2 + m_2^2}}$

product of perpendicular distance =  $\frac{|n_1|}{\sqrt{l_1^2 + m_1^2}} \times \frac{|n_2|}{\sqrt{l_2^2 + m_2^2}}$

$$= \frac{|n_1n_2|}{\sqrt{(l_1^2 + m_1^2)(l_2^2 + m_2^2)}}$$

$$= \frac{|n_1n_2|}{\sqrt{l_1^2l_2^2 + l_1^2m_2^2 + l_2^2m_1^2 + m_1^2m_2^2}}$$

$$= \frac{|n_1n_2|}{\sqrt{(l_1l_2 - m_1m_2)^2 + 2l_1l_2m_1m_2 + (l_1m_2 + l_2m_1)^2 - 2l_1m_2l_2m_1}}$$

$$= \frac{|n_1n_2|}{\sqrt{(l_1l_2 - m_1m_2)^2 + (l_1m_2 + l_2m_1)^2}}$$

$$= \frac{|c|}{\sqrt{(a-b)^2 + (2h)^2}}$$

$$= \frac{|c|}{\sqrt{(a-b)^2 + 4h^2}}$$

10. Find the values of k, if the lines joining the origin to the points of intersection of the curve  $2x^2 - 2xy + 3y^2 + 2x - y - 1 = 0$  and the line  $x+2y=k$  are mutually perpendicular

**Sol:** Given straight line equation  $x + 2y = k \Rightarrow \frac{x+2y}{k} = 1$  ..... (1)

Given curve equation  $2x^2 - 2xy + 3y^2 + 2x - y - 1 = 0$  ..... (2)

To get the required equation by homogenising (2) with the help of equation (1)

$$\therefore 2x^2 - 2xy + 3y^2 + (2x - y)(1) - 1(1^2) = 0$$

$$\Rightarrow 2x^2 - 2xy + 3y^2 + (2x - y)\frac{(x+2y)}{k} - 1\frac{(x+2y)^2}{k^2} = 0$$

$$\Rightarrow k^2(2x^2 - 2xy + 3y^2) + k(2x^2 + 4xy - xy - 2y^2) - (x^2 + 4y^2 + 4xy) = 0$$

$$\Rightarrow x^2(2k^2 + 2k - 1) + xy(-2k^2 + 3k - 4) + y^2(3k^2 - 2k - 4) = 0$$

If the two lines in above equation are perpendicular then

$$\text{coeff. } x^2 + \text{coeff. } y^2 = 0$$


$$\Rightarrow 2k^2 + 2k - 1 + 3k^2 - 2k - 4 = 0$$

$$\Rightarrow 5k^2 - 5 = 0$$

$$\Rightarrow 5k^2 = 5$$

$$\Rightarrow k^2 = 1$$

$$\Rightarrow k^2 = \pm 1$$



11. Find the angle between the lines joining the origin to the points of intersection of the curve  $x^2 + 2xy + y^2 + 2x + 2y - 5 = 0$  and the line  $3x - y + 1 = 0$

**Sol:** Given straight line equation  $3x - y + 1 = 0 \Rightarrow y - 3x = 1$  ..... (1)

Given curve equation  $x^2 + 2xy + y^2 + 2x + 2y - 5 = 0$  ..... (2)

to get the equations of the lines OA, OB is obtained by homogenising the equation (2) by equation (1)

$$\therefore x^2 + 2xy + y^2 + 2x + 2y - 5(1^2) = 0$$

$$\Rightarrow x^2 + 2xy + y^2 + (2x + 2y)(y - 3x) - 5(y - 3x)^2 = 0$$

$$\Rightarrow x^2 + 2xy + y^2 + 2xy - 6x^2 + 2y^2 - 6xy - 5(y^2 + 9x^2 - 6xy) = 0$$

$$\Rightarrow x^2 + 2xy + y^2 + 2xy - 6x^2 + 2y^2 - 6xy - 5y^2 - 45x^2 + 30xy = 0$$

$$\Rightarrow -50x^2 + 28xy - 2y^2 = 0$$

$$\Rightarrow 25x^2 - 14xy + y^2 = 0$$


If  $\theta$  is the angle between the lines in the above equation then

$$\cos\theta = \frac{|a+b|}{\sqrt{(a-b)^2 + 4h^2}}$$

$$a = 25, h = -7, b = 1$$

$$\therefore \cos\theta = \frac{|25+1|}{\sqrt{(25-1)^2 + 4(-7)^2}}$$

$$= \frac{26}{\sqrt{276+196}}$$



$$= \frac{26}{\sqrt{4 \times 196}} = \frac{13}{\sqrt{193}}$$

$$\therefore \theta = \cos^{-1}\left(\frac{13}{\sqrt{193}}\right)$$

**12. Find the condition for the chord  $lx + my = 1$  of the circle  $x^2 + y^2 = a^2$  (whose centre is the origin) to subtend a right angle at the origin.**

**Sol:** Given circle equation  $x^2 + y^2 - a^2 = 0$  ..... (1)

Equation of the chord  $lx + my = 1$  ..... (2)

to get the required equation by homogenising (1) by (2) we get

$$x^2 + y^2 - a^2 (lx + my)^2 = 0$$

$$\Rightarrow x^2 + y^2 - a^2 (l^2 x^2 + m^2 y^2 + 2lmxy) = 0$$

$$\Rightarrow x^2(1 - a^2 l^2) - 2a^2 lmxy + y^2(1 - a^2 m^2) = 0$$

If two lines in above equation are perpendicular then

$$\text{coeff. of } x^2 + \text{coeff. of } y^2 = 0$$

$$\Rightarrow 1 - a^2 l^2 + 1 - a^2 m^2 = 0$$

$$\Rightarrow 2 = a^2 l^2 + a^2 m^2$$

$$\Rightarrow a^2 (l^2 + m^2) = 2$$

**13. Find the condition for the lines joining the origin to the points of intersection of the circle  $x^2 + y^2 = a^2$  and the line  $lx + my = 1$  to coincide.**

**Sol:** Given circle equation  $x^2 + y^2 - a^2 = 0$  ..... (1)

Equation of the chord  $lx + my = 1$  ..... (2)

to get the required equation by homogenising (1) by (2) we get

$$x^2 + y^2 - a^2 (lx + my)^2 = 0$$

$$\Rightarrow x^2 + y^2 - a^2 (l^2 x^2 + m^2 y^2 + 2lmxy) = 0$$

$$\Rightarrow x^2(1 - a^2 l^2) - 2a^2 lmxy + y^2(1 - a^2 m^2) = 0$$

[If two lines in  $ax^2 + 2hxy + by^2 = 0$  are coincide then  $h^2 = ab$ ]

$$\therefore (-a^2 lm)^2 = (1 - a^2 l^2)(1 - a^2 m^2)$$

$$\Rightarrow a^4 l^2 m^2 = 1 - a^2 m^2 - a^2 l^2 + a^4 l^2 m^2$$

$$\Rightarrow a^2 m^2 + a^2 l^2 = 1$$

$$\Rightarrow a^2 (l^2 + m^2) = 1$$

\*\*\*\*\*

## Three Dimensional Coordinates

### Key concepts

→ If  $P(x_1, y_1, z_1)$ ,  $Q(x_2, y_2, z_2)$  are two points in the space then distance between P and Q

$$PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

→ The point dividing the segment  $\overline{AB}$  where  $A(x_1, y_1, z_1)$ ,  $B(x_2, y_2, z_2)$  in the ratio  $m:n$  is

(i)  $\left( \frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right)$

(ii) Mid point of  $\overline{AB}$

$$\left( \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right)$$

→ Centroid of a triangle with vertices  $(x_1, y_1, z_1)$ ,  $(x_2, y_2, z_2)$ ,  $(x_3, y_3, z_3)$

$$\left( \frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right)$$

→ Centroid of a tetrahedron whose vertices are

$$(x_1, y_1, z_1), (x_2, y_2, z_2), (x_3, y_3, z_3), (x_4, y_4, z_4)$$

$$\left( \frac{x_1 + x_2 + x_3 + x_4}{4}, \frac{y_1 + y_2 + y_3 + y_4}{4}, \frac{z_1 + z_2 + z_3 + z_4}{4} \right)$$

### Very Short Answer Questions (2 Marks)

1. Find  $x$  if the distance between  $(5, -1, 7)$  and  $(x, 5, 1)$  is 9 units.

**Sol:** Given points  $(5, -1, 7)$ ,  $(x, 5, 1)$

from problem we have  $AB = 9$

$$\Rightarrow AB^2 = 81$$

$$\Rightarrow (x-5)^2 + (5+1)^2 + (1-7)^2 = 81$$

$$\Rightarrow (x-5)^2 + 36 + 36 = 81$$

$$\Rightarrow (x-5)^2 = 9$$

$$\Rightarrow (x-5) = \pm 3$$

$$\Rightarrow x = 5 \pm 3$$

$$\Rightarrow x = 8, 2$$

**2. Show that the points (1, 2, 3), (2, 3, 1), (3, 1, 2) form an equilateral triangle.**

**Sol:** Let given points be A(1, 2, 3), B(2, 3, 1), C(3, 1, 2)

$$AB = \sqrt{(2-1)^2 + (3-2)^2 + (1-3)^2} = \sqrt{1^2 + 1^2 + (-2)^2} = \sqrt{6}$$

$$BC = \sqrt{(3-2)^2 + (1-3)^2 + (2-1)^2} = \sqrt{1^2 + (-2)^2 + 1^2} = \sqrt{6}$$

$$CA = \sqrt{(1-3)^2 + (2-1)^2 + (3-2)^2} = \sqrt{(-2)^2 + 1^2 + 1^2} = \sqrt{6}$$

$$AB = BC = CA$$

$\therefore \Delta ABC$  is an equilateral triangle

**3. find the centroid of the triangle whose vertices are (5, 4, 6), (1, -1, 3), and (4, 3, 2)**

**Sol:** Let given points be A ( $x_1, y_1, z_1$ ) = (5, 4, 6)

$$B (x_2, y_2, z_2) = (1, -1, 3)$$

$$C (x_3, y_3, z_3) = (4, 3, 2)$$

$$\begin{aligned} \text{Centroid of } \Delta ABC &= \left( \frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right) \\ &= \left( \frac{5+1+4}{3}, \frac{4-1+3}{3}, \frac{6+3+2}{3} \right) \\ &= \left( \frac{10}{3}, 2, \frac{11}{3} \right) \end{aligned}$$

**4. Find the centroid of the tetrahedron whose vertices are (2, 3, -4), (-3, 3, -2) (-1, 4, 2) and (3, 5, 1)**

**Sol:** Let given points of a tetrahedron

$$A (x_1, y_1, z_1) = (2, 3, -4)$$

$$B (x_2, y_2, z_2) = (-3, 3, -2)$$

$$C (x_3, y_3, z_3) = (-1, 4, 2)$$

$$D (x_4, y_4, z_4) = (3, 5, 1)$$

Centroid of a tetrahedron

$$\begin{aligned} &\left( \frac{x_1 + x_2 + x_3 + x_4}{4}, \frac{y_1 + y_2 + y_3 + y_4}{4}, \frac{z_1 + z_2 + z_3 + z_4}{4} \right) \\ &= \left( \frac{2-3-1+3}{4}, \frac{3+3+4+5}{4}, \frac{-4-2+2+1}{4} \right) = \left( \frac{1}{4}, \frac{15}{4}, \frac{-3}{4} \right) \end{aligned}$$

5. Find the fourth vertex of the parallelogram whose consecutive vertices are (2, 4, -1), (3, 6, -1) and (4, 5, 1)

**Sol:** Let ABCD be parallelogram where A(2, 4, -1), B(3, 6, -1), C(4, 5, 1) and D (a, b, c)

$\therefore$  mid point of AC = mid point of BD

$$\Rightarrow \left( \frac{2+4}{2}, \frac{4+5}{2}, \frac{-1+1}{2} \right) = \left( \frac{3+a}{2}, \frac{6+b}{2}, \frac{-1+c}{2} \right)$$

$$\Rightarrow \left( \frac{6}{2}, \frac{9}{2}, 0 \right) = \left( \frac{a+3}{2}, \frac{b+6}{2}, \frac{c-1}{2} \right)$$

$$\Rightarrow \frac{a+3}{2} = \frac{6}{2}, \frac{b+6}{2} = \frac{9}{2}, \frac{c-1}{2} = 0$$

$\therefore a = 3, b = 3, c = 1 \Rightarrow 4^{\text{th}}$  vertex D(a, b, c) = (3, 3, 1)

6. Find the coordinates of the vertex 'C' of  $\Delta$  ABC if its centroid is the origin and the vertices A, B are (1, 1, 1) and (-2, 4, 1) respectively.

**Sol:** Let given points A(1, 1, 1), B(-2, 4, 1), C(x, y, z)

Given that centroid of  $\Delta$  ABC = (0, 0, 0)

$$\Rightarrow \left( \frac{1-2+x}{3}, \frac{1+4+y}{3}, \frac{1+1+z}{3} \right) = (0, 0, 0)$$

$$\Rightarrow \left( \frac{x-1}{3}, \frac{y+5}{3}, \frac{z+2}{3} \right) = (0, 0, 0)$$

$$\Rightarrow \frac{a+3}{2} = \frac{6}{2}, \frac{b+6}{2} = \frac{9}{2}, \frac{c-1}{2} = 0$$

$$\Rightarrow -1+x=0, 5+y=0, z+2=0$$

$$\Rightarrow x=1, y=-5, z=-2 \Rightarrow \text{third vertex } C(x, y, z) = (1, -5, -2)$$

7. If (3, 2, -1), (4, 1, 1), (6, 2, 5) and (4, 2, 2) is the centroid of a tetrahedron, find the fourth vertex.

**Sol:** Let given points A(3, 2, -1), B(4, 1, 1), C(4, 1, 1), D(x, y, z)

Given that centroid of ABCD tetrahedron = (4, 2, 2)

$$\Rightarrow \left( \frac{3+4+6+x}{4}, \frac{2+1+2+y}{4}, \frac{-1+1+5+z}{4} \right) = (4, 2, 2)$$

$$\Rightarrow \left( \frac{13+x}{4}, \frac{5+y}{4}, \frac{5+z}{4} \right) = (4, 2, 2) \Rightarrow \frac{13+x}{4} = 4, \frac{5+y}{4} = 2, \frac{5+z}{4} = 2$$

$$\Rightarrow x=3, y=3, z=3 \Rightarrow 4^{\text{th}}$$
 vertex D(x, y, z) = (3, 3, 3)

\*\*\*\*\*



## Direction Cosines and Direction Ratios

### Key concepts

- A ray  $\overline{OP}$  passing through origin  $O$  and making angles  $\alpha, \beta, \gamma$  respectively with  $\overline{OX}, \overline{OY}, \overline{OZ}$  then the numbers  $\cos\alpha, \cos\beta, \cos\gamma$  are called the direction cosines of the ray  $\overline{OP}$ . Usually they are denoted by  $(l, m, n)$  where  $l = \cos\alpha, m = \cos\beta, n = \cos\gamma$
- If  $(l, m, n)$  are direction cosines of  $\overline{OP}$ . then direction cosines of  $\overline{PO}$  are  $(-l, -m, -n)$
- If  $(l, m, n)$  are direction cosines of a line then  $l^2 + m^2 + n^2 = 1$
- Triad of numbers  $(a, b, c)$  proportional to direction cosines of a line are called its direction ratios.
- Direction cosines of a line whose direction ratios  $(a, b, c)$  are
- $$\pm \left( \frac{a}{\sqrt{a^2 + b^2 + c^2}}, \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \frac{c}{\sqrt{a^2 + b^2 + c^2}} \right)$$
- Direction ratios of the line joining  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  are  $(x_2 - x_1, y_2 - y_1, z_2 - z_1)$  and its direction cosines are
- $$\left( \frac{x_2 - x_1}{\sqrt{\Sigma(x_2 - x_1)^2}}, \frac{y_2 - y_1}{\sqrt{\Sigma(x_2 - x_1)^2}}, \frac{z_2 - z_1}{\sqrt{\Sigma(x_2 - x_1)^2}} \right)$$
- If  $\theta$  is angle between two lines whose direction cosines are  $(l_1, m_1, n_1), (l_2, m_2, n_2)$  then  $\cos\theta = |l_1l_2 + m_1m_2 + n_1n_2|$
- If  $\theta$  is angle between two lines whose direction ratios are  $(a_1, b_1, c_1), (a_2, b_2, c_2)$  then

$$\cos\theta = \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

### Long Answer Questions (7 Marks)

1. Find the direction cosines of two lines which are connected by the relations  $l + m + n = 0$  and  $mn - 2nl - 2lm = 0$

**Sol:** Given that  $l + m + n = 0$  ....(1)

$$mn - 2nl - 2lm = 0 \quad \dots(2)$$

From (1)  $l = -m - n$  substituting  $l$  value in (2) we get

$$mn - 2n(-m - n) - 2(-m - n)m = 0$$

$$\Rightarrow mn + 2mn + 2n^2 + 2m^2 + 2mn = 0$$

$$\Rightarrow 2m^2 + 4mn + mn + 2n^2 = 0$$

$$\Rightarrow 2m(m + 2n) + n(m + 2n) = 0$$

$$\Rightarrow (m + 2n)(2m + n) = 0$$

$$\Rightarrow m + 2n = 0 \text{ or } 2m + n = 0$$

Now  $2m + n = 0 \Rightarrow n = -2m$

from (1)  $l = -m - (-2m) = m$

$\therefore l : m : n = m : m : -2m = 1 : 1 : -2$

directional cosines  $\left( \frac{1}{\sqrt{1^2 + 1^2 + (-2)^2}}, \frac{1}{\sqrt{1^2 + 1^2 + (-2)^2}}, \frac{-2}{\sqrt{1^2 + 1^2 + (-2)^2}} \right)$

$$= \left( \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, \frac{-2}{\sqrt{6}} \right)$$

Now  $m + 2n = 0 \Rightarrow m = -2n$

from (1)  $l = -m - n = 2n - n = n$

$\therefore l : m : n = n : -2n : n = 1 : -2 : 1$

Direction cosines of a line is  $\left( \frac{1}{\sqrt{1^2 + (-2)^2 + 1^2}}, \frac{-2}{\sqrt{1^2 + (-2)^2 + 1^2}}, \frac{1}{\sqrt{1^2 + (-2)^2 + 1^2}} \right)$

$$= \left( \frac{1}{\sqrt{6}}, \frac{-2}{\sqrt{6}}, \frac{1}{\sqrt{6}} \right)$$

Thus the d.c's of the two lines are  $= \left( \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, \frac{-2}{\sqrt{6}} \right) = \left( \frac{1}{\sqrt{6}}, \frac{-2}{\sqrt{6}}, \frac{1}{\sqrt{6}} \right)$

2. Find the direction cosines of two lines which are connected by the relations  $l - 5m + 3n = 0$  and  $7l^2 + 5m^2 - 3n^2 = 0$

**Sol:** Given that  $l - 5m + 3n = 0$  ....(1)

$$7l^2 + 5m^2 - 3n^2 = 0 \quad \dots(2)$$

From (1)  $l = 5m - 3n$  substitute this value in (2)

$$\begin{aligned} 7(5m - 3n)^2 + 5m^2 - 3n^2 &= 0 \\ \Rightarrow 7(25m^2 + 9n^2 - 30mn) + 5m^2 - 3n^2 &= 0 \\ \Rightarrow 175m^2 + 63n^2 - 210mn + 5m^2 - 3n^2 &= 0 \\ \Rightarrow 180m^2 - 210mn + 60n^2 &= 0 \\ \Rightarrow 6m^2 - 7mn + 2n^2 &= 0 \\ \Rightarrow (3m - 2n)(2m - n) &= 0 \\ \Rightarrow 3m - 2n = 0 \text{ ຫຼື } 2m - n = 0 \end{aligned}$$

Now  $3m - 2n = 0 \Rightarrow m = \frac{2}{3}n$

From (1)  $l = 5m - 3n = \frac{10}{3}n - 3n = \frac{10n - 9n}{3} = \frac{n}{3}$

$\therefore l : m : n = \frac{n}{3} : \frac{2}{3}n : n = \frac{1}{3} : \frac{2}{3} : 1 = 1 : 2 : 3$

Direction cosines  $\left( \frac{1}{\sqrt{1^2 + 2^2 + 3^2}}, \frac{2}{\sqrt{1^2 + 2^2 + 3^2}}, \frac{3}{\sqrt{1^2 + 2^2 + 3^2}} \right)$   
 $= \left( \frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}} \right)$

similarly  $2m - n = 0 \Rightarrow n = 2m$

(1) ສອດ້  $l = 5m - 3(2m) = 5m - 6m = -m$

$\therefore l : m : n = -m : m : 2m = -1 : 1 : 2$

Direction cosines of another line  $\left( \frac{-1}{\sqrt{(-1)^2 + 1^2 + 2^2}}, \frac{1}{\sqrt{(-1)^2 + 1^2 + 2^2}}, \frac{2}{\sqrt{(-1)^2 + 1^2 + 2^2}} \right)$   
 $= \left( \frac{-1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, \frac{2}{\sqrt{6}} \right)$

**3. Find the angle between the lines whose direction cosines satisfy the equations**

$l + m + n = 0, l^2 + m^2 - n^2 = 0$

**Sol:** Given equations  $l + m + n = 0$  ....(1)

$l^2 + m^2 - n^2 = 0$  ....(2)

From (1)  $l = -m - n$  substitute this value in (2)

$(-m - n)^2 + m^2 - n^2 = 0$

$$\Rightarrow m^2 + n^2 + 2mn + m^2 - n^2 = 0$$

$$\Rightarrow 2m^2 + 2mn = 0$$

$$\Rightarrow 2m(m+n) = 0$$

$$\Rightarrow m = 0 \text{ or } m + n = 0$$

Now if  $m = 0$  From (1)  $l = 0 - n = -n$

$$\therefore l : m : n = -n : 0 : n = -1 : 0 : 1$$

direction ratios of a line  $(a_1, b_1, c_1) = (-1, 0, 1)$

$$m + n = 0 \Rightarrow m = -n$$

$$l = -m - n = -(-n) - n = n - n = 0$$

$$\therefore l : m : n = 0 : -n : n = 0 : -1 : 1$$

direction ratios of another line  $(a_2, b_2, c_2) = (0, -1, 1)$

Let  $\theta$  be the angle between two lines then

$$\cos \theta = \frac{|a_1 a_2 + b_1 b_2 + c_1 c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} = \frac{|0 \times (-1) + 0 \times (-1) + 1 \times 1|}{\sqrt{1+0+1} \sqrt{1+0+1}} = \frac{1}{\sqrt{2}\sqrt{2}} = \frac{1}{2}$$

$$\therefore \theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ = \frac{\pi}{3}$$

4. Find the angle between the lines whose direction cosines are given by the equations  $3l + m + 5n = 0$  and  $6mn - 2nl + 5lm = 0$

**Sol:** Given equation  $3l + m + 5n = 0$  ....(1)

$$6mn - 2nl + 5lm = 0 \quad \dots(2)$$

From (1)  $m = -3l - 5n$  substitute this value in (2)

$$6(-3l - 5n)n - 2nl + 5l(-3l - 5n) = 0$$

$$\Rightarrow -18ln - 30n^2 - 2nl - 15l^2 - 25ln = 0$$

$$\Rightarrow -15l^2 - 45ln - 30n^2 = 0$$

$$\Rightarrow l^2 + 3ln + 2n^2 = 0$$

$$\Rightarrow (l+2n)(l+n) = 0$$

$$\Rightarrow l + 2n = 0 \text{ or } l + n = 0$$

Now  $l + 2n = 0 \Rightarrow l = -2n$

From (1)  $m = -3l - 5n = 6n - 5n = n$

$$\therefore l : m : n = -2n : n : n = -2 : 1 : 1$$

Direction ratios of a line  $(a_1, b_1, c_1) = (-2, 1, 1)$

also  $l + n = 0 \Rightarrow l = -n$

from (1)  $m = -3l - 5n = -3(-n) - 5n = 3n - 5n = -2n$

$\therefore l : m : n = -n : -2n : n = -1 : -2 : 1 = 1 : 2 : -1$

direction ratios of another line  $(a_2, b_2, c_2) = (1, 2, -1)$

If  $\theta$  is angle between two lines then

$$\begin{aligned} \cos\theta &= \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} = \frac{|(-2) \times 1 + 1 \times 2 + 1 \times (-1)|}{\sqrt{(-2)^2 + 1^2 + 1^2} \sqrt{1^2 + 2^2 + (-1)^2}} \\ &= \frac{|-1|}{\sqrt{6}\sqrt{6}} = \frac{1}{6} \end{aligned}$$

$$\therefore \theta = \cos^{-1}\left(\frac{1}{6}\right)$$

**5. Find the angle between two diagonals of a cube**

**Sol:** Let 'O', One of the vertices of the cube, be taken as the origin and the three coterminous edges  $\overline{OA}, \overline{OB}, \overline{OC}$  as coordinate axes.

let  $\overline{OA} = \overline{OB} = \overline{OC} = a$

Coordinates of vertices of cube  $O(0, 0, 0), A(a, 0, 0), B(0, a, 0)$

$C(0, 0, a), D(a, a, 0), E(a, 0, a), F(a, a, a), G(0, a, a)$

Also four diagonals of cube are  $\overline{OF}, \overline{AG}, \overline{BE}, \overline{CD}$


Now direction ratio's of diagonal  $\overline{OF}$  are  $(a - 0, a - 0, a - 0) = (a, a, a)$

direction ratio's of diagonal  $\overline{AG}$   $(0 - a, a - 0, a - 0) = (-a, a, a)$

If  $\theta$  is the angle between  $\overline{OF}, \overline{AG}$  then by

$$\begin{aligned} \cos\theta &= \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} = \frac{|a(-a) + a(a) + a(a)|}{\sqrt{a^2 + a^2 + a^2} \sqrt{a^2 + a^2 + a^2}} \\ &= \frac{a^2}{\sqrt{3}a \cdot \sqrt{3}a} = \frac{1}{3} \end{aligned}$$

$$\therefore \theta = \cos^{-1}\left(\frac{1}{3}\right)$$



6. If a ray makes angles  $\alpha, \beta, \gamma, \delta$  with the four diagonals of a cube

find  $\cos^2\alpha + \cos^2\beta + \cos^2\gamma + \cos^2\delta$ .


**Sol:** Let 'O', One of the vertices of the cube, be taken as the origin and the three coterminous edges  $\overline{OA}, \overline{OB}, \overline{OC}$  as coordinate axes.

let  $\overline{OA} = \overline{OB} = \overline{OC} = a$

Coordinates of vertices of cube  $O(0, 0, 0), A(a, 0, 0), B(0, a, 0),$

$C(0, 0, a), D(a, a, 0), E(a, 0, a), F(a, a, a), G(0, a, a)$

Also four diagonals of cube are  $\overline{OF}, \overline{AG}, \overline{BE}, \overline{CD}$



Now direction ratio's of diagonal  $\overline{OF}$  are  $(a - 0, a - 0, a - 0) = (a, a, a)$

direction ratio's of diagonal  $\overline{AG}$   $(0 - a, a - 0, a - 0) = (-a, a, a)$

direction ratio's of diagonal  $\overline{BE}$   $(a - 0, a - 0, a - 0) = (a, -a, a)$

direction ratio's of diagonal  $\overline{CD}$   $(a - 0, a - 0, 0 - a) = (a, a, -a)$

If any ray having direction ratio's  $(l, m, n)$  and makes an angle with four diagonals  $\overline{OF}, \overline{AG}, \overline{BE}, \overline{CD}$  are  $\alpha, \beta, \gamma, \delta$  then

$$\cos\alpha = \frac{|al + am + an|}{\sqrt{a^2 + a^2 + a^2} \sqrt{l^2 + m^2 + n^2}} = \frac{|a(l + m + n)|}{\sqrt{3a^2} \sqrt{l^2 + m^2 + n^2}}$$

$$\therefore \cos^2\alpha = \frac{a^2(l + m + n)^2}{3a^2(l^2 + m^2 + n^2)} = \frac{(l + m + n)^2}{3(l^2 + m^2 + n^2)}$$

similarly

$$\cos^2\beta = \frac{(-l + m + n)^2}{3(l^2 + m^2 + n^2)}, \cos^2\gamma = \frac{(l - m + n)^2}{3(l^2 + m^2 + n^2)}, \cos^2\delta = \frac{(l + m - n)^2}{3(l^2 + m^2 + n^2)}$$

$$\therefore \cos^2\alpha + \cos^2\beta + \cos^2\gamma + \cos^2\delta = \frac{(-l + m + n)^2 + (l - m + n)^2 + (l + m - n)^2}{3(l^2 + m^2 + n^2)}$$

simplification

$$= \frac{4l^2 + 4m^2 + 4n^2}{3(l^2 + m^2 + n^2)} = \frac{4(l^2 + m^2 + n^2)}{3(l^2 + m^2 + n^2)} = \frac{4}{3}$$

\*\*\*\*\*

## The Plane

### Key concepts

- A plane is a surface with at least three non collinear points such that the line joining any two points on the surface lies entirely on it.
- Equation of the plane in normal form is  $lx + my + nz = p$  where  $(l, m, n)$  are direction cosines of the normal to the plane and 'p' is the perpendicular distance to the plane from the origin.
- General equation of the plane is  $ax + by + cz + d = 0$  where  $(a, b, c)$  are direction ratios of the normal to the plane.
- Equation of the plane in intercept form  $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$  where  $a, b, c$  are intercepts on X, Y, Z – axes respectively.
- Angle between two planes is the angle between their normals. If  $\theta$  is the angle between the planes  $a_1x + b_1y + c_1z + d_1 = 0$  and  $a_2x + b_2y + c_2z + d_2 = 0$  then

$$\cos \theta = \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

- (i) If  $a_1a_2 + b_1b_2 + c_1c_2 = 0$  then planes are perpendicular.
- (ii) If  $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$  then the planes are parallel.

### Very Short Answer Questions (2 Marks)

1. Reduce the equation  $x + 2y - 3z - 6 = 0$  of the plane to the normal form.

**Sol:** Given equation of the plane  $x + 2y - 3z - 6 = 0$

dividing both sides  $\sqrt{1^2 + 2^2 + (-3)^2} = \sqrt{14}$

$$\frac{x + 2y - 3z}{\sqrt{14}} = \frac{6}{\sqrt{14}}$$

$$\Rightarrow \frac{1}{\sqrt{14}}x + \frac{2}{\sqrt{14}}y - \frac{3}{\sqrt{14}}z = \frac{6}{\sqrt{14}}$$

2. Find the equation of the plane whose intercepts on X, Y, Z – axes are 1, 2, 4 respectively.

**Sol:** Let the intercepts on X, Y, Z – axes

are  $a = 1, b = 2, c = 4$  respectively then equation of the plane in intercept form is

$$\begin{aligned}\frac{x}{a} + \frac{y}{b} + \frac{z}{c} &= 1 \\ \Rightarrow \frac{x}{1} + \frac{y}{2} + \frac{z}{4} &= 1 \\ \Rightarrow 4x + 2y + z &= 4\end{aligned}$$

3. Write the equation of the plane  $4x - 4y + 2z + 5 = 0$  in the intercept form.

**Sol:** Given equation of the plane  $4x - 4y + 2z + 5 = 0$

$$\begin{aligned}\Rightarrow 4x - 4y + 2z &= -5 \\ \Rightarrow \frac{4x - 4y + 2z}{5} &= \frac{-5}{5} \\ \Rightarrow \frac{-4}{5}x + \frac{4}{5}y - \frac{2}{5}z &= 1 \\ \Rightarrow \frac{x}{-5/4} + \frac{y}{5/4} + \frac{z}{-5/2} &= 1\end{aligned}$$

4. Find the angle between the planes  $x + 2y + 2z - 5 = 0$  and  $3x + 3y + 2z - 8 = 0$

**Sol:** Given equation of the plane  $x + 2y + 2z - 5 = 0$  .....(1)

$$a_1 = 1, b_1 = 2, c_1 = 2$$

and given another plane equation  $3x + 3y + 2z - 8 = 0$  .....(2)

$$a_2 = 3, b_2 = 3, c_2 = 2$$

Let  $\theta$  be the angle between the planes (1) and (2)

$$\begin{aligned}\text{then } \cos\theta &= \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} \\ &= \frac{|1 \times 3 + 2 \times 3 + 2 \times 2|}{\sqrt{1^2 + 2^2 + 2^2} \sqrt{3^2 + 3^2 + 2^2}} \\ &= \frac{|3 + 6 + 4|}{\sqrt{1 + 4 + 4} \sqrt{9 + 9 + 4}} = \frac{13}{3\sqrt{22}} \\ \therefore \theta &= \cos^{-1}\left(\frac{13}{3\sqrt{22}}\right)\end{aligned}$$

\*\*\*\*\*



## Limits

### Key concepts

$$* \quad \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}$$

$$* \quad \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \quad (\text{x is in radians})$$

$$* \quad \lim_{x \rightarrow 0} \frac{\tan x}{x} = 1 \quad (\text{x is in radians})$$

$$* \quad \lim_{x \rightarrow 0} (1+x)^{1/x} = e$$

$$* \quad \lim_{x \rightarrow 0} \left(1 + \frac{1}{x}\right)^x = e$$

$$* \quad \lim_{x \rightarrow 0} \left(\frac{a^x - 1}{x}\right) = \log_e a$$

$$* \quad \lim_{x \rightarrow 0} \frac{x^n - a^n}{x^m - a^m} = \frac{n}{m} a^{n-m}$$

$$* \quad \lim_{x \rightarrow 0} \frac{\sin ax}{x} = a \quad (\text{x is in radians})$$

$$* \quad \lim_{x \rightarrow 0} \frac{\tan ax}{x} = a \quad (\text{x is in radians})$$

$$* \quad \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$$

### Very Short Answer Questions (2 Marks)

1. 
$$\lim_{x \rightarrow 0} \frac{e^{x+3} - e^3}{x}$$

A. 
$$= \lim_{x \rightarrow 0} \frac{e^x \cdot e^3 - e^3}{x} \quad \boxed{a^{m+n} = a^m \cdot a^n}$$

$$= \lim_{x \rightarrow 0} e^3 \frac{(e^x - 1)}{x}$$

$$= e^3 \lim_{x \rightarrow 0} \frac{e^x - 1}{x}$$

$$= e^3 \cdot 1 = e^3$$

2. 
$$\lim_{x \rightarrow 0} \frac{e^x - \sin x - 1}{x}$$

A. 
$$\lim_{x \rightarrow 0} \left( \frac{e^x - 1}{x} - \frac{\sin x}{x} \right)$$

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = \lim_{x \rightarrow 0} \frac{\sin x}{x}$$

$$1 - 1 = 0$$

3.  $\lim_{x \rightarrow 0} \frac{\sin ax}{x \cos x}$

A.  $\lim_{x \rightarrow 0} \frac{\sin ax}{x \cos x} = \frac{\lim_{ax \rightarrow 0} \frac{\sin ax}{ax} \cdot a}{\lim_{x \rightarrow 0} \cos x} = \frac{a}{1} = a$

4.  $\lim_{x \rightarrow \infty} \frac{11x^3 - 3x + 4}{13x^3 - 5x^2 - 7}$

A.  $\lim_{x \rightarrow \infty} \frac{11x^3 - 3x + 4}{13x^3 - 5x^2 - 7}$

$$= \lim_{x \rightarrow \infty} \frac{x^3 \left( 11 - \frac{3}{x^2} + \frac{4}{x^3} \right)}{x^3 \left( 13 - \frac{5}{x} - \frac{7}{x^3} \right)}$$

As  $x \rightarrow \infty$ ,  $\frac{1}{x}$ ,  $\frac{1}{x^2}$  and  $\frac{1}{x^3} \rightarrow 0$

$$= \lim_{x \rightarrow \infty} \frac{11 - \frac{3}{x^2} + \frac{4}{x^3}}{13 - \frac{5}{x} - \frac{7}{x^3}} = \frac{11 - 0 + 0}{13 - 0 - 0} = \frac{11}{13}$$

5.  $\lim_{x \rightarrow 1} \frac{\sin(x-1)}{(x^2-1)}$

A.  $\lim_{x \rightarrow 1} \frac{\sin(x-1)}{(x^2-1)}$

$$\lim_{x \rightarrow 1} \frac{\sin(x-1)}{(x-1)} \cdot \lim_{x \rightarrow 1} \frac{1}{x+1}$$

Put  $y = x - 1$  so that as  $x \rightarrow 1, y \rightarrow 0$

$$\lim_{x \rightarrow 1} \frac{\sin(x-1)}{(x-1)} = \lim_{y \rightarrow 0} \frac{\sin y}{y} = 1$$

$$= 1 \cdot \frac{1}{1+1} = \frac{1}{2}$$

6.  $\lim_{x \rightarrow 0} \frac{\sin ax}{\sin bx} \quad b \neq 0, a \neq b$

A.  $\lim_{x \rightarrow 0} \frac{\frac{\sin ax}{ax} \times ax}{\frac{\sin bx}{bx} \times bx}$

$$\frac{\lim_{ax \rightarrow 0} \frac{\sin ax}{ax} \cdot a}{\lim_{bx \rightarrow 0} \frac{\sin bx}{bx} \cdot b}$$

$$\frac{1 \cdot a}{1 \cdot b} = \frac{a}{b}$$

7.  $\lim_{x \rightarrow \infty} \frac{8|x|+3x}{3|x|-2x}$

A.  $x \rightarrow \infty \Rightarrow x > 0 \therefore |x| = x$

$$\lim_{x \rightarrow \infty} \frac{8|x|+3x}{3|x|-2x} = \lim_{x \rightarrow \infty} \frac{8x+3x}{3x-2x}$$

$$\lim_{x \rightarrow \infty} \frac{11x}{x} = 11$$

8.  $\lim_{x \rightarrow 0} \frac{a^x - 1}{b^x - 1}$

A.  $\lim_{x \rightarrow 0} \frac{\frac{a^x - 1}{x}}{\frac{b^x - 1}{x}}$

$$\frac{\lim_{x \rightarrow 0} \frac{a^x - 1}{x}}{\lim_{x \rightarrow 0} \frac{b^x - 1}{x}}$$

$$= \frac{\log_e a}{\log_e b}$$

$$= \log_b a$$

$$9. \quad \lim_{x \rightarrow 0} \frac{e^{7x} - 1}{x}$$

A. As  $x \rightarrow 0$

$$\Rightarrow 7x \rightarrow 0$$

$$\lim_{7x \rightarrow 0} \frac{e^{7x} - 1}{7x} \times 7 = 7 \left( \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1 \right)$$

$$10. \quad \lim_{x \rightarrow 0} \left( \frac{\sqrt{1+x} - 1}{x} \right)$$

A. For  $0 < |x| < 1$ , we have

$$\frac{\sqrt{x+1} - 1}{x} = \frac{\sqrt{1+x} - 1}{x} \cdot \frac{\sqrt{1+x} + 1}{\sqrt{1+x} + 1}$$

$$= \frac{1+x-1}{(\sqrt{1+x}+1)x} = \frac{x}{x(\sqrt{1+x}+1)}$$

$$\Rightarrow \lim_{x \rightarrow 0} \frac{\sqrt{1+x} - 1}{x} = \lim_{x \rightarrow 0} \frac{1}{\sqrt{1+x} + 1}$$

$$= \frac{1}{1+1} = \frac{1}{2}$$

$$11. \quad \lim_{x \rightarrow 0} \frac{e^{\sin x} - 1}{x}$$

$$A. \quad \lim_{x \rightarrow 0} \frac{\frac{e^{\sin x} - 1}{\sin x}}{\frac{x}{x \sin x}}$$

$$\lim_{x \rightarrow 0} \frac{e^{\sin x} - 1}{\sin x} \quad \lim_{x \rightarrow 0} \frac{\sin x}{x}$$

$$\lim_{\sin x \rightarrow 0} \frac{e^{\sin x} - 1}{\sin x} \cdot 1$$

$$1 \cdot 1 = 1$$

$$12. \quad \lim_{x \rightarrow 0} \frac{\log(1+5x)}{x}$$

$$A. \quad \lim_{x \rightarrow 0} \log(1+5x)^{\frac{1}{5x}} \quad \boxed{m \log x = \log x^m}$$

$$\log \left( \lim_{5x \rightarrow 0} (1+5x)^{\frac{1}{5x}} \right)^5$$

$$\begin{aligned} & \log_e e^5 \\ &= 5 \log_e e \quad \boxed{\log_e e = 1} \\ &= 5 \end{aligned}$$

13. Show that  $\lim_{x \rightarrow 0} \left( \frac{2|x|}{x} + x + 1 \right) = 3$

A.  $x \rightarrow 0+ \Rightarrow x > 0$

$$|x| = x$$

$$\lim_{x \rightarrow 0} \left( \frac{2|x|}{x} + x + 1 \right)$$

$$= \lim_{x \rightarrow 0} \left( \frac{2x}{x} + x + 1 \right)$$

$$= 2 + 0 + 1 = 3$$

Short Answer Questions (4 Marks)

1.  $\lim_{x \rightarrow 3} \frac{x^2 - 8x + 15}{x^2 - 9}$

A.  $\lim_{x \rightarrow 3} \frac{x^2 - 5x - 3x + 15}{(x-3)(x+3)}$

$$\lim_{x \rightarrow 3} \frac{(x-5)(\cancel{x-3})}{(\cancel{x-3})(x+3)}$$

$$= \frac{-2}{6}$$

$$= \frac{-1}{3}$$

2. Compute  $\lim_{x \rightarrow 0} \left[ \frac{(1+x)^{\frac{1}{3}} - (1-x)^{\frac{1}{3}}}{x} \right]$

A.  $\lim_{x \rightarrow 0} \frac{(1+x)^{\frac{1}{3}} - (1-x)^{\frac{1}{3}}}{x}$

$$= \lim_{(1+x) \rightarrow 1} \frac{(1+x)^{\frac{1}{3}} - 1}{(1+x) - 1} + \lim_{(1-x) \rightarrow 1} \frac{(1-x)^{\frac{1}{3}} - 1^{\frac{1}{3}}}{(1-x) - 1}$$

$$= \frac{1}{3} 1^{-2/3} + \frac{1}{3} 1^{-2/3} = \frac{1}{3} + \frac{1}{3} = \frac{2}{3}$$

$$3. \quad \lim_{x \rightarrow a} \frac{\sqrt{a+2x} - \sqrt{3x}}{\sqrt{3a+x} - 2\sqrt{x}}$$

$$A. \quad \lim_{x \rightarrow a} \frac{\sqrt{a+2x} - \sqrt{3x}}{\sqrt{3a+x} - 2\sqrt{x}} \times \frac{\sqrt{a+2x} + \sqrt{3x}}{\sqrt{a+2x} + \sqrt{3x}} \times \frac{\sqrt{3a+x} + 2\sqrt{x}}{\sqrt{3a+x} + 2\sqrt{x}}$$

$$\lim_{x \rightarrow a} \frac{(a+2x-3x)}{3a+x-4x} \cdot \frac{\sqrt{a+2x} + \sqrt{3x}}{\sqrt{3a+x} + 2\sqrt{x}}$$

$$\lim_{x \rightarrow a} \frac{a-x}{3(a-x)} \quad \lim_{x \rightarrow a} \frac{\sqrt{a+2x} + \sqrt{3x}}{\sqrt{3a+x} + 2\sqrt{x}}$$

$$\frac{1}{3} \cdot \frac{\cancel{\sqrt{3a}}}{\cancel{\sqrt{a}}} = \frac{1}{2\sqrt{3}}$$

$$4. \quad \lim_{x \rightarrow 0} \frac{\cos ax - \cos bx}{x^2}$$

$$A. \quad \lim_{x \rightarrow 0} \frac{-2 \sin \frac{ax+bx}{2} \sin \frac{ax-bx}{2}}{x^2}$$

$$\cos C - \cos D = -2 \sin \frac{C+D}{2} \sin \frac{C-D}{2}$$

$$-2 \lim_{x \rightarrow 0} \frac{\sin(a+b)x}{x} \cdot \frac{\sin(a-b)x}{x}$$

$$-2 \lim_{x \rightarrow 0} \frac{\sin(a+b)x}{\frac{(a+b)}{2}x} \left( \frac{a+b}{2} \right) \lim_{x \rightarrow 0} \frac{\sin(a-b)x}{\frac{(a-b)}{2}x} \left( \frac{a-b}{2} \right)$$

$$-2 \frac{a+b}{2} \lim_{x \rightarrow 0} \frac{\sin \left( \frac{a+b}{2} \right) x}{\left( \frac{a+b}{2} \right) x} \frac{(a+b)}{2} \frac{a-b}{2} \lim_{x \rightarrow 0} \frac{\sin \left( \frac{a-b}{2} \right) x}{\left( \frac{a-b}{2} \right) x} \left( \frac{a-b}{2} \right)$$

$$-2 \cdot 1 \left( \frac{a+b}{2} \right) \cdot 1 \frac{(a-b)}{2}$$

$$= \frac{(a+b)(a-b)}{2}$$

$$= \frac{b^2 - a^2}{2}$$

5.  $\lim_{x \rightarrow 0} \frac{1 - \cos 2mx}{\sin^2 nx} (m, n \in \mathbb{Z})$

A.  $\lim_{x \rightarrow 0} \frac{1 - \cos 2mx}{\sin^2 nx} = \lim_{x \rightarrow 0} \frac{2 \sin^2 mx}{\sin^2 nx}$

$$= 2 \frac{\left( \lim_{x \rightarrow 0} \frac{\sin mx}{mx} \right)^2}{\left( \lim_{x \rightarrow 0} \frac{\sin nx}{nx} \right)^2} = \frac{2m^2}{n^2}$$

6.  $\lim_{x \rightarrow 0} \frac{\sin(a + bx) - \sin(a - bx)}{x}$

A.  $\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$

$$\lim_{x \rightarrow 0} \frac{2 \cos \frac{a+bx+a-bx}{2} \sin \frac{(a+bx)-(a-bx)}{2}}{x}$$

$$2 \cos \frac{2a}{2} \lim_{x \rightarrow 0} \frac{\sin bx}{bx} \cdot b$$

$$2 \cos a \times b = 2b \cos a$$

## Differentiation

### Key concepts

- |  |   |
|--|---|
| <p>* <math>\frac{d}{dx}(uv) = u \frac{du}{dx} + v \frac{dv}{dx}</math></p> <p>* <math>\frac{d}{dx}(x^n) = nx^{n-1}</math></p> <p>* <math>\frac{d}{dx}(\sin x) = \cos x</math></p> <p>* <math>\frac{d}{dx}(\tan x) = \sec^2 x</math></p> <p>* <math>\frac{d}{dx}(\sec x) = \sec x \tan x</math></p> <p>* <math>\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}</math></p> <p>* <math>\frac{d}{dx}(\sec^{-1} x) = \frac{1}{ x \sqrt{x^2-1}}</math></p> <p>* <math>\frac{d}{dx}(e^x) = e^x</math></p> <p>* <math>\frac{d}{dx}(\log x) = \frac{1}{x}</math></p> <p>* <math>\frac{d}{dx}(\cosh x) = \sinh x</math></p> <p>* <math>\frac{d}{dx}(\coth x) = -\operatorname{cosech}^2 x</math></p> <p>* <math>\frac{d}{dx}(\operatorname{cosech} x) = -\operatorname{cosech} x \coth x</math></p> <p>* <math>\frac{d}{dx}(\cosh^{-1} x) = \frac{1}{\sqrt{x^2-1}}</math></p> <p>* <math>\frac{d}{dx}(\coth^{-1} x) = \frac{1}{1-x^2}</math></p> <p>* <math>\frac{d}{dx}(\operatorname{cosech}^{-1} x) = \frac{-1}{ x \sqrt{x^2+1}}</math></p> | <p>* <math>\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}</math></p> <p>* <math>\frac{d}{dx}(x) = 1</math></p> <p>* <math>\frac{d}{dx}(\cos x) = -\sin x</math></p> <p>* <math>\frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x</math></p> <p>* <math>\frac{d}{dx}(\operatorname{cosec} x) = -\operatorname{cosec} x \cot x</math></p> <p>* <math>\frac{d}{dx}(\cot^{-1} x) = \frac{-1}{1+x^2}</math></p> <p>* <math>\frac{d}{dx}(\operatorname{cosec}^{-1} x) = \frac{-1}{ x \sqrt{x^2-1}}</math></p> <p>* <math>\frac{d}{dx}(a^x) = a^x \log a</math></p> <p>* <math>\frac{d}{dx}(\sinh x) = \cosh x</math></p> <p>* <math>\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x</math></p> <p>* <math>\frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \tanh x</math></p> <p>* <math>\frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{1+x^2}}</math></p> <p>* <math>\frac{d}{dx}(\tanh^{-1} x) = \frac{1}{1-x^2}</math></p> <p>* <math>\frac{d}{dx}(\operatorname{sech}^{-1} x) = \frac{-1}{ x \sqrt{1-x^2}}</math></p> |
|--|---|



## Very Short Answer Questions (2 Marks)

1.  $y = \log(\sin(\log x))$ , then find  $\frac{dy}{dx}$

A. Differentiate both sides with respect to 'x'

$$v = \log x, u = \sin v \quad y = \log u$$

$$\frac{dy}{du} = \frac{1}{u}; \frac{du}{dv} = \cos v; \frac{dv}{dx} = \frac{1}{x}$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}$$

$$= \frac{1}{\sin(\log x)} \cdot \cos(\log x) \cdot \frac{1}{x} = \frac{\cot(\log x)}{x}$$

2.  $f(x) = \log(\sec x + \tan x)$ , then find  $f'(x)$

A. Differentiate both sides with respect to 'x'

$$u = \sec x + \tan x \quad \text{and} \quad y = \log u$$

$$\frac{dy}{du} = \frac{1}{u}, \frac{du}{dx} = \sec x \cdot \tan x + \sec^2 x$$

$$= \sec x(\sec x + \tan x)$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

$$= \frac{1}{\sec x + \tan x} \cdot \sec x(\sec x + \tan x) = \sec x$$

3. Find the derivatives of the following functions.

A. (i)  $\frac{d}{dx} x \tan^{-1} x = x \frac{d}{dx} \tan^{-1} x + \tan^{-1} x \frac{d}{dx} x$

$$= \frac{x}{1+x^2} + \tan^{-1} x$$

(ii)  $\frac{d}{dx} \tan^{-1}(\log x)$

$$= \frac{1}{1+(\log x)^2} \cdot \frac{d}{dx}(\log x)$$

$$= \frac{1}{x[1+(\log x)^2]}$$

(iii)  $\frac{d}{dx} e^{\sin^{-1} x}$

$$= e^{\sin^{-1} x} \frac{d}{dx} \sin^{-1} x$$

$$= \frac{e^{\sin^{-1} x}}{\sqrt{1-x^2}}$$

$$(iv) x = e^{\sinh y}$$

$$\frac{dx}{dy} = \frac{d}{dy} e^{\sinh y}$$

$$\frac{dx}{dy} = e^{\sinh y} \frac{d}{dy} \cosh y$$

$$\frac{dx}{dy} = e^{\sinh y} \cosh y$$

$$\Rightarrow \frac{dx}{dy} = \frac{1}{e^{\sinh y} \cosh y}$$

$$(v) \frac{d}{dx} \sin(\cos(x^2))$$

$$\cos(\cos(x^2)) \frac{d}{dx} \cos x^2$$

$$\cos(\cos(x^2)) (-\sin x^2) \frac{d}{dx} x^2$$

$$\cos(\cos(x^2)) \sin(x^2) 2x$$

4.  $f(x) = 1 + x + x^2 + \dots + x^{100}$ , then find  $f'(1)$

A. Differentiate both sides with respect to 'x'

$$f'(x) = 1 + 2x + 3x^2 \dots + 100x^{99}$$

$$f'(1) = 1 + 2 + 3 \dots + 100$$

$$= \frac{100 \times 101}{2} = 5050 \left( \sum x = \frac{x(x+1)}{2} \right)$$

5.  $f(x) = xe^x \sin x$  then find  $f'(x)$

A. Differentiate both sides with respect to 'x'

$$\frac{d}{dx} f(x) = \frac{d}{dx} xe^x \sin x$$

$$f'(x) = xe^x \frac{d}{dx} \sin x + x \sin x \frac{d}{dx} e^x \sin x \frac{dx}{dx}$$

$$= xe^x \cos x + x \sin x e^x + e^x \sin x$$

6.  $f(x) = e^x, g(x) = \sqrt{x}, g(x)$  Find the derivatives of the  $f(x)$  with respect to  $g(x)$

A.  $f(x) = e^x, g(x) = 5x$

$$\Rightarrow \frac{d}{dx} f(x) = \frac{d}{dx} e^x$$

$$\frac{d}{dx}g(x) = \frac{d}{dx}\sqrt{x} = \frac{1}{2\sqrt{x}}$$

$$\therefore \frac{df(x)}{dg(x)} = \frac{\frac{df(x)}{dx}}{\frac{dg(x)}{dx}} = \frac{e^x}{\frac{1}{2\sqrt{x}}} = 2\sqrt{x}e^x$$

7.  $\frac{2}{x^3} + y^2 = a^2$  then show that  $\frac{dy}{dx} = -\sqrt[3]{\frac{y}{x}}$

A. Differentiate both sides with respect to 'x'

$$\frac{2}{3}x^{\frac{2}{3}-1} + \frac{2}{3}y^{\frac{2}{3}-1} \frac{dy}{dx} = 0$$

$$x^{-\frac{1}{3}} + y^{-1/3} \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\frac{x^{-1/3}}{y^{-1/3}} = -\sqrt[3]{\frac{y}{x}}$$

8.  $y = \sin^{-1}\sqrt{x}$  then find  $\frac{dy}{dx}$

A.  $\frac{dy}{dx} = \frac{1}{\sqrt{1-x}} \cdot \frac{1}{2\sqrt{x}}$

$$\frac{dy}{dx} = \frac{1}{2\sqrt{x(1-x)}}$$

9.  $x = a\cos^3 t$ ,  $y = a\sin^3 t$  then find  $\frac{dy}{dx}$

A. Differentiate both sides with respect to 'x'

$$\frac{dx}{dt} = -3a\cos^2 t \sin t \quad \frac{dy}{dt} = 3\sin^2 t \cos t$$

$$\frac{dy}{dx} = \frac{dy}{dt} \frac{dt}{dx}$$

$$= \frac{3a\sin^2 t \cos t}{-3a\cos^2 t \sin t}$$

$$= -\tan t$$

## Short Answer Questions (4 Marks)

1. Find the derivatives of the following functions from the first principles.

A. (i)  $f(x) = \sin 2x$

$$f(x+h) = \sin 2(x+h) = \sin(2x+2h)$$

First principle

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\sin(2x+2h) - \sin 2x}{h}$$

$$\boxed{\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}}$$

$$= \lim_{h \rightarrow 0} 2 \frac{\cos \left( \frac{2x+2h+2h}{2} \right) \sin \left( \frac{2x+2h-2x}{2} \right)}{h}$$

$$= 2 \lim_{h \rightarrow 0} \cos \frac{4x+2h}{2} \lim_{h \rightarrow 0} \frac{\sin \frac{2h}{2}}{h}$$

$$= 2 \cos \frac{4x}{2} \cdot 1$$

$$\therefore \frac{d}{dx} \sin 2x = 2 \cos 2x$$

(ii)  $f(x) = \tan 2x$

$$f(x+h) = \tan 2(x+h) = \tan(2x+2h)$$

First principle

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\tan(2x+2h) - \tan 2x}{h}$$

$$\lim_{h \rightarrow 0} \frac{\frac{\sin(2x+2h)}{\cos(2x+2h)} - \frac{\sin 2x}{\cos 2x}}{h}$$

$$\lim_{h \rightarrow 0} \frac{\frac{\sin(2x+2h)\cos 2x - \cos(2x+2h)\sin 2x}{\cos(2x+2h)\cos 2x}}{h}$$

$$\lim_{h \rightarrow 0} \frac{\sin[2x + 2h - 2h]}{h \cos(2x + 2h) \cos 2x}$$

$$\lim_{2h \rightarrow 0} \frac{\sin 2h}{2h} \times 2 \lim_{h \rightarrow 0} \frac{1}{\cos(2x + 2h) \cos 2x}$$

$$1 \times 2 \frac{1}{\cos^2 2x} \qquad 2 \sec^2 2x$$

(iii)  $f(x) = ax^2 + bx + c$

First principle

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{a[x+h]^2 + b(x+h) - (ax^2 + bx + c)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{a[x^2 + 2hx + h^2] + b[x+h] - (ax^2 + bx + c)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{ax^2 + 2hax + ah^2 + bx + bh + c - ax^2 - bx - c}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h(2x + b + ah)}{h} \qquad \therefore \frac{d}{dx} f(x) = 2ax + b$$

2.  $x^3 + y^3 - 3axy = 0$  then find  $\frac{dy}{dx}$

A. Differentiate both sides with respect to 'x'

$$\frac{dy}{dx} [x^3 + y^3 - 3axy] = 0$$

$$3x^2 + 3y^2 \frac{dy}{dx} - 3a \left[ x \frac{dy}{dx} + y \frac{d}{dx} x \right] = 0$$

$$x^2 + y^2 \frac{dy}{dx} - ax \frac{dy}{dx} - ay = 0$$

$$(y^2 - ax) \frac{dy}{dx} = ay - x^2$$

$$\therefore \frac{dy}{dx} = \frac{ay - x^2}{y^2 - ax}$$

3.  $y = e^{a \sin^{-1} x}$  then show that  $\frac{dy}{dx} = \frac{ay}{\sqrt{1-y^2}}$

A.  $y = e^{a \sin^{-1} x} \dots\dots(1)$

Differentiate both sides with respect to 'x'

$$\frac{dy}{dx} = \frac{d}{dx} e^{a \sin^{-1} x}$$

$$= e^{a \sin^{-1} x} \frac{d}{dx} a \sin^{-1} x$$

$$\frac{dy}{dx} = e^{a \sin^{-1} x} \frac{a}{\sqrt{1-x^2}}$$

$$\frac{dy}{dx} = \frac{ya}{\sqrt{1+x^2}} \text{ from (1)}$$

4. Find the derivative of  $\tan^{-1} \left( \frac{a-x}{1+ax} \right)$

A. put  $a = \tan A \Rightarrow A = \tan^{-1} a$

$$= \tan B \Rightarrow B = \tan^{-1} x$$

$$\frac{d}{dx} \left( \tan^{-1} \left( \frac{\tan A - \tan B}{1 + \tan A \tan B} \right) \right)$$

$$\frac{d}{dx} \left( \tan^{-1} (\tan(A-B)) \right)$$

$$\frac{d}{dx} (A-B)$$

$$\frac{d}{dx} (\tan^{-1} a - \tan^{-1} x)$$

$$0 - \frac{1}{1+x^2}$$

$$= -\frac{1}{1+x^2}$$

5.  $x^y = e^{x-y}$  then show that  $\frac{dy}{dx} = \frac{\log x}{(1+\log x)^2}$

A.  $x^y = e^{x-y}$

take log both sides

$$\log_e x^y = \log_e e^{x-y}$$

$$y \log x = (x - y) \log_e e$$

$$x = y + y \log x$$

$$x = y(1 + \log x)$$

$$y = \frac{x}{(1 + \log x)}$$

Differentiate both sides with respect to 'x'

$$\frac{dy}{dx} = \frac{d}{dx} \frac{x}{1 + \log x}$$

$$= \frac{(1 + \log x)1 - x \left( \frac{1}{x} \right)}{(1 + \log x)^2}$$

$$\therefore \frac{dy}{dx} = \frac{1 + \log x - 1}{(1 + \log x)^2}$$

6.  $\sin y = x \sin(a + y)$  then show that  $\frac{dy}{dx} = \frac{\sin^2(a + y)}{\sin a}$

A.  $x = \frac{\sin y}{\sin(a + y)}$

Differentiate both sides with respect to 'y'

$$\frac{dx}{dy} = \frac{\sin(a + y) \cos y - \sin y \cos(a + y)}{\sin^2(a + y)}$$

$$\frac{dx}{dy} = \frac{\sin[a + y - y]}{\sin^2(a + y)}$$

$$\therefore \frac{dy}{dx} = \frac{\sin^2(a + y)}{\sin a}$$

### Long Answer Questions (8 Marks)

1.  $x = \frac{3at}{1 + t^3}$   $y = \frac{3a t^2}{1 + t^3}$  then find  $\frac{dy}{dx}$

A. Differentiate both sides with respect to 't'

$$\frac{d}{dt} x = 3a \frac{d}{dt} \left[ \frac{t}{1 + t^3} \right]$$

$$= 3a \left[ \frac{(1 + t^3) \frac{dt}{dt} - t \frac{d}{dt} (1 + t^3)}{(1 + t^3)^2} \right]$$

$$\begin{aligned}
&= 3a \left[ \frac{(1+t^3) - t \cdot 3t^2}{(1+t^3)^2} \right] \\
&= 3a \left[ \frac{1-2t^3}{(1+t^3)^2} \right] \\
\frac{dy}{dt} &= 3a \left[ \frac{(1+t^3) \frac{d}{dt} t^2 - t^2 \frac{d}{dt} (1+t^3)}{(1+t^3)^2} \right] \\
&= 3a \left[ \frac{(1+t^3)2t - t^2 \cdot 3t^2}{(1+t^3)^2} \right] \\
&= 3a \left[ \frac{2t + 2t^4 - 3t^4}{(1+t^3)^2} \right] \\
&= 3a \left[ \frac{2t - t^4}{(1+t^3)^2} \right] \\
\therefore \frac{dy}{dx} &= \frac{dy}{dt} \cdot \frac{dt}{dx} \\
&= 3a \left[ \frac{2t - t^4}{(1+t^3)^2} \right] \frac{(1+t^3)^2}{3a(1-2t^3)} \\
&= \frac{t(2-t^3)}{3(1-2t^3)}
\end{aligned}$$

2.  $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$ , then show that  $\frac{dy}{dx} = \frac{\sqrt{1-y^2}}{\sqrt{1-x^2}}$

A. Let  $x = \sin A$      $y = \sin B \Rightarrow A = \sin^{-1} x \Rightarrow B = \sin^{-1} y$

$$\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$$

Put  $x = \sin \theta, y = \sin \phi$

$$\therefore \sqrt{1-\sin^2 \theta} + \sqrt{1-\sin^2 \phi} = a(\sin \theta - \sin \phi)$$

$$\cos \theta + \cos \phi = a(\sin \theta - \sin \phi)$$

$$2 \cos \frac{\theta + \phi}{2} \cdot \cos \frac{\theta - \phi}{2}$$

$$= \left[ 2 \cos \frac{\theta + \phi}{2} \sin \frac{\theta - \phi}{2} \right]$$



$$\therefore \cos \frac{\theta - \phi}{2} = a \cdot \sin \frac{\theta - \phi}{2}$$

$$\tan \frac{\theta - \phi}{2} = \frac{1}{2}; \frac{\theta - \phi}{2} = \tan^{-1} \left( \frac{1}{2} \right)$$

$$\phi = \theta - 2 \tan^{-1} \left( \frac{1}{a} \right);$$

$$\sin^{-1} y = \sin^{-1} x - 2 \tan^{-1} \left( \frac{1}{a} \right)$$

Differentiating w.r.to x

$$\frac{1}{\sqrt{1-y^2}} \cdot \frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}} \Rightarrow \frac{dy}{dx} = \frac{\sqrt{1-y^2}}{\sqrt{1-x^2}}$$

3.  $\tan^{-1} \left( \frac{2x}{1-x^2} \right) + \tan^{-1} \frac{3x-x^3}{1-3x^2} - \tan^{-1} \left( \frac{4x-4x^3}{1-6x^2+x^4} \right)$  then show that  $\frac{dy}{dx} = \frac{1}{1+x^2}$

A. Let  $x = \tan \theta \Rightarrow \theta = \tan^{-1} x$

$$y = \tan^{-1} \left( \frac{2 \tan \theta}{1 - \tan^2 \theta} \right) + \tan^{-1} \left( \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \right)$$

$$- \tan^{-1} \left( \frac{4 \tan \theta - 4 \tan^3 \theta}{1 - 6 \tan^2 \theta + \tan^4 \theta} \right)$$

$$= \tan^{-1} (\tan 2\theta) + \tan^{-1} (\tan 3\theta)$$

$$y = 2\theta + 3\theta - 4\theta$$

$$y = \theta$$

$$y = \tan^{-1} x$$

Differentiate both sides with respect to 'x'

$$\frac{dy}{dx} = \frac{1}{1+x^2}$$

4.  $y = \frac{x^3 \sqrt{2+3x}}{(2+x)(1-x)}$  then find  $\frac{dy}{dx}$

A. Take 'log' both sides

$$\log y = \log \frac{x^3 \sqrt{2+3x}}{(2+x)(1-x)}$$

$$= \log x^3 + \log \sqrt{2+3x} - \log(2+x) - \log(1-x)$$

$$\log y = 3 \log x + \frac{1}{2} (2+3x) - \log(2+x) - \log(1-x)$$

Differentiate both sides with respect to 'x'

$$\frac{1}{y} \frac{dy}{dx} = \frac{3}{x} + \frac{3}{2(2+3x)} - \frac{1}{2+x} + \frac{1}{1+x}$$

$$\therefore \frac{dy}{dx} = y \left[ \frac{3}{x} + \frac{3}{2(2+3x)} - \frac{1}{2+x} + \frac{1}{1+x} \right]$$

5.  $y = x\sqrt{a^2 + x^2} + a^2 \log(x + \sqrt{a^2 + x^2})$  then show that  $\frac{dy}{dx} = 2\sqrt{a^2 + x^2}$

A. Differentiate both sides with respect to 'x'

$$\frac{dy}{dx} = x \frac{1}{2\sqrt{a^2 + x^2}} \cdot 2x + \sqrt{a^2 + x^2} + \frac{a^2 \left[ 1 + \frac{1}{2\sqrt{a^2 + x^2}} \cdot 2x \right]}{x + \sqrt{a^2 + x^2}}$$

$$= \frac{x^2}{\sqrt{a^2 + x^2}} + \frac{a^2 (\sqrt{a^2 + x^2} + x)}{(\sqrt{a^2 + x^2} + x) \sqrt{a^2 + x^2}}$$

$$= \frac{x^2 + a^2}{\sqrt{a^2 + x^2}} + \sqrt{a^2 + x^2}$$

$$= \sqrt{a^2 + x^2} + \sqrt{a^2 + x^2}$$

$$= 2\sqrt{a^2 + x^2}$$

6.  $x^{\log y} = \log x$  then show that  $\frac{dy}{dx} = \frac{y}{x} \left[ \frac{1 - \log x \log y}{(\log x)^2} \right]$

A. Take 'log' both sides

$$\log x^{\log y} = \log(\log x)$$

$$\log y \log x = \log(\log x)$$

Differentiate both sides with respect to 'x'

$$(\log y) \frac{1}{x} + \frac{\log x}{y} \frac{dy}{dx} = \frac{1}{\log x} \cdot \frac{1}{x}$$

$$\frac{\log x}{y} \frac{dy}{dx} = \frac{1}{x \log x} - \frac{\log y}{x}$$

$$\frac{\log x}{y} \frac{dy}{dx} = \frac{1 - \log x \log y}{x(\log x)}$$

$$\therefore \frac{dy}{dx} = \frac{y}{x} \left[ \frac{1 - \log x \log y}{(\log x)^2} \right]$$

## Errors and approximations

### Key concepts

1. A small change in  $x$  is  $\Delta x$
2. If  $x$  is changed as  $x + \Delta x$  then change in  $y$   $\Delta y = f(x + \Delta x) - f(x)$
3. Differential in  $y$   $dy = f'(x) \cdot \Delta x$
4. Relative error in  $y = \frac{\Delta y}{y}$
5. Percentage error in  $y = \frac{\Delta y}{y} \times 100$

### Short Answer Questions (4 Marks)

1. Find  $dy, \Delta y$  for the following  $y = x^2 + 3x + 6, x = 10, \Delta x = 0.01$

**Sol:**  $\Delta y = f(x + \Delta x) - f(x)$

$$\Delta y = f(10 + 0.01) - f(10)$$

$$\Delta y = f(10.01) - f(10)$$

$$= (10.01)^2 + 3 \times 10.01 + 6 - (10^2 + 3 \times 10 + 6)$$

$$= 100.2001 + 30.03 + 6 - 100 - 30 - 6$$

$$= 130.2301 - 130$$

$$= 0.2301$$

$$\Delta y = f'(x) \Delta x$$

$$= (2x + 3) \Delta x$$

$$= (2 \times 10 + 3) \times 0.01$$

$$= 23 \times 0.01$$

$$= 0.23$$

2. Find  $dy, \Delta y$  for the following function  $y = \cos x$   $x = 60^\circ$ ,  $\Delta x = 1^\circ$

Sol:  $x = 60^\circ$ ,  $\Delta x = 1^\circ$

$$\Delta y = f(x + \Delta x) - f(x)$$

$$\Delta y = f(60^\circ + 1^\circ) - f(60^\circ)$$

$$= f(61^\circ) - f(60^\circ)$$

$$= \cos 61^\circ - \cos 60^\circ$$

$$= 0.4848 - 0.5 = -0.0152$$

$$\Delta y = f'(x)\Delta x$$

$$= -\sin x \cdot \Delta x$$

$$= -\sin 60^\circ \times 1^\circ$$

$$= -0.866 \times 0.0174 \quad (1^\circ = 0.0174 \text{ ರೇಡಿಯನ್})$$

$$= -0.0150$$

3. Find  $dy, \Delta y$  for the following function  $y = e^x + x$ ,  $x = 5$ ,  $\Delta x = 0.02$

Sol:  $\Delta y = f(x + \Delta x) - f(x)$

$$= f(5 + 0.02) - f(5)$$

$$= f(5.02) - f(5)$$

$$= e^{5.02} + 5.02 - e^5 - 5$$

$$= e^{5.02} - e^5 + 0.02$$

$$\Delta y = f'(x)\Delta x$$

$$= (e^x + 1) \cdot \Delta x$$

$$= (e^5 + 1)(0.02)$$

4. Find  $dy, \Delta y$  for the following function  $y = 5x^2 + 6x + 6$ ,  $x = 2$ ,  $\Delta x = 0.001$

Sol:  $\Delta y = f(x + \Delta x) - f(x)$

$$= f(2 + 0.001) - f(2)$$

$$= f(2.001) - f(2)$$

$$= 5(2.001)^2 + 6 \times 2.001 + 6 - (5 \times 2^2 + 6 \times 2 + 6)$$

$$= 5(4.004001) + 12.006 + 6 - 20 - 12 - 6$$

$$= 0.026005$$

$$\Delta y = f'(x)\Delta x$$

$$= (10x + 6) \cdot \Delta x$$

$$= (10 \times 2 + 6)(0.001)$$

$$= 26 \times 0.001 = 0.026$$

5. If the increase in the side of a square is 2%, find the change in the area of the square.

**Sol:** Let side of the square =  $x$

Area of the square  $A = x^2$

$$\text{given } \frac{\Delta x}{x} \times 100 = 2 \quad \dots\dots(1)$$

$$\Rightarrow \frac{\Delta A}{dx} = 2x \Rightarrow \frac{\Delta A}{A} = \frac{2x}{x^2}$$

$$\Rightarrow \frac{\Delta A}{A} \times 100 = 2 \times \left( \frac{\Delta x}{dx} \times 100 \right)$$

$$= 2 \times 2 \text{ from (1)}$$

$$= 4$$

6. If the increase in the side of the square is 4% find the change in the area of the square.

**Sol:** Let side of the square =  $x$

Area of the square  $A = x^2$

$$\frac{\Delta x}{x} \times 100 = 4 \text{ (given)}$$

$$A = x^2 \Rightarrow \frac{\Delta A}{dx} = 2x$$

$$\Rightarrow \frac{\Delta A}{A} \times 100 \times \Delta x = \frac{2x}{x^2} \times 100 \times \Delta x$$

$$= 2 \times \frac{\Delta x}{x} \times 100$$

$$= 2 \times 4 \text{ from (1)}$$

$$= 8$$

7. The radius of the sphere is measured is 14 c.m., Later it was found that there is an error 0.02cm in measuring the radius, find the approximate error in surface area of the sphere.

**Sol:** Let radius of the sphere =  $r$

$$\text{Surface Area } A = 4\pi r^2 \Rightarrow \frac{dA}{dr} = 4\pi \times 2r$$

$$dA = \frac{dA}{dr} \times \Delta r$$

$$\begin{aligned}
 &= 4\pi \times 2r \times \Delta r \\
 &= 8 \times \frac{22}{7} \times 14 \times 0.02 \\
 &= 7.04 \text{ sq.cm.}
 \end{aligned}$$

### Geometrical interpretation of the derivative

#### Key concepts

1. Slope of the tangent at point  $(x, y)$  on the curve  $y = f(x)$  is  $\frac{dy}{dx}$
2. Equation of tangent at point  $(a, b)$  on the curve  $y = f(x)$  is  $y - b = m(x - a)$   $\left(m = \frac{dy}{dx}\right)$
3. Slope of the normal =  $-\frac{1}{m}$
3. Equation of the normal  $y - b = -\frac{1}{m}(x - a)$

1. Find the slope of the tangents to the curve  $y = 3x^2 - x^3$ , where it meets the X-axis.

**Sol:**

$$y = 3x^2 - x^3 \text{ --- (1)}$$

$$\text{Equation of } x\text{-axis } y = 0 \text{ ---- (2)}$$

from (1) and (2)

$$3x^2 - x^3 = 0 \quad (\because y = 0)$$

$$x^2(3 - x) = 0$$

$$x^2 = 0 \quad \text{or} \quad 3 - x = 0$$

$$x = 0 \quad \text{or} \quad x = 3$$

$\therefore$  The given curve intersecting x-axis at points  $(0, 0)$ ,  $(3, 0)$

$$\begin{aligned}
 m &= \frac{dy}{dx}_{(0,0)} \\
 &= 6x - 3x^2 \\
 &= 6 \times 0 - 3 \times 0 \\
 &= 0
 \end{aligned}$$

Equation of tangent at  $(0, 0)$

$$\begin{aligned}
 y - 0 &= 0(x - 0) \\
 \Rightarrow y &= 0
 \end{aligned}$$

$$\begin{aligned}
 \text{Slope of tangent } M &= \frac{dy}{dx}_{(3,0)} \\
 &= 6x - 3 \times 2 \\
 &= 6 \times 3 - 3 \times 3^2 \\
 &= 18 - 27 = -9
 \end{aligned}$$

Equation of tangent at (3, 0)

$$\begin{aligned}
 y - 0 &= -9(x - 3) \\
 y &= -9x + 27 \\
 9x + y - 27 &= 0
 \end{aligned}$$

2. Show that the area of the triangle formed by the tangent at any point on the curve  $xy = C$  ( $C \neq 0$ ) with the co-ordinate axes is constant.

**Sol:** Let  $P(x_1, y_1)$  be the point on the curve  $xy = c$  and  $x_1 \neq 0, y_1 \neq 0$ .

$$y = \frac{c}{x}$$

Defferentiate both sides

$$\frac{dy}{dx} = -\frac{c}{x^2}$$

$$\text{Slope of tangent } m = -\frac{c}{x_1^2} \text{ (at } P(x_1, y_1))$$

$P(x_1, y_1)$  Equation of the tangent at P

$$y - y_1 = -\frac{c}{x_1^2} (x - x_1)$$

$$y \cdot x_1^2 - y_1 x_1^2 = -cx + cx_1$$

$$cx + yx_1^2 = cx_1 + y_1 x_1^2$$

$$cx + yx_1^2 = cx_1 + y_1 x_1 x_1$$

$$cx + yx_1^2 = cx_1 + cx_1 \quad (\because x_1 y_1 = c)$$

$$cx + yx_1^2 = 2cx_1$$

$$cx + yx_1^2 - 2cx_1 = 0$$

$$\text{area of the triangle} = \frac{1}{2} \left| \frac{c^2}{ab} \right|$$

$$= \frac{1}{2} \left| \frac{(-2c_1)^2}{c \times x_1^2} \right|$$

$$= \frac{1}{2} \left| \frac{4c^2 x_1^2}{c^2 \times x_1^2} \right| = 2c$$

= which is constant

**3. Find the slope of the tangent to the curve  $y = 3x^4 - 4x$  at  $x = 4$**

**Sol:**  $y = 3x^4 - 4x$

Differentiate both sides

$$\frac{dy}{dx} = 3 \times 4x^3 - 4$$

$$\begin{aligned} \text{slope of the tangent at } x = 4 \quad m &= 12 \times 4^3 - 4 \\ &= 12 \times 64 - 4 \\ &= 768 - 4 = 764 \end{aligned}$$

**4(i). Find the slope of the tangent to the curve  $y = x^3 - 3x + 2$  at the point whose  $x$  coordinate is 3.**

**Sol:**  $y = x^3 - 3x + 2$

Differentiate both sides

$$\frac{dy}{dx} = 3x^2 - 3$$

$$\begin{aligned} \text{slope of the tangent at } x = 3 \quad m &= 3 \times 3^2 - 3 \\ &= 27 - 3 = 24 \end{aligned}$$

**4(ii). Find the slope of the normal to the curve  $x = a \cos^3 \theta$ ,  $y = a \sin^3 \theta$  at  $\theta = \frac{\pi}{4}$**

**Sol:**  $x = a \cos^3 \theta$ ,  $y = a \sin^3 \theta$

Differentiating with respect to  $\theta$

$$\frac{dx}{d\theta} = a3 \cos^2 \theta (-\sin \theta)$$



$$\frac{dy}{d\theta} = a \times 3 \sin^2 \theta (\cos \theta)$$

$$\text{slope of the tangent } (m) \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{3a \sin^2 \theta \cos \theta}{-3a \cos^2 \theta \cdot \sin \theta} = -\tan \theta = -\tan \frac{\pi}{4}$$

$$m = -1$$

$$\therefore \text{slope of the normal} = -\frac{1}{m} = -\frac{-1}{-1} = 1$$

5. Find the slope of the normal to the curve  $x = 1 - a \sin \theta$ ,  $y = b \cos^3 \theta$ , at  $\theta = \frac{\pi}{4}$

**Sol:**  $x = 1 - a \sin \theta$ ,  $y = b \cos^3 \theta$

differentiating with respect to  $\theta$

$$\frac{dx}{d\theta} = -a \cos \theta, \quad \frac{dy}{d\theta} = 2b \cos \theta (-\sin \theta)$$

$$\text{slope of the tangent (at } \theta = \frac{\pi}{2} \text{)} \quad m = \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{-2b \cos \theta \sin \theta}{-a \cos \theta}$$

$$= \frac{2b}{a} \sin \frac{\pi}{2}$$

$$m = \frac{2b}{a}$$

$$\text{slope of the normal at } \theta = \frac{\pi}{2} = -\frac{1}{m}$$

$$= -\frac{1}{2b/a}$$

$$= \frac{-a}{2b}$$

6. Find the equation of tangent and normal to the curve  $y = x^2 - 4x + 2$  at point (4, 2)

**Sol:**  $y = x^2 - 4x + 2$

differentiating with respect to  $x$

$$\frac{dy}{dx} = 2x - 4$$

$$\text{slope of the tangent at } (4, 2) \quad m = 2 \times 4 - 4 = 4$$

equation of the tangent

$$y - y_1 = m(x - x_1)$$

$$y - 2 = 4(x - 4)$$

$$y - 2 = 4x - 16$$

$$4x - y - 16 + 2 = 0$$

$$4x - y - 14 = 0$$

$$\text{slope of the normal} = -\frac{1}{m} = -\frac{1}{4}$$

$$\text{equation of the normal } y - 2 = -\frac{1}{4}(x - 4)$$

$$4y - 8 = -x + 4$$

$$x + 4y - 8 - 4 = 0$$

$$x + 4y - 12 = 8.$$

7. Find the equation of tangent and normal to the curve  $y = x^3 + 4x^2$  at  $(-1, 3)$

**Sol:**  $y = x^3 + 4x^2$

differentiating with respect to x

$$\frac{dy}{dx} = 3x^2 + 8x$$

$$\begin{aligned} \text{slope of the tangent at } (-1, 3) \quad m &= 3 \times (-1)^2 + 8(-1) \\ &= 3 - 8 \\ &= -5 \end{aligned}$$

$$\text{equation of the tangent} = y - 3 = -5(x + 1)$$

$$y - 3 = -5x - 5$$

$$5x + y + 2 = 0$$

$$\text{slope of the normal} = -\frac{1}{m}$$

$$= \frac{-1}{-5} = \frac{1}{5}$$

$$\text{equation of the normal } y - 3 = \frac{1}{5}(x + 1)$$

$$5y - 15 = x + 1$$

$$x - 5y - 16 = 0$$

8. Find the tangent and normal to the curve  $y = 2e^{-\frac{x}{3}}$  at the point where the curve meets y-axis.

Sol: Equation of y-axis  $x = 0$

$$y = 2e^{-\frac{x}{3}}, \text{ the curve meets y-axis at } (0, 2) \quad [\because y = 2e^{-\frac{x}{3}} = 2]$$

$$y = 2e^{-\frac{x}{3}}$$

differentiating with respect to x

$$\begin{aligned} \frac{dy}{dx} &= 2e^{-x/3} \times \left(-\frac{1}{3}\right) \\ &= -\frac{2}{3} \cdot e^{-x/3} \end{aligned}$$

$$\text{slope of the tangent (at } x = 0) \quad m = -\frac{2}{3} \cdot e^{-0/3} = -\frac{2}{3}$$

equation of the tangent at (0, 2)

$$y - 2 = -\frac{2}{3}(x - 0)$$

$$3y - 6 = -2x$$

$$2x + 3y - 6 = 0$$

$$\text{slope of the normal} = -\frac{1}{m} = \frac{-1}{-\frac{2}{3}} = \frac{3}{2}$$

$$\text{equation of the normal } y - 2 = \frac{3}{2}(x - 0)$$

$$2y - 4 = 3x$$

$$3x - 2y + 4 = 0$$

9. If the tangent at any point on the curve  $x^{2/3} + y^{2/3} = a^{2/3}$  intersects the co-ordinate axes in A and B then show that the length of AB is constant.

Sol: P  $(x_1, y_1)$  be the any point on the curve  $x^{2/3} + y^{2/3} = a^{2/3}$

$$x_1^{2/3} + y_1^{2/3} = a^{2/3} \quad \dots (1)$$

$$x^{2/3} + y^{2/3} = a^{2/3}$$

differentiating both sides with respect to 'x'

$$\frac{2}{3} \cdot x^{-\frac{1}{3}} + \frac{2}{3} \cdot y^{-\frac{1}{3}} \cdot \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\frac{y^{1/3}}{x^{1/3}}$$

slope of the tangent at P(x<sub>1</sub>, y<sub>1</sub>)

$$m = -\left(\frac{y_1}{x_1}\right)^{1/3}$$

equation of the tangent at P(x<sub>1</sub>, y<sub>1</sub>)

$$y - y_1 = -\left(\frac{y_1}{x_1}\right)^{1/3} \cdot (x - x_1)$$

$$y \cdot y_1 - y_1 \cdot x_1^{1/3} = -x \cdot y_1^{1/3} (y_1)^{1/3} \cdot x_1$$

$$\frac{x \cdot y_1^{1/3}}{y_1^{1/3} x_1^{1/3}} + \frac{y \cdot x_1^{1/3}}{y_1^{1/3} \cdot x_1^{1/3}} = \frac{x_1 y_1^{1/3}}{y_1^{1/3} x_1^{1/3}} + \frac{x_1^{1/3} \cdot y_1}{y_1^{1/3} \cdot x_1^{1/3}}$$

$$= \frac{x}{x_1^{1/3}} + \frac{y}{y_1^{1/3}} = x_1^{2/3} + y_1^{2/3}$$

$$= \frac{x}{x_1^{1/3}} + \frac{y}{y_1^{1/3}} = a^{2/3} \quad (\text{from eqn-1})$$

above equation intersects x-axis at A (a<sup>2/3</sup> · x<sub>1</sub><sup>1/3</sup>, 0), y-axis at B(0, a<sup>2/3</sup> · y<sub>1</sub><sup>1/3</sup>)

$$AB = \sqrt{(a^{2/3} \cdot x_1^{2/3})^2 + (a^{2/3} \cdot y_1^{1/3})^2}$$

$$= \sqrt{(a^{2/3})^2 (x_1^{2/3} + y_1^{2/3})^2}$$

$$= \sqrt{a^{4/3} \cdot a^{2/3}} \quad (\text{from eqn-1})$$

$$= \sqrt{a^{6/3}}$$

$$= \sqrt{a^2}$$

$$= a = \text{constant}$$

### Lengths of tangent, normal, subtangent and subnormal

$$1. \quad \text{Length of tangent} = \left| \frac{y^1 \cdot \sqrt{1+(y^1)^2}}{y^1} \right|$$

$$2. \quad \text{Length of tangent} = \left| y^1 \cdot \sqrt{1+(y^1)^2} \right|$$

$$3. \quad \text{Length of the subtangent} = \left| \frac{y}{y^1} \right|$$

$$4. \quad \text{Length of the subnormal} = \left| y \cdot y^1 \right|$$

1. **Show that the length of the subtangent at any point on the curve  $y = a^x$  ( $a > 0$ ) is a constant.**

**Sol:** Differentiating  $y = a^x$  with respect to 'x'

We have  $y^1 = a^x \cdot \log a$

$$\text{Length of the subtangent} = \left| \frac{y}{y^1} \right|$$

$$= \left| \frac{a^x}{a^x \log a} \right|$$

$$= \left| \frac{1}{\log a} \right| \text{ it is constant}$$

2. **Find the lengths of subtangent and subnormal at a point on the curve  $y = b \sin \frac{x}{a}$**

**Sol:**  $y = b \sin \frac{x}{a}$

Differentiating with respect to 'x'

$$y^1 = b \cos \frac{x}{a} \cdot \left(\frac{1}{a}\right)$$

$$y^1 = \frac{b}{a} \times \cos \frac{x}{a}$$

$$\text{Length of the subtangent} = \left| \frac{y}{y^1} \right|$$

$$= \left| \frac{b \sin \frac{x}{a}}{\frac{b}{a} \cdot \cos \frac{x}{a}} \right|$$

$$= a \cdot \tan \frac{x}{a}$$

$$\text{Length of the subnormal} = |y \cdot y^1|$$

$$= \left| b \sin \frac{x}{a} \cdot \frac{b}{a} \cdot \cos \frac{x}{a} \right|$$

$$= \left| \frac{b^2}{2a} \times 2 \sin \frac{x}{a} \cdot \cos \frac{x}{a} \right|$$

$$= \left| \frac{b^2}{2a} \cdot \sin \frac{2x}{a} \right|$$

3. Show that at any point (x,y) on the curve  $y = be^{x/a}$ , the length of the subtangent is a constant and the length of the subnormal is  $\frac{y^2}{a}$

**Sol:**  $y = be^{x/a}$

Differentiating with respect to 'x'

$$y^1 = be^{x/a} \cdot \frac{1}{a} = \frac{b}{a} \cdot e^{x/a}$$

$$\text{Length of the subtangent} = \left| \frac{y}{y^1} \right| = \left| \frac{be^{x/a}}{\frac{b}{a} \cdot e^{x/a}} \right| = |a| = \text{is constant}$$

$$\text{Length of the subnormal} = |y y^1| = \left| b e^{x/a} \cdot \frac{b}{a} \cdot e^{x/a} \right| = \left| \frac{(b e^{x/a})^2}{a} \right| = \frac{y^2}{a}$$

4. At any point 't' on the curve  $x = a(t + \sin t)$ ,  $y = a(1 - \cos t)$  Find the lengths of tangent, normal, and subtangent and subnormal.

Sol:  $x = a(t + \sin t)$ ,  $y = a(1 - \cos t)$

$$y = a^x$$

differentiating both sides with respect to 't'

$$\frac{dx}{dt} = a(1 + \cos t), \quad \frac{dy}{dt} = a \sin t$$

$$y^1 = \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{a \sin t}{a(1 + \cos t)} = \frac{2 \sin t/2 \cdot \cos t/2}{2 \cos^2 t/2}$$

$$= \frac{\sin t/2}{\cos t/2} = \tan t/2$$

$$\text{length of the tangent} = \left| \frac{y \cdot \sqrt{1 + (y^1)^2}}{y^1} \right| = \left| \frac{a(1 - \cos t) \sqrt{1 + \tan^2 t/2}}{\tan t/2} \right|$$

$$= \left| \frac{a \cdot 2 \sin^{2t/2} \cdot \sqrt{\sec^{2t/2}}}{\tan^{t/2}} \right|$$

$$= \left| \frac{a \cdot 2 \sin^{2t/2}}{\frac{\sin^{t/2}}{\cos^{t/2}}} \times \frac{1}{\cos^{t/2}} \right| = 2a \sin^{t/2}$$

$$\text{Length of the normal} = \left| y \sqrt{1 + (y^1)^2} \right|$$

$$= \left| a(1 - \cos t) \cdot \sqrt{1 + \tan^2 t/2} \right|$$

$$= \left| 2a \cdot \sin^{2t/2} \cdot \sqrt{\sec^{2t/2}} \right|$$

$$= \left| 2a \cdot \sin^{2t/2} \cdot \frac{\sin^{2t/2}}{\cos^{2t/2}} \right|$$

$$= \left| 2a \sin^{2t/2} \cdot \tan^{t/2} \right|$$

$$\text{length of the subtangent} = \left| \frac{y}{y^1} \right| = \left| \frac{a(1 - \cos t)}{\tan^{t/2}} \right|$$

$$= \left| \frac{a(1 - \cos t)}{\frac{\sin^{t/2}}{\cos^{t/2}}} \right| = 2a \sin^{t/2} \cdot \cos^{t/2}$$

$$\begin{aligned} \text{length of the subnormal} &= \left| y \cdot y^1 \right| = \left| 2(1 - \cos t) \cdot \tan^{t/2} \right| \\ &= \left| 2a \sin^{2t/2} \cdot \tan^{t/2} \right| \end{aligned}$$

5. At any point 't' on the curve  $x = a(\cos t + t \sin t)$ ,  $y = a(\sin t - t \cos t)$  Find the lengths of tangent, normal, and subtangent and subnormal.

**Sol:**  $x = a(\cos t + t \sin t)$

differentiating both sides with respect to 't'

$$\frac{dx}{dy} = a(-\sin t + \sin t + t \cos t)$$

$$= a t \cos t.$$

$$y = a(-\sin t - t \cos t)$$

differentiating both sides with respect to 't'

$$\frac{dx}{dy} = a(\cos t - \cos t + t \sin t)$$

$$= a t \sin t.$$

$$y^1 = \frac{dx}{dy} = \frac{dy/dt}{dx/dt} = \frac{at \sin t}{at \cos t} = \tan t$$



$$\begin{aligned} \text{length of the subtangent} &= \left| \frac{y}{y^2} \right| = \left| \frac{a(\sin t - t \cos t)}{\tan t} \right| \\ &= (ac \sin t - t \cos t) \cdot \cot t \end{aligned}$$

$$\text{length of the subnormal} = |y \cdot y^1| = |a(\sin t - t \cos t) \cdot \tan t|$$

### Angle between two curves and condition for orthogonality of curves

1. Angle between tangents drawn at intersecting points of two curves  $C_1$  and  $C_2$  is angle between curves  $C_1$  and  $C_2$
2. If slopes of tangents drawn at intersecting points of curves is  $m_1$  and  $m_2$  and angle between the curves ' $\theta$ ' then  $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$
3. If  $m_1 = m_2$ , the curves have common tangents, and touch each other.
4. If  $m_1 \cdot m_2 = -1$  the curves intersect orthogonally.

1. Show that the curves  $y^2 = 4(x+1)$ ,  $y^2 = 36(9-x)$  intersect orthogonally.

**Sol:**  $y^2 = 4(x+1)$ ,  $y^2 = 36(9-x)$

$$\therefore 4(x+1) = 36(9-x)$$

$$x + 1 = 9(9-x)$$

$$x + 1 = 81 - 9x$$

$$x + 9x = 81 - 1$$

$$10x = 80$$

$$x = 8$$

$$y^2 = 4(x+1)$$

$$y^2 = 4(8+1)$$

$$y^2 = 4 \times 9$$

$$y = \pm 6$$

intersecting points of given curves (8, 6), (8, -6)

slope of tangent at 'p' to the curve  $y^2 = 4(x+1)$

$$2y \cdot \frac{dy}{dx} = 4$$

$$= \frac{dy}{dx} = \frac{4}{2y} = \frac{2}{y} \text{ (first curve)}$$

$$y^2 = 36(9+x) \Rightarrow 2y \cdot \frac{dy}{dx} = -36$$

$$= \frac{dy}{dx} = -\frac{36}{2y} = -\frac{18}{y} \text{ (second curve)}$$

$$\text{slope of the curve at } (2, 6) \text{ } y^2 = 4(x+1) \text{ } m_1 = \frac{2}{6} = \frac{1}{3}$$

$$\text{slope of the curve at } (2, 6) \text{ } y^2 = 36(9+x) \text{ } m_2 = \frac{-18}{6} = -3$$

$$m_1 \times m_2 = \frac{1}{3} \times (-3) = -1$$

The given curves intersect orthogonally similarly we can prove at point Q(8,-6)

2. Show that the condition for the orthogonality of curves  $ax^2 + bx = 1$ ,  $a, x^2 + b, y^2 = 1$  is

$$\frac{1}{a} - \frac{1}{b} = \frac{1}{a_1} - \frac{1}{b_1}$$

**Sol:** Let intersecting point of given curves  $ax^2 + bx = 1$ ,  $a, x^2 + b, y^2 = 1$

$P(x_1, y_1)$

$$\therefore ax_1^2 + by_1^2 - 1 = 0, \quad a_1x_1^2 + b_1y_1^2 - 1 = 0$$

$$\begin{array}{cccc} x_1^2 & y_1^2 & 1 & \\ b & -1 & a & b \\ b_1 & -1 & a_1 & b_1 \end{array}$$

$$\frac{x_1^2}{-b+b_1} = \frac{y_1^2}{-a_1+a} = \frac{1}{ab_1-a_1b} \quad \dots (1)$$

$\therefore$  slope of the tangent to the curve at  $P(x_1, y_1)$   $ax^2 + by^2 = 1$

$$2ax + 2by^2 \cdot y^1 = 0$$

$$\Rightarrow y_1 = \frac{-2ax}{2by} = \frac{-ax}{by}$$

$$\therefore m_1 = \frac{ax_1}{by_1}$$

slope of the tangent to the curve at  $P(x_1, y_1)$   $a_1x^2 + b_1y^2 = 1$

$$m_1 = \frac{a_1x_1}{b_1y_1}$$

condition for orthogonality is  $m_1 m_2 = -1$

$$-\frac{ax_1}{by_1} \times \frac{a_1x_1}{b_1y_1} = -1$$

$$\frac{aa_1}{bb_1} \cdot \frac{x_1^2}{y_1^2} = -1$$

$$\frac{x_1^2}{y_1^2} = -\frac{bb_1}{aa_1}$$

$$\frac{b_1 - b}{a - a_1} = \frac{-bb_1}{aa_1} \quad (\text{from eqn---(1)})$$

$$b_1 \cdot aa_1 - b \cdot aa_1 = -abb_1 + ba_1b_1$$

$$a a_1 b_1 - b a_1 b_1 = a b a_1 - a b b_1$$

$$a_1 b_1 (a - b) = ab (a_1 - b_1)$$

$$\frac{a - b}{ab} = \frac{a_1 - b_1}{a_1 b_1}$$

$$\frac{a}{ab} - \frac{b}{ab} = \frac{a_1}{a_1 b_1} - \frac{b_1}{a_1 b_1}$$

$$\frac{1}{b} - \frac{1}{a} = \frac{1}{b_1} - \frac{1}{a_1}$$

$$\therefore \frac{1}{a} - \frac{1}{b} = \frac{1}{a_1} - \frac{1}{b_1}$$

3. Find the angle between the curves  $x + y + 2 = 0$ ,  $x^2 + y^2 - 10y = 0$

**Sol:**  $x + y + 2 = 0$

$$\Rightarrow x = -(y + 2)$$

$$x^2 + y^2 - 10y = 0$$

$$(-(y + 2))^2 + y^2 - 10y = 0$$

$$y^2 + 4y + 4 + y^2 - 10y = 0$$

$$2y^2 - 6y + 4 = 0$$

$$y^2 - 3y + 2 = 0$$

$$(y - 1)(y - 2)$$

$$y = +1 \text{ or } y = 2$$

$$y = 1 \quad x = -(1 + 2) = -3$$

$$y = 2x = -(2 + 2) = -4$$

Intersecting points are  $(-3, 1)$ ,  $(-4, 2)$

slope of the tangent at  $(-3, 1)$  to the curve  $x + y + 2 = 0$

$$1 + y^1 = 0$$

$$y^1 = -1$$

$$m_1 = -1$$

$$x^2 + y^2 - 10y = 0$$

differentiating both sides with respect to 'x'

$$2x + 2y \cdot y^1 - 10 \cdot y^1 = 0.$$

$$y^1 = \frac{-2x}{2y - 10}$$

$$m_2 = \frac{-2 \times (-3)}{2(+1) + 10}$$

$$= \frac{6}{2 - 10} = \frac{6}{-8} = \frac{-3}{4}$$

angle between two curves

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

$$= \left| \frac{-1 + 3/4}{1 + (-1)(-3/4)} \right|$$

$$= \left| \frac{\frac{-4 + 3}{4}}{\frac{4 + 3}{4}} \right|$$

$$= \left| \frac{-1}{7} \right|$$

$$= \frac{1}{7}$$

$$\theta = \text{Tan}^{-1} \left( \frac{1}{7} \right)$$

4. Find the angle between the curves  $y^2 = 4x$ ,  $x^2 + y^2 = 5$

**Sol:**  $y^2 = 4x$ ,  $x^2 + y^2 = 5$

$$\Rightarrow x^2 + 4x - 5 = 0$$

$$(x + 5)(x - 1) = 0$$

$$x = -5 \text{ or } 1$$

$$x = 1 \Rightarrow y^2 = 4 \times 1$$

$$\Rightarrow y^2 = 4$$

$$y = \pm 2$$

intersecting points of two curves (1, 2), (1, -2)

$$y^2 = 4x$$

differentiating both sides with respect to 'x'

$$2y \cdot y^1 = 4$$

$$y^1 = \frac{4}{2y} = \frac{2}{y}$$

slope of the first curve at (1, 2) =  $\frac{2}{2} = 1$  ( $m_1$ )

$$x^2 + y^2 = 5$$

differentiating both sides with respect to 'x'

$$2x + 2y \cdot y^1 = 0 \Rightarrow y^1 = \frac{-x}{y}$$

slope of the second curve at (1, 2)

$$m_2 = -\frac{1}{2}$$

angle between two curves ' $\theta$ '

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

$$\tan \theta = \left| \frac{1 + \frac{1}{2}}{1 + (1)\left(-\frac{1}{2}\right)} \right|$$

$$= \left| \frac{\frac{3}{2}}{\frac{1}{2}} \right| = 3$$

$$\tan \theta = 3$$

$$\theta = \tan^{-1}(3)$$

5. Find the angle between two curves  $x^2 = 2(y+1)$ ;  $y = \frac{8}{x^2+4}$

Sol:  $x^2 = 2(y+1)$ ,  $y = \frac{8}{x^2+4}$

$$\Rightarrow x^2 + 4 = \frac{8}{y}$$

$$x^2 = \frac{8}{y} - 4$$

$$\therefore \frac{8}{y} - 4 = 2(y+1)$$

$$\frac{8}{y} = 2y + 2 + 4$$

$$\frac{8}{y} = 2y + 6$$

$$\frac{4}{y} = y + 3$$

$$4 = y^2 + 3y$$

$$y^2 + 3y - 4 = 0$$

$$(y+4)(y-1) = 0$$

$$y = -4 \text{ or } y = 1$$

$$x^2 = 2(1+1)$$

$$x^2 = 2 \times 2$$

$$x = \pm 2$$

intersecting points (2,1) and (2, -1)

$$x^2 = 2(y+1)$$

differentiating both sides with respect to 'x'

$$2x = 2y^1$$

$$y^1 = x$$

slope of the curve at (2, 1)  $m_1 = 2$

$$y = \frac{8}{x^2 + 4}$$

differentiating both sides with respect to 'x'

$$y^1 = -\frac{8}{(x^2 + 4)^2} \cdot 2x$$

slope of the second curve at (2, 1) to the curve  $y = \frac{8}{x^2 + 4}$

$$m_2 = \frac{-16 \times 2}{(2^2 + 4)^2} = \frac{-32}{64} = -\frac{1}{2}$$

$$m_1 \times m_2 = 2 \times \left(-\frac{1}{2}\right) = -1$$

$\therefore$  the given curves intersect orthogonally.

6. Show that the curves  $6x^2 - 5x + 2y = 0$ ,  $4x^2 + 8y^2 = 3$  touch each other at points

$$\left(\frac{1}{2}, \frac{1}{2}\right)$$

**Sol:**  $6x^2 - 5x + 2y = 0$

differentiating both sides with respect to 'x'

$$6 \times 2x - 5 + 2y^1 = 0$$

$$2y^1 = 5 - 12x$$

$$y^1 = \frac{5 - 12x}{2}$$

slope of the tangent at  $\left(\frac{1}{2}, \frac{1}{2}\right)$  to the first curve is  $m_1 = \frac{5 - 12 \times \frac{1}{2}}{2}$

$$= \frac{5 - 6}{2}$$

$$= -\frac{1}{2}$$

$$4x^2 + 8y^2 = 3$$

differentiating both sides with respect to 'x'

$$4 \times 2x + 8 \times 2y \cdot y^1 = 0$$

$$y^1 = \frac{-8x}{16y} = \frac{-x}{2y}$$

slope of the tangent at  $\left(\frac{1}{2}, \frac{1}{2}\right)$  to the second curve is  $m_2 = \frac{-\frac{1}{2}}{2 \times \frac{1}{2}} = -\frac{1}{2}$

$$\therefore m_1 = m_2$$

$\therefore$  the given curves touch each other at  $\left(\frac{1}{3}, \frac{1}{2}\right)$

## Maxima and Minima

### Key concepts

#### 1. 1<sup>st</sup> derivative test :

Let  $f$  be a differential function on an interval  $D$ ,  $c \in D$ , and  $f$  is defined in some neighbourhood of  $c$ , suppose,  $c$  is a stationary point of  $f$  such that  $(c - \delta, c + \delta)$  does not contain any other stationary point for some  $\delta > 0$  then.

- (i)  $c$  is a point of local maximum, if  $f'(x)$  changes sign from positive to negative at  $x = c$
- (ii)  $c$  is a point of local minimum, if  $f'(x)$  changes sign from negative to positive at  $x = c$
- (iii)  $c$  is neither a point of local maximum nor a points of local minimum  $f'(x)$  does not change sign at  $x = c$

#### 2. 2<sup>nd</sup> derivative test :

- (i)  $x=c$  is a point of local maximum of 'f' if  $f'(c)=0$  and  $f''(c) < 0$ , and local maximum value of 'f' is  $f(c)$
- (ii)  $x=c$  is a point of local minimum of 'f' if  $f'(c) = 0$  and  $f''(c) > 0$  and local minimum value 'f' is  $f(c)$
- (iii) the test fails if  $f'(c)=0$  and  $f''(c)=0$




1. Find the maximum area of the rectangle that can be formed with fixed perimeter 20.

Sol: Let  $x$  and  $y$  denote the length and the breadth of a rectangle respectively . Given that the perimeter of the rectangle is 20.

$$\therefore 2(x + y) = 20 \dots\dots (1)$$

$$\Rightarrow x + y = 10 \dots\dots\dots(2)$$



area of the rectangle  $A = x.y \dots\dots\dots (3)$

$$A = x \cdot (10 - x) \quad \because x + y = 10$$

$$A = 10x - x^2 \dots\dots\dots (4)$$

differentiating both sides with respect to 'x'

$$\frac{dA}{dx} = 10 - 2x \dots\dots (5)$$

$$\frac{dA}{dx} = 0 \Rightarrow 10 - 2x = 0$$

$$\Rightarrow 2x = 10$$

$$\Rightarrow x = 5$$

$\therefore x = 5$  is the stationary point

differentiating (5) with respect to 'x'

$$\frac{d^2A}{dx^2} = -2$$

$$\frac{d^2A}{dx^2} < 0$$

which is negative, therefore by second derivative test the area  $A$  is maximized at  $x = 5$  and hence  $y = 10 - 5 = 5$ , and the maximum area is  $A = 5(5) = 25$

2. The profit function  $P(x)$  of a company selling  $x$  items per day is given by  $P(x) = (150 - x)x - 1600$ . Find the number of items that the company should sell to get maximum profit. Also find the maximum profit.

Sol: 
$$P(x) = (150 - x)x - 1600$$

$$= 150x - x^2 - 1600$$

differentiating both sides with respect to 'x'

$$\frac{dP}{dx} = 150 - 2x \dots\dots (1)$$

$$\frac{dP}{dx} = 0$$

$$\Rightarrow 150 - 2x = 0$$

$$2x = 150$$

$$x = 75$$

Again differentiating Eqn .... (1) with respect to 'x'

$$\frac{d^2P}{dx^2} = -2$$

$$\frac{d^2P}{dx^2} < -2$$

∴ The profit P (x) is maximum for x = 75

∴ The company should sell 75 items a day to make maximum profit.

the maximum profit P (75) = (150 - 75) · 75 - 1600

$$= 75 \times 75 - 1600$$

$$= 5625 - 1600$$

$$= 4025.$$

3. From a rectangular sheet of dimensions 30cm × 80cm four equal squares of side x cm are removed at the corners, and the sides are then turned up so as to form an open rectangular box. Find the value of x, so that the volume of the box is the greatest.

Sol: length of the box = 80 - 2x


breadth of the box = 30 - 2x

height of the box = x

volume = (80 - 2x) (30 - 2x) · x

$$= (2400 - 160x - 60x + 4x^2) \cdot x.$$

$$= 4x^3 - 220x^2 + 2400x$$



differentiating both sides with respect to 'x'

$$\frac{dv}{dx} = 12x^2 - 440x + 2400 \quad \dots\dots 91)$$

$$\frac{dv}{dx} = 0$$

$$12x^2 - 440x + 2400 = 0$$

$$3x^2 - 110x + 600 = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{110 \pm \sqrt{(110)^2 - 4 \times 3 \times 600}}{2 \times 3}$$

$$x = \frac{110 \pm \sqrt{12100 - 7200}}{6}$$

$$= \frac{110 \pm \sqrt{4900}}{6}$$

$$= \frac{110 \pm 70}{6}$$

$$= \frac{110+70}{6} \quad = \frac{110-70}{6}$$

$$= \frac{180}{6} \quad = \frac{40}{6}$$

$$= 30 \quad = \frac{20}{3}$$

$$x = 30 \quad b = 30 - 2 \times 30$$

$$= -30$$

$$b < 0$$

$$\therefore x \neq 30$$

$$\therefore x = \frac{20}{3}$$

Again differentiating Eqn .... (1) with respect to 'x'

$$\frac{d^2v}{dx^2} = 24x - 440$$

$$x = \frac{20}{3} \quad \& \quad \frac{d^2v}{dx^2} = 24 \times \frac{20}{3} - 440$$

$$= 160 - 440$$

$$= -280$$

$$\frac{d^2v}{dx^2} < 0$$

Volume of box is maximum at  $x = \frac{20}{3}$

4. A window is in the shape of a rectangle surrounded by a semicircle. If the perimeter of the window is 20 ft. Find the maximum area.

Sol: The perimeter =  $22 + 2y = \pi x = 20$

$$2y = 20 - (\pi + 2)x.$$

$$y = 10 - \frac{\pi + 2}{2} \cdot x$$

$$\text{area} = 2xy + \frac{\pi x^2}{2}$$

$$A = (20 - (\pi + 2) \cdot x) + \frac{\pi x^2}{2}$$

$$= 20 - (\pi + 2)x^2 + \frac{\pi x^2}{2}$$

differintating both sides with respect to 'x'

$$\frac{dA}{dx} = 20 - (\pi + 2) \cdot 2x + \frac{\pi}{2} \times 2x \quad \dots 91)$$

$$\frac{dA}{dx} = 0$$

$$20 - (\pi + 2)x = 0$$

$$x(\pi - 2\pi - 4) = -20$$

$$x(-\pi - 4) = -20$$

$$x(\pi + 4) = 20$$


$$x = \frac{20}{\pi + 4}$$

Again differintating Eqn .... (1) with respect to 'x'

$$\frac{d^2A}{dx^2} = -(\pi + 2) \times 2 + \pi$$

$$= -2\pi - 4 + \pi$$

$$= -\pi - 4$$



$$\frac{d^2A}{dx^2} < 0$$

area of the window maximum at  $x = \frac{20}{\pi + 4}$

$$A = 2 \times y + \frac{\pi}{2} \cdot x^2$$

$$2y = 20 - (\pi + 2)x$$

$$2y = 20 - (\pi + 2) \times \frac{20}{\pi + 4}$$

$$2y = \frac{20\pi + 80 - 20\pi - 40}{\pi + 4}$$

$$= \frac{40}{\pi + 4}$$

$$A = 2xy + \frac{\pi}{2} \cdot x^2$$

$$= \frac{20}{\pi + 4} \cdot \frac{40}{\pi + 4} + \frac{\pi}{2} \left( \frac{20}{\pi + 4} \right)^2$$

$$= \frac{1600 + 400 \cdot \pi}{2(\pi + 4)^2}$$

$$= \frac{200(4 + \pi)}{(\pi + 4)2}$$

$$= \frac{200}{\pi + 4}$$


5. If the curved surface of right circular cylinder inscribed in a sphere of radius  $r$  is maximum, show that the height of the cylinder is  $\sqrt{2} r$ .

**Sol:** From  $\Delta OAB$

$$OA^2 + AB = OB^2$$

$$R^2 + \left(\frac{h}{2}\right)^2 = r^2$$

$$R^2 = r^2 - \frac{h^2}{4}$$



lateral surface area of cylinder =  $2\pi rh$

$$= 2\pi \cdot \sqrt{r^2 - \frac{h^2}{4}} \cdot h$$

$$A = \frac{2\pi \cdot h}{2} \cdot \sqrt{4r^2 - h^2}$$

$$A = \pi \cdot h \sqrt{4r^2 - h^2}$$

differintating both sides with respect to 'h'

$$\frac{dA}{dh} = \pi \left\{ 1 \cdot \sqrt{4r^2 - h^2} + \frac{h \cdot 1 \cdot (\cancel{2}h)}{\cancel{2}\sqrt{4r^2 - h^2}} \right\}$$

$$\frac{dA}{dh} = \pi \cdot \left\{ \frac{4r^2 - h^2 - h^2}{\sqrt{4r^2 - h^2}} \right\}$$

$$= \pi \cdot \left\{ \frac{4r^2 - 2h^2}{\sqrt{4r^2 - h^2}} \right\}$$

$$\frac{dA}{dh} = 0$$

$$\pi \cdot \left\{ \frac{4r^2 - 2h^2}{\sqrt{4r^2 - h^2}} \right\} = 0 \quad 4r^2 - 2h^2 = 0$$

$$h = \sqrt{2} \cdot r.$$

Again differintating Eqn .... (1) with respect to 'h'

$$\frac{d^2A}{dh^2} = 2\pi \cdot \left\{ \frac{\sqrt{4R^2 - h^2} \cdot (-2h) - \frac{(2R^2 - h^2)}{2\sqrt{4R^2 - h^2}} \cdot (-2h)}{4e^2 - h^2} \right\}$$

$$= 2\pi \cdot \left\{ \frac{-8R^2h + 2h^3 + 2R^2h - h^3}{(4e^2 - h^2)(\sqrt{4R^2 - 4^2})} \right\}$$

$$= 2\pi \cdot \left\{ \frac{-8R^2h + 2h^3 + 2R^2h - h^3}{4R^2 - h^2 \cdot \sqrt{4R^2 - h^2}} \right\}$$

$$= \frac{-4\pi h}{\sqrt{4R^2 - h^2}} < 0, \text{ at } h = \sqrt{2}R \quad \frac{d^2A}{dh^2} < 0$$

∴ lateral surface area of cylinder is maximum when  $h = \sqrt{2}R$

**6. A wire of length  $l$  is cut into two parts which are bent respectively in the form of a square and a circle. What are the lengths of the pieces of the wire respectively so that the sum of the areas is the least.**

**Sol:** Let the square formed with the wire of length  $x$

Let the circle formed with wire of length is  $l-x$

side of the square =  $\frac{x}{4}$

area =  $\left(\frac{x}{4}\right)^2$

circumfrance of the circle  $2\pi r = l - x$

$$r = \frac{l-x}{2\pi}$$

area of the circle =  $\pi r^2 = \pi \cdot \left(\frac{l-x}{2\pi}\right)^2$

some of the areas  $A = \frac{x^2}{16} + \frac{(l-x)^2}{4\pi}$

differintating both sides with respect to 'x'

$$\frac{dA}{dx} = \frac{2x}{16} + \frac{2(l-x)}{4\pi} \cdot (-1) \quad \dots (1)$$

$$\frac{dA}{dx} = 0$$

$$= \frac{2x}{16} - \frac{2(l-x)}{4\pi} = 0 \Rightarrow \frac{x}{8} - \frac{l-x}{2\pi} = 0$$

$$= \frac{\pi \cdot x - 4l + 4x}{8\pi} = 0 \Rightarrow x(\pi + 4) = 4l$$

$$x = \frac{4l}{\pi + 4}$$

Again differentiating Eqn .... (1) with respect to 'x'

$$\begin{aligned} \frac{dA^2}{dx^2} &= \frac{1}{8} - \frac{(-1)}{2\pi} \\ &= \frac{1}{8} + \frac{1}{2\pi} > 0 \end{aligned}$$

∴ At  $x = \frac{4l}{\pi + 4}$  we get minimum value

$$l - x = l - \frac{4l}{\pi + 4} = \frac{\pi l + 4l - 4l}{\pi + 4} = \frac{\pi l}{\pi + 4}$$

differentiating equation (1) with respect to 'x'

$$\frac{d^2A}{dx^2} = \frac{1}{8} - \frac{(-1)}{2\pi} = \frac{1}{8} + \frac{1}{2\pi} > 0$$

sum of areas is minimum at  $x = \frac{4l}{\pi + 4}$

$$l - x = l - \frac{4l}{\pi + 4} = \frac{\pi l + 4l - 4l}{\pi + 4} = \frac{\pi l}{\pi + 4}$$

\*\*\*\*\*



