

# REDOX REACTION

## Table of Contents

➤	Theory .....	2
➤	Solved Examples .....	21
➤	Exercise - 1 : Basic Objective Questions .....	23
➤	Exercise - 2 : Previous Year JEE Mains Questions .....	27
➤	Exercise - 3 : Advanced Objective Questions .....	29
➤	Exercise - 4 : Previous Year JEE Advanced Questions .....	35
➤	Answer Key .....	38

# THEORY

## Oxidation & Reduction

Let us do a comparative study of oxidation and reduction :

Oxidation	Reduction
1. Addition of Oxygen e.g. $2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$	1. Removal of Oxygen e.g. $\text{CuO} + \text{C} \rightarrow \text{Cu} + \text{CO}$
2. Removal of Hydrogen e.g. $\text{H}_2\text{S} + \text{Cl}_2 \rightarrow 2\text{HCl} + \text{S}$	2. Addition of Hydrogen e.g. $\text{S} + \text{H}_2 \rightarrow \text{H}_2\text{S}$
3. Increase in positive charge e.g. $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-$	3. Decrease in positive charge e.g. $\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+}$
4. Increase in oxidation number (+2)            (+4) e.g. $\text{SnCl}_2 \rightarrow \text{SnCl}_4$	4. Decrease in oxidation number (+7)            (+2) e.g. $\text{MnO}_4^- \rightarrow \text{Mn}^{2+}$
5. Removal of electron e.g. $\text{Sn}^{2+} \rightarrow \text{Sn}^{4+} + 2\text{e}^-$	5. Addition of electron e.g. $\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+}$

## Oxidation Number

- It is an imaginary or apparent charge developed over atom of an element when it goes from its elemental free state to combined state in molecules.
- It is calculated on basis of an arbitrary set of rules.
- It is a relative charge in a particular bonded state.
- In order to keep track of electron-shifts in chemical reactions involving formation of compounds, a more practical method of using oxidation number has been developed.
- In this method, it is always assumed that there is a complete transfer of electron from a less electronegative atom to a more electronegative atom.

## Rules governing oxidation number


The following rules are helpful in calculating oxidation number of the elements in their different compounds. it is to be remembered that the basis of these rule is the electronegativity of the element.

- **Fluorine atom :**  
Fluorine is most electronegative atom (known). It always has oxidation number equal to  $-1$  in all its compounds
- **Oxygen atom :**  
In general and as well as in its oxides, oxygen atom has oxidation number equal to  $-2$ .

### In case of

- (i) peroxide (e.g.  $\text{H}_2\text{O}_2$ ,  $\text{Na}_2\text{O}_2$ ) is  $-1$ ,
- (ii) super oxide (e.g.  $\text{KO}_2$ ) is  $-1/2$
- (iii) ozonide (e.g.  $\text{KO}_3$ ) is  $-1/3$
- (iv) in  $\text{OF}_2$  is  $+2$  & in  $\text{O}_2\text{F}_2$  is  $+1$
- **Hydrogen atom :**  
In general, H atom has oxidation number equal to  $+1$ . But in metallic hydrides (e.g.  $\text{NaH}$ ,  $\text{KH}$ ), it is  $-1$ .
- **Halogen atom :**  
In general, all halogen atoms (Cl, Br, I) have oxidation number equal to  $-1$ .

But if halogen atom is attached with a more electronegative atom than halogen atom, then it will show positive oxidation numbers.




### ○ Metals :

- (a) Alkali metal (Li, Na, K, Rb, ..... ) always have oxidation number +1
- (b) Alkaline earth metal (Be, Mg, Ca ..... ) always have oxidation number +2.
- (c) Aluminium always has +3 oxidation number


**Note :** Metal may have negative or zero oxidation number

- Oxidation number of an element in free state or in allotropic forms is always zero



- Sum of the oxidation numbers of atoms of all elements in a molecule is zero.
- Sum of the oxidation numbers of atoms of all elements in an ion is equal to the charge on the ion.
- If the group number of an element in modern periodic table is n, then its oxidation number may vary from (n – 10) to (n – 18) (but it is mainly applicable for p-block elements)

e.g. N-atom belongs to 15th group in the periodic table, therefore as per rule, its oxidation number may vary from





- The maximum possible oxidation number of any element in a compound is never more than the number of electrons in valence shell. (but it is mainly applicable for p-block elements)

### Calculation of average oxidation number :

#### Solved Examples

#### Example-1 :

Calculate oxidation number of underlined element :



**Sol.** (a) Let oxidation number of S-atom is x. Now work accordingly with the rules given before.

$$(+1) \times 2 + (x) \times 2 + (-2) \times 3 = 0$$

$$x = +2$$

(b) Let oxidation number of S-atom is x

$$\therefore (+1) \times 2 + (x) \times 4 + (-2) \times 6 = 0$$

$$x = +2.5$$

- It is important to note here that  $\text{Na}_2\underline{\text{S}}_2\text{O}_3$  have two S-atoms and there are four S-atom in  $\text{Na}_2\underline{\text{S}}_4\text{O}_6$ . However none of the sulphur atoms in both the compounds have +2 or +2.5 oxidation number, it is the average of oxidation number, which reside on each sulphur atom. Therefore, we should work to calculate the individual oxidation number of each sulphur atom in these compounds.

### Calculation of individual oxidation number

It is important to note that to calculate individual oxidation number of the element in its compound one should know the structure of the compound and use the following guidelines.

#### Formula :


Oxidation Number = Number of electrons in the valence shell - Number of electrons taken up after bonding

**Guidelines :** It is based on electronegativity of elements.

1. If there is a bond between similar type of atom and each atom has same type of hybridisation, then bonded pair electrons are equally shared by each element.

#### Example :

Calculate oxidation number of each Cl-atom in  $\text{Cl}_2$  molecule



I :      Number of electrons in the valence shell = 7

Number of electrons taken up after bonding = 7.

$\therefore$  oxidation number =  $7 - 7 = 0$ .

II :      similarly, oxidation number =  $7 - 7 = 0$


2. If there is a bond between different type of atoms :

e.g.      A – B (if B is more electronegative than A)

Then after bonding, bonded pair of electrons are counted with B-atom

#### Example :

Calculate oxidation number of each atom in HCl molecule



**Note :** Electron of H-atom is now counted with Cl-atom, because Cl-atom is more electronegative than H-atom

H :      Number of electrons in the valence shell = 1

Number of electrons taken up after bonding = 0

Oxidation number of H =  $1 - 0 = +1$

Cl :      Number of electrons in the valence shell = 7


Number of electrons taken up after bonding = 8

Oxidation number of Cl =  $7 - 8 = -1$

## Solved Examples

### Example - 2

Calculate individual oxidation number of each S-atom in  $\text{Na}_2\text{S}_2\text{O}_3$  (sodium thiosulphate) with the help of its structure.



**Note :** I (central S-atom) is  $\text{sp}^3$  hybridised (25% s-character) and II (terminal S-atom) is  $\text{sp}^2$  hybridised (33% s-character). Therefore, terminal sulphur atom is more electronegative than central sulphur atom. Now, the shared pair of electrons are counted with terminal S-atom.

$\therefore$  I, S-atom : Number of electrons in the valence shell = 6

Number of electrons left after bonding = 0

Oxidation number of central S-atom =  $6 - 0 = +6$

II, S-atom : Number of electrons in the valence shell = 6


Number of electrons left after bonding = 8

Oxidation number of terminal S-atom =  $6 - 8 = -2$

Now, you can also calculate Average Oxidation number of S =  $\frac{6 + (-2)}{2} = +2$  (as we have calculated before)

### Miscellaneous Examples :


In order to determine the exact or individual oxidation number we need to take help from the structures of the molecules. Some special cases are discussed as follows :



From the structure, it is evident that in  $\text{CrO}_5$  there are two peroxide linkages and one double bond. The contribution of each peroxide linkage is  $-2$ . Let the oxidation number of Cr is  $x$ .

$\therefore x + (-2) \times 2 + (-2) = 0$  or  $x = 6$

$\therefore$  Oxidation number of Cr =  $+6$     **Ans.**



From the structure, it is evident that in  $\text{H}_2\text{SO}_5$ , there is one peroxide linkage, two sulphur-oxygen double bonds and one OH group. Let the oxidation number of S =  $x$ .

$$\therefore (+1) + (-2) + x + (-2) 2 + (-2) + 1 = 0$$


$$\text{or } x + 2 - 8 \quad \text{or } x - 6 = 0 \quad \text{or } x = 6$$

$\therefore$  Oxidation number of S in  $\text{H}_2\text{SO}_5$  is + 6 Ans.

### Paradox of fractional oxidation number


Fractional oxidation number is the average of oxidation state of all atoms of element under examination and the structural parameters reveal that the atoms of element for whom fractional oxidation state is realised are actually present in different oxidation states. Structure of the species  $\text{C}_3\text{O}_2$ ,  $\text{Br}_3\text{O}_8$  and  $\text{S}_4\text{O}_6^{2-}$  reveal the following bonding situations :

- The element marked with asterisk (\*) in each species is exhibiting different oxidation number from rest of the atoms of the same element in each of the species. This reveals that in  $\text{C}_3\text{O}_2$ , two carbon atoms are present in +2 oxidation state each whereas the third one is present in zero oxidation state and the average is +4/3. However, the realistic picture is +2 for two terminal carbons and zero for the middle carbon.




Structure of  $\text{C}_3\text{O}_2$  (Carbon suboxide)

- Likewise in  $\text{Br}_3\text{O}_8$ , each of the two terminal bromine atoms are present in +6 oxidation state and the middle bromine\* is present in +4 oxidation state. Once again the average, that is different from reality, is + 16/3.



Structure of  $\text{Br}_3\text{O}_8$  (Tribromooctaoxide)

- In the same fashion, in the species  $\text{S}_4\text{O}_6^{2-}$ , average oxidation number of S is + 2.5, whereas the reality being + 5, 0\*, 0\* and +5 oxidation number respectively for respective sulphur atoms.



Structure of  $\text{S}_4\text{O}_6^{2-}$ , (tetrathionate ion)

In general, the conclusion is that the idea of fractional oxidation state should be taken with care and the reality is revealed by the structures only.

### Oxidising and reducing agent

- **Oxidising agent or Oxidant :**

Oxidising agents are those compounds which can oxidise others and reduce itself during the chemical reaction. Those reagents in which for an element, oxidation number decreases or which undergoes gain of electrons in a redox reaction are termed as oxidants.

e.g.  $\text{KMnO}_4$ ,  $\text{K}_2\text{Cr}_2\text{O}_7$ ,  $\text{HNO}_3$ , conc.  $\text{H}_2\text{SO}_4$  etc are powerful oxidising agents.

### ○ Reducing agent or Reductant :


Reducing agents are those compounds which can reduce other and oxidise itself during the chemical reaction. Those reagents in which for an element, oxidation number increases or which undergoes loss of electrons in a redox reaction are termed as reductants.

e.g. KI,  $\text{Na}_2\text{S}_2\text{O}_3$  etc are the powerful reducing agents.

**Note :** There are some compounds also which can work both as oxidising agent and reducing agent

e.g.  $\text{H}_2\text{O}_2$ ,  $\text{NO}_2^-$


### HOW TO IDENTIFY WHETHER A PARTICULAR SUBSTANCE IS AN OXIDISING OR A REDUCING AGENT



### Redox reaction

A reaction in which oxidation and reduction simultaneously take place is called a redox reaction



In all redox reactions, the total increase in oxidation number must be equal to the total decrease in oxidation number.




### Disproportionation Reaction :

A redox reaction in which same element present in a particular compound in a definite oxidation state is oxidized as well as reduced simultaneously is a disproportionation reaction.


Disproportionation reactions are a special type of redox reactions. One of the reactants in a disproportionation reaction always contains **an element that can exist in at least three oxidation states**. The element in the form of reacting substance is in the intermediate oxidation state and both higher and lower oxidation states of that element are formed in the reaction. For example :




### Consider the following reactions :



$\text{KClO}_3$  plays a role of oxidant and reductant both. Here, Cl present in  $\text{KClO}_3$  is reduced and O present in  $\text{KClO}_3$  is oxidized. Since same element is not oxidized and reduced, so it is not a disproportionation reaction, although it looks like one.



Nitrogen in this compound has -3 and +3 oxidation number, which is not a definite value. So it is not a disproportionation reaction. It is an example of comproportionation reaction, which is a class of redox reaction in which an element from two different oxidation state gets converted into a single oxidation state.











It is a case of disproportionation reaction and Cl atom is disproportionating.



**List of some important disproportionation reactions**

1.  $\text{H}_2\text{O}_2 \longrightarrow \text{H}_2\text{O} + \text{O}_2$
2.  $\text{X}_2 + \text{OH}^- (\text{dil.}) \longrightarrow \text{X}^- + \text{XO}^- (\text{X} = \text{Cl, Br, I})$
3.  $\text{X}_2 + \text{OH}^- (\text{conc.}) \longrightarrow \text{X}^- + \text{XO}_3^-$





**F<sub>2</sub> does not undergo disproportionation as it is the most electronegative element.**




9. Oxyacids of Phosphorus (+1, +3 oxidation number)



10. Oxyacids of Chlorine (Halogens) (+1, +3, +5 Oxidation number)



- Reverse of disproportionation is called Comproportionation. In some of the disproportionation reactions, by changing the medium (from acidic to basic or reverse), the reaction goes in backward direction and can be taken as an example of Comproportionation reaction.





## Balancing of redox reactions

All balanced equations must satisfy two criteria.

### 1. Atom balance (mass balance) :

There should be the same number of atoms of each kind on reactant and product side.

### 2. Charge balance :

The sum of actual charges on both sides of the equation must be equal.

There are two methods for balancing the redox equations :

1. Oxidation – number change method
2. Ion electron method or half cell method

- Since First method is not very much fruitful for the balancing of redox reactions, students are advised to use second method (Ion electron method) to balance the redox reactions

### Ion electron method :

By this method redox equations are balanced in two different medium.

- (a) Acidic medium (b) Basic medium


### ○ Balancing in acidic medium

Students are advised to follow the following steps to balance the redox reactions by ion electron method in acidic medium


## Solved Examples

### Example - 3 :

Balance the following redox reaction :




**Sol. Step-I** Assign the oxidation number to each element present in the reaction




### Step II :

Now convert the reaction in Ionic form by eliminating the elements or species, which are not undergoing either oxidation or reduction.



### Step III :

Now identify the oxidation/reduction occurring in the reaction




### Step IV :

Split the Ionic reaction in two half, one for oxidation and other for reduction.



**Step V :**

Balance the atom other than oxygen and hydrogen atom in both half reactions




Fe & Mn atoms are balanced on both side.

**Step VI :**

Now balance O & H atom by  $\text{H}_2\text{O}$  &  $\text{H}^+$  respectively by the following way : For one excess oxygen atom, add one  $\text{H}_2\text{O}$  on the other side and two  $\text{H}^+$  on the same side.


**Step VII :**

Equation (i) & (ii) are balanced atomwise. Now balance both equations chargewise. To balance the charge, add electrons to the electrically positive side.

**Step VIII :**

The number of electrons gained and lost in each half-reaction are equalised by multiplying both the half reactions with a suitable factor and finally the half reactions are added to give the overall balanced reaction.

Here, we multiply equation (1) by 5 and (2) by 1 and add them :





(Here, at this stage, you will get balanced redox reaction in Ionic form)

**Step IX :**

Now convert the ionic reaction into molecular form by adding the elements or species, which are removed in step (2).

Now, by some manipulation, you will get :




○ **Balancing in basic medium :**

In this case, except step VI, all the steps are same. We can understand it by the following example :


## Solved Examples

### Example - 4 :


Balance the following redox reaction in basic medium :




**Sol.** By using upto step V, we will get :



Now, students are advised to follow step VI to balance 'O' and 'H' atom.




○ Now, since we are balancing in basic medium, therefore add as many as  $\text{OH}^-$  on both side of equation as there are  $\text{H}^+$  ions in the equation.



Now see equation (i) and (ii) in which O and H atoms are balanced by  $\text{OH}^-$  and  $\text{H}_2\text{O}$


Now from step VIII



## Concept of equivalents

### Equivalent mass of element

Number of parts by mass of an element which reacts or displaces from a compound 1.008 parts by mass of hydrogen, 8 parts by mass of oxygen and 35.5 parts by mass of chlorine, is known as the equivalent weight of the element.



$$48\text{g} \quad 32\text{g}$$

$$12\text{g} \quad 8\text{g}$$

$\therefore$  32g of  $\text{O}_2$  reacts with 48 g of Mg


$$\therefore 8 \text{ g of } \text{O}_2 = \frac{48 \times 8}{32} = 12\text{g}$$

$\therefore$  Equivalent weight of Mg = 12



$$65.5 \text{ g} \quad 32.75$$

$$\therefore \text{Equivalent weight of Zn} = \frac{65.5}{2} = 32.75\text{g}$$



$$27\text{ g} \quad \frac{3}{2} \times 71\text{ g}$$

$\therefore$  111.5 g chlorine reacts with 27 g of Al.

$$\therefore 35.5\text{ g chlorine reacts with } \frac{27 \times 35.5}{111.5} = 9.0\text{ g of Al}$$

$$\therefore \text{Equivalent weight of aluminium} = \frac{27}{3} = 9.0$$

As we can see from the above examples that equivalent weight is the ratio of atomic weight and a factor (say n-factor or valency factor) which is in above three cases is their respective valencies.

#### Equivalent weight (E) :

$$\text{In general, Eq. wt. (E)} = \frac{\text{Atomic weight or molecular weight}}{\text{valency factor (v.f)}} = \frac{\text{Mol. wt.}}{\text{n - factor}} = \frac{M}{x}$$

$$\text{Number of Equivalents} = \frac{\text{mass of species}}{\text{eq. wt. of that species}}$$

For a solution, Number of equivalents =  $N_1 V_1$ , where N is the normality and V is the volume in litres

- Equivalent mass is a pure number which, when expressed in gram, is called gram equivalent mass.
- The equivalent mass of substance may have different values under different conditions.
- There is no hard and fast rule that equivalent weight will be always less than the molecular mass.

#### Valency factor calculation :

##### ○ For Elements :

Valency factor = valency of the element.

##### ○ For Acids :

Valency factor = number of replaceable  $\text{H}^+$  ions per acid molecule

#### Solved Examples

**Example - 5 :** NaOH, KOH


**Sol.** v.f.  $\rightarrow$  1 1

$$\text{Eq. wt.} \quad \frac{M}{1} \quad \frac{M}{1}$$

- Bases may be defined as the substances in which OH group is/are directly attached with group I elements (Li, Na, K, Rb, Cs), group II elements (Be, Mg, Ca, Ba) or group III elements (Al, Ga, In, Tl), transition metals, non-metallic cations like  $\text{PH}_4^+$ ,  $\text{NH}_4^+$  etc.
- **Acid - base reaction :**  
In case of acid base reaction, the valence factor is the actual number of  $\text{H}^+$  or  $\text{OH}^-$  replaced in the reaction. The acid or base may contain more number of replaceable  $\text{H}^+$  or  $\text{OH}^-$  than actually replaced in reaction.
- **v. f.** for base is the number of  $\text{H}^+$  ion from the acid replaced by each molecule of the base

## Solved Examples

### Example - 6 :



Base      Acid


**Sol.** Valency factor of base = 1

Here, two molecule of NaOH replaced  $2\text{H}^+$  ion from the  $\text{H}_2\text{SO}_4$ . Therefore, each molecule of NaOH replaced only one  $\text{H}^+$  ion of acid, so v.f. = 1.

○ v. f. for acid is the number of  $\text{OH}^-$  replaced from the base by each molecule of acid

## Solved Examples

### Example - 7 :



Base      Acid

**Sol.** Valency factor of acid = 1

Here, one of molecule of  $\text{H}_2\text{SO}_4$  replaced one  $\text{OH}^-$  from NaOH. Therefore, valency factor for  $\text{H}_2\text{SO}_4$  is one

$$\therefore \text{Eq. wt. of } \text{H}_2\text{SO}_4 = \frac{\text{Mol. wt}}{1}$$

○ **Salts :**

(a) **In non-reacting condition**

○ **Valency factor** = Total number of positive charge or negative charge present in the compound.

## Solved Examples


### Example - 8 :

		$\text{Na}_2\text{CO}_3$ ,	$\text{Fe}_2(\text{SO}_4)_3$	$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$
<b>Sol.</b>	v.f.	2	$2 \times 3 = 6$	2
	Eq. wt.	$\frac{M}{2}$	$\frac{M}{6}$	$\frac{M}{2}$

(b) **In reacting condition**

## Solved Examples

### Example - 9 :



Base      Acid


**Sol.** It is an acid base reaction, therefore valency factor for  $\text{Na}_2\text{CO}_3$  is one while in non-reacting condition, it will be two.

(c) **Equivalent weight of oxidising / reducing agents in a redox reaction**

In case of redox change, v.f. = Total change in oxidation number per molecule.

## Solved Examples

### Example - 10 :



**Sol.** Mn in  $\text{KMnO}_4$  is going from +7 to +2, so change in oxidation number per molecule of  $\text{KMnO}_4$  is 5. So the valency factor of

$\text{KMnO}_4$  is 5 and equivalent weight is  $\frac{M}{5}$ .

### Normality :

Normality of a solution is defined as the number of equivalents of solute present in one litre (1000 mL) solution.

Let V mL of a solution is prepared by dissolving W g of solute of equivalent weight E in water.

○ Number of equivalents of solute =  $\frac{W}{E}$

V mL of solution contain  $\frac{W}{E}$  equivalents of solute

∴ 1000 mL solution will contain  $\frac{W \times 1000}{E \times V}$  equivalents of solute.

○ **Normality (N)** =  $\frac{W \times 1000}{E \times V}$

○ **Normality (N) = Molarity × Valency factor**

$$N \times V \text{ (in mL)} = M \times V \text{ (in mL)} \times n$$

or

○ milliequivalents = millimoles × n

## Solved Examples

### Example 11 :

Calculate the normality of a solution containing 15.8 g of  $\text{KMnO}_4$  in 50 mL acidic solution.

**Sol :** **Normality (N)** =  $\frac{W \times 1000}{E \times V}$

Here  $W = 15.8 \text{ g}$ ,  $V = 50 \text{ mL}$   $E = \frac{\text{molar mass of } \text{KMnO}_4}{\text{Valency factor}} = 158/5 = 31.6$

So, normality = 10 N

### Example 12 :


Calculate the normality of a solution containing 50 mL of 5 M solution of  $\text{K}_2\text{Cr}_2\text{O}_7$  in acidic medium.

**Sol :** Normality (N) Molarity × valency factor =  $5 \times 6 = 30 \text{ N}$


## Law of Equivalence

The law states that one equivalent of an element combine with one equivalent of the other. In a chemical reaction, equivalents and milli equivalents of reactants react in equal amount to give same number of equivalents or milli equivalents of products separately.

### Accordingly



$$\text{meq of A} = \text{meq of B} = \text{meq of M} = \text{m.eq. of N}$$




$$\text{meq of } M_xN_y = \text{meq of M} = \text{meq of N}$$

## Solved Examples

### Example 13 :

Find the number of moles of  $\text{KMnO}_4$  needed to oxidise one mole  $\text{Cu}_2\text{S}$  in acidic medium.



**Sol.** From law of equivalence,

equivalents of  $\text{Cu}_2\text{S} = \text{equivalents of } \text{KMnO}_4$

moles of  $\text{Cu}_2\text{S} \times \text{v.f.} = \text{moles of } \text{KMnO}_4 \times \text{v.f.}$

$$1 \times 8 = \text{moles of } \text{KMnO}_4 \times 5 \quad \Rightarrow \text{moles of } \text{KMnO}_4 = 8/5$$

$$(\therefore \text{v.f. of } \text{Cu}_2\text{S} = 2(2 - 1) + 1(4 - (-2)) = 8 \text{ and v.f. of } \text{KMnO}_4 = 1(7 - 2) = 5)$$

### Example 14 :

The number of moles of oxalate ions oxidized by one mole of  $\text{MnO}_4^-$  ion in acidic medium are :

(a)  $\frac{5}{2}$

(b)  $\frac{2}{5}$

(c)  $\frac{3}{5}$

(d)  $\frac{5}{3}$

**Sol.** Equivalents of  $\text{C}_2\text{O}_4^{2-} = \text{equivalents of } \text{MnO}_4^-$

$$x (\text{mole}) \times 2 = 1 \times 5$$

$$(\therefore \text{v.f. of } \text{C}_2\text{O}_4^{2-} = 2(4 - 3) = 2 \text{ and v.f. of } \text{MnO}_4^- = 1(7 - 2) = 5.$$

$$x = \frac{5}{2} \text{ mole of } \text{C}_2\text{O}_4^{2-} \text{ ions.}$$

## Solved Examples

### Example : 15

How many millilitres of 0.02 M  $\text{KMnO}_4$  solution would be required to exactly titrate 25 mL of 0.2 M  $\text{Fe}(\text{NO}_3)_2$  solution in acidic medium ?

**Sol. Method -1 : Mole concept method**


Starting with 25 mL of 0.2 M  $\text{Fe}^{2+}$ , we can write :

$$\text{Millimoles of } \text{Fe}^{2+} = 25 \times 0.2 \quad \dots\dots (1)$$

and in volume V (in milliliters) of the  $\text{KMnO}_4$ ,

$$\text{Millimoles of } \text{MnO}_4^- = V(0.02) \quad \dots\dots\dots (2)$$

The balanced reaction is :



This requires that at the equivalent point,

$$\frac{\text{m.moles of MnO}_4^-}{1} = \frac{\text{m.moles of Fe}^{2+}}{5}$$

$$\therefore \frac{V(0.02)}{1} = \frac{(25)(0.2)}{5} \text{ (from (1) \& (2))}$$

$$\therefore V = 50 \text{ mL.}$$

### Method -2 : Equivalent Method :

At the equivalence point,

milliequivalents of  $\text{MnO}_4^-$  = milliequivalents of  $\text{Fe}^{2+}$

$$M_1 \times v f_1 \times V_1 = M_2 \times v f_2 \times V_2$$

$$0.02 \times 5 \times V_1 = 0.2 \times 1 \times 25 \quad (\because \text{MnO}_4^- \rightarrow \text{Mn}^{2+}; v.f. = 5, \text{Fe}^{2+} \rightarrow \text{Fe}^{3+}; v.f. = 1)$$

$$\therefore V_1 = 50 \text{ mL.}$$

## Titration

Titration is procedure for determining the concentration of a solution by allowing a carefully measured volume to react with a standard solution of another substance, whose concentration is known.

**Standard solution** - It is a solution whose concentration is known and is taken in burette. It is also called **Titrant**.

There are two type of **titrants** :

- **Primary titrants/standard** - These reagents can be accurately weighed and their solutions are not to be standardised before use.

Ex : Oxalic acid,  $\text{K}_2\text{Cr}_2\text{O}_7$ ,  $\text{AgNO}_3$ ,  $\text{CuSO}_4$ , ferrous ammonium sulphate, hypo etc.

- **Secondarily titrants/standard** : These reagents cannot be accurately weighed and their solutions are to be standardised before use.

Ex :  $\text{NaOH}$ ,  $\text{KOH}$ ,  $\text{HCl}$ ,  $\text{H}_2\text{SO}_4$ ,  $\text{I}_2$ ,  $\text{KMnO}_4$ , etc.

**Titrate** : Solution consisting of substance to be estimated, generally taken in a beaker.

**Equivalence point** : It is the point when number of equivalents of titrant added becomes equal to number of equivalents of titrate.

**At equivalence point** :

$$n_1 V_1 M_1 = n_2 V_2 M_2$$

**Indicator** : An auxiliary substance added for physical detection of completion of titration at equivalence point. It generally show colour change on completion of titration.

**Types of Titrations** :

- Acid-base titrations (to be studied in Ionic equilibrium)



## ○ Redox Titrations

### Some Common Redox Titrations

**Table of Redox Titrations : (Excluding Iodometric/Iodimetric titrations)**

	Estimation of	By titrating with	Reactions	Relation *between OA and RA
1.	$\text{Fe}^{2+}$	$\text{MnO}_4^-$	$\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-$ $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$	$5\text{Fe}^{2+} \equiv \text{MnO}_4^-$ Eq. wt. of $\text{Fe}^{2+} = \text{M}/1$
2.	$\text{Fe}^{2+}$	$\text{Cr}_2\text{O}_7^{2-}$	$\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-$ $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	$6\text{Fe}^{2+} \equiv \text{Cr}_2\text{O}_7^{2-}$ Eq. wt. of $\text{Cr}_2\text{O}_7^{2-} = \text{M}/6$
3.	$\text{C}_2\text{O}_4^{2-}$	$\text{MnO}_4^-$	$\text{C}_2\text{O}_4^{2-} \rightarrow 2\text{CO}_2 + 2\text{e}^-$ $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$	$5\text{C}_2\text{O}_4^{2-} \equiv 2\text{MnO}_4^-$ Eq. wt. of $\text{C}_2\text{O}_4^{2-} = \text{M}/2$
4.	$\text{H}_2\text{O}_2$	$\text{MnO}_4^-$	$\text{H}_2\text{O}_2 \rightarrow 2\text{H}^+ + \text{O}_2 + 2\text{e}^-$ $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$	$5\text{H}_2\text{O}_2 \equiv 2\text{MnO}_4^-$ Eq. wt. of $\text{H}_2\text{O}_2 = \text{M}/2$
5.	$\text{As}_2\text{O}_3$	$\text{MnO}_4^-$	$\text{As}_2\text{O}_3 + 5\text{H}_2\text{O} \rightarrow 2\text{AsO}_4^{3-} + 10\text{H}^+ + 4\text{e}^-$ $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$	Eq. wt. of $\text{As}_2\text{O}_3 = \text{M}/4$
6.	$\text{AsO}_3^{3-}$	$\text{BrO}_3^-$	$\text{AsO}_3^{3-} + \text{H}_2\text{O} \rightarrow \text{AsO}_4^{3-} + 2\text{H}^+ + 2\text{e}^-$ $\text{BrO}_3^- + 6\text{H}^+ + 6\text{e}^- \rightarrow \text{Br}^- + 3\text{H}_2\text{O}$	Eq. wt. of $\text{AsO}_3^{3-} = \text{M}/2$ Eq. wt. of $\text{BrO}_3^- = \text{M}/6$



### Permanganate Titrations :

- $\text{KMnO}_4$  is generally used as oxidising agent in acidic medium, generally provided by dilute  $\text{H}_2\text{SO}_4$ .
- $\text{KMnO}_4$  works as self indicator persistent pink color is indication of end point.
- Mainly used for estimation of  $\text{Fe}^{2+}$ , oxalic acid, oxalates,  $\text{H}_2\text{O}_2$  etc.

### Solved Examples

#### Example 16 :



Write the balanced reaction of titration of  $\text{KMnO}_4$  Vs oxalic acid in presence of  $\text{H}_2\text{SO}_4$ .



Indicator :  $\text{KMnO}_4$  acts as self indicator.

**Example : 17**


Write the balanced reaction of titration of  $\text{KMnO}_4$  Vs ferrous ammonium sulphate in presence of  $\text{H}_2\text{SO}_4$ .




**Indicator :**  $\text{KMnO}_4$  acts as self indicator.

**Hydrogen peroxide ( $\text{H}_2\text{O}_2$ )**


$\text{H}_2\text{O}_2$  can behave both like oxidising and reducing agent in both the mediums (acidic and basic).



○ **Oxidising agent : ( $\text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{O}$ )**



$$\text{v.f} = 2$$




$$\text{v.f} = 2$$

○ **Reducing agent : ( $\text{H}_2\text{O}_2 \rightarrow \text{O}_2$ )**



$$\text{v.f} = 2$$




$$\text{v.f} = 2$$

**Note :** Valency factor of  $\text{H}_2\text{O}_2$  is always equal to 2.

**Volume strength of  $\text{H}_2\text{O}_2$  :** Strength of  $\text{H}_2\text{O}_2$  is represented as 10V, 20V, 30V etc.

**20V  $\text{H}_2\text{O}_2$**  means **one litre** of this sample of  $\text{H}_2\text{O}_2$  on decomposition gives **20L of  $\text{O}_2$**  gas of **STP**.

Decomposition of  $\text{H}_2\text{O}_2$  is given as :



$$1 \text{ mole} \quad \frac{1}{2} \times 22.4 \text{ L O}_2 \text{ at STP}$$

$$= 34\text{g} \quad = 11.2 \text{ L O}_2 \text{ at STP}$$

To obtain 11.2 litre O<sub>2</sub> at STP, at least 34 g H<sub>2</sub>O<sub>2</sub> must be decomposed.

$$\text{For 20 L O}_2, \text{ we should decompose atleast } \frac{34}{11.2} \times 20 \text{ g H}_2\text{O}_2$$

$$\therefore \quad 1\text{L solution of H}_2\text{O}_2 \text{ contains } \frac{34}{11.2} \times 20 \text{ g H}_2\text{O}_2$$

$$\therefore \quad 1\text{L solution of H}_2\text{O}_2 \text{ contains } \frac{34}{11.2} \times \frac{20}{17} \text{ equivalents of H}_2\text{O}_2 \quad (E_{\text{H}_2\text{O}_2} = \frac{M}{2} = \frac{34}{2} = 17)$$

$$\text{Normality of H}_2\text{O}_2 = \frac{34}{11.2} \times \frac{20}{17} = \frac{20}{5.6}$$

$$\bigcirc \quad \text{Normality of H}_2\text{O}_2 \text{ (N)} = \frac{\text{Volume strength of H}_2\text{O}_2}{5.6}$$

$$\therefore \quad M_{\text{H}_2\text{O}_2} = \frac{N_{\text{H}_2\text{O}_2}}{\text{v.f.}} = \frac{N_{\text{H}_2\text{O}_2}}{2}$$

$$\bigcirc \quad \text{Molarity of H}_2\text{O}_2 \text{ (M)} = \frac{\text{Volume strength of H}_2\text{O}_2}{11.2}$$

**Strength (in g/L) :** Denoted by S

$$\text{Strength} = \text{Molarity} \times \text{Mol. wt} = \text{Molarity} \times 34$$

$$\text{Strength} = \text{Normality} \times \text{Eq. weight} = \text{Normality} \times 17$$

### Solved Examples

#### Example 18 :

20 mL of H<sub>2</sub>O<sub>2</sub> after acidification with dilute H<sub>2</sub>SO<sub>4</sub> required 30 mL of  $\frac{N}{12}$  KMnO<sub>4</sub> for complete oxidation. Find the strength of H<sub>2</sub>O<sub>2</sub> solution. [Molar mass of H<sub>2</sub>O<sub>2</sub> = 34]

**Sol.** meq. of KMnO<sub>4</sub> = meq. of H<sub>2</sub>O<sub>2</sub>

$$30 \times \frac{1}{12} = 20 \times N'$$

$$N' = \frac{30}{12 \times 20} = \frac{1}{8} N$$

$$\therefore \quad \text{strength} = N' \times \text{equivalent mass} = \frac{1}{8} \times 17 = 2.12 \text{ g/L.}$$

### Hardness of water (Hard water does not give lather with soap)

Temporary hardness - due to bicarbonates of Ca & Mg

Permanent hardness - due to chlorides & sulphates of Ca & Mg. There are some method by which we can soften the water sample.

- (a) By boiling :  $2\text{HCO}_3^- \longrightarrow \text{H}_2\text{O} + \text{CO}_2 + \text{CO}_3^{2-}$  or  
 By Slaked lime :  $\text{Ca}(\text{HCO}_3)_2 + \text{Ca}(\text{OH})_2 \longrightarrow \text{CaCO}_3 + 2\text{H}_2\text{O}$   
 $\text{Ca}^{2+} + \text{CO}_3^{2-} \longrightarrow \text{CaCO}_3$
- (b) By Washing Soda :  $\text{CaCl}_2 + \text{Na}_2\text{CO}_3 \longrightarrow \text{CaCO}_3 + 2\text{NaCl}$
- (c) By ion exchange resins :  $\text{Na}_2\text{R} + \text{Ca}^{2+} \longrightarrow \text{CaR} + 2\text{Na}^+$
- (d) By adding chelating agents like  $(\text{PO}_3^-)_3$  etc.

### Parts Per Million (ppm)

When the solute is present in very less amount, then this concentration term is used. It is defined as the number of parts of the solute present in every 1 million parts of the solution. ppm can both be in terms of mass or in terms of moles. If nothing has been specified, we take ppm to be in terms of mass. Hence, a 100 ppm solution means that 100 g of solute is present in every 1000000g of solution.

$$\text{ppm}_A = \frac{\text{mass of A}}{\text{Total mass}} \times 10^6 = \text{mass fraction} \times 10^6$$

### Measurement of Hardness :

Hardness is measured in terms of ppm (parts per million) of  $\text{CaCO}_3$  or equivalent to it.

$$\text{Hardness in ppm} = \frac{\text{mass of CaCO}_3}{\text{Total mass of solution}} \times 10^6$$

### Solved Examples

#### Example 19 :

0.00012%  $\text{MgSO}_4$  and 0.000111%  $\text{CaCl}_2$  is present in water. What is the measured hardness of water and millimoles of washing soda required to purify water 1000 L water ?

**Sol.** Basis of calculation = 100 g hard water


$$\text{MgSO}_4 = 0.00012\text{g} = \frac{0.00012}{120} \text{ mole}$$

$$\text{CaCl}_2 = 0.000111\text{g} = \frac{0.000111}{111} \text{ mole}$$

$$\therefore \text{equivalent moles of CaCO}_3 = \left( \frac{0.00012}{120} + \frac{0.000111}{111} \right) \text{ mole}$$

$$\therefore \text{mass of CaCO}_3 = \left( \frac{0.00012}{120} + \frac{0.000111}{111} \right) \times 100 = 2 \times 10^{-4} \text{ g}$$

$$\text{Hardness (in terms of ppm of CaCO}_3) = \frac{2 \times 10^{-4}}{100} \times 10^6 = 2 \text{ ppm}$$





$$\therefore \text{Required Na}_2\text{CO}_3 \text{ for 100g of water} = \left( \frac{0.00012}{120} + \frac{0.000111}{111} \right) \text{mole}$$

$$= 2 \times 10^{-6} \text{ mole}$$


$$\therefore \text{Required Na}_2\text{CO}_3 \text{ for 1000 litre water} = \frac{2 \times 10^{-6}}{100} \times 10^6 = \frac{2}{100} \text{ mole} \quad (\because d = 1 \text{ g/mL})$$

$$= \frac{20}{1000} \text{ mole} = 20 \text{ m mole}$$

## SOLVED EXAMPLES

### Example – 1

Calculate individual oxidation number of each S-atom in  $\text{Na}_2\text{S}_4\text{O}_6$  (sodium tetrathionate) with the help of its structure.



### Example – 2

Find the average and individual oxidation number of Fe & Pb in  $\text{Fe}_3\text{O}_4$  &  $\text{Pb}_3\text{O}_4$ , which are mixed oxides.

Sol. (i)  $\text{Fe}_3\text{O}_4$  is mixture of  $\text{FeO}$  &  $\text{Fe}_2\text{O}_3$  in 1 : 1 ratio  
so, individual oxidation number of Fe = +2 & +3

$$\& \text{average oxidation number} = \frac{1(+2) + 2(+3)}{3} = 8/3$$






(ii)  $\text{Pb}_3\text{O}_4$  is a mixture of  $\text{PbO}$  &  $\text{PbO}_2$  in 2 : 1 molar ratio  
so, individual oxidation number of Pb are +2 & +4

$$\& \text{average oxidation number of Pb} = \frac{2(+2) + 1(+4)}{3} = 8/3$$

### Example – 3

Balance the following equations :

- $\text{H}_2\text{O}_2 + \text{MnO}_4^- \longrightarrow \text{Mn}^{+2} + \text{O}_2$  (acidic medium)
- $\text{Zn} + \text{HNO}_3 \text{ (dil)} \longrightarrow \text{Zn}(\text{NO}_3)_2 + \text{H}_2\text{O} + \text{NH}_4\text{NO}_3$
- $\text{CrI}_3 + \text{KOH} + \text{Cl}_2 \longrightarrow \text{K}_2\text{CrO}_4 + \text{KIO}_4 + \text{KCl} + \text{H}_2\text{O}$
- $\text{P}_2\text{H}_4 \longrightarrow \text{PH}_3 + \text{P}_4$
- $\text{Ca}_3(\text{PO}_4)_2 + \text{SiO}_2 + \text{C} \longrightarrow \text{CaSiO}_3 + \text{P}_4 + \text{CO}$



**Example – 4**

Calculate the normality of a solution obtained by mixing 50 mL of 5M solution of  $K_2Cr_2O_7$  and 50 mL of 2 M  $K_2Cr_2O_7$  in acidic medium.

**Sol.** v.f. of  $K_2Cr_2O_7 = 6$

$$\text{so } N_f = \frac{N_1 V_1 + N_2 V_2}{V_1 + V_2}$$

$$= \frac{5 \times 6 \times 50 + 2 \times 6 \times 50}{50 + 50} = 21 \text{ N}$$

**Example – 5**

Calculate the normality of a solution containing 13.4 g of sodium oxalate in 100 mL Sol.

**Sol.** Normality =  $\frac{\text{wt. in g/eq. wt}}{\text{vol of solution in litre}}$

Here, eq. wt. of  $Na_2C_2O_4 = 134/2 = 67$

$$\text{so } N = \frac{13.4/67}{100/1000} = 2 \text{ N}$$

**Example – 6**

The number of moles of ferrous oxalate oxidised by one mole of  $KMnO_4$  in acidic medium is :

- (a)  $\frac{5}{2}$  (b)  $\frac{2}{5}$   
(c)  $\frac{3}{5}$  (d)  $\frac{5}{3}$

**Sol.** Eq. of  $FeC_2O_4 = \text{Eq. of } KMnO_4$   
moles of  $FeC_2O_4 \times 3 = \text{moles of } KMnO_4 \times 5$   
so, moles of  $FeC_2O_4 = 5/3$  **Ans. (d)**

**Example – 7**

How many moles of  $KMnO_4$  are needed to oxidise a mixture of 1 mole of each  $FeSO_4$  &  $FeC_2O_4$  in acidic medium ?

- (a)  $\frac{4}{5}$  (b)  $\frac{5}{4}$   
(c)  $\frac{3}{4}$  (d)  $\frac{5}{3}$

**Sol.** Eq. of  $KMnO_4 = \text{Eq. of } FeSO_4 + \text{Eq. of } FeC_2O_4$   
moles of  $KMnO_4 \times 5 = \text{moles of } FeSO_4 \times 1 + \text{moles of } FeC_2O_4 \times 3$   
 $\therefore$  moles of  $KMnO_4 = 4/5$  **Ans. (a)**

**Example – 8**

A fresh  $H_2O_2$  solution is labelled 11.2 V. This solution has the same concentration as a solution which is :

- (a) 3.4 % (w/w) (b) 3.4% (v/v)  
(c) 3.4% (w/v) (d) None of these

**Sol.** Molarity of  $H_2O_2 = \frac{\text{vol. strength}}{11.2} = \frac{11.2}{11.2} = 1$

$$\text{Now, \% (w/v)} = \frac{\text{wt. of solute in g}}{\text{wt. of solution in mL}} \times 100$$

$$= \text{Molarity} \times \text{Mol. wt. of solute} \times \frac{1}{10}$$

$$= 1 \times 34 \times \frac{1}{10} = 3.4\% \quad \text{Ans. (c)}$$

**Example – 9**

100 mL each of 1N  $H_2O_2$  and 11.2 V  $H_2O_2$  solution are mixed, then the final solution is equivalent to :

- (a) 3 M  $H_2O_2$  solution (b) 0.5 N  $H_2O_2$  solution  
(c) 25.5 g/L  $H_2O_2$  solution  
(d) 2.55 g/L  $H_2O_2$  Sol.

**Sol.**  $N_{\text{final}} = \frac{N_1 V_1 + N_2 V_2}{V_1 + V_2} = \frac{1 \times 100 + \left(\frac{11.2}{5.6}\right) \times 100}{100 + 100} = 3/2 = 1.5 \text{ N}$

$$\text{So, Molarity} = \frac{\text{Normality}}{\text{v.f.}} = \frac{1.5}{2} = 0.75 \text{ M}$$

Strength of solution in g/L = Molarity  $\times$  Mol. wt. =  $0.75 \times 34 = 25.5 \text{ g/L}$  **Ans. (c)**

## EXERCISE - 1 : BASIC OBJECTIVE QUESTIONS

### Oxidizing/Reducing Agents

1. The compound that can work both as an oxidising as well as a reducing agent is :  
 (a)  $\text{KMnO}_4$  (b)  $\text{H}_2\text{O}_2$   
 (c)  $\text{Fe}_2(\text{SO}_4)_3$  (d)  $\text{K}_2\text{Cr}_2\text{O}_7$
2. Which of the following behaves as both oxidising and reducing agents?  
 (a)  $\text{H}_2\text{SO}_4$  (b)  $\text{SO}_2$   
 (c)  $\text{H}_2\text{S}$  (d)  $\text{HNO}_3$
3. Which of the following is not a redox reaction?  
 (a)  $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$  (b)  $\text{O}_2 + 2\text{H}_2 \rightarrow 2\text{H}_2\text{O}$   
 (c)  $\text{Na} + \text{H}_2\text{O} \rightarrow \text{NaOH} + 1/2 \text{H}_2$   
 (d)  $\text{MnCl}_3 \rightarrow \text{MnCl}_2 + 1/2 \text{Cl}_2$
4. Which substance serves as reducing agent in the following reaction?  
 $14\text{H}^+ + \text{Cr}_2\text{O}_7^{2-} + 3\text{Ni} \rightarrow 2\text{Cr}^{3+} + 7 \text{H}_2\text{O} + 3\text{Ni}^{2+}$   
 (a)  $\text{H}_2\text{O}$  (b)  $\text{Ni}$   
 (c)  $\text{H}^+$  (d)  $\text{Cr}_2\text{O}_7^{2-}$
5. Which of the following reactions depicts the oxidising property of  $\text{SO}_2$ ?  
 (a)  $\text{SO}_2 + \text{H}_2\text{O} \longrightarrow \text{H}_2\text{SO}_3$   
 (b)  $2\text{H}_2\text{S} + \text{SO}_2 \longrightarrow 3\text{S} + 2\text{H}_2\text{O}$   
 (c)  $\text{Cl}_2 + \text{SO}_2 \longrightarrow \text{SO}_2\text{Cl}_2$   
 (d)  $2\text{MnO}_4^- + 5\text{SO}_2 + 2\text{H}_2\text{O} \longrightarrow 5\text{SO}_4^{2-} + 2\text{Mn}^{2+} + 4\text{H}^+$
6. In which of the following reactions, there is no change in valency?  
 (a)  $4\text{KClO}_3 \longrightarrow 3\text{KClO}_4 + \text{KCl}$   
 (b)  $\text{SO}_2 + 2\text{H}_2\text{S} \longrightarrow 2\text{H}_2\text{O} + 3\text{S}$   
 (c)  $\text{BaO}_2 + \text{H}_2\text{SO}_4 \longrightarrow \text{BaSO}_4 + \text{H}_2\text{O}_2$   
 (d)  $2\text{BaO} + \text{O}_2 \longrightarrow 2\text{BaO}_2$
7. Nitric oxide acts as a reducing agent in the reaction  
 (a)  $4\text{NH}_3 + 5\text{O}_2 \longrightarrow 4\text{NO} + 6\text{H}_2\text{O}$   
 (b)  $2\text{NO} + 3\text{I}_2 + 4\text{H}_2\text{O} \longrightarrow 2\text{NO}_3^- + 6\text{I}^- + 8\text{H}^+$   
 (c)  $2\text{NO} + \text{H}_2\text{SO}_3 \longrightarrow \text{N}_2\text{O} + \text{H}_2\text{SO}_4$   
 (d)  $2\text{NO} + \text{H}_2\text{S} \longrightarrow \text{N}_2\text{O} + \text{S} + \text{H}_2\text{O}$

### Oxidation Number

8. The oxidation state of osmium (Os) in  $\text{OsO}_4$  is  
 (a) +7 (b) +5  
 (c) +4 (d) +8
9. Oxidation number of nitrogen in  $(\text{NH}_4)_2\text{SO}_4$  is  
 (a)  $-1/3$  (b)  $-1$   
 (c) +1 (d)  $-3$
10. In which of the following compounds, the oxidation number of iodine is fractional?  
 (a)  $\text{IF}_7$  (b)  $\text{I}_3^-$   
 (c)  $\text{IF}_5$  (d)  $\text{IF}_3$
11. The oxidation number of Phosphorus in  $\text{Mg}_2\text{P}_2\text{O}_7$  is :  
 (a) +3 (b) +2  
 (c) +5 (d)  $-3$
12. In which of the following compounds, nitrogen has an oxidation state of  $-1$ ?  
 (a)  $\text{N}_2\text{O}$  (b)  $\text{NO}_2^-$   
 (c)  $\text{NH}_2\text{OH}$  (d)  $\text{N}_2\text{H}_4$
13. In which of the following reactions is there a change in the oxidation number of nitrogen atom?  
 (a)  $2 \text{NO}_2 \rightarrow \text{N}_2\text{O}_4$   
 (b)  $\text{NH}_3 + \text{H}_2\text{O} \rightarrow \text{NH}_4^+ + \text{OH}^-$   
 (c)  $\text{N}_2\text{O}_5 + \text{H}_2\text{O} \rightarrow 2\text{HNO}_3$   
 (d) None of these
14. When  $\text{SO}_2$  is passed through an acidified solution of potassium dichromate the oxidation state of S changes from:  
 (a) +4 to 0 (b) +4 to +2  
 (c) +4 to +6 (d) +6 to +4
15. Oxidation state of nitrogen is correctly given for

Compound	Oxidation state
(a) $[\text{Co}(\text{NH}_3)_5 \text{Cl}]\text{Cl}_2$	0
(b) $\text{NH}_2\text{OH}$	+1
(c) $(\text{N}_2\text{H}_5)_2\text{SO}_4$	+2
(d) $\text{Mg}_3\text{N}_2$	$-3$


16. The oxidation state of chromium in  $\text{Cr}(\text{CO})_6$  is :  
 (a) 0 (b) +2  
 (c) -2 (d) +6
17. In which of the following pairs, there is greatest difference in the oxidation number of the under lined elements?  
 (a)  $\underline{\text{N}}\text{O}_2$  and  $\underline{\text{N}}_2\text{O}_4$  (b)  $\underline{\text{P}}_2\text{O}_5$  and  $\underline{\text{P}}_4\text{O}_{10}$   
 (c)  $\underline{\text{N}}_2\text{O}$  and  $\underline{\text{N}}\text{O}$  (d)  $\underline{\text{S}}\text{O}_2$  and  $\underline{\text{S}}\text{O}_3$
18. In which of the compounds does manganese exhibit highest oxidation number?  
 (a)  $\text{MnO}_2$  (b)  $\text{Mn}_3\text{O}_4$   
 (c)  $\text{K}_2\text{MnO}_4$  (d)  $\text{MnSO}_4$
19. Which of the following elements has least oxidation number ?  
 (a)  $\text{Ni}(\text{CN})_4$  (b)  $\text{Ni}(\text{CO})_4$   
 (c)  $\text{Fe}_2\text{O}_3$  (d)  $\text{SF}_6$
20. The oxidation number of sulphur in  $\text{S}_8$ ,  $\text{S}_2\text{F}_2$ ,  $\text{H}_2\text{S}$  respectively are :  
 (a) 0, +1 and -2 (b) +2, +1 and -2  
 (c) 0, +1 and +2 (d) -2, +1 and -2
21. A metal ion  $\text{M}^{3+}$  loses 3 electrons, its oxidation number will be :  
 (a) +3 (b) +6  
 (c) 0 (d) -3

### Balancing of Redox Reactions

22. For the redox reaction  $\text{MnO}_4^- + \text{C}_2\text{O}_4^{2-} + \text{H}^+ \rightarrow \text{Mn}^{2+} + \text{CO}_2 + \text{H}_2\text{O}$ , The correct stoichiometric coefficients of  $\text{MnO}_4^-$ ,  $\text{C}_2\text{O}_4^{2-}$  and  $\text{H}^+$  are respectively:  
 (a) 2,5,16 (b) 16,5,2  
 (c) 5,16,2 (d) 2,16,5
23. In the chemical reaction,  $\text{K}_2\text{Cr}_2\text{O}_7 + \text{XH}_2\text{SO}_4 + \text{YSO}_2 \rightarrow \text{K}_2\text{SO}_4 + \text{Cr}_2(\text{SO}_4)_3 + \text{ZH}_2\text{O}$ , X, Y and Z are  
 (a) 1,3,1 (b) 4,1,4  
 (c) 3,2,3 (d) 2,1,2
24. For the redox reaction,  
 $\text{MnO}_4^- + \text{C}_2\text{O}_4^{2-} + \text{H}^+ \longrightarrow \text{Mn}^{2+} + \text{CO}_2 + \text{H}_2\text{O}$  the correct coefficients of the reactants for the balanced reaction are:
- |                  |                             |              |
|------------------|-----------------------------|--------------|
| $\text{MnO}_4^-$ | $\text{C}_2\text{O}_4^{2-}$ | $\text{H}^+$ |
| (a) 2            | 5                           | 16           |
| (b) 16           | 5                           | 2            |


- |       |    |   |
|-------|----|---|
| (c) 5 | 16 | 2 |
| (d) 2 | 16 | 5 |

25. What is the coefficient of oxalate ion in the following reaction?



- |       |       |
|-------|-------|
| (a) 4 | (b) 2 |
| (c) 3 | (d) 5 |


26. In the equation



n stands for

- |       |       |
|-------|-------|
| (a) 1 | (b) 2 |
| (c) 3 | (d) 4 |

27. In the chemical reaction,



x, y and z are

- |             |             |
|-------------|-------------|
| (a) 1, 3, 1 | (b) 4, 1, 4 |
| (c) 3, 2, 3 | (d) 2, 1, 2 |

### Equivalent Weight

28. In the conversion  $\text{NH}_2\text{OH} \rightarrow \text{N}_2\text{O}$ , the equivalent weight of  $\text{NH}_2\text{OH}$  will be :

- |                  |                  |
|------------------|------------------|
| (a) $\text{M}/4$ | (b) $\text{M}/2$ |
| (c) $\text{M}/5$ | (d) $\text{M}/1$ |

(M = molecular weight of  $\text{NH}_2\text{OH}$ )


29. The mass of oxalic acid crystals ( $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ ) required to prepare 50 mL of a 0.2 N solution is :

- |            |            |
|------------|------------|
| (a) 4.5 g  | (b) 6.3 g  |
| (c) 0.63 g | (d) 0.45 g |

30. M is molecular weight of  $\text{KMnO}_4$ . The equivalent weight of  $\text{KMnO}_4$  when it is converted into  $\text{K}_2\text{MnO}_4$  is :

- |                  |                  |
|------------------|------------------|
| (a) M            | (b) $\text{M}/3$ |
| (c) $\text{M}/5$ | (d) $\text{M}/7$ |


31. The equivalent mass of oxidising agent in the following reaction is



- |        |        |
|--------|--------|
| (a) 32 | (b) 64 |
| (c) 16 | (d) 8  |



32. In the following reaction :



the equivalent weight of  $\text{As}_2\text{S}_3$  (with molecular weight M) is :

(a)  $\frac{3M}{28}$

(b)  $\frac{M}{4}$

(c)  $\frac{M}{24}$

(d)  $\frac{M}{28}$

33. Equivalent weight of  $\text{H}_3\text{PO}_2$  (molecular weight = M) when it disproportionates into  $\text{PH}_3$  and  $\text{H}_3\text{PO}_3$  is

(a) M

(b)  $\frac{M}{2}$

(c)  $\frac{M}{4}$

(d)  $\frac{3M}{4}$

34. Equivalent weight of  $\text{MnO}_4^-$  in acidic, neutral and basic media are in ratio of :

(a) 3 : 5 : 15

(b) 5 : 3 : 1

(c) 5 : 1 : 13

(d) 3 : 15 : 5

35.  $\text{K}_2\text{CrO}_4$  oxidises KI in the presence of HCl to  $\text{I}_2$ . The equivalent weight of the  $\text{K}_2\text{CrO}_4$  is

(a)  $\frac{\text{Mw}}{2}$

(b)  $\text{Mw} \times \frac{2}{3}$

(c)  $\frac{\text{Mw}}{3}$

(d)  $\frac{\text{Mw}}{6}$

36. The equivalent weight of  $\text{MnSO}_4$  is half its molecular weight when it is converted to



37. In the reaction  $\text{VO} + \text{Fe}_2\text{O}_3 \rightarrow \text{FeO} + \text{V}_2\text{O}_5$  the eq. wt. of  $\text{V}_2\text{O}_5$  is equal to its

(a) mol. wt.

(b) mol. wt./8

(c) mol. wt./6

(d) none of these

38. The eq. wt. of  $\text{K}_2\text{CrO}_4$  as an oxidising agent in acid medium is

(a) mol. wt./2





(b)  $\frac{2 \times \text{mol. wt.}}{3}$

(c)  $\frac{\text{mol. wt.}}{3}$


(d)  $\frac{\text{mol. wt.}}{6}$

## Disproportionation

39. Which of the following is a disproportionation reaction ?



40. In the balanced chemical reaction



a, b, c and d respectively, correspond to

(a) 5, 6, 3, 3

(b) 5, 3, 6, 3

(c) 3, 5, 3, 6

(d) 5, 6, 5, 5

## Titration

41. If equal volumes of 0.1 M  $\text{KMnO}_4$  and 0.1 M  $\text{K}_2\text{Cr}_2\text{O}_7$  solutions are allowed to oxidise  $\text{Fe}^{2+}$  to  $\text{Fe}^{3+}$  in acidic medium, then  $\text{Fe}^{2+}$  oxidised will be :

(a) More by  $\text{KMnO}_4$

(b) More by  $\text{K}_2\text{Cr}_2\text{O}_7$

(c) Equal in both cases

(d) Cannot be determined.

42. Volume  $V_1$  mL of 0.1M  $\text{K}_2\text{Cr}_2\text{O}_7$  is needed for complete oxidation of 0.678 g  $\text{N}_2\text{H}_4$  in acidic medium. The volume of 0.3 M  $\text{KMnO}_4$  needed for same oxidation in acidic medium will be :


(a)  $2/5 V_1$

(b)  $5/2 V_1$

(c)  $113 V_1$

(d) Can not be determined

43. In the chemical reaction,



(a) 1, 3, 1

(b) 4, 1, 4

(c) 3, 2, 3

(d) 2, 1, 2

44. 80 mL of  $\text{KMnO}_4$  solution reacts with 3.4 g of  $\text{Na}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$  in acidic medium. The molarity of the  $\text{KMnO}_4$  solution is


(a) 0.5 M

(b) 0.1 M

(c) 5 M

(d) 1 M

45. What weight of  $\text{NaHSO}_3$  is required to react with 100 mL of solution containing 0.33 g of  $\text{NaIO}_3$  according to the following reaction








- (a) 0.52 g (b) 5.2 g  
(c) 1.04 g (d) 10.4 g
46. How many moles of  $\text{MnO}_4^-$  ion will react with 1 mol of ferrous oxalate in acidic medium ?
- (a)  $\frac{1}{5}$  (b)  $\frac{2}{5}$   
(c)  $\frac{3}{5}$  (d)  $\frac{5}{3}$
47. What volume of 0.05 M  $\text{K}_2\text{Cr}_2\text{O}_7$  in acidic medium is needed for complete oxidation of 200 mL of 0.6 M  $\text{FeC}_2\text{O}_4$  solution?
- (a) 1.2 mL (b) 1.2 L  
(c) 120 mL (d) 800 mL
48. KI reacts with  $\text{H}_2\text{SO}_4$  producing  $\text{I}_2$  and  $\text{H}_2\text{S}$ . The volume of 0.2 M  $\text{H}_2\text{SO}_4$  required to produce 0.1 mol of  $\text{H}_2\text{S}$  is
- (a) 4 L (b) 2.5 L  
(c) 3.8 L (d) 5 L

### Daniell Cell

49. Which of the following statement is correct for a galvanic cell ?
- (a) Reduction occurs at cathode  
(b) Oxidation occurs at anode  
(c) Electrons flow from anode to cathode  
(d) All the statements are correct

### Electrode Potential : Feasibility



50. Consider the following standard reduction potentials:



Which of the following metals is the strongest REDUCING AGENT?





- (a) Ca (b) Pb  
(c) Cu (d) Hg

51. Consider the following electrodes potentials:



Which of the following reactions will proceed spontaneously from left to right?

- (a)  $\text{Mg}^{2+} + \text{V} \rightarrow \text{Mg} + \text{V}^{2+}$   
(b)  $\text{Mg}^{2+} + 2\text{Cu}^+ \rightarrow \text{Mg} + 2\text{Cu}^{2+}$   
(c)  $\text{V}^{2+} + 2\text{Cu}^+ \rightarrow \text{V} + 2\text{Cu}^{2+}$   
(d)  $\text{V} + 2\text{Cu}^{2+} \rightarrow \text{V}^{2+} + 2\text{Cu}^+$
52. The standard reduction potentials at 298 K are given against each of the following half cell reactions :



Based on the above data, state which of the following is the strongest reducing agent ?

- (a) Zn (b) Cr  
(c)  $\text{H}_2$  (d) Fe
53. The correct order of reactivity of K, Mg, Zn and Cu with water according to the electrochemical series is
- (a)  $\text{K} > \text{Mg} > \text{Zn} > \text{Cu}$  (b)  $\text{Mg} > \text{Zn} > \text{Cu} > \text{K}$   
(c)  $\text{K} > \text{Zn} > \text{Mg} > \text{Cu}$  (d)  $\text{Cu} > \text{Zn} > \text{Mg} > \text{K}$

### $\text{H}_2\text{O}_2$

54. 34 g of  $\text{H}_2\text{O}_2$  is present in 1120 mL of solution. This solution is called
- (a) 10 vol solution (b) 20 vol solution  
(c) 34 vol solution (d) 32 vol solution
55. A 5.0 mL solution of  $\text{H}_2\text{O}_2$  liberates 1.27 g of iodine from an acidified KI solution. The percentage strength of  $\text{H}_2\text{O}_2$  is
- (a) 11.2 (b) 5.6  
(c) 1.7 (d) 3.4
56. The volume strength of 1.5 N  $\text{H}_2\text{O}_2$  solution is
- (a) 4.8 (b) 8.4  
(c) 3.0 (d) 8.0

## EXERCISE - 2 : PREVIOUS YEAR JEE MAINS QUESTIONS

1. Which of the following reaction is possible at anode? (2002)
  - (a)  $F_2 + 2e^- \longrightarrow 2F^-$
  - (b)  $2H^+ + \frac{1}{2}O_2 + 2e^- \longrightarrow H_2O$
  - (c)  $2Cr_2^{3+} + 7H_2O \longrightarrow Cr_2O_7^{2-} + 14H^+ + 6e^-$
  - (d)  $Fe^{2+} \longrightarrow Fe^{3+} + e^-$
2. Which of the following is a redox reaction (2002)
  - (a)  $NaCl + KNO_3 \longrightarrow NaNO_3 + KCl$
  - (b)  $CaC_2O_4 + 2HCl \longrightarrow CaCl_2 + H_2C_2O_4$
  - (c)  $Ca(OH)_2 + 2NH_4Cl \longrightarrow CaCl_2 + 2NH_3 + 2H_2O$
  - (d)  $2K[Ag(CN)_2] + Zn \longrightarrow 2Ag + K_2[Zn(CN)_4]$
3.  $MnO_4^-$  is a good oxidising agent in different medium changing to
 

$MnO_4^- \longrightarrow \left\{ \begin{array}{l} Mn^{2+} \\ MnO_4^{2-} \\ MnO_2 \\ Mn_2O_3 \end{array} \right.$

Changes in oxidation number respectively, are (2002)


  - (a) 1, 3, 4, 5
  - (b) 5, 4, 3, 2
  - (c) 5, 1, 3, 4
  - (d) 2, 6, 4, 3
4. Oxidation number of Cl in  $CaOCl_2$  (bleaching powder) is (2002)
  - (a) zero, since it contains  $Cl_2$
  - (b) -1, since it contains  $Cl^-$
  - (c) +1, since it contains  $ClO^-$
  - (d) +1 and -1, since it contains  $ClO^-$  and  $Cl^-$
5. The oxidation state of chromium in the final product formed by the reaction between KI and acidified potassium dichromate solution is (2005)
  - (a) +3
  - (b) +2
  - (c) +6
  - (d) +4
6. The oxidation state of Cr in  $[Cr(NH_3)_4Cl_2]^+$  is (2005)
  - (a) 0
  - (b) +1
  - (c) +2
  - (d) +3
7. Which of the following chemical reactions depicts the oxidising behaviour of  $H_2SO_4$ ? (2006)
  - (a)  $2HI + H_2SO_4 \longrightarrow I_2 + SO_2 + 2H_2O$
  - (b)  $Ca(OH)_2 + H_2SO_4 \longrightarrow CaSO_4 + 2H_2O$
  - (c)  $NaCl + H_2SO_4 \longrightarrow NaHSO_4 + HCl$
  - (d)  $2PCl_5 + H_2SO_4 \longrightarrow 2POCl_3 + 2HCl + SO_2Cl_2$
8. Consider the following reaction,
 
$$xMnO_4^- + yC_2O_4^{2-} + zH^+ \rightarrow xMn^{2+} + 2yCO_2 + \frac{z}{2}H_2O$$

The values of x, y and z in the reaction are, respectively (2013)

  - (a) 5, 2 and 16
  - (b) 2, 5 and 8
  - (c) 2, 5 and 16
  - (d) 5, 2 and 8
9. Given,  $E_{Cr^{3+}/Cr}^\circ = -0.74 \text{ V}$ ;  $E_{MnO_4^-/Mn^{2+}}^\circ = 1.51 \text{ V}$   
 $E_{Cr_2O_7^{2-}/Cr^{3+}}^\circ = 1.33 \text{ V}$ ;  $E_{Cl/Cl^-}^\circ = 1.36 \text{ V}$   
 Based on the data given above strongest oxidising agent will be (2013)
  - (a) Cl
  - (b)  $Cr^{3+}$
  - (c)  $Mn^{2+}$
  - (d)  $MnO_4^-$
10. Which of the following reactions is an example of a redox reaction? (2017)
  - (a)  $XeF_2 + PF_5 \rightarrow [XeF]^+ PF_6^-$
  - (b)  $XeF_6 + H_2O \rightarrow XeOF_4 + 2HF$
  - (c)  $XeF_6 + 2H_2O \rightarrow XeO_2F_2 + 4HF$
  - (d)  $XeF_4 + O_2F_2 \rightarrow XeF_6 + O_2$

## JEE MAINS ONLINE QUESTION


1. Consider the reaction :



Which of the following statements is correct?

**Online 2014 SET (4)**

- (a)  $\text{H}_2\text{SO}_3$  is the reducing agent because it undergoes reduction  
(b)  $\text{Sn}^{4+}$  is the reducing agent because it undergoes oxidation  
(c)  $\text{H}_2\text{SO}_3$  is the reducing agent because it undergoes oxidation  
(d)  $\text{Sn}^{4+}$  is the oxidizing agent because it undergoes oxidation
2. How many electrons are involved in the following redox reaction ?



**Online 2014 SET (4)**


- (a) 3 (b) 4  
(c) 5 (d) 6

3. What is the oxidation number of sulphur in  $\text{Na}_2\text{S}_4\text{O}_6$  ?

**Online 2016 SET (1)**

- (a) 2/3 (b) 3/2  
(c) 3/5 (d) 5/2

4. The value of n in



**Online 2017 SET (1)**

- (a) 5 (b) 4  
(c) 2 (d) 3

5. In  $\text{KO}_2$ , the nature of oxygen species and the oxidation state of oxygen atom are, respectively :

**Online 2018 SET (2)**

- (a) Oxide and -2  
(b) Superoxide and  $-\frac{1}{2}$   
(c) Peroxide and  
(d) Superoxide and -1

## EXERCISE - 3 : ADVANCED OBJECTIVE QUESTIONS

1. All questions marked “S” are single choice questions
2. All questions marked “M” are multiple choice questions
3. All questions marked “C” are comprehension based questions
4. All questions marked “A” are assertion–reason type questions
 

(A) If both assertion and reason are correct and reason is the correct explanation of assertion.

(B) If both assertion and reason are true but reason is not the correct explanation of assertion.

(C) If assertion is true but reason is false.

(D) If reason is true but assertion is false.
5. All questions marked “X” are matrix–match type questions
6. All questions marked “I” are integer type questions

### Oxidizing/Reducing Agents

1. (S) In the reaction,  $2\text{FeCl}_3 + \text{H}_2\text{S} \longrightarrow 2\text{FeCl}_2 + 2\text{HCl} + \text{S}$ 
  - (a)  $\text{FeCl}_3$  acts as an oxidizing agent
  - (b) Both  $\text{H}_2\text{S}$  and  $\text{FeCl}_3$  are oxidized
  - (c)  $\text{FeCl}_3$  is oxidised while  $\text{H}_2\text{S}$  is reduced
  - (d)  $\text{H}_2\text{S}$  acts as an oxidizing agent
2. (S) When  $\text{KMnO}_4$  reacts with acidified  $\text{FeSO}_4$ 
  - (a) Only  $\text{FeSO}_4$  is oxidised
  - (b) Only  $\text{KMnO}_4$  is oxidised
  - (c)  $\text{FeSO}_4$  is oxidised and  $\text{KMnO}_4$  is reduced
  - (d) None of the above
3. (S) Which of the following is a redox reaction?
  - (a)  $\text{H}_2\text{SO}_4$  with  $\text{NaOH}$
  - (b) In atmosphere,  $\text{O}_3$  from  $\text{O}_2$  by lightning
  - (c) Nitrogen oxides from nitrogen and oxygen by lightning
  - (d) Evaporation of  $\text{H}_2\text{O}$
4. (M) Which of the following reactions do not involve oxidation or reduction:
  - (a)  $2\text{Rb} + 2\text{H}_2\text{O} \longrightarrow 2\text{RbOH} + \text{H}_2$
  - (b)  $2\text{CuI}_2 \longrightarrow 2\text{CuI} + \text{I}_2$
  - (c)  $\text{NH}_4\text{Cl} + \text{NaOH} \longrightarrow \text{NaCl} + \text{NH}_3 + \text{H}_2\text{O}$
  - (d)  $4\text{KCN} + \text{Fe}(\text{CN})_2 \longrightarrow \text{K}_4[\text{Fe}(\text{CN})_6]$

5. (M) Which of the following can act both as an oxidising as well as reducing agent :

- |                          |                            |
|--------------------------|----------------------------|
| (a) $\text{HNO}_2$       | (b) $\text{H}_2\text{O}_2$ |
| (c) $\text{H}_2\text{S}$ | (d) $\text{SO}_2$          |

### Oxidation Number

6. (A) **Assertion :**  $\text{MnO}_2$  can act as an oxidizing agent as well as reducing agent.  
**Reason :** Oxidation state of Mn lies between highest and lowest oxidation state.
 

(a) A	(b) B
(c) C	(d) D
7. (S) The oxidation number of cobalt in  $\text{K}_3[\text{Co}(\text{NO}_2)_6]$  is
 

(a) 0	(b) + 4
(c) + 3	(d) + 6
8. (S) Phosphorus has the oxidation state of + 3 in
 

(a) Phosphorous acid	(b) Orthophosphoric
(c) Hypophosphorous acid	(d) Metaphosphoric acid
9. (I) It requires 40 mL of 1 M  $\text{Ce}^{4+}$  to titrate 20 mL of 1M  $\text{Sn}^{2+}$  to  $\text{Sn}^{4+}$ . What is the oxidation state of the Cerium in the product?
10. (M) When  $\text{Cl}_2$  reacts with aqueous  $\text{NaOH}$  in cold condition then oxidation number of chlorine changes from 0 to:
 

(a) -1	(b) +1
(c) -2	(d) +2

## Comprehension

The valency of carbon is generally 4, but its oxidation state may be -4, -2, 0, 2, -1 etc. In the compounds containing C, H and O, the oxidation number of C is calculated as

$$\text{Oxidation number of C} = \frac{2n_{\text{O}} - n_{\text{H}}}{n_{\text{C}}}$$

where  $n_{\text{O}}$ ,  $n_{\text{H}}$ ,  $n_{\text{C}}$  are the number of oxygen, hydrogen, and carbon atoms, respectively.

11. (C) In which of the following compounds is the valency of C two?

- (a) Ketenes (b) Alkenes  
(c) Allenes (d) Carbenes

12. (C) In which of the following compounds is the oxidation state of C highest?

- (a) HCOOH (b) HCHO  
(c) CH<sub>3</sub>OH (d) CH<sub>4</sub>

13. (C) In which of the following compounds is the oxidation state of C a fraction?

- (a) CO (b) CO<sub>2</sub>  
(c) Carbon suboxide (d) All

14. (A) **Assertion :** KO<sub>2</sub> is superoxide.


**Reason :** Oxidation state of oxygen of KO<sub>2</sub> is -2.

- (a) A (b) B  
(c) C (d) D

15. (I) Oxidation number of chlorine in NOClO<sub>4</sub> is .....


## Balancing of Redox Reactions

16. (I) In the redox reaction,




What is the value of  $\frac{x}{z}$ ?

17. Dichromate ion in acidic medium oxidizes stannous ion as:



- (a) the value of x:y is 1 : 3 (b) the value of x+y+z is 18  
(c) a:b is 3 : 2 (d) the value of z-c is 7


18. (S) For the redox reaction,



the correct coefficients of the reactions for the balanced reaction are respectively  $\text{MnO}_4^-$ ,  $\text{C}_2\text{O}_4^{2-}$ ,  $\text{H}^+$  :

- (a) 2, 5, 16 (b) 16, 3, 12  
(c) 15, 16, 12 (d) 2, 16, 5

19. (S) In a chemical reaction



the value of x, y and z respectively are:

- (a) x = 1, y = 3, z = 1 (b) x = 4, y = 1, z = 4  
(c) x = 3, y = 2, z = 1 (d) x = 2, y = 2, z = 1

20. (X)

### Column I

(Redox reaction)

### Column II

(molar ratio of reducing to oxidising agent)

- (A)  $\text{ClO}^- + \text{Fe}(\text{OH})_2 \rightarrow \text{Cl}^- + \text{Fe}(\text{OH})_3$  (p) 3 : 2  
(B)  $\text{Cr}_2\text{O}_7^{2-} + \text{FeC}_2\text{O}_4 \rightarrow \text{Cr}^{3+} + \text{CO}_2 + \text{Fe}^{3+}$  (q) 2 : 1  
(C)  $\text{H}_2\text{O}_2 + \text{Cr}(\text{OH})_3 \rightarrow \text{CrO}_4^{2-} + \text{H}_2\text{O}$  (r) 1 : 3  
(D)  $\text{N}_2\text{H}_4 + \text{Cu}(\text{OH})_2 \rightarrow \text{N}_2\text{O} + \text{Cu}$  (s) 2 : 3  
(E)  $\text{MnO}_4^- + \text{C}_2\text{O}_4^{2-} \rightarrow \text{MnO}_2 + \text{CO}_2$

21. (X) Match the reaction in column I with the coefficients x and y given in column II.

### Column I

(Reaction)

### Column II

(The coefficients of x and y are)

- (A)  $x\text{Cu} + y\text{HNO}_3 \rightarrow \text{Cu}(\text{NO}_3)_2 + \text{NO} + \text{NO}_2 + \text{H}_2\text{O}$  (p) 2 and 6  
(B)  $x\text{KI} + y\text{BaCrO}_4 \xrightarrow{\text{H}^+} \text{I}_2 + \text{CrCl}_3$  (q) 6 and 2  
(C)  $x\text{As}_2\text{S}_3 + y\text{NO}_3^- \rightarrow \text{AsO}_4^{3-} + \text{NO} + \text{SO}_4^{2-}$  (r) 3 and 28  
(D)  $4\text{P} + 3\text{OH}^- + 3\text{H}_2\text{O} \rightarrow x\text{PH}_3 + y\text{H}_2\text{PO}_2$  (s) 1 and 3  
(E)  $x\text{KI} + y\text{H}_2\text{SO}_4 \rightarrow \text{I}_2 + \text{H}_2\text{S} + \text{K}_2\text{SO}_4$  (t) 8 and 5

22. (M) In the reaction



which of the following statements is/are correct?

- (a) The coefficients of  $\text{OH}^-$  and  $\text{I}^-$  in the given in balanced equation are, respectively, 6 and 5.
- (b) The coefficients of  $\text{OH}^-$  and  $\text{I}^-$  in the given balanced equation are, respectively, 5 and 6.
- (c)  $\text{C}_2\text{H}_5\text{OH}$  is oxidised to  $\text{CHI}_3$  and  $\text{HCOO}^-$
- (d) The number of electrons in the conversion of  $\text{C}_2\text{H}_5\text{OH}$  to  $\text{CHI}_3$  and  $\text{HCOO}^-$  is 8.

### Equivalent Weight

23. (I) When  $\text{BrO}_3^-$  ion react with  $\text{Br}^-$  ion in acidic medium,  $\text{Br}_2$  is liberated. Calculate the ratio of molecular weight and equivalent weight of  $\text{KBrO}_3$ .

24. (I) A volume of 12.5 mL of 0.05 M  $\text{SeO}_2$  reacts with 25 mL of 0.1M  $\text{CrSO}_4$  which is oxidised to  $\text{Cr}^{3+}$ . To what oxidation state was the selenium converted by the reaction?

25. (X) 

Column I	Column II
----------	-----------

(A) Eq. wt. =  $\frac{\text{Molecular weight}}{33}$  (p) When  $\text{CrI}_3$  oxidises

into  $\text{Cr}_2\text{O}_7^{2-}$  and  $\text{IO}_4^-$

(B) Eq. wt. =  $\frac{\text{Molecular weight}}{27}$  (q) When  $\text{Fe}(\text{SCN})_2$  oxidises

into  $\text{Fe}^{3+}$ ,  $\text{SO}_4^{2-}$ ,  $\text{CO}_3^{2-}$ , and  $\text{NO}_3^-$

(C) Eq. wt. =  $\frac{\text{Molecular weight}}{28}$  (r) When  $\text{NH}_4\text{SCN}$  oxidizes

into  $\text{SO}_4^{2-}$ ,  $\text{CO}_3^{2-}$  and  $\text{NO}_3^-$

(D) Eq. wt. =  $\frac{\text{Molecular weight}}{24}$  (s) When  $\text{As}_2\text{S}_3$  oxidises

into  $\text{AsO}_3^-$  and  $\text{SO}_4^{2-}$

26. (X) 

Column I	Column II
----------	-----------

(A)  $\text{P}_2\text{H}_4 \rightarrow \text{PH}_3 + \text{P}_4\text{H}_2$

(p)  $E = \frac{3M}{4}$

(B)  $\text{I}_2 \rightarrow \text{I}^- + \text{IO}_3^-$

(q)  $E = \frac{3M}{5}$

(C)  $\text{MnO}_4^- + \text{Mn}^{2+} + \text{H}_2\text{O} \rightarrow \text{Mn}_3\text{O}_4 + \text{H}^+$

(r)  $E = \frac{15M}{26}$

(D)  $\text{H}_3\text{PO}_2 \rightarrow \text{PH}_3 + \text{H}_3\text{PO}_3$

(s)  $E = \frac{5M}{6}$

27. (M) When a equimolar mixture of  $\text{Cu}_2\text{S}$  and  $\text{CuS}$  is titrated with  $\text{Ba}(\text{MnO}_4)_2$  in acidic medium, the final product's contains  $\text{Cu}^{2+}$ ,  $\text{SO}_2$  and  $\text{Mn}^{2+}$ . If the mol. wt. of  $\text{Cu}_2\text{S}$ ,  $\text{CuS}$  and  $\text{Ba}(\text{MnO}_4)_2$  are  $M_1$ ,  $M_2$  and  $M_3$  respectively then:

(a) eq. wt. of  $\text{Cu}_2\text{S}$  is  $\frac{M_1}{8}$

(b) eq. wt. of  $\text{CuS}$  is  $\frac{M_2}{6}$

(c) eq. wt. of  $\text{Ba}(\text{MnO}_4)_2$  is  $\frac{M_3}{5}$

(d)  $\text{Cu}_2\text{S}$  and  $\text{CuS}$  both have same equivalents in mixture

### Comprehension


$$\text{Equivalent weight} = \frac{\text{Molecular weight / Atomic weight}}{n - \text{factor}}$$

n-factor is very important in redox as well as non-redox reactions. With the help of n-factor we can predicts the molar ratio of the reactant species taking part in reactions. The reciprocal of n-factor's ratio of the reactants is the molar ratio of the reactants.

In general n-factor of acid/base is number of moles of  $\text{H}^+/\text{OH}^-$  furnished per mole of acid/base. n-factor of a reactant is no. of moles of electrons lost or gained per mole of reactant.

### Example 1 :

- In acidic medium :  $\text{KMnO}_4$  ( $n = 5$ )  $\rightarrow \text{Mn}^{2+}$
- In neutral medium :  $\text{KMnO}_4$  ( $n = 3$ )  $\rightarrow \text{Mn}^{2+}$
- In basic medium :  $\text{KMnO}_4$  ( $n = 1$ )  $\rightarrow \text{Mn}^{6+}$

**Example 2 :**

Total no. of moles of  $\text{e}^-$  lost by 1 mole of  $\text{FeC}_2\text{O}_4$


$$= 1 + 1 \times 2 \Rightarrow 3$$

$$\therefore \text{ n-factor of } \text{FeC}_2\text{O}_4 = 3$$

**28. (C)** n-factor of  $\text{Ba}(\text{MnO}_4)_2$  in acidic medium is

- (a) 2 (b) 6  
(c) 10 (d) None of these

**29. (C)** For the reaction,



What is the equivalent weight of  $\text{H}_3\text{PO}_2$ ? (mol. wt. is M)

- (a) M (b)  $M/2$   
(c)  $M/3$  (d) None of these

**30. (C)** For the reaction,  $\text{Fe}_{0.95}\text{O}$  (molar mass: M)  $\rightarrow \text{Fe}_2\text{O}_3$ . What is the eq. wt. of  $\text{Fe}_{0.95}\text{O}$ ?

- (a)  $\frac{M}{0.85}$  (b)  $\frac{M}{0.95}$   
(c)  $\frac{M}{0.8075}$  (d) None of these

**31. (S)** In the reaction between  $\text{SO}_2$  and  $\text{O}_3$ , the equivalent weight of ozone is :

- (a) The same as its molecular weight  
(b) Half the molecular weight  
(c) One-third of the molecular weight  
(d) One-fourth of the molecular weight

**32. (I)** In the reaction :  $\text{Na}_2\text{S}_2\text{O}_3 + 4\text{Cl}_2 + 5\text{H}_2\text{O} \longrightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{SO}_4 + 8\text{HCl}$ , the equivalent weight of  $\text{Na}_2\text{S}_2\text{O}_3$  will be : (M = molecular weight of  $\text{Na}_2\text{S}_2\text{O}_3$ )

**Disproportionation**

**33. (A)** **Assertion :**  $\text{F}_2$  does not undergo disproportionation reactions.

**Reason :** Fluorine shows only 0 and -1 oxidation states.

- (a) A (b) B  
(c) C (d) D

**34. (M)** Identify the correct statements with reference to the given reaction,  $\text{P}_4 + 3\text{OH}^- + \text{H}_2\text{O} \rightarrow \text{PH}_3 + 3\text{H}_2\text{PO}_2^-$

- (a) Phosphorus is undergoing reduction only  
(b) Phosphorus is undergoing oxidation only  
(c) Phosphorus is undergoing oxidation as well as reduction.  
(d) Hydrogen is undergoing neither oxidation nor reduction

**35. (A)** **Assertion :** White phosphorous reacts with aqueous caustic soda to form  $\text{PH}_3$  and  $\text{NaH}_2\text{PO}_2$ . It is disproportionation reaction.

**Reason :** In the reaction of disproportionation, same substance is oxidised as well as reduced simultaneously.

- (a) A (b) B  
(c) C (d) D

**Titration**

**36. (S)** 1 mole of equimolar mixture of ferric oxalate and ferrous oxalate will require x mole of  $\text{KMnO}_4$  in acidic medium for complete oxidation, x is:

- (a) 0.5 mole (b) 0.9 mole  
(c) 1.2 mole (d) 4.5 mole

**37. (S)** 20 mL of 0.2 M  $\text{NaOH}$  (aq) solution is mixed with 35 mL of 0.1 M  $\text{NaOH}$  (aq) solution and the resultant solution is diluted to 100 mL. 40 mL of this diluted solution reacted with 10% impure sample of oxalic acid ( $\text{H}_2\text{C}_2\text{O}_4$ ). The weight of impure sample is:

- (a) 0.15 gram (b) 0.135 gram  
(c) 0.59 gram (d) None of these

**38. (S)** A mixture of  $\text{FeO}$  and  $\text{Fe}_2\text{O}_3$  is completely reacted with 100 mL of 0.25 M acidified  $\text{KMnO}_4$  solution. The resultant solution was then titrated with Zn dust which converted  $\text{Fe}^{3+}$  of the solution of  $\text{Fe}^{2+}$ . The  $\text{Fe}^{2+}$  required 1000 mL of 0.10 M  $\text{K}_2\text{Cr}_2\text{O}_7$  solution. Find out the weight %  $\text{Fe}_2\text{O}_3$  in the mixture.

- (a) 80.85 (b) 19.15  
(c) 50 (d) 89.41

**39. (S)** 2 mole, equimolar mixture of  $\text{Na}_2\text{C}_2\text{O}_4$  and  $\text{H}_2\text{C}_2\text{O}_4$  required  $V_1$  L of 0.1 M  $\text{KMnO}_4$  in acidic medium for complete oxidation. The same amount of the mixture required  $V_2$  L of 0.2 M  $\text{NaOH}$  for neutralization. The ratio of  $V_1$  to  $V_2$  is:

- (a) 1 : 2 (b) 2 : 1  
(c) 4 : 5 (d) 5 : 4



40. (S) 32 g of a sample of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  were dissolved in dilute sulphuric acid and water and its volume was made up to 1 litre, 25 mL of this solution required 20 mL of 0.02 M  $\text{KMnO}_4$  solution for complete oxidation. Calculate the weight % of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  in the sample.
- (a) 34.75 (b) 69.5  
(c) 89.5 (d) None of these
41. (S) 125 mL of 63% (w/v)  $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$  solution is made to react with 125 mL of a 40% (w/v) NaOH solution. The resulting solution is : (ignoring hydrolysis of ions)
- (a) Neutral (b) Acidic  
(c) Strongly acidic (d) Alkaline
42. (S) An element A in a compound ABD has oxidation number  $-n$ . It is oxidised by  $\text{Cr}_2\text{O}_7^{2-}$  in acidic medium. In the experiment,  $1.68 \times 10^{-3}$  moles of  $\text{K}_2\text{Cr}_2\text{O}_7$  were used for  $3.36 \times 10^{-3}$  moles of ABD. The new oxidation number of A after oxidation is :
- (a) 3 (b)  $3 - n$   
(c)  $n - 3$  (d)  $+n$
43. (M) 25 mL of 0.5 M  $\text{H}_2\text{O}_2$  solution is added to 50 mL of 0.2 M  $\text{KMnO}_4$  in acid solution. Which of the following statements is false :
- (a) 0.010 mole of oxygen gas is liberated.  
(b) 0.005 mole of  $\text{KMnO}_4$  is left.  
(c) 0.030 g of oxygen gas is evolved.  
(d) 0.0025 mole  $\text{H}_2\text{O}_2$  does not react with  $\text{KMnO}_4$ .
44. (M) Which of the following solutions will exactly oxidize 25 mL of an acid solution of 0.1 M iron (II) oxalate :
- (a) 10 mL of 0.25 M  $\text{KMnO}_4$   
(b) 25 mL of 0.2 M  $\text{KMnO}_4$   
(c) 25 mL of 0.6 M  $\text{KMnO}_4$   
(d) 15 mL of 0.1 M  $\text{KMnO}_4$
45. (M) In the following reaction :  $\text{Cr}(\text{OH})_3 + \text{OH}^- + \text{IO}_3^- \rightarrow \text{CrO}_4^{2-} + \text{H}_2\text{O} + \text{I}^-$
- (a)  $\text{IO}_3^-$  is oxidising agent  
(b)  $\text{Cr}(\text{OH})_3$  is oxidised  
(c)  $6e^-$  are being taken per iodine atom  
(d) None of these
46. (S) In the reaction,  $8\text{Al} + 3\text{Fe}_3\text{O}_4 \rightarrow 4\text{Al}_2\text{O}_3 + 9\text{Fe}$ , the number of electrons transferred from reductant to oxidant is:
- (a) 8 (b) 4  
(c) 7 (d) 24
47. (S) In the reaction of sodium thiosulphate with  $\text{I}_2$  in aqueous medium the equivalent weight of sodium thiosulphate is equal to:
- (a) molar mass of sodium thiosulphate  
(b) the average molar masses of  $\text{Na}_2\text{S}_2\text{O}_3$  and  $\text{I}_2$   
(c) half the molar masses of sodium thiosulphate  
(d) twice of molar mass of sodium thiosulphate
48. (A) **Assertion :** If 1.50 mol of  $\text{KMnO}_4$  is required for oxidation in acidic medium, 2.50 mol of  $\text{KMnO}_4$  would be required for same oxidation but in basic medium.
- Reason :** In acidic medium, oxidation state of Mn changes from +7 to +2, while in basic medium, it changes from +7 to +4.
- (a) A (b) B  
(c) C (d) D
49. (M) 500 mL of a 0.05 M Mohr salt solution required the same volume of permanganate solution for complete oxidation. Which of the followings is (are) true regarding the above redox reaction?
- (a) The molarity of  $\text{KMnO}_4$  must be 0.01 M if the medium is acidic  
(b) The molarity of  $\text{KMnO}_4$  must be 0.0167 M if the medium is basic.  
(c) The medium has no role to play in redox reaction.  
(d) In the above reaction Fe(II) is oxidized to Fe(III) irrespective of medium.

### Daniell Cell




50. (M) In an electrochemical cell, a salt bridge is used:
- (a) to avoid mechanical contact of electrolyte solution in cathodic and anodic half-cell  
(b) to maintain electrical neutrality in the cell  
(c) to avoid liquid junction potential  
(d) to mix the solution of anodic and cathodic half-cell

### Electrode Potential : Feasibility

51. (S) Based on the following information arrange four metals A, B, C and D in order of decreasing ability to act as reducing agents:
- (I) Only A, B and C react with 1 M HCl to give  $\text{H}_2$  (g)  
(II) When C is added to solutions of the other metal ions, metallic B and D are formed  
(III) Metal C does not reduce  $\text{A}^{n+}$ .
- (a)  $\text{C} > \text{A} > \text{B} > \text{D}$  (b)  $\text{C} > \text{A} > \text{D} > \text{B}$   
(c)  $\text{A} > \text{C} > \text{D} > \text{B}$  (d)  $\text{A} > \text{C} > \text{B} > \text{D}$

## Comprehension

The next two questions deal with an experiment. An unknown metal 'X' is found to react spontaneously with 1.0M solution of  $\text{CuSO}_4$  plating out  $\text{Cu(s)}$ . X does not react with a solution of 1.0M  $\text{Zn(NO}_3)_2$ . The half-reactions for these metals are:



52. (C) What is the correct order for listing the metals according to decreasing strength as reducing agent ?

- (a) X, Cu, Zn                      (b) Cu, Zn, X  
(c) Cu, X, Zn                      (d) Zn, X, Cu

53. (C) Another metal Y displaces  $\text{Cu}^{2+}$  from its aqueous solution but can't displace  $\text{Zn}^{2+}$  from its aqueous solution. Which of the following statements regarding X and Y is correct?

- (a) X is stronger reducing agent than Y  
(b) X is weaker reducing agent than Y  
(c) Both X and Y are weaker reducing agents than Cu  
(d) X can be either stronger or weaker reducing agent than Y.

54. (S) If  $\text{Sn}^{2+} + 2\text{e}^- \rightarrow \text{Sn}$ ;  $E^\circ = -0.14 \text{ V}$



- (a)  $\text{Sn}^{2+}$  will disproportionate to  $\text{Sn}^{4+}$  and Sn  
(b)  $\text{Sn}^{2+}$  is stable and it will not disproportionate  
(c)  $\text{Sn}^{4+}$  is easily reduced to Sn  
(d) none of the above

55. (M) Given that,  $E^\circ_{\text{Ni}^{2+}/\text{Ni}} = -0.25 \text{ V}$ ,  $E^\circ_{\text{Cu}^{2+}/\text{Cu}} = +0.34 \text{ V}$



Which of the following redox processes will not take place in specified direction?

- (a)  $\text{Ni}^{2+}(\text{aq}) + \text{Cu(s)} \rightarrow \text{Ni(s)} + \text{Cu}^{2+}(\text{aq})$   
(b)  $\text{Cu(s)} + 2\text{Ag}^+(\text{aq}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{Ag(s)}$   
(c)  $\text{Cu(s)} + 2\text{H}^+(\text{aq}) \rightarrow \text{Cu}^{2+}(\text{aq}) + \text{H}_2(\text{g})$   
(d)  $\text{Zn(s)} + 2\text{H}^+(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + \text{H}_2(\text{g})$

## $\text{H}_2\text{O}_2$

56. (I) 20 mL of  $\text{H}_2\text{O}_2$  after acidification with dilute  $\text{H}_2\text{SO}_4$  required 30 mL of N/12  $\text{KMnO}_4$  for complete oxidation. The strength of  $\text{H}_2\text{O}_2$  solution is in g/L : [Molar mass of  $\text{H}_2\text{O}_2 = 34$ ]

57. (I) A sample of 28 mL of  $\text{H}_2\text{O}_2(\text{aq})$  solution required 10 mL of 0.1  $\text{KMnO}_4(\text{aq})$  solution for complete reaction in acidic medium. What is the volume strength of  $\text{H}_2\text{O}_2$ ?

58. (S) A bottle of  $\text{H}_2\text{O}_2$  is labelled as 10 vol  $\text{H}_2\text{O}_2$ . 112 mL of this solution of  $\text{H}_2\text{O}_2$  is titrated against 0.04 M acidified solution of  $\text{KMnO}_4$ . The volume of  $\text{KMnO}_4$  in litre is

- (a) 1 L                                      (b) 2 L  
(c) 3 L                                      (d) 4 L

## EXERCISE - 4 : PREVIOUS YEAR JEE ADVANCED QUESTION

1. The equivalent weight of  $\text{MnSO}_4$  is half of its molecular weight when it converts to (1988)  
 (a)  $\text{Mn}_2\text{O}_3$  (b)  $\text{MnO}_2$   
 (c)  $\text{MnO}_4^-$  (d)  $\text{MnO}_4^{2-}$
2. The oxidation number of phosphorus in  $\text{Ba}(\text{H}_2\text{PO}_2)_2$  is (1988)  
 (a) +3 (b) +2  
 (c) +1 (d) -1
3. The volume strength of 1.5 N  $\text{H}_2\text{O}_2$  is (1990)  
 (a) 4.8 (b) 8.4  
 (c) 3.0 (d) 8.0
4. For the redox reaction  

$$\text{MnO}_4^- + \text{C}_2\text{O}_4^{2-} + \text{H}^+ \rightarrow \text{Mn}^{2+} + \text{CO}_2 + \text{H}_2\text{O}$$
 The correct coefficients of the reactants for the balanced reaction are  

$\text{MnO}_4^-$	$\text{C}_2\text{O}_4^{2-}$	$\text{H}^+$	(1992)
(a) 2	5	16	
(b) 16	5	2	
(c) 5	16	2	
(d) 2	16	5	
5. The number of moles of  $\text{KMnO}_4$  that will be needed to react with one mole of sulphite ion in acidic solution  
 (a)  $\frac{2}{5}$  (b)  $\frac{3}{5}$  (1997)  
 (c)  $\frac{4}{5}$  (d) 1
6. The number of moles of  $\text{KMnO}_4$  that will be needed to react completely with one mole of ferrous oxalate in acidic medium is (1997)  
 (a)  $\frac{2}{5}$  (b)  $\frac{3}{5}$   
 (c)  $\frac{4}{5}$  (d) 1
7. The normality of 0.3 M phosphorus acid ( $\text{H}_3\text{PO}_3$ ) is  
 (a) 0.1 (b) 0.9 (1999)  
 (c) 0.3 (d) 0.6
8. The oxidation number of sulphur in  $\text{S}_8$ ,  $\text{S}_2\text{F}_2$ ,  $\text{H}_2\text{S}$  respectively, are (1999)  
 (a) 0, +1 and -2 (b) +2, +1 and -2  
 (c) 0, +1 and +2 (d) -2, +1 and -2
9. Among the following, the species in which the oxidation number of an element is +6 (2000)  
 (a)  $\text{MnO}_4^-$  (b)  $\text{Cr}(\text{CN})_6^{3-}$   
 (c)  $\text{NiF}_6^{2-}$  (d)  $\text{CrO}_2\text{Cl}_2$
10. An aqueous solution of 6.3 g oxalic acid dihydrate is made up to 250 mL. The volume of 0.1 N NaOH required to completely neutralize 10 mL of this solution is (2000)  
 (a) 40 mL (b) 20 mL  
 (c) 10 mL (d) 4 mL
11. The reaction,  $3\text{ClO}(\text{aq})^- \longrightarrow \text{ClO}_3(\text{aq})^- + 2\text{Cl}(\text{aq})^-$  is an example of (2001)  
 (a) oxidation reaction  
 (b) reduction reaction  
 (c) disproportionation reaction  
 (d) decomposition reaction
12. In the standardization of  $\text{Na}_2\text{S}_2\text{O}_3$  and  $\text{K}_2\text{Cr}_2\text{O}_7$  by iodometry, the equivalent weight of  $\text{K}_2\text{Cr}_2\text{O}_7$  is (2000)  
 (a) (molecular weight)/2 (b) (molecular weight)/6  
 (c) (molecular weight)/3 (d) same as molecular weight
13. Consider a titration of potassium dichromate solution with acidified Mohr's salt solution using diphenylamine as indicator. The number of moles of Mohr's salt required per mole of dichromate is (2007)  
 (a) 3 (b) 4  
 (c) 5 (d) 6

14. Which ordering of compounds is according to the decreasing order of the oxidation state of nitrogen ?

(2012)

- (a)  $\text{HNO}_3, \text{NO}, \text{NH}_4\text{Cl}, \text{N}_2$
- (b)  $\text{HNO}_3, \text{NO}, \text{N}_2, \text{NH}_4\text{Cl}$
- (c)  $\text{HNO}_3, \text{NH}_4\text{Cl}, \text{NO}, \text{N}_2$
- (d)  $\text{NO}, \text{HNO}_3, \text{NH}_4\text{Cl}, \text{N}_2$


15. The reaction of white phosphorus with aqueous NaOH gives phosphine along with another phosphorus containing compound. The reaction type; the oxidation states of phosphorus in phosphine and the other product are respectively

(2013)

- (a) redox reaction;  $-3$  and  $-5$
- (b) redox reaction;  $3$  and  $+5$
- (c) disproportionation reaction;  $-3$  and  $+5$
- (d) disproportionation reaction;  $-3$  and  $+3$

16. For the reaction :

(2016)



The correct statement(s) in the balanced equation is/are:

- (a) Stoichiometric coefficient of  $\text{HSO}_4^-$  is 6.
- (b) Iodide is oxidized
- (c) Sulphur is reduced
- (d)  $\text{H}_2\text{O}$  is one of the products.

### Assertion and Reason

17. **Assertion :** In the titration of  $\text{Na}_2\text{CO}_3$  with HCl using methyl orange indicator, the volume required at the equivalence point is twice that of the acid required using phenolphthalein indicator.

**Reason :** Two moles of HCl are required for the complete neutralization of one mole of  $\text{Na}_2\text{CO}_3$ .

(1991)

- (a) Assertion is true; Reason is true; Reason is the correct explanation of Assertion.
- (b) Assertion is true; Reason is true; Reason is not the correct explanation of Assertion.
- (c) Assertion is true; Reason is false.
- (d) Assertion is false; Reason is true.

### Fill in the Blanks

18. The compound  $\text{YBa}_2\text{Cu}_3\text{O}_7$ , which shows super conductivity, has copper in oxidation state..... Assume that the rare earth element yttrium is in its usual  $+3$  oxidation state.

(1994)

### Integer Type Questions

19. A student performs a titration with different burettes and finds titre values of 25.2 mL, 25.25 mL, and 25.0 mL. The number of significant figures in the average titre value is

(2010)

20. Among the following, the number of elements showing only one non-zero oxidation state is O, Cl, F, N, P, Sn, Tl, Na, Ti

(2010)

21. The difference in the oxidation numbers of the two types of sulphur atoms in  $\text{Na}_2\text{S}_4\text{O}_6$  is

(2011)

### Subjective Questions

22. The density of a 3 M sodium thiosulphate solution ( $\text{Na}_2\text{S}_2\text{O}_3$ ) is 1.25 g per mL. Calculate (i) the percentage by weight of sodium thiosulphate (ii) the mole fraction of sodium thiosulphate and (iii) the molalities of  $\text{Na}^+$  and  $\text{S}_2\text{O}_3^{2-}$  ions.

(1983)

23. 4.08 g of a mixture of BaO and unknown carbonate  $\text{MCO}_3$  was heated strongly. The residue weighed 3.64 g. This was dissolved in 100 mL of 1 N HCl. The excess acid required 16 mL of 2.5 N NaOH solution for complete neutralization. Identify the metal M.

(1983)

24.  $2.68 \times 10^{-3}$  moles of a solution containing an ion  $\text{A}^{n+}$  require  $1.61 \times 10^{-3}$  moles of  $\text{MnO}_4^-$  for the oxidation of  $\text{A}^{n+}$  to  $\text{AO}_3^-$  in acidic medium. What is the value of n?



(1984)

25. 5 mL of 8 N nitric acid, 4.8 mL of 5 N hydrochloric acid and a certain volume of 17 M sulphuric acid are mixed together and made up to 2 L. 30 mL of this acid mixture exactly neutralise 42.9 mL of sodium carbonate solution containing one gram of  $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$  in 100 mL of water. Calculate the amount in gram of the sulphate ions in solution.

(1985)

26. A sample of hydrazine sulphate ( $\text{N}_2\text{H}_6\text{SO}_4$ ) was dissolved in 100 mL of water, 10 mL of this solution was reacted with excess of ferric chloride solution and warmed to complete the reaction. Ferrous ion formed was estimated and it, required 20 mL of M/50 potassium permanganate solution. Estimate the amount of hydrazine sulphate in one litre of the solution

### Reaction:




27. An equal volume of a reducing agent is titrated separately with 1 M  $\text{KMnO}_4$  in acid, neutral and alkaline medium. The volumes of  $\text{KMnO}_4$  required are 20 mL in acid, 33.3 mL in neutral and 100 mL in alkaline media. Find out the oxidation state of manganese in each reduction product. Give the balanced equations for all the three half reaction. Find out the volume of 1 M  $\text{K}_2\text{Cr}_2\text{O}_7$  consumed, if the same volume of the reducing agent is titrated in acid medium.

(1989)

28. An organic compound X on analysis gives 24.24 per cent carbon and 4.04 per cent hydrogen. Further, sodium extract of 1.0 g of X gives 2.90 g of silver chloride with acidified silver nitrate solution. The compound X may be represented by two isomeric structures Y and Z. Y on treatment with aqueous potassium hydroxide solution gives a dihydroxy compound while Z on similar treatment gives ethanal. Find out the molecular formula of X and gives the structure of Y and Z. (1989)
29. A mixture of  $\text{H}_2\text{C}_2\text{O}_4$  (oxalic acid) and  $\text{NaHC}_2\text{O}_4$  weighing 2.02 g was dissolved in water and the solution made up to one litre. Ten millilitres of the solution required 3.0 mL of 0.1 N sodium hydroxide solution for complete neutralization. In another experiment, 10.0 mL of the same solution, in hot dilute sulphuric acid medium, required 4.0 mL of 0.1 N potassium permanganate solution for complete reaction. Calculate the amount of  $\text{H}_2\text{C}_2\text{O}_4$  and  $\text{NaHC}_2\text{O}_4$  in the mixture. (1990)
30. A solution of 0.2 g of a compound containing  $\text{Cu}^{2+}$  and  $\text{C}_2\text{O}_4^{2-}$  ions on titration with 0.02 M  $\text{KMnO}_4$  in presence of  $\text{H}_2\text{SO}_4$  consumes 22.6 mL of the oxidant. The resultant solution is neutralized with  $\text{Na}_2\text{CO}_3$ , acidified with dilute acetic acid and treated with excess KI. The liberated iodine requires 11.3 mL of 0.05 M  $\text{Na}_2\text{S}_2\text{O}_3$  solution for complete reduction. Find out the mole ratio of  $\text{Cu}^{2+}$  to  $\text{C}_2\text{O}_4^{2-}$  in the compound. Write down the balanced redox reactions involved in the above titrations. (1991)
31. A 1.0 g sample of  $\text{Fe}_2\text{O}_3$  solid of 55.2% purity is dissolved in acid and reduced by heating the solution with zinc dust. The resultant solution is cooled and made up to 100.0 mL. An aliquot of 25.0 mL of this solution requires for titration. Calculate the number of electrons taken up by the oxidant in the reaction of the above titration. (1991)
32. A 2.0 g sample of a mixture containing sodium carbonate, sodium bicarbonate and sodium sulphate is gently heated till the evolution of  $\text{CO}_2$  ceases. The volume of  $\text{CO}_2$  at 750 mm Hg pressure and at 298 K is measured to be 123.9 mL. A 1.5 g of the same sample requires 150 mL of (M/10) HCl for complete neutralization. Calculate the percentage composition of the components of the mixture. (1992)
33. One gram of commercial  $\text{AgNO}_3$  is dissolved in 50 mL of water. It is treated with 50 mL of a KI solution. The silver iodide thus precipitated is filtered off. Excess of KI in the filtrate is titrated with (M/10)  $\text{KIO}_3$  solution in presence of 6 M HCl till all  $\text{I}^-$  ions are converted into  $\text{ICl}$ . It requires 50 mL of (M/10)  $\text{KIO}_3$  solution. 20 mL of the same stock solution of KI requires 30 mL of (M/10)  $\text{KIO}_3$  under similar conditions. Calculate the percentage of  $\text{AgNO}_3$  in the sample.
34. A 5.0  $\text{cm}^3$  solution of  $\text{H}_2\text{O}_2$  liberates 0.508 g of iodine from an acidified KI solution. Calculate the strength of  $\text{H}_2\text{O}_2$  solution in terms of volume strength at STP (1995)
35. A 20.0  $\text{cm}^3$  mixture of CO,  $\text{CH}_4$  and He gases is exploded by an electric discharge at room temperature with excess of oxygen. The volume contraction is found to be 13.0  $\text{cm}^3$ . A further contraction of 14.0  $\text{cm}^3$  occurs when the residual gas is treated with KOH solution. Find out the composition of the gaseous mixture in terms of volume percentage. (1995)
36. A 3.00 g sample containing  $\text{Fe}_3\text{O}_4$ ,  $\text{Fe}_2\text{O}_3$  and an inert impure substance, is treated with excess of KI solution in presence of dilute  $\text{H}_2\text{SO}_4$ . The entire iron is converted into  $\text{Fe}^{2+}$  along with liberation of iodine. The resulting solution is diluted to 100 mL. A 20 mL of the dilute solution requires 11.0 mL of 0.5 M  $\text{Na}_2\text{S}_2\text{O}_3$  solution to reduce the iodine present. A 50 mL of the dilute solution, after complete extraction of the iodine required 12.80 mL of 0.25 M  $\text{KMnO}_4$  solution in dilute  $\text{H}_2\text{SO}_4$  medium for the oxidation of  $\text{Fe}^{2+}$ . Calculate the percentage of  $\text{Fe}_2\text{O}_3$  and  $\text{Fe}_3\text{O}_4$  in the original sample. (1996)
37. To a 25 mL  $\text{H}_2\text{O}_2$  solution, excess of acidified solution of potassium iodide was added. The iodine liberated required 20 mL of 0.3 N sodium thiosulphate solution. Calculate the volume strength of  $\text{H}_2\text{O}_2$  solution. (1997)
38. An aqueous solution containing 0.10 g  $\text{KIO}_3$  (formula weight = 214.0) was treated with an excess of KI solution. The solution was acidified with HCl. The liberated  $\text{I}_2$  consumed 45.0 mL of thiosulphate solution decolourise the blue starch-iodine complex. Calculate the molarity of the sodium thiosulphate solution. (1998)
39. How many millilitres of 0.5 M  $\text{H}_2\text{SO}_4$  are needed to dissolve 0.5 g of copper (II) carbonate? (1999)
40. Hydrogen peroxide solution (20 mL) reacts quantitatively with a solution of  $\text{KMnO}_4$  (20 mL) acidified with dilute  $\text{H}_2\text{SO}_4$ . The same volume of the  $\text{KMnO}_4$  solution is just decolourised by 10 mL of  $\text{MnSO}_4$  in neutral medium simultaneously forming a dark brown precipitate of hydrated  $\text{MnO}_2$ . The brown precipitate is dissolved in 10 mL of 0.2 M sodium oxalate under boiling condition in the presence of dilute  $\text{H}_2\text{SO}_4$ . Write the balanced equation involved in the reactions and calculate the molarity of  $\text{H}_2\text{O}_2$  (2001)
41. Calculate the amount of calcium oxide required when it reacts with 852 g of  $\text{P}_4\text{O}_{10}$ . (2005)

**Reaction :**



## ANSWER KEY

### Exercise - 1 : (Basic Objective Questions)

1. (b)	2. (b)	3. (a)	4. (b)	5. (b)	6. (c)	7. (b)	8. (d)	9. (d)	10. (b)
11. (c)	12. (c)	13. (d)	14. (c)	15. (d)	16. (a)	17. (d)	18. (c)	19. (b)	20. (a)
21. (b)	22. (a)	23. (a)	24. (a)	25. (d)	26. (b)	27. (a)	28. (b)	29. (c)	30. (a)
31. (c)	32. (d)	33. (d)	34. (a)	35. (c)	36. (b)	37. (c)	38. (c)	39. (a)	40. (a)
41. (b)	42. (a)	43. (a)	44. (b)	45. (a)	46. (c)	47. (b)	48. (b)	49. (d)	50. (a)
51. (d)	52. (a)	53. (a)	54. (a)	55. (d)	56. (b)				

### Exercise - 2 : (Previous Year JEE Mains Questions)

1. (cd)	2. (d)	3. (c)	4. (d)	5. (a)	6. (d)	7. (a)	8. (c)	9. (d)	10. (d)
---------	--------	--------	--------	--------	--------	--------	--------	--------	---------

### JEE Mains Online

1. (c)	2. (d)	3. (d)	4. (a)	5. (b)
--------	--------	--------	--------	--------

### Exercise - 3 : (Advanced Objective Questions)

1. (a)	2. (c)	3. (c)	4. (cd)	5. (abd)	6. (a)	7. (c)	8. (a)	9. + 3	10. (ab)
11. (d)	12. (a)	13. (c)	14. (c)	15. (0007)	16. (0007)	17. (bcd)	18. (a)	19. (a)	
20. $A \rightarrow q; B \rightarrow q; C \rightarrow s; D \rightarrow r; E \rightarrow p$	21. $A \rightarrow p; B \rightarrow q; C \rightarrow r; D \rightarrow s; E \rightarrow t$	22. (acd)	23. (0005)						
24. (0000)	25. $A \rightarrow q; B \rightarrow p; C \rightarrow s; D \rightarrow r$	26. $A \rightarrow s; B \rightarrow q; C \rightarrow r; D \rightarrow p$	27. (ab)	28. (c)					
29. (a)	30. (a)	31. (b)	32. M/8	33. (a)	34. (cd)	35. (a)	36. (b)	37. (a)	38. (a)
39. (c)	40. (b)	41. (a)	42. (b)	43. (acd)	44. (d)	45. (abc)	46. (d)	47. (a)	48. (a)
49. (abd)	50. (abc)	51. (d)	52. (d)	53. (d)	54. (b)	55. (ac)	56. 2 g/L	57. (0001)	
58. (a)									

### Exercise - 4 : (Previous Year JEE Advanced Questions)

1. (b)	2. (c)	3. (b)	4. (a)	5. (a)	6. (b)	7. (d)	8. (a)	9. (d)	10. (a)
11. (c)	12. (b)	13. (d)	14. (b)	15. (c)	16. (b)	17. (a,b,d)	18. $\frac{7}{3}$	19. (0003)	20. (0002)
21. (0005)	22. (i) 37.92, (ii) 0.065, (iii) 7.73 and 3.86	23. Calcium	24. 2	25. 6.5g	26. 6.5g/L	27. 16.67 mL			
28. Leave	29. 1.12g	30. 1 : 2	31. $1.04 \times 10^{21}$	32. 26.5%	33. 85%	34. 4.48%	35. 50%, 20%, 30%		
36. 49.33%, 34.8%	37. 1.33V	38. 0.06	39. 8.1 mL	40. 0.1 M	41. 10.08g				