

Chapter 26

Aldehydes, Ketones and Carboxylic acids

(A) ALDEHYDES AND KETONES

Structure of carbonyl group



Hybridisation state of carbon in carbonyl group is sp^2 and sp^2 orbital overlaps with p -orbital of oxygen and forms C-O, sp^2-p , σ -bond.

The carbonyl group is polar in nature. The π -bond cloud is unsymmetrical because of the greater electronegativity of oxygen (3.5) than the carbon (2.5).


PREPARATION OF ALDEHYDES AND KETONES

I. For Both Aldehydes and Ketones

(i) By Oxidation of Alcohols





Aldehydes are quite susceptible to further oxidation to acids.



Thus oxidation of primary alcohols is made at the temperature much above the B.P. of aldehyde and thus aldehydes are vapourised out and prevented from being oxidised.


Note : Collins reagent ($CrO_3 \cdot 2C_5H_5N$) and pyridinium chlorochromate ($C_5H_5NH^+CrO_3Cl^-$) are mild oxidising agents thereby preventing the further oxidation of aldehydes to carboxylic acids.

(ii) By Dehydrogenation of Alcohols




(iii) From Hydrocarbons


(a) By ozonolysis of alkenes



To prevent further oxidation of carbonyl compound by H_2O_2 , we add zinc in the reaction to destroy H_2O_2



(b) By hydration of alkynes





This is called Kucherov reaction.

Note:



- Formaldehyde can't be prepared by this method
- Only ethyne gives acetaldehyde while other alkynes give ketones.






(c) By Wacker's process

Reagent : Acidified aqueous solution of $PdCl_2$ and $CuCl_2$






(iv) Oxidation of Nitroalkane (NEF Reaction)





(v) By Hydrolysis of Oximes:**(vi) By Dry Distillation of Calcium Salts of Fatty Acids****II. For Aldehydes Only****(i) From Acyl/Acid Chloride**

Note: Formaldehyde can not be prepared by this method because, the corresponding acid chloride (HCOCl) is unstable.


In the above reaction BaSO_4 acts as catalyst poison and decreases the catalytic efficiency of Pd, due to which further reduction of aldehyde to alcohol can't take place.

(ii) From Nitriles and Esters**(iii) From Hydrocarbons**


(a) By oxidation of methyl benzene






(b) By side-chain chlorination followed by hydrolysis




(c) By Gatterman-Koch reaction





(d) By Gattermann reaction

**III. For Ketones Only****(i) From Acyl Chloride**



Dialkyl cadmium can be obtained by




This method is very important for synthesis of Ketones.


(ii) From Nitriles

Note: If instead of alkyl or aryl nitriles, HCN is used aldehydes are formed.



(iii) By Friedal-Crafts Acylation Reaction

Note: Oppenauer oxidation:

**PHYSICAL PROPERTIES OF ALDEHYDES AND KETONES****(i) Physical State**

Formaldehyde is a gas. All other aldehydes and ketones upto C_{11} are colourless volatile liquids. Higher members are solids at room temperature.

(ii) Odour

Lower aldehydes have an unpleasant odour. Higher aldehydes and ketones have a pleasant odour.


(iii) Boiling and melting points:

Boiling point and melting point \propto Molecular weight \propto

$$\frac{1}{\text{Branching}}$$

1. Nucleophilic Addition Reactions

Mechanism of nucleophilic addition to carbonyl group.



Aldehydes are more reactive than ketones due to:

- (i) **Inductive effect:** Alkyl group has +I effect which decreases polarity of carbonyl group. Thus ketones containing two alkyl groups are more reactive in comparison to one alkyl group containing aldehydes.
- (ii) **Steric effects:** As the number and size of alkyl groups increases the attack of the nucleophile on carbonyl group C becomes more and more difficult due to steric hindrance.

Boiling point of aldehydes and ketones are higher than those of hydrocarbons and ethers. This is due to dipole-dipole interactions between the opposite ends of >C=O dipoles.

Ketones have higher Boiling point than isomeric aldehydes. This is due to presence of two electrons donating alkyl groups around >C=O group which makes them more polar.

Boiling point of alcohols and carboxylic acids are higher than aldehydes and ketones. This is due to intermolecular H-bonding in alcohols and acids which is much stronger than dipole-dipole interactions in aldehydes and ketones.

(iv) Solubility :


Lower aldehydes and ketones are soluble in water due to H-bonding between the polar carbonyl group and the water molecules. Solubility in H_2O decreases with increase in size of alkyl group. Due to larger hydrocarbon part, aromatic aldehydes and ketones are much less soluble than aliphatic analogues. All aldehydes and ketones are fairly soluble in organic solvents.

CHEMICAL PROPERTIES OF ALDEHYDES AND KETONES







The reaction of aldehydes and ketones can be divided into the following categories :

1. Nucleophilic addition reactions.
2. Reduction reactions.
3. Oxidation reactions
4. Reactions with alkalis.
5. Miscellaneous reactions.

The various nucleophilic addition reactions are:



Some ammonia-substituted derivatives of aldehydes and ketones


Z	Reagent Name	Carbonyl derivative	Product name
- H	Ammonia	>C=NH	Imine
- R	Amine	>C=NR	Substituted imine (Schiff's base)
- OH	Hydroxylamine	>C=N-OH	Oxime
- NH ₂	Hydrazine	>C=N-NH_2	Hydrazone
- HN- 	Phenylhydrazine	>C=N-NH- 	Phenylhydrazone
- HN- 	2, 4 - Dinitrophenylhydrazine	>C=N-NH- 	2, 4-Dinitrophenylhydrazone
- NH- 	Semicarbazide	>C=N-NH- 	Semicarbazone

Note: (i) Benzaldehyde forms NaHSO₃ adduct but acetophenone does not.

(ii) In addition of alcohols if dihydric alcohol is used a cyclic acetal or ketal is formed.

(iii) In addition of NH₃ derivatives resulting compounds formed are crystalline solids with sharp melting points. Thus these derivatives are used for identification and characterization of aldehydes and ketones.


2. Reduction Reactions



Note: Aldehydes and ketones can be also reduced to corresponding hydrocarbons by heating with HI in presence of red P at 423K.

3. Oxidation Reactions


Oxidation of Aldehydes




Aldehydes get easily oxidised by strong oxidising agents like HNO_3 , $\text{K}_2\text{Cr}_2\text{O}_7$, KMnO_4 etc. as well as by weak oxidising agents like Fehling's and Tollen's reagents.

Oxidation of Ketones



Strong oxidising agent and elevated temperatures needed. In case of symmetrical ketones mixture of two acids is always obtained.





In case of unsymmetrical ketones ($\text{R} \neq \text{R}'$), keto group stays with smaller alkyl group (Popoff's rule).



For example:



Note: Aldehydes and ketones having one methyl group linked to carbonyl C-atom, i.e., methyl ketones, show haloform reaction



This is **Iodoform** test and is used for characterising compounds with CH_3CO group or $\text{CH}_3\text{CH}(\text{OH})$ group.

4. Reactions with Alkalies


Due to strong electron withdrawing nature ($-\text{I}$ effect) of the carbonyl group and resonance stabilisation of the conjugate base, α -H-atoms of carbonyl compounds are acidic in nature. This acidic nature of α -H gives rise to a large number of reactions of carbonyl compounds.

(i) Aldol Condensation

Two molecules of aldehyde or ketone containing atleast one α -H atom condense in presence of a dilute alkali to form a β -hydroxyaldehyde or a β -hydroxyketone respectively.

- (ii) **Cross-aldol condensation:** An aldol condensation between two different aldehydes or two ketones or between one aldehyde and one ketone.

For example,




Note: Cross aldol condensation between an aromatic aldehyde and an aliphatic aldehyde or a ketone is called Claisen-Schmidt condensation.

- (iii) **Cannizzaro reaction:** Those aldehydes, which do not contain α -hydrogen atoms give this reaction with conc. alkali like conc. NaOH or KOH. In this reaction two molecules of aldehyde react in such a way that one molecule gets oxidised and the other gets reduced. This type of reaction is called disproportionation reaction.

When two molecules in a reaction are same it is called simple Cannizzaro reaction. In case of different molecules, it is a mixed cross Cannizzaro reaction. In mixed cross Cannizzaro reaction

More reactive aldehyde \longrightarrow Gets oxidised


Less reactive aldehyde \longrightarrow Gets reduced




Miscellaneous Reactions

- (i) **Electrophilic substitution reactions:** Carbonyl group is a deactivating and meta-directing group.


(a) **Nitration :**



(b) **Sulphonation :**





- (ii) **Benzoïn condensation :**



(iii) Cold dilute Schiff's reagent (colour less) + Aldehyde \longrightarrow Pink coloured solution

(iv) **With ammonia :**



Hydrobenzamide (amide without $-\text{CONH}_2$ group)




DISTINCTION BETWEEN ALDEHYDES AND KETONES

Test	Aldehydes	Ketones
1. With Tollen's reagent	give silver mirror $\text{RCHO} + 2 [\text{Ag}(\text{NH}_3)_2]^+ + 3\text{OH}^- \longrightarrow \text{RCOO}^- + 2\text{Ag} + 2\text{H}_2\text{O} + 4\text{NH}_3$	No action
2. With Fehling's solution	give red ppt of Cu_2O (only aliphatic aldehydes)	No action
3. Reduction with LiAlH_4	1° alcohols are formed	2° alcohols are formed
4. Action with alcohols in presence of dry HCl gas	Form acetals easily	Do not form ketals easily
5. Action with ammonia	Give aldehyde-ammonia adducts	Form complex condensation products
6. With Schiff's reagent	Restore pink colour	No action


(B) CARBOXYLIC ACIDS

PREPARATION OF CARBOXYLIC ACIDS







(i) **From 1° Alcohols, Aldehydes and Ketones.**



This gives poor yield of carboxylic acids.

(ii) From alkylbenzenes.

Note : Presence of electron withdrawing groups in benzene ring increases the reactivity of the benzylic hydrogens. Thus p-nitrotoluene is much more easily oxidised than toluene.

(iii) From Nitriles and Amides.**(iv) From Grignard Reagent.****(v) From Hydrolysis of Acid Derivatives:****PHYSICAL PROPERTIES OF CARBOXYLIC ACIDS**

(i) Melting point (M.P.) : Melting point of carboxylic acid do not vary smoothly from one molecule to another

For first ten members



Reason : In acids of even number of carbon atoms, terminal $-\text{CH}_3$ and $-\text{COOH}$ groups lie on the opposite sides of the carbon chain which results in more effective packing of the molecules in the lattice. While in acids of odd number of carbon atoms they lie on the same sides on the carbon chain.

(ii) Boiling points: Boiling point \propto Molecular weight




The boiling points of carboxylic acids are higher than hydrocarbons, aldehydes, ketones and even alcohols of comparable molecular masses.

This is due to extensive H-bonding resulting in association of carboxylic acid molecules. The H-bonds formed in carboxylic acids are even more stronger than alcohols. Carboxylic acids exist as cyclic dimer in vapour phase or in aprotic solvents.


(iii) Solubility: Simple aliphatic acids (upto 4 C-atoms) are soluble in H_2O due to H-bonding. Higher acids are insoluble in H_2O due to increased hydrophobic interaction of hydrocarbon part. Aromatic acids are also insoluble due to their large hydrocarbon part. Carboxylic acids are soluble in organic solvents like C_6H_6 , ether, CHCl_3 , etc.

CHEMICAL PROPERTIES OF CARBOXYLIC ACIDS**I. Reactions Due to Cleavage of O-H Bond:**


Acid strength of acids



Note: Acids liberate CO_2 (gas) with NaHCO_3 . This reaction is used to distinguish carboxylic acids from phenols which do not react with NaHCO_3 .



In water, carboxylic acids dissociate as :



$$\Rightarrow K_a = K_{\text{eq}} [\text{H}_2\text{O}] = \frac{[\text{H}_3\text{O}^+][\text{RCOO}^-]}{[\text{RCOOH}]}$$


and $\text{p}K_a = -\log K_a$

\therefore smaller the $\text{p}K_a$, stronger the acid.

The acidity of carboxylic acids, phenols and alcohols follows the order:

Carboxylic acid > phenol > alcohol

Consider the resonating structures of carboxylate ion and phenoxide ion.




This is because the carboxylate ion is stabilised by two equivalent resonance structures with $-ve$ charge at more electronegative O-atom. On the other hand, phenoxide ion has non-equivalent structures with $-ve$ charge on less electronegative C-atom. Thus, contribution of resonating structures of phenoxide ion towards resonance is very small.

Therefore, carboxylate ion is more stable than phenoxide ion so carboxylic acids are more acidic than phenols.

EWG increase the acidity of acid by stabilising the carboxylate ion whereas EDG decrease the acidity by destabilising the ion.

Note:

- More will be the electron withdrawing nature of substituent greater will be acidic strength.
- More will be the number of electron withdrawing substituents greater will be acidic strength.
- As the distance between electron withdrawing substituent and $-\text{COOH}$ group increases acidic strength decreases.
- Direct attachment of phenyl or vinyl groups increase the acidity of acid due to resonance:



In general, acidity increases as the electronegativity of the carbon atom directly attached to $-\text{COOH}$ group increases or the hybridization of the carbon atom directly attached to $-\text{COOH}$ changes from $sp^3 \rightarrow sp^2 \rightarrow sp$.


The order of acidity of various acids is as follows:



Note: o -Substituted benzoic acids are usually stronger acids than benzoic acid regardless of the nature of the substituent.


II. Cleavage of C – OH Bond

(i) Anhydride Formation



Acid anhydrides can also be obtained by treating acid chlorides with carboxylic acids.



(ii) Esterification



Note: As the number and size of the substituents around the $-\text{COOH}$ or $-\text{OH}$ group increases, the rate of esterification decreases.


(iii) Formation of Acid Chlorides.

Reaction with PCl_5 , PCl_3 and SOCl_2



Note: In this reaction, SO_2 being a gas escapes out and HCl gets absorbed by basic pyridine and pure acyl chloride is left. So it is best method for preparing acyl chloride from acid.


(iv) Reaction with NH_3



III. Reaction Involving $-\text{COOH}$ Group


(i) Reduction:

(a) Reduction to alkanes: RCOOH




NaBH_4 does not reduce the carboxyl group B_2H_6 does not reduce groups like ester, $-\text{NO}_2$, halo, etc.



(b) Reduction to alkanes:




(ii) Decarboxylation




(iii) Kolbe's electrolytic synthesis :



(iv) Hunsdiecker reaction :




(v) Formation of carbonyl compounds [Dry Distillation of calcium salt] :



IV. Substitution in Hydrocarbon Part

(i) Halogenation




α - Halocarboxylic acid (X = Cl, Br)

This is **Hell-Volhard-Zelinsky (HVZ)** reaction

Acids which do not have α -hydrogens do not show HVZ reaction

(ii) Ring Substitution :



Note: – COOH group is m-directing. Aromatic acids do not undergo Friedal Crafts reaction.


DISTINCTION BETWEEN PHENOL AND CARBOXYLIC ACID

Test	Carboxylic acid	Phenol
1. Sodium bicarbonate test	R – COOH with NaHCO ₃ , give brisk effervescence due to evolution of CO ₂ gas	No reaction
2. Ferric chloride test	R – COOH give coloured ppt with neutral FeCl ₃ sol Ex: $3\text{CH}_3\text{COOH} + \text{FeCl}_3 \longrightarrow (\text{CH}_3\text{COO})_3\text{Fe} + 3\text{HCl}$ Buff coloured	C ₆ H ₅ OH give violet coloured ppt. with FeCl ₃ solution $3\text{C}_6\text{H}_5\text{OH} + \text{FeCl}_3 \longrightarrow (\text{C}_6\text{H}_5\text{O})_3\text{Fe} + 3\text{HCl}$ ferric phenoxide (violet coloured)


ALDEHYDES, KETONES AND CARBOXYLIC ACIDS

Aldehydes and Ketones


Preparation of aldehydes and ketones



Preparation of aldehydes only





Preparation of ketones only




Properties of aldehydes and ketones


- Aldehydes are more reactive than ketones in nucleophilic addition reactions. These reactions are following



Clemmensen reduction



Wolff-Kishner reduction



- Aldehydes gives positive test with Tollen's reagent and Fehling solution while ketones do not give such test

Aldol condensation


Condensation of aldehydes and ketones having atleast one α -H atom.

Cannizzaro reaction

Aldehydes with no α -hydrogen undergoes, self oxidation and reduction.

Carboxylic acids


Preparation



Properties

- Gives CO_2 with carbonates and NaHCO_3
- 2 Ethanoic acid $\xrightarrow[\text{or P}_2\text{O}_5, \Delta]{\text{H}^+, \Delta} \text{Ethanoic anhydride}$

Esterification








Hell-Volhard Zelinsky reaction



EXERCISE - 1

Conceptual Questions


- Benzaldehyde and alcohol KCN reacts to produce which of the following :
 (a) ethyl benzoate (b) benzyl alcohol
 (c) benzoin (d) benzyl benzoate
 - Reaction of phenylacetylene with dil. H_2SO_4 and HgSO_4 gives
 (a) acetophenone (b) 2-phenylethanol
 (c) phenylacetaldehyde (d) phenylacetic acid
 - A new C — C bond is formed in :
 (a) Cannizzaro's reaction
 (b) Reimer Tiemann reaction
 (c) Friedel Crafts reaction
 (d) both (b) and (c)
 - Two molecules of acetic acid on heating with phosphorous pentoxide produce :
 (a) ethyl alcohol
 (b) acetic anhydride
 (c) acetyl pentaoxide
 (d) pentaoxide of acetic acid
 - Which of the following gases does not form a ketone on treatment with dil H_2SO_4 and 1% HgSO_4 ?
 (a) C_4H_6 (b) C_2H_2
 (c) C_3H_4 (d) All of these
 - A compound X of formula $\text{C}_3\text{H}_8\text{O}$ yields a ketone $\text{C}_3\text{H}_6\text{O}$ on oxidation. To which of the following class of compounds would X belong?
 (a) Aldehyde (b) Tertiary alcohol
 (c) Secondary alcohol (d) Alkene
 - Clemmensen's reduction of ketones is carried out in :
 (a) LiAlH_4 in H_2O (b) glycol and KOH
 (c) Zn-Hg and HCl (d) H_2 and Pd catalyst
 - Suggest appropriate structures for the missing final compound. (The number of carbon atom remains the same throughout the reaction)
- 
- 

- 

- An unknown compound 'D' first oxidised to aldehyde and then acetic acid by a dilute solution of $\text{K}_2\text{Cr}_2\text{O}_7$ and H_2SO_4 . The compound 'D' is :
 (a) CH_3OH (b) $\text{C}_2\text{H}_5\text{OH}$
 (c) $\text{CH}_3\text{CH}_2\text{COOH}$ (d) $\text{CH}_3\text{CH}_2\text{CHO}$
 - Identify the product C in the series

$$\text{CH}_3\text{CN} \xrightarrow{\text{Na/C}_2\text{H}_5\text{OH}} \text{A} \xrightarrow{\text{HNO}_2} \text{B} \xrightarrow{\text{Cu/573K}} \text{C}$$
 (a) CH_3COOH (b) $\text{CH}_3\text{CH}_2\text{NHOH}$
 (c) CH_3CONH_2 (d) CH_3CHO
 - Benzaldehyde can be prepared by oxidation of toluene by
 (a) Acidic KMnO_4 (b) $\text{K}_2\text{Cr}_2\text{O}_7/\text{H}^+$
 (c) CrO_2Cl_2 (d) All of these
 - Formalin is an aqueous solution of
 (a) fluorescein (b) formic acid
 (c) formaldehyde (d) furfuraldehyde
 - The reagent (s) which can be used to distinguish acetophenone from benzophenone is (are)
 (a) 2,4- dinitrophenylhydrazine
 (b) aqueous solution of NaHSO_3
 (c) benedict reagent
 (d) I_2 and Na_2CO_3 .
 - The test used for the distinction of HCOOH and CH_3COOH is treatment with :
 (a) sodium (b) sodium hydroxide
 (c) Tollen's reagent (d) alcohol
 - Sodium formate on heating yields
 (a) oxalic acid and H_2 (b) sodium oxalate and H_2
 (c) CO_2 and NaOH (d) sodium oxalate.
 - $(\text{CH}_3)_2\text{C}=\text{CHCOCH}_3$ can be oxidized to $(\text{CH}_3)_2\text{C}=\text{CHCOOH}$ by
 (a) Chromic acid (b) NaOI
 (c) Cu at 300°C (d) KMnO_4 .
 - Pinacolone is
 (a) 2, 3-dimethyl-2 3-butanediol
 (b) 3, 3-dimethyl-2 butanone
 (c) 1-phenyl-2propanone
 (d) 1,1-diphenyl-2-ethandiol.
 - Urotropine is obtained when ammonia reacts with
 (a) HCHO (b) CH_3COCH_3
 (c) CH_3CHO (d) $\text{C}_6\text{H}_5\text{OH}$
 - Benzaldehyde reacts with ethanoic KCN to give
 (a) $\text{C}_6\text{H}_5\text{CHOHCN}$
 (b) $\text{C}_6\text{H}_5\text{CHOHCOC}_6\text{H}_5$
 (c) $\text{C}_6\text{H}_5\text{CHOHCOOH}$
 (d) $\text{C}_6\text{H}_5\text{CHOHCHOHC}_6\text{H}_5$
 - Aldehydes and ketones will not form crystalline derivatives with
 (a) sodium bisulphite
 (b) phenylhydrazine
 (c) semicarbazide hydrochloride
 (d) dihydrogen sodium phosphate.


21. When formaldehyde is polymerised in the presence of dilute H_2SO_4 , the compound obtained is
 (a) paraldehyde (b) paraformaldehyde
 (c) metaldehyde (d) trioxo formaldehyde
22. $(\text{CH}_3)_3\text{C}-\text{CHO}$ does not undergo Aldol condensation due to
 (a) three electron donating methyl groups
 (b) cleavage taking place between $-\text{C}-\text{CHO}$ bond
 (c) absence of alpha hydrogen atom in the molecule
 (d) bulky $(\text{CH}_3)_3\text{C}-$ group
23. In the reaction





$$\text{CH}_3\text{CN} + 2\text{H} \xrightarrow[\text{Ether}]{\text{HCl}} \text{X} \xrightarrow[\text{H}_2\text{O}]{\text{Boiling}} \text{Y}; \text{ the term Y is :}$$


 (a) acetone (b) ethyl amine
 (c) acetaldehyde (d) dimethyl amine
24. Phenylmethyl ketone can be converted into ethylbenzene in one step by which of the following reagents?
 (a) LiAlH_4 (b) Zn-Hg/HCl
 (c) NaBH_4 (d) CH_3MgI
25. Which one of the following is reduced with zinc and hydrochloric acid to give the corresponding hydrocarbon?
 (a) Acetamide (b) Acetic acid
 (c) Ethyl acetate (d) Butan-2-one
26. Acetone reacts with bleaching powder to give
 (a) CCl_4
 (b) CHCl_3
 (c) $\text{CCl}_3-\text{COCH}_3$
 (d) none of these.
27. Appropriate reducing agent for the following conversion is—




- (a) $\text{LiAlH}_4/\text{H}_2\text{O}$ (b) $\text{NaBH}_4/\text{H}_2\text{O}$
 (c) $\text{Na} + \text{C}_2\text{H}_5\text{OH}$ (d) $\text{B}_2\text{H}_6/\text{H}^+$
28. A compound called chloral whose formula is $\text{C}_2\text{Cl}_3\text{OH}$ is a/an
 (a) aldehyde (b) alcohol
 (c) ketone (d) alkanoyl chloride



- (a)  (b) 
 (c)  (d) 





- (a) $\text{C}_6\text{H}_5\overset{\text{OH}}{\underset{|}{\text{CH}}} - \text{CH}_2\text{CHO}$ (b) $\text{C}_6\text{H}_5\text{CH} = \text{CH} - \text{CHO}$
 (c) $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CHO}$ (d) Both (b) & (c)



because



- (a) carbon is bounded by 3 heavy methyl group
 (b) due to absence of α -hydrogen
 (c) it is a neutral molecule
 (d) due to hindrance created by methyl for carbanion.
32. Which of the following is best method for reducing 3-bromopropanal to 1-bromopropane —
 (a) Wolf-Kishner reduction
 (b) Clemmenson reduction
 (c) Either (a) or (b)
 (d) Stephen's reduction
33. Cannizzaro reaction occurs with
 (a) $\text{CH}_3 - \text{CH}_2\text{OH}$ (b) $\text{C}_6\text{H}_5\text{CHO}$
 (c) CH_3CHO (d) $\text{CH}_3 - \text{CO} - \text{CH}_3$
34. In an alkaline medium, acetaldehyde undergoes:
 (a) benzoin condensation (b) aldol condensation
 (c) polymerisation (d) Cannizzaro reaction
35. Which of the following compound will show positive silver mirror test?
 (a) HCOOH (b) $\text{CH}_3(\text{CHOH})_3\text{CHO}$
 (c) $\text{CH}_3\text{CO}(\text{CHOH})\text{CH}_3$ (d) Both (a) and (b)
36. Oxalic acid on heating with conc. H_2SO_4 produces
 (a) $\text{CO} + \text{CO}_2$ (b) $\text{CO}_2 + \text{H}_2$
 (c) $\text{C} + \text{H}_2\text{O}$ (d) $\text{CO} + \text{H}_2$
37. Aldehydes and ketones are generally reduced by :
 (a) Clemmensen reduction (b) H_2S
 (c) H_2/Ni (d) None of these
38. The product obtained by the reaction of an aldehyde and hydroxylamine is
 (a) hydrazone (b) aldoxime
 (c) primary amine (d) alcohol
39. In which reaction, $>\text{C}=\text{O}$ can be reduced to $>\text{CH}_2$?
 (a) Wolf-Kishner reaction
 (b) Reimer-Tiemann reaction
 (c) Wurtz reaction
 (d) None of these

40. In benzilic acid rearrangement,
 (a) benzoin is converted into benzilic acid
 (b) benzaldehyde is converted into benzoin
 (c) benzil is converted into benzilic acid
 (d) benzilic acid is converted into benzil
41. Which of the following does not show Cannizzaro reaction?
 (a) CH_3CHO (b) HCHO
 (c) $\text{C}_6\text{H}_5\text{CHO}$ (d) $(\text{CH}_3)_3\text{C}\cdot\text{CHO}$
42. A compound does not react with 2, 4-dinitrophenylhydrazine, the compound is :
 (a) Acetone (b) Acetaldehyde
 (c) CH_3OH (d) $\text{CH}_3\text{CH}_2\text{COCH}_3$
43. Which gives lactic acid on hydrolysis after reacting with HCN ?
 (a) HCHO (b) CH_3CHO
 (c) $\text{C}_6\text{H}_5\text{CHO}$ (d) CH_3COCH_3
44. Ketones react with Mg-Hg over water to give :
 (a) pinacolone (b) pinacols
 (c) alcohols (d) none of these
45. Which of the following functional groups cannot be reduced to alcohol using NaBH_4 in ethanolic solution ?
 (a) $\text{R}-\text{O}-\text{R}$ (b) RCOCl
 (c) $\text{R}-\text{COOH}$ (d) $\text{R}-\text{CHO}$
46. What is the final product of the following reaction ?




47. 2-pentanone and 3-pentanone can be distinguished by :
 (a) Cannizzaro's reaction
 (b) Aldol condensation
 (c) Iodoform reaction
 (d) Clemmensen's reduction
48. When dihydroxyacetone reacts with HIO_4 , the product is/ are :
 (a) HCHO
 (b) HCOOH
 (c) HCHO and HCOOH
 (d) HCHO and CO_2

49. Which of the following is disproportionation reaction?





- (d) Both (a) & (b)


50. Formaldehyde reacts with ammonia to give urotropine is
 (a) $(\text{CH}_2)_6\text{N}_4$ (b) $(\text{CH}_2)_4\text{N}_3$
 (c) $(\text{CH}_2)_6\text{N}_6$ (d) $(\text{CH}_2)_3\text{N}_3$
51. The property which distinguishes formic acid from acetic acid is
 (a) only ammonium salt of formic acid on heating gives amide
 (b) when heated with alcohol/ H_2SO_4 only acetic acid forms ester
 (c) only acetic acid forms salts with alkali
 (d) only formic acid reduces Fehling's solution
52. Which of the following is an example of aldol condensation?






53. The reagent which does not give acid chloride on treating with a carboxylic acid is
 (a) PCl_5 (b) Cl_2
 (c) SOCl_2 (d) PCl_3
54. The reactant (X) in the reaction



55. Main product obtained from the reaction of ammonia and formaldehyde is
 (a) formic acid (b) methylamine
 (c) methanol (d) urotropine
56. Which of the products is formed when acetone is reacted with barium hydroxide solution?





57. Hydride ion transfer takes place in
 (a) Frankland method (b) Wurtz reaction
 (c) Cannizzaro's reaction (d) Wolf-Kishner reduction
58. Ketone upon treatment with Grignard Reagent gives
 (a) primary alcohol (b) secondary alcohol
 (c) tertiary alcohol (d) aldehyde
59. Which of the following compounds would be the main product of an aldol condensation of acetaldehyde and acetone?
 (a) $\text{CH}_3\text{CH}=\text{CH}\cdot\text{CHO}$ (b) $\text{CH}_3\text{CH}=\text{CHCOCH}_3$
 (c) $(\text{CH}_3)_2\text{C}=\text{CH}\cdot\text{CHO}$ (d) $(\text{CH}_3)_2\text{C}=\text{CHCOCH}_3$
60. What is formed, when acetonitrile is hydrolysed partially with cold concentrated HCl?
 (a) Acetamide (b) Acetic acid
 (c) Methyl cyanide (d) Acetic anhydride
61. Among acetic acid, phenol and *n*-hexanol, which of the following compounds will react with NaHCO_3 solution to give sodium salt and carbon dioxide?
 (a) acetic acid (b) *n*-Hexanol
 (c) acetic acid and phenol (d) phenol.
62. Formic acid is obtained when
 (a) calcium acetate is heated with conc. H_2SO_4
 (b) calcium formate is heated with calcium acetate
 (c) glycerol is heated with oxalic acid at 373 K
 (d) acetaldehyde is oxidised with $\text{K}_2\text{Cr}_2\text{O}_7$ and H_2SO_4 .
63. Schotten-Baumann reaction is a reaction of phenols with
 (a) benzoyl chloride and sodium hydroxide
 (b) acetyl chloride and sodium hydroxide
 (c) salicylic acid and conc. H_2SO_4
 (d) acetyl chloride and conc H_2SO_4
64. Which of the following products is formed when benzaldehyde is treated with CH_3MgBr and the addition product so obtained is subjected to acid hydrolysis?
 (a) A secondary alcohol
 (b) A primary alcohol
 (c) Phenol
 (d) *tert*-Butyl alcohol
65. An ester is boiled with KOH. The product is cooled and acidified with concentrated HCl. A white crystalline acid separates. The ester is
 (a) methyl acetate (b) ethyl acetate
 (c) ethyl formate (d) ethyl benzoate

66. In the Friedel Craft's acylation reaction, the effective electrophile is
 (a) RCOCl^\oplus (b) AlCl_3
 (c) RCOCl (d) RCO^\oplus
67. The compound which is not formed during the dry distillation of a mixture of calcium formate and calcium acetate is
 (a) methanal (b) propanal
 (c) propanone (d) ethanal
68. Which one of the following esters cannot undergo Claisen self-condensation?
 (a) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{COOC}_2\text{H}_5$
 (b) $\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5$
 (c) $\text{C}_6\text{H}_5\text{CH}_2\text{COOC}_2\text{H}_5$
 (d) $\text{C}_6\text{H}_{11}\text{CH}_2\text{COOC}_2\text{H}_5$
69. The cyanohydrin of a compound on hydrolysis gives an optically active α -hydroxy acid. The compound is
 (a) diethyl ketone (b) formaldehyde
 (c) acetaldehyde (d) acetone
70. The product of following reaction is
 $2\text{CH}_3\text{MgBr} + \text{CH}_2\text{O} \longrightarrow$
 (a) CH_3OH (b) $\text{C}_2\text{H}_5\text{OH}$
 (c) CH_4 (d) C_2H_6
71. $\text{CH}_3\text{CH}_2\text{COOH} \xrightarrow[\text{red P}]{\text{Cl}_2} A \xrightarrow{\text{alc. KOH}} B$. What is B?
 (a) $\text{CH}_3\text{CH}_2\text{COCl}$ (b) $\text{CH}_3\text{CH}_2\text{CHO}$
 (c) $\text{CH}_2=\text{CHCOOH}$ (d) $\text{ClCH}_2\text{CH}_2\text{COOH}$
72. When dihydroxy acetone reacts with HIO_4 , the product is/are
 (a) HCHO
 (b) HCOOH
 (c) HCHO and HCOOH
 (d) HCHO and CO_2
73. When $\text{CH}_2=\text{CH}-\text{COOH}$ is reduced with LiAlH_4 , the compound obtained will be
 (a) $\text{CH}_2=\text{CH}-\text{CH}_2\text{OH}$ (b) $\text{CH}_3-\text{CH}_2-\text{CH}_2\text{OH}$
 (c) $\text{CH}_3-\text{CH}_2-\text{CHO}$ (d) $\text{CH}_3-\text{CH}_2-\text{COOH}$
74. On mixing ethyl acetate with aqueous sodium chloride, the composition of the resultant solution is
 (a) $\text{CH}_3\text{COCl} + \text{C}_2\text{H}_5\text{OH} + \text{NaOH}$
 (b) $\text{CH}_3\text{COONa} + \text{C}_2\text{H}_5\text{OH}$
 (c) $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{NaCl}$
 (d) $\text{CH}_3\text{Cl} + \text{C}_2\text{H}_5\text{COONa}$
75. Acetyl bromide reacts with excess of CH_3MgI followed by treatment with a saturated solution of NH_4Cl gives
 (a) 2-methyl-2-propanol (b) acetamide
 (c) acetone (d) acetyl iodide
76. Acetic anhydride reacts with diethyl ether in the presence of anhydrous AlCl_3 to give :
 (a) $\text{CH}_3\text{CH}_2\text{COOH}$ (b) $\text{CH}_3\text{CH}_2\text{COOC}_2\text{H}_5$
 (c) $\text{CH}_3\text{COOCH}_3$ (d) $\text{CH}_3\text{COOC}_2\text{H}_5$
77. An important reaction of acetone is auto condensation in presence of concentrated sulphuric acid to give the aromatic compound.....
 (a) Mesitylene (b) Mesityl oxide
 (c) Trioxan (d) Phorone

78. The acid(s) which do not contain $-\text{COOH}$ groups is/are :

- (a) palmitic acid
- (b) lactic acid
- (c) ethanoic acid
- (d) picric acid and *p*-toluene sulphonic acid

79. The best combination of reagents for carrying out the conversion $\text{RCH}_2\text{CH}_2\text{OH} \rightarrow \text{RCH}_2\text{CH}_2\text{COOH}$ is

- (a) $\text{PBr}_3, \text{KCN}, \text{H}_3\text{O}^+$
- (b) $\text{PBr}_3, \text{KCN}, \text{H}_2/\text{Pt}$
- (c) $\text{KCN}, \text{H}_3\text{O}^+$
- (d) $\text{PBr}_3, \text{H}_3\text{O}^+$

80. $\text{CH}_3\text{COOH} \xrightarrow{\text{A}} \text{CH}_3\text{COCl}$. What is A ?

- (a) PCl_5
- (b) Cl_2
- (c) HCl
- (d) COCl_2

81. Ethyl acetate on reaction with Grignard's reagent, forms a/an

- (a) alcohol
- (b) ketone
- (c) ether
- (d) hydrocarbon

82. The compound not soluble in acetic acid is :

- (a) CaCO_3
- (b) CaO
- (c) CaC_2O_4
- (d) $\text{Ca}(\text{OH})_2$


83. Among the following, the most acidic is :

- (a) CH_3COOH
- (b) ClCH_2COOH
- (c) Cl_2CHCOOH
- (d) $\text{Cl}_2\text{CHCH}_2\text{COOH}$

84. The compound that neither forms semicarbazone for oxime is

- (a) HCHO
- (b) $\text{CH}_3\text{COCH}_2\text{Cl}$
- (c) CH_3CHO
- (d) $\text{CH}_3\text{CONHCH}_3$


85. Consider the following transformations :



The molecular formula of C is

- (a) $\text{CH}_3-\overset{\text{OH}}{\underset{\text{I}}{\text{C}}}-\text{CH}_3$
- (b) $\text{ICH}_2-\text{COCH}_3$
- (c) CHI_3
- (d) CH_3I

86. In the following reaction



The major compounds X and Y are

- (a) RCHBrCONH_2 ; $\text{RCH}(\text{NH}_2)\text{COOH}$
- (b) RCHBrCOOH ; $\text{RCH}(\text{NH}_2)\text{COOH}$
- (c) RCH_2COBr ; $\text{RCH}_2\text{COONH}_4$
- (d) RCHBrCOOH ; $\text{RCH}_2\text{CONH}_2$

87. An organic compound $\text{C}_3\text{H}_6\text{O}$ neither gives precipitate with semicarbazide nor reacts with sodium. It could be

- (a) $\text{CH}_3\text{CH}_2\text{CHO}$
- (b) CH_3COCH_3
- (c) $\text{CH}_2=\text{CH}-\text{CH}_2-\text{OH}$
- (d) $\text{CH}_2=\text{CH}-\text{O}-\text{CH}_3$


88. Which one of the following compounds will not react with CH_3MgBr ?

- (a) Ethyl acetate
- (b) Acetone
- (c) Dimethyl ether
- (d) Ethanol

89. Some carboxylic acids and their IUPAC names are given below. Which of the following is not correctly matched?

- (a) Formic acid — Methanoic acid
- (b) Acetic acid — Ethanoic acid
- (c) *n*-Butyric acid — Butanoic acid
- (d) *Iso*-butyric acid — 2-methylbutanoic acid

90. Silver benzoate will react with bromine in CCl_4 to form :



91. Which of the following pairs can be distinguished by sodium hypoiodite?

- (a) CH_3CHO and CH_3COCH_3
- (b) $\text{CH}_3\text{CH}_2\text{CHO}$ and CH_3COCH_3
- (c) $\text{CH}_3\text{CH}_2\text{OH}$ and $\text{CH}_3\text{CH}_2\text{CHOHCH}_3$
- (d) CH_3OH and $\text{CH}_3\text{CH}_2\text{CHO}$

92. Reduction of aldehydes and ketones into hydrocarbons using zinc amalgam and conc. HCl is called

- (a) Cope reduction
- (b) Dow reduction
- (c) Wolf-Kishner reduction
- (d) Clemmensen reduction.

93. Which of the following represents the correct order of the acidity in the given compounds?

- (a) $\text{FCH}_2\text{COOH} > \text{CH}_3\text{COOH} > \text{BrCH}_2\text{COOH} > \text{ClCH}_2\text{COOH}$
- (b) $\text{BrCH}_2\text{COOH} > \text{ClCH}_2\text{COOH} > \text{FCH}_2\text{COOH} > \text{CH}_3\text{COOH}$
- (c) $\text{FCH}_2\text{COOH} > \text{ClCH}_2\text{COOH} > \text{BrCH}_2\text{COOH} > \text{CH}_3\text{COOH}$
- (d) $\text{CH}_3\text{COOH} > \text{BrCH}_2\text{COOH} > \text{ClCH}_2\text{COOH} > \text{FCH}_2\text{COOH}$

94. Which one of the following on treatment with 50% aqueous sodium hydroxide yields the corresponding alcohol and acid?


- (a) $\text{C}_6\text{H}_5\text{CHO}$
- (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$

- (c) $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$
- (d) $\text{C}_6\text{H}_5\text{CH}_2\text{CHO}$

95. The product formed in Aldol condensation is

- (a) a beta-hydroxy aldehyde or a beta-hydroxy ketone
- (b) an alpha-hydroxy aldehyde or ketone
- (c) an alpha, beta unsaturated ester
- (d) a beta-hydroxy acid

96. Propionic acid with Br_2/P yields a dibromo product. Its structure would be:



97. Which of the following reactions will not result in the formation of carbon-carbon bonds?

- (a) Reimer-Tiemann reaction (b) Cannizzaro reaction
(c) Wurtz reaction (d) Friedel-Crafts acylation

98. Following compounds are given:

- (i) $\text{CH}_3\text{CH}_2\text{OH}$ (ii) CH_3COCH_3
(iii) $\begin{array}{c} \text{CH}_3-\text{CHOH} \\ | \\ \text{CH}_3 \end{array}$ (iv) CH_3OH

Which of the above compound(s), on being warmed with iodine solution and NaOH, will give iodoform?

- (a) (i) and (ii) (b) (i), (iii) and (iv)
(c) Only (ii) (d) (i), (ii) and (iii)

99. Clemmensen reduction of a ketone is carried out in the presence of which of the following?

- (a) Glycol with KOH (b) Zn-Hg with HCl
(c) LiAlH_4 (d) H_2 and Pt as catalyst

100. Match the compounds given in List-I with List-II and select the suitable option using the code given below:

List I

- (A) Benzaldehyde
(B) Phthalic anhydride
(C) Phenyl benzoate
(D) Methyl salicylate

List-II

- (i) Phenolphthalein
(ii) Benzoin condensation
(iii) Oil of wintergreen
(iv) Fries rearrangement

Code :

- | | | | |
|----------|-------|-------|-------|
| (A) | (B) | (C) | (D) |
| (a) (iv) | (i) | (iii) | (ii) |
| (b) (iv) | (ii) | (iii) | (i) |
| (c) (ii) | (iii) | (iv) | (i) |
| (d) (ii) | (i) | (iv) | (iii) |

EXERCISE - 2

Applied Questions

1. The catalyst used in the Wacker process of oxidation of ethyne to ethanal is


- (a) PdCl_2 (b) V_2O_5
(c) Nickel complexes (d) TiCl_4 and $\text{Al}(\text{CH}_3)_3$

2. Which one of the following reagents enables the purification of benzaldehydes?

- (a) HCN (b) CH_3MgBr
(c) NH_2OH (d) NaHSO_3

3. Which of the following is the correct decreasing order of acidic strength of

- (i) Methanoic acid (ii) Ethanoic acid
(iii) Propanoic acid (iv) Butanoic acid
(a) (i) > (ii) > (iii) > (iv) (b) (ii) > (iii) > (iv) > (i)
(c) (i) > (iv) > (iii) > (ii) (d) (iv) > (i) > (iii) > (ii)




The above shown polymer is obtained when a carbonyl compound is allowed to stand. It is a white solid. The polymer is

- (a) Trioxane (b) Formose
(c) Paraformaldehyde (d) Metaldehyde

5. Which of the following gives reddish brown precipitate on heating with a solution of CuSO_4 and alkaline sodium potassium tartarate?

- (a) Ethanol (b) Ethanal
(c) Propanone (d) Ethanoic acid



The above compound describes a condensation polymer which can be obtained in two ways: either treating 3 molecules of acetone (CH_3COCH_3) with conc. H_2SO_4 or passing propyne ($\text{CH}_3\text{C}\equiv\text{CH}$) through a red hot tube. The polymer is

- (a) Phorone (b) Mesityl oxide
(c) Deacetonyl alcohol (d) Mesitylene

7. In which of the following, the number of carbon atoms does not remain same when carboxylic acid is obtained by oxidation

- (a) CH_3COCH_3 (b) $\text{CCl}_3\text{CH}_2\text{CHO}$
(c) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ (d) $\text{CH}_3\text{CH}_2\text{CHO}$

8. Aldehydes that do not undergo aldol condensation are

1. propanal 2. trichloroethanal
3. methanal 4. ethanal
5. benzaldehyde

- (a) 3 and 4 only (b) 3 and 5 only
(c) 1, 2 and 3 only (d) 2, 3 and 5 only

9. Benzoic acid gives benzene on being heated with X and phenol gives benzene on being heated with Y. Therefore X and Y are respectively

- (a) Soda-lime and copper (b) Zn dust and NaOH
(c) Zn dust and soda-lime (d) Soda-lime and zinc dust

10. An ester (A) with molecular formula, $\text{C}_9\text{H}_{10}\text{O}_2$ was treated with excess of CH_3MgBr and the complex so formed was treated with H_2SO_4 to give an olefin (B). Ozonolysis of (B) gave a ketone with molecular formula $\text{C}_8\text{H}_8\text{O}$ which shows positive iodoform test. The structure of (A) is

- (a) $\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5$
(b) $\text{C}_2\text{H}_5\text{COOC}_6\text{H}_5$
(c) $\text{H}_3\text{COCH}_2\text{COC}_6\text{H}_5$
(d) $\text{p}-\text{H}_3\text{CO}-\text{C}_6\text{H}_4-\text{COCH}_3$

11. Polarization of electrons in acrolein may be written as:

- (a) $\overset{\delta-}{\text{C}}\text{H}_2 = \text{CH} - \overset{\delta+}{\text{CH}} = \text{O}$
 (b) $\overset{\delta+}{\text{C}}\text{H}_2 = \overset{\delta-}{\text{C}}\text{H} - \text{CH} = \text{O}$
 (c) $\overset{\delta+}{\text{C}}\text{H}_2 = \text{CH} - \overset{\delta-}{\text{CH}} = \text{O}$
 (d) $\overset{\delta-}{\text{C}}\text{H}_2 = \text{CH} - \overset{\delta+}{\text{C}}\text{H} = \text{O}$

12. Which compound among the following gives positive iodoform test?

1. Ethanol 2. Ethanal
 3. 1-Butanol 4. 2-Butanol
 5. Phenyl ethanol
 (a) 1, 2 and 5 (b) 1, 3 and 4
 (c) 1, 2 and 3 (d) 1, 2 and 4

13. Reaction of cyclohexanone with dimethylamine in the presence of catalytic amount of an acid forms a compound if water during the reaction is continuously removed. The compound formed is generally known as

- (a) an amine (b) an imine
 (c) an enamine (d) a schiff's base

14. The increasing order of the rate of HCN addition to compound A – D is

- (A) HCHO (B) CH_3COCH_3
 (C) PhCOCH_3 (D) PhCOPh
 (a) $\text{D} < \text{C} < \text{B} < \text{A}$ (b) $\text{C} < \text{D} < \text{B} < \text{A}$
 (c) $\text{A} < \text{B} < \text{C} < \text{D}$ (d) $\text{D} < \text{B} < \text{C} < \text{A}$

15. Reaction of butanone with methylmagnesium bromide followed by hydrolysis gives

- (a) 2-methyl-2-butanol
 (b) 2-butanol
 (c) 3-methyl-2-butanol
 (d) 2, 2-dimethyl-1-butanol


16. The correct order of increasing acid strength of the compounds

- (A) $\text{CH}_3\text{CO}_2\text{H}$ (B) $\text{MeOCH}_2\text{CO}_2\text{H}$
 (C) $\text{CF}_3\text{CO}_2\text{H}$ (D) $\text{Me}_2\text{CHCO}_2\text{H}$

is

- (a) $\text{D} < \text{A} < \text{B} < \text{C}$ (b) $\text{A} < \text{D} < \text{B} < \text{C}$
 (c) $\text{B} < \text{D} < \text{A} < \text{C}$ (d) $\text{D} < \text{A} < \text{C} < \text{B}$


17. How many cross aldol products (without counting stereoisomers) are produced in the following reaction?



- (a) 5 (b) 2
 (c) 3 (d) 1


18. Monocarboxylic acids are functional isomers of:

- (a) Ethers (b) Amines
 (c) Esters (d) Alcohols







In the above sequence of reaction X and Y are respectively

- (a) H_2/Pt ; Br_2 (b) KMnO_4 ; H_2/Pt
 (c) KMnO_4 (aq); HI/P (d) $\text{NH}_2 - \text{NH}_2/\text{KOH}$, HI/P







The product formed in the reaction is –


- (a)  (b) 
 (c)  (d) 

21. An aromatic compound 'X' with molecular formula $\text{C}_9\text{H}_{10}\text{O}$ gives the following chemical tests. It





- (i) forms 2, 4-DNP derivative
 (ii) reduces Tollen's reagent
 (iii) undergoes Cannizzaro reaction and
 (iv) on vigorous oxidation 1, 2-benzenedicarboxylic acid is obtained.

X is


- (a)  (b) 
 (c)  (d) 












Product (X) of the reaction is–

- (a)  (b) 
 (c)  (d) 




23. In a set of reactions acetic acid yielded a product D.








The structure of (D) would be –







25. $\text{CH}_3\text{CO}_2\text{C}_2\text{H}_5$ on reaction with sodium ethoxide in ethanol gives A, which on heating in the presence of acid gives B. Compound B is :







What is Z ?




The final product is –





28. *o*-Toluic acid on reaction with $\text{Br}_2 + \text{Fe}$ gives




29. Which of the following compound will not give benzoic acid on oxidation with $\text{KMnO}_4/\text{OH}^-/\Delta$ –




30. The final product (III) obtained in the reaction sequence –




31. Calculate number of molecules of Grignard reagent consumed by 1 molecule of following compound.



- (a) 5 (b) 2
(c) 3 (d) 1
32. Cannizzaro's reaction is not given by:




33. Which of the following reagent(s) used for the conversion?




- (a) glycol/ $\text{LiAlH}_4/\text{H}_3\text{O}^+$ (b) glycol/ $\text{NaH}/\text{H}_3\text{O}^+$
(c) LiAlH_4 (d) NaBH_4


34. In a set of reactions, ethylbenzene yield a product D.




Identify D:



35. Which one of the following compounds will be most readily dehydrated?



36. The order of reactivity of phenyl magnesium bromide (PhMgBr) with the following compounds




- (a) $\text{III} > \text{II} > \text{I}$ (b) $\text{II} > \text{I} > \text{III}$
(c) $\text{I} > \text{III} > \text{II}$ (d) $\text{I} > \text{II} > \text{III}$

37. An organic compound 'A' on treatment with NH_3 gives 'B' which on heating gives 'C', 'C' when treated with Br_2 in the presence of KOH produces ethylamine. Compound 'A' is:

- (a) CH_3COOH (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$
 (c) $\text{CH}_3 - \underset{\text{CH}_3}{\text{CHCOOH}}$ (d) $\text{CH}_3\text{CH}_2\text{COOH}$


38. In Cannizzaro reaction given below



the slowest step is :


- (a) the transfer of hydride to the carbonyl group
 (b) the abstraction of proton from the carboxylic group
 (c) the deprotonation of PhCH_2OH
 (d) the attack of $:\text{OH}^-$ at the carboxyl group

39. In the following reaction sequence, the correct structures of E, F and G are






- (a) $\text{E} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{CH}_3$ $\text{F} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{O}^- \text{Na}^+$ $\text{G} = \text{CHI}_3$
 (b) $\text{E} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{CH}_3$ $\text{F} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{O}^- \text{Na}^+$ $\text{G} = \text{CHI}_3$
 (c) $\text{E} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{CH}_3$ $\text{F} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}^- \text{Na}^+$ $\text{G} = \overset{*}{\text{C}}\text{HI}_3$
 (d) $\text{E} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{CH}_3$ $\text{F} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}^- \text{Na}^+$ $\text{G} = \overset{*}{\text{C}}\text{H}_3\text{I}$

40. In the reaction




the structure of the product T is :





- (a) $\text{H}_3\text{C}-\text{C}_6\text{H}_4-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{C}_6\text{H}_5$




41. The compounds P, Q and S



were separately subjected to nitration using $\text{HNO}_3/\text{H}_2\text{SO}_4$ mixture. The major product formed in each case respectively, is :




42. The number of aldol reaction(s) that occurs in the given transformation is :




- (a) 1 (b) 2
 (c) 3 (d) 4

43. The major product **H** of the given reaction sequence is



- (a) $\text{CH}_3 - \text{CH} = \underset{\text{CH}_3}{\text{C}} - \text{COOH}$
- (b) $\text{CH}_3 - \text{CH} = \underset{\text{CH}_3}{\text{C}} - \text{CN}$
- (c) $\text{CH}_3 - \text{CH}_2 - \underset{\text{CH}_3}{\overset{\text{OH}}{\text{C}}} - \text{COOH}$
- (d) $\text{CH}_3 - \text{CH} = \underset{\text{CH}_3}{\text{C}} - \text{CO} - \text{NH}_2$


44. The compound that undergoes decarboxylation most readily under mild condition is





45. An organic compound A upon reacting with NH_3 gives B. On heating B gives C. C in presence of KOH reacts with Br_2 to give $\text{CH}_3\text{CH}_2\text{NH}_2$. A is :

- (a) CH_3COOH (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$
- (c) $\text{CH}_3 - \underset{\text{CH}_3}{\text{CH}} - \text{COOH}$ (d) $\text{CH}_3\text{CH}_2\text{COOH}$

46. Predict the product in the given reaction.



- (a)
- (b)



47. Acetone is treated with excess of ethanol in the presence of hydrochloric acid. The product obtained is :

- (a) $\text{CH}_3\text{CH}_2\text{CH}_2 - \overset{\text{O}}{\parallel} \text{C} - \text{CH}_3$
- (b) $\text{CH}_3\text{CH}_2\text{CH}_2 - \overset{\text{O}}{\parallel} \text{C} - \text{CH}_2\text{CH}_2\text{CH}_3$
- (c) $(\text{CH}_3)_2\text{C} \begin{matrix} \nearrow \text{OH} \\ \searrow \text{OC}_2\text{H}_5 \end{matrix}$
- (d) $(\text{CH}_3)_2\text{C} \begin{matrix} \nearrow \text{OC}_2\text{H}_5 \\ \searrow \text{OC}_2\text{H}_5 \end{matrix}$

DIRECTIONS for Qs. 48 to 50 : These are Assertion-Reason type questions. Each of these question contains two statements : **Statement-1 (Assertion)** and **Statement-2 (Reason)**. Answer these questions from the following four options.

- (a) Statement-1 is true, Statement-2 is true, Statement-2 is a correct explanation for Statement -1
- (b) Statement-1 is True, Statement-2 is True ; Statement-2 is NOT a correct explanation for Statement-1
- (c) Statement-1 is True, Statement-2 is False
- (d) Statement-1 is False, Statement-2 is True

48. **Statement-1 :** Acetic acid does not undergo haloform reaction.

Statement-2 : Acetic acid has no alpha hydrogens.

49. **Statement-1 :** Benzaldehyde is more reactive than ethanol towards nucleophilic attack.

Statement-2 : The overall effect of $-I$ and $+R$ effect of phenyl group decreases the electron density on the carbon atom of $>\text{C}=\text{O}$ group in benzaldehyde.

50. **Statement-1 :** Lower aldehyde and ketones are soluble in water but the solubility decreases as molecular mass increases.





Statement-2 : Aldehydes and ketones can be distinguished by Tollen's reagent.

EXERCISE - 3





Exemplar & Past Years NEET/AIPMT Questions

Exemplar Questions

1. Addition of water to alkynes occurs in acidic medium and in the presence of Hg^{2+} ions as a catalyst. Which of the following products will be formed on addition of water to but - 1-yne under these conditions?



2. Which of the following compounds is most reactive towards nucleophilic addition reactions?



3. The correct order of increasing acidic strength is

- (a) phenol < ethanol < chloroacetic acid < acetic acid
 (b) ethanol < phenol < chloroacetic acid < acetic acid
 (c) ethanol < phenol < acetic acid < chloroacetic acid
 (d) chloroacetic acid < acetic acid < phenol < ethanol


4. Compound $\text{Ph} - \text{O} - \overset{\text{O}}{\parallel} \text{C} - \text{Ph}$ can be prepared by the reactions of

- (a) phenol and benzoic acid in the presence of NaOH
 (b) phenol and benzoyl chloride in the presence of pyridine
 (c) phenol and benzoyl chloride in the presence of ZnCl_2
 (d) phenol and benzaldehyde in the presence of palladium

5. The reagent which does not react with both, acetone and benzaldehyde?






- (a) Sodium hydrogen sulphite
 (b) Phenyl hydrazine
 (c) Fehling's solution
 (d) Grignard reagent

6. Cannizzaro's reaction is not given by



7. Which product is formed when the compound


is treated with concentrated aqueous KOH solution?



structure of 'A' and type of isomerism in the above reaction are respectively


- (a) Prop-1-en-2-ol, metamerism
 (b) Prop-1-en-1-ol, tautomerism
 (c) Prop-2-en-2-ol, geometrical isomerism
 (d) Prop-1-en-2-ol, tautomerism

9. Compounds A and C in following reaction are



- (a) identical (b) positional isomers
 (c) functional isomers (d) optical isomers

10. Which is the most suitable reagent for the following conversion?



- (a) Tollen's reagent (b) Benzoyl peroxide
 (c) I_2 and NaOH solution (d) Sn and NaOH solution



11. Which of the following compounds will give butanone on oxidation with alkaline KMnO_4 solution?

- (a) Butan-1-ol (b) Butan-2-ol
 (c) Both (a) and (b) (d) None of these

12. In Clemmensen reduction, carbonyl compounds are treated with
- zinc amalgam + HCl
 - sodium amalgam + HCl
 - zinc amalgam + nitric acid
 - sodium amalgam + HNO_3

NEET/AIPMT (2013-2017) Questions





13. The order of stability of the following tautomeric compounds is:



[2013]





- III > II > I
 - II > I > III
 - II > III > I
 - I > II > III
14. Reaction by which Benzaldehyde cannot be prepared:

[2013]



15. Which one is most reactive towards Nucleophilic addition reaction?


[2014]



16. An organic compound 'X' having molecular formula $\text{C}_5\text{H}_{10}\text{O}$ yields phenyl hydrazone and gives negative response to the Iodoform test and Tollen's test. It produces n-pentane on reduction. 'X' could be :-

[2015]


- 2-pentanone
- 3-pentanone
- n-amyl alcohol
- pentanal

17. Treatment of cyclopentanone  with methyl lithium gives which of the following species?

[2015]

- Cyclopentanonyl cation
- Cyclopentanonyl radical
- Cyclopentanonyl biradical
- Cyclopentanonyl anion





18. The enolic form of ethyl acetoacetate as below has: [2015]



- 16 sigma bonds and 1 pi - bond
- 9 sigma bonds and 2 pi - bonds
- 9 sigma bonds and 1 pi - bond
- 18 sigma bonds and 2 pi - bonds

19. Which one of the following esters gets hydrolysed most easily under alkaline conditions?

[2015 RS]




20. Reaction of a carbonyl compound with one of the following reagents involves nucleophilic addition followed by elimination of water. The reagent is:

[2015 RS]

- a Grignard reagent
- hydrazine in presence of feebly acidic solution
- hydrocyanic acid
- sodium hydrogen sulphite


21. The correct statement regarding a carbonyl compound with a hydrogen atom on its α -carbon, is : [2016]
- a carbonyl compound with a hydrogen atom on its α -carbon never equilibrates with its corresponding enol.
 - a carbonyl compound with a hydrogen atom on its α -carbon rapidly equilibrates with its corresponding enol and this process is known as aldehyde-ketone equilibration.
 - a carbonyl compound with a hydrogen atom on its α -carbon rapidly equilibrates with its corresponding enol and this process is known as carbonylation.
 - a carbonyl compound with a hydrogen atom on its α -carbon rapidly equilibrates with its corresponding enol and this process is known as keto-enol tautomerism.
22. The product formed by the reaction of an aldehyde with a primary amine is [2016]
- Schiff base
 - Ketone
 - Carboxylic acid
 - Aromatic acid
23. Consider the reactions :- [2017]



Identify A, X, Y and Z

- (a) A-Methoxymethane, X-Ethanol, Y-Ethanoic acid, Z-Semicarbazide.


- A - Ethanal, X-Ethanol, Y - But - 2-enal, Z-Semicarbazone
 - A-Ethanol, X-Acetaldehyde, Y - Butanone, Z-Hydrazone
 - A-Methoxymethane, X-Ethanoic acid, Y-Acetate ion, Z-hydrazine.
24. Of the following, which is the product formed when cyclohexanone undergoes aldol condensation followed by heating ? [2017]




Hints & Solutions

EXERCISE - 1

1. (c) Benzaldehyde and alcoholic KCN react to form Benzoin. This is known as Benzoin condensation.




3. (d) (a) Cannizzaro reaction




No new C—C bond is formed

- (b) Reimer - Tiemann reaction :




New C—C bond is formed

- (c) Friedel-Craft reaction :





New C—C bond is formed

- (d) Clemmensen reduction




4. (b) Acetic Anhydride is produced due to dehydration of two molecules of acetic acid by P_2O_5





5. (b) 
- Calcium acetate Acetone

6. (c) $\text{C}_3\text{H}_8\text{O} \xrightarrow[\text{(x)}]{[\text{O}]} \text{C}_3\text{H}_6\text{O}$ (Ketone)




Ketones are oxidation products of 2° alcohols.


7. (c) Clemmensen's Reduction of ketones is carried out in Zn-Hg and HCl.




9. (b) The compound is $\text{C}_2\text{H}_5\text{OH}$.



10. (d) $\text{CH}_3\text{CN} \xrightarrow[\text{[A]}]{\text{Na/C}_2\text{H}_5\text{OH}} \text{CH}_3\text{CH}_2\text{NH}_2 \xrightarrow{\text{HNO}_2}$



11. (c)




Acidic KMnO_4 and $\text{K}_2\text{Cr}_2\text{O}_7$ oxidise toluene to benzoic acid but CrO_2Cl_2 oxidises it to Benzaldehyde.


12. (c) Formalin is an aqueous solution (40%) of formaldehyde.
13. (d) I_2 and Na_2CO_3 react with acetophenone ($\text{C}_6\text{H}_5\text{COCH}_3$) to give yellow ppt. of CHI_3 but benzophenone ($\text{C}_6\text{H}_5\text{COC}_6\text{H}_5$) does not and hence can be used to distinguish between them.
14. (c) The test used for the distinction of HCOOH and CH_3COOH is treatment with Tollen's reagent because formic acid can reduce Tollen's reagent but acetic acid can not.

15. (b) $2\text{HCOONa} \xrightarrow[360^\circ]{\Delta} \begin{array}{c} \text{COONa} \\ | \\ \text{COONa} \end{array} + \text{H}_2$


16. (b) Chromic acid and KMnO_4 will cleave the molecule at the site of double bond while Cu at 300°C cannot oxidise $\text{COCH}_3 \longrightarrow \text{COOH}$. The only reagent suitable for this conversion is NaOI or $\text{NaOH} + \text{I}_2$ (iodoform test):




17. (b) Pinacolone is 3,3-dimethyl-2-butanone.







18. (a) Formaldehyde reacts with ammonia to form hexamethylene tetramine which is also known as urotropine. It is used as urinary tract antiseptic



19. (b) When benzaldehyde is refluxed with aqueous alcoholic potassium cyanide, two molecules of benzaldehyde condense together to form benzoin








20. (d) Dihydrogen sodium phosphate (NaH_2PO_4) does not have a lone pair of electrons on the P atom. As such it can not act as a nucleophile and hence does not react with aldehydes and ketones.
21. (d) The compound formed is trioxo formaldehyde.
22. (c) Aldol condensation is given by the compounds which contain α hydrogen atom. As the given compound does not contain α hydrogen atom. Hence it does not undergo aldol condensation.
23. (c) Acetaldehyde is formed




This reaction is known as Clemmensen's reduction.

25. (d) It is Clemmensen's reduction




28. (a) Chloral is CCl_3CHO so, it is an aldehyde.
29. (c) In the presence of base, cyclohexanone show aldol condensation.




30. (b) Aldol formed in aromatic aldehydes itself loses water molecule without heating because double bond formed is more stable due to conjugation with benzene ring. $\text{C}_6\text{H}_5\text{CH}=\text{CHCHO}$ is commonly known as cinnamaldehyde.
31. (b) $(\text{CH}_3)_3\text{C} - \text{CHO}$ does not show aldol condensation because it does not contain α -hydrogen atom.
32. (b) In Wolf-Kishner reduction $\text{NH}_2\text{NH}_2/\text{OH}^-$ is used. Br^- can be replaced by OH^- .
33. (b) Cannizzaro reaction is given by aldehydes and ketones which do not have α -hydrogen atom. Benzaldehyde ($\text{C}_6\text{H}_5\text{CHO}$) does not have α -H atom and hence gives Cannizzaro reaction.
34. (b) Acetaldehyde has α -H atoms. In alkaline medium it will undergo aldol condensation to produce aldol (β -hydroxyaldehyde).


35. (d) Compounds having $-\text{CHO}$ group reduce Tollen's reagent to silver mirror. It is called silver mirror test.



Both (a) and (b) have $-\text{C}-\text{H}$ group so both of them give positive silver mirror test.






37. (a) Aldehydes and ketones are reduced to alkanes by Clemmensen reduction.




38. (b)

39. (a) Wolf-Kishner reduction



Benzilic acid.


41. (a) Aldehydes having α -hydrogen atom do not show Cannizzaro reaction.



(a)



(b)





(c)


(d)

Thus only CH_3CHO has α -hydrogen atom and hence it will not show Cannizzaro reaction.

42. (c) Only aldehydes and ketones react with 2,4-dinitrophenylhydrazine.



44. (b) Example:




Pinacol



(Pinacols have adjacent hydroxyl groups)

45. (a) Compounds having $-\text{C}(=\text{O})-$ moiety in their structure are reduced to alcohols using NaBH_4 in ethanolic solution.


Thus $\text{R}-\text{C}(=\text{O})-\text{Cl}$, $\text{R}-\text{C}(=\text{O})-\text{OH}$ and $\text{R}-\text{C}(=\text{O})-\text{H}$ are reduced while, $\text{R}-\text{O}-\text{R}$ (ethers) are inert and can't be reduced by NaBH_4 .




47. (c) Iodoform test is given by compounds which have CH_3CO group.



\therefore 2-pentanone has CH_3CO group, so it gives iodoform test, while 3-pentanone does not have CH_3CO group, so it does not give iodoform test.





49. (a) In this reaction, one molecule is oxidised and other is reduced simultaneously.



61. (a) among acetic acid, phenol and n-hexanol only CH_3COOH reacts with NaHCO_3 to evolve CO_2 gas.

$$\text{CH}_3\text{COOH} + \text{NaHCO}_3 \longrightarrow \text{CH}_3\text{COONa} + \text{CO}_2 + \text{H}_2\text{O}$$


62. (c) When glycerol is heated with oxalic acid following reaction occurs.




The function of NaOH is

- (i) To convert phenol to more stronger nucleophile PhO^-
 (ii) To neutralize the acid formed

64. (a) Aldehydes, other than formaldehyde, when treated with RMgX give 2° alcohols.
 65. (d) Methyl acetate and ethyl acetate on hydrolysis give CH_3COOH which is a liquid. Similarly ethyl formate on hydrolysis will give formic acid which is also a liquid. Only ethyl benzoate on hydrolysis will give benzoic acid which is a solid.




67. (b) Propanal is not formed during the dry distillation of a mixture of calcium formate and calcium acetate. While methanal, propanone and ethanal are formed as follows:






68. (b) The ester having α hydrogen atom show Claisen condensation reaction.

We know that ethyl benzoate ($\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5$) does not contain α -hydrogen. Therefore $\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5$ does not undergo Claisen self condensation.



(As it has a chiral C-atom thus it is optically active)





72. (d)



73. (a) LiAlH_4 can reduce COOH group and not the double bond.



74. (c) There is no reaction hence the resultant mixture contains $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{NaCl}$.




77. (a) When distilled with conc. H_2SO_4 , three molecules of acetone condense to form mesitylene, i.e. 1, 3, 5-trimethylbenzene.




Other three acids contain $-\text{COOH}$ group.






Palmitic acid



Lactic acid








Ethanoic acid



82. (c) CaC_2O_4 is the only compound of Ca which is not soluble in acetic acid. This property of Ca is used in inorganic salt analysis.

83. (c) Cl_2CHCOOH is most acidic because it has two chlorine at α -position.




100. (d)




5. (b) Only aldehydes give reddish brown ppt. with Fehling's solution.
6. (d) Acetone or Propyne form mesitylene (1,3,5-trimethyl benzene) on distillation with conc. H_2SO_4 .
7. (a) Ketones on oxidation give carboxylic acids with lesser number of carbon atoms, i.e.,



8. (d) Structures of given aldehydes




Trichloroethanal, methanal and benzaldehyde do not undergo aldol condensation. Aldol condensation is not given by aldehydes and ketones which do not contain α -hydrogen atom(s).








EXERCISE - 2

1. (a) Ethyne get converted to aldehydes and ketones when treated with an acidified aqueous solution of palladium chloride (PdCl_2) and cupric chloride (CuCl_2) in presence of air or oxygen. This method is known as Wacker's process.




2. (d) Carbonyl compound form crystalline addition product with NaHSO_3 which on treatment with dilute acid or alkali regenerates the carbonyl compound.
3. (a) An electron releasing substituent (+I) intensify the negative charge on the anion resulting in the decrease of stability and thus decreases the acidity of the acid. Hence acid character decreases as the + I-effect of the alkyl group increases as
 $\text{CH}_3^- < \text{CH}_3\text{CH}_2^- < \text{CH}_3\text{CH}_2\text{CH}_2^- < \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2^-$
Hence the order becomes : (i) > (ii) > (iii) > (iv)










11. (c) In $\text{CH}_2 = \text{CH} - \text{CHO}$ due to $-\text{M}$ effect of $-\text{CHO}$ group polarization of electron takes place as follows




Hence partial polarization is represented as




12. (d) Structures of given compounds




Compounds containing $\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} -$ or $\text{CH}_3 - \text{CHOH} -$ group give positive iodoform test.



14. (a) Addition of HCN to carbonyl compounds is nucleophilic addition reaction. The order of reactivity of carbonyl compounds is Aldehydes (smaller to higher) Ketones (smaller to higher), Then




The lower reactivity of Ketones is due to presence of two alkyl group which shows $+\text{I}$ effect. The reactivity of Ketones decreases as the size of alkyl group increases.



16. (a) The correct order of increasing acid strength




Electron withdrawing groups increase the acid strength and electron donating groups decrease the acid strength.




Two products are formed.



18. (c) Mono-carboxylic acids are functional isomers of esters.




19. (c) KMnO_4 converts $-\text{CH}_3$ group of toluene into $-\text{COOH}$ while HI reduces $-\text{COOH}$ group into $-\text{CH}_3$ group.







21. (a)








24. (a) It is an example of intramolecular cannizzaro reaction.




(B)

β -Keto acids, on heating, undergo decarboxylation.







28. (c) In *o*-toluic acid, $-\text{CH}_3$ group is *ortho-para* directing



and $-\text{COOH}$ group is *meta*-directing. So, the resulting product will be (c) in which Br is attached at *para* to $-\text{CH}_3$ and *meta* to $-\text{COOH}$ group.


29. (b) Generally benzene ring is very resistant to oxidation thus side chain is always oxidised to $-\text{COOH}$ group. But if the side chain is tert-alkyl group, oxidation is difficult, but on vigorous oxidation benzene ring is oxidised.






32. (c) Only those aldehydes which do not have α -H atom undergo Cannizzaro's reaction. Hence CH_3CHO will not undergo Cannizzaro's reaction as it has 3 α H atoms.

33. (a)




34. (d)












35. (d) The intermediate is carbocation which is destabilised by C = O group in the first three cases. In (d), α -hydrogen is more acidic which can be removed as water. Moreover, the positive charge on the intermediate carbocation is relatively away from the C = O group.

36. (d) The reactivity of the carbonyl group toward the nucleophilic addition reactions depend upon the magnitude of the positive charge on the carbonyl carbon atom (electronic factor) and also on the crowding around the carbonyl carbon atom in the transition state (steric factor). Both these factors predict the following order






(due to steric crowding).


37. (d) Since, C when heated with Br_2 in presence of KOH produces ethylamine, hence it must be propanamide and hence the organic compound (A) will be propanoic acid. The reactions follows.




40. (c)




(OH group is activating)




(OCH₃ group is more activating)




(Benzene ring having -O- is activated)




44. (b) β -Ketoacids undergo decarboxylation easily.




Reaction (III) is a Hofmann bromamide reaction. Now formation of CH₃CH₂NH₂ is possible only from a compound CH₃CH₂CONH₂ (C) which can be obtained from the compound CH₃CH₂COO⁻NH₄⁺ (B). Thus (A) should be CH₃CH₂COOH



46. (c) Cannizzaro reaction - when an aldehyde containing no α -H undergo reaction in presence of 50% KOH. It disproportionates to form a molecule of carboxylic acid and a molecule of alcohol.



47. (d) Anhydrous alcohols add to the carbonyl group of aldehydes in the presence of anhydrous hydrogen chloride to form acetals via hemiacetals.



48. (c)


49. (d) Benzaldehyde is less reactive than ethanol towards nucleophilic attack. The combined effect of -I and +R effect of phenyl group is electron donating which increases the electron density on the carbon atom of the $>\text{C}=\text{O}$ in benzaldehyde.

50. (b) It is true that lower aldehyde and ketones are soluble in water but as the molecular mass increases their solubility decreases. On adding Tollen's reagent to a solution of Carbonyl compound if silver mirror is obtained then it is aldehyde. Therefore Tollen's reagent is used for the identification of aldehydes and ketones. Here, Statement-1 and Statement-2 both are true but the Statement-2 is not the correct explanation of Statement-1.

EXERCISE - 3

Exemplar Questions

1. (b) But-1-yne on reaction with water in presence of Hg^{2+} ions as a catalyst produces butan-2-one.




2. (a) The carbonyl group in ketones being influenced by two alkyl group is less reactive than in aldehydes where the carbonyl group is under the influence of one alkyl group only. As the number of alkyl group increases both the +I effect and the steric hinderance get increases preventing the attack of nucleophile.

Now among benzaldehyde and acetaldehyde former is less electrophilic than carbon atom of carbonyl group present in ethanal. The polarity of carbonyl group is reduced in benzaldehyde due to resonance hence it is less reactive than ethanal.

3. (c) Due to the electron withdrawing characteristic of $-\text{Cl}$, chloroacetic acid is more acidic than acetic acid further, in phenol phenoxide ion obtained on ionisation of phenol is stabilised due to resonance where as no such stabilisation occurs in case of ethanol. Hence phenol is stronger acid than ethanol. Thus acidic strength increases in the order.


ethanol < phenol < acetic acid < chloro acetic acid

4. (b) Compound $\text{Ph} - \text{COO} - \text{Ph}$ can be prepared by the reaction of phenol & benzoyl chloride




This an example of **Schotten-Baumann reaction**.



5. (c) Acetone and benzaldehyde both do not react with Fehling's solution.
6. (d) Cannizzaro reaction is given by those aldehydes which have absence of α -hydrogen atom. So, CH_3CHO will not give Cannizzaro reaction.
7. (b) Benzaldehyde having no α -hydrogen on reaction with aqueous KOH solution undergo Cannizzaro reaction and produces benzyl alcohol and potassium benzoate.





8. (d) Chemical reaction can be shown as




9. (b) Chemical reaction can be shown as




10. (c)



11. (b) 2° alcohols on oxidation with alkaline KMnO_4 solution produce ketones.



12. (a) Clemmensen reduction is used to convert carbonyl group as follows



Zinc amalgam and HCl act as reagent in this reaction.


NEET/AIPMT (2013-2017) Questions

13. (a) Enolic form predominates in compounds containing two carbonyl groups separated by a $-\text{CH}_2$ group. This is due to following two factors.
- Presence of conjugation which increases stability.
 - Formation of intramolecular hydrogen bond between enolic hydroxyl group and second



14. (c) Zn/Hg and conc. HCl reduce carboxyl group to methylene group (Clemmensen reduction).
15. (d) Any substituent in the carbonyl compound that increases the positive charge on the carbonyl carbon will increase reactivity towards nucleophilic addition.

O=Cc1ccc([N+](=O)[O-])cc1

c1ccc(cc1)[N+](=O)[O-] is most reactive towards nucleophilic addition reaction.



$$\text{(CH}_3\text{—C(=O)—}$$

On reduction




$$\text{H}_3\text{C}-\text{CH}_2-\underset{\text{O}}{\underset{\parallel}{\text{C}}}-\text{CH}_2-\text{CH}_3 \xrightarrow[\text{or NH}_3\text{, -NH}_2\text{/OH}^-]{\text{reduction, Zn-Hg/HCl}} \text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_3$$


Cyclopentanovl anion

18. (d) Enolic form of ethyl acetoacetate has 18 sigma and 2 pi-bonds as shown below:





19. (a) Among the substituent attached to the benzene ring, NO_2 group is the most electron withdrawing, thus withdraws electron density from carbonyl carbon thus facilitate the attack of OH^\ominus ion.




While in all other case no elimination take place.

21. (d) Keto-enol tautomerism is possible only in those aldehydes and ketones which have at least one α -hydrogen atom, which can convert the ketonic group to the enolic group. e.g.



22. (a) Schiff base is formed when 1° amine reacts with aldehydes.



23. (b) Since 'A' gives positive silver mirror test therefore, it must be an aldehyde of α -Hydroxyketone. Also, reaction with OH^- i.e., aldol condensation (by assuming alkali to be dilute) indicates that A is aldehyde as aldol reaction of ketones is reversible and carried out in special apparatus. It indicates that A is an aldehyde

