- The chemical reactions which takes place in both directions are called reversible reactions
- Equilibrium is the end state of a reversible reaction.
- Gaseas Equilibrium is established only in a closed container.
- At equilibrium, the rate of forward and backward
- At equilibrium, the concentration of reactants &



Q.For the equilibrium A B, the variation of the rate of the forward (a) & reverse (b) reaction with time is given by



3. value of k depends only on temperature.

- 4. If K for the reaction  $aA + bB \implies cC+dD$  is K, then K for the reaction cC+dD  $\implies$  aA+ bB will be  $\frac{1}{2}$
- 5 If K for the reaction  $aA + bB \implies cC+dD$  is K, then K for the reaction  $naA + nbB \implies ncC + ndD$  will be (K)<sup>n</sup>
- 6. During the addition of two reactions having equilibrium constants  $K_1 \& K_2$ , then the net Constant  $K = K_1 \times K_2$
- 7. During the subtraction of a reaction having constant  $K_2$  from a reaction having constant  $K_1$ , then the net constant  $K = K_1/K_2$
- 8. If Q<K, the reaction will proceed in forward direction
- 9. If Q>K, the reaction will proceed in backward direction
- 10. If Q=K, the system is in equilibrium.
- 11. If  $K \ge 10^3$ , the reaction is almost complete in forward direction.
- 12. If  $k < 10^{-3}$ , the reaction is mostly backward.
- 13. If K is in blw 10<sup>3</sup> & 10<sup>-3</sup>, almost same reaction takes place

≜ ●

Q.At a given temperature, the equilibrium constants for the reactions,  $NO_{(q)} + \frac{1}{2}O_{2(q)} \longrightarrow NO_{2(q)} \& 2NO_{2(q)} \longrightarrow 2NO_{(q)} + O_{2(q)}$ are  $K_1$  and  $K_2$  respectively. If K, is 4 x 10<sup>-3</sup>. then  $K_2$  will be (A) 8x10<sup>-3</sup> (B) 16x10<sup>-3</sup> (C) 6.25x10<sup>4</sup> (D) 6.25x10<sup>6</sup>

- Such equilibrium is established in physical reactions.
- It is dynamic in nature.

Here vapour pressure is constant at a constant temp

- Established only at a constant temperature (m.p of solid ie, freeging point of liquid)
- Ice-water equilibrium is established at o°c at 1 atm.

- eg: Saturated sugar solution.
- Sugar (dissolved) <u>Sugar</u> (undissolved)

- Here solubility depends upon pressure (Henry's law)
- eg: Soda water
- $CO_2$  (dissolved)  $\leftarrow$   $CO_2$  (undissolved)

Q.Which of the given statements does not elucidate the equilibrium state

(A) The equilibrium can be approached from either direction.

(B) The equilibrium can be attained only if the system is an isolated system. (C) The free energy change at constant pressure and temperature is zero. (D) It is dynamic in nature.

# chemical equilibrium



- Chemical equilibrium approaches from both forward & backward direction
- Chemical equilibrium is dynamic in nature.

## EQUILIBRIUM CONSTANT (K)

For a general reversible reaction aA + bB ← \_\_\_\_ cC+ dD

$$K = \frac{[C]^{C} [D]^{d}}{[A]^{\alpha} [B]^{b}}$$

## **REACTION OUOTIENT (O)**

- At any time during the reaction  $aA+bB \longrightarrow cC+dD$
- At equilibrium Q = K

- Value of k does not depend upon initial concentration of reactants and products.
- 2. Value of k does not depend upon the direction from which equilibrium is attained.

## Q. In the given reaction: A+2B 2C, 2 moles each of A & B present in 10 L of solution combine to form 1 mole of C. Calculate K for the reaction.

ž 🗖

(A) 1.5 (B) 6.67 (C) 0.15 (D) 2.3

According to Le-chatelier's principle, if a system at equilibrium is subjected to a change in concentration, temperature or pressure,

- Conc. of reactant increases → shift towards forward reaction.
- 2. Conc. of product decreases  $\rightarrow$  shift towards forward reaction.
- Conc. of reactant decreases→shift towards backward reaction.
- $\triangle$  Conc. of product increases  $\rightarrow$  shift towards backward reaction
- 5. Pressure increases → shift towards lesser number of aaseous moles
- Pressure decreases → shift towards higher number of gaseous moles
- 8. If temperature increases shift towards endothermic
- 10. Catalyst helps to attain eqm state easily. After the establishment of eqm, catalyst has no effect.
- 12. Addition of inert gas at constant pressure  $\rightarrow$  shift towards

Q.Which one of the following conditions will favour maximum formation of the product in the reaction  $A_{2(a)} + B_{2(a)} \longrightarrow X_{2(a)}$ ,  $\Delta_r H = -XkJ/mol$ ?

(A) Low temperature and high pressure (B) High temperature and high pressure

(C) Low temperature and low pressure (D) High temperature and low pressure