

# 26. VECTORS

## 1. INTRODUCTION TO VECTOR ALGEBRA

### 1.1 Scalars and Vectors

**Scalar:** A scalar is a quantity that has only magnitude but no direction. Scalar quantity is expressed as a single number, followed by appropriate unit, e.g. length, area, mass, etc. In linear algebra, real numbers are called scalars.

**Vector:** A vector is a quantity that has both magnitude and direction, e.g. displacement, velocity, etc.

### 1.2 Representation of Vectors

- (a) A vector is represented diagrammatically by a directed line segment or an arrow. A directed line segment has both magnitude (length) and direction. The length is denoted by  $|V|$ .
- (b) If P and Q are the given two points, then the vector from P to Q is denoted by  $\overrightarrow{PQ}$ , where P is called the tail and Q is called the nose of the vector.

### 1.3 Vector Components

In a two-dimensional coordinate system, any vector can be resolved into x-component and y-component

$$\vec{v} = \langle v_x, v_y \rangle$$

Let us consider the figure shown (adjacent) here. In this figure, the components can be quickly read. The vector in the component form is  $\vec{v} = \langle 4, 5 \rangle$ .

The relation between magnitude of the vector and the components of the vector can be calculated by using trigonometric ratios.

$$\cos \theta = \frac{\text{Adjacent side}}{\text{Hypotenuse}} = \frac{v_x}{v}$$

$$\sin \theta = \frac{\text{Opposite side}}{\text{Hypotenuse}} = \frac{v_y}{v}$$

$$v_x = v \cos \theta; v_y = v \sin \theta$$




Figure 26.1

If  $v_x$  and  $v_y$  are the known lengths of a right triangle, then the length of the hypotenuse,  $V$ , is calculated by using the Pythagorean theorem

$$|V| = \sqrt{v_x^2 + v_y^2}$$

## 2. TYPE OF VECTORS

### 2.1 Null Vector/Zero Vector

A zero vector or null vector is a vector that has zero magnitude, i.e. initial and terminal points are coincident, so that its direction is in indeterminate form. It is denoted by  $\phi$ .

### 2.2 Unit Vector

A unit vector is a vector of unit length. A unit vector is sometimes denoted by replacing the arrow on a vector with " $\wedge$ ".

Unit vectors parallel to x-axis, y-axis and z-axis are denoted by  $\hat{i}$ ,  $\hat{j}$  and  $\hat{k}$ , respectively.

Unit vector  $\hat{U}$  parallel to  $\vec{V}$  can be obtained as  $\hat{U} = \frac{\vec{V}}{|\vec{V}|}$ .

**Illustration 1:** Find unit vector of  $\vec{i} - 2\vec{j} + 3\vec{k}$

(JEE MAIN)

**Sol:** Here unit vector of  $\vec{a}$  is given by  $\hat{a} = \frac{\vec{a}}{|\vec{a}|}$ .

$$\vec{a} = \vec{i} - 2\vec{j} + 3\vec{k}$$

If  $\vec{a} = a_x\hat{i} + a_y\hat{j} + a_z\hat{k}$  then it's magnitude  $|\vec{a}| = \sqrt{a_x^2 + a_y^2 + a_z^2} \Rightarrow |\vec{a}| = \sqrt{14} \Rightarrow \hat{a} = \frac{\vec{a}}{\sqrt{14}} = \frac{1}{\sqrt{14}}(\vec{i} - 2\vec{j} + 3\vec{k})$

### 2.3 Collinear or Parallel Vectors

Two or more vectors are said to be collinear, when they are along the same lines or parallel lines irrespective of their magnitudes and directions.

### 2.4 Like and Unlike Vectors

Vectors having the same direction are called like vectors. Any two vectors parallel to one another, having unequal magnitudes and acting in opposite directions are called unlike vectors.

### 2.5 Co-Initial Vectors

All those vectors whose terminal points are same, are called co-terminal vectors.

### 2.6 Co-Terminal Vectors

Vectors that have the same initial points are called co-initial vectors.

**Illustration 2:** Which are co-initial and equal vectors in the given rectangle diagram?

(JEE MAIN)

**Sol:** By following above mentioned conditions we can obtain co-initial and equal vectors.




Figure 26.2

Here,  $\vec{a}$  and  $\vec{b}$  are co-initial vectors,  $\vec{b}$  and  $\vec{c}$ ,  $\vec{a}$  and  $\vec{d}$  are equal vectors.

## 2.7 Coplanar Vectors

Vectors lie on the same plane are called coplanar.

## 2.8 Negative vector

A vector that points to a direction opposite to that of the given vector is called a negative vector.

## 2.9 Reciprocal of a Vector

A vector having the same direction as that of a given vector  $\vec{a}$ , but magnitude equal to the reciprocal of the given vector is known as the reciprocal of  $\vec{a}$  and is denoted by  $\vec{a}^{-1}$ .

## 2.10 Localized and Free vectors

A vector drawn parallel to a given vector through a specified point unlike free vector in space is called a localized vector. For example, the effect of force acting on a rigid body depends not only on the magnitude and direction but also on the line of action of the force. A vector that depends only on its length and direction and not on its position in the space is called a free vector, e.g. gravity. In this chapter, we will deal with free vectors, unless otherwise stated. Thus a free vector can be determined in space by choosing an arbitrary initial point.

**Illustration 3:** Let  $\vec{a} = \hat{i} + 2\hat{j}$  and  $\vec{b} = 2\hat{i} + \hat{j}$ . Is  $|\vec{a}| = |\vec{b}|$ ? Are the vectors  $\vec{a}$  and  $\vec{b}$  equal? **(JEE MAIN)**

**Sol:** Two vectors are equal if their modulus and corresponding components both are equal.

We have  $|\vec{a}| = \sqrt{1^2 + 2^2} = \sqrt{5}$  and  $|\vec{b}| = \sqrt{2^2 + 1^2} = \sqrt{5}$ . So,  $|\vec{a}| = |\vec{b}|$ . But, the two vectors are not equal, since their corresponding components are distinct.

**Illustration 4:** Find a vector of magnitude 5 units which is parallel to the vector  $2\hat{i} - \hat{j}$ . **(JEE MAIN)**

**Sol:** As we know  $\hat{a} = \frac{\vec{a}}{|\vec{a}|}$ , therefore required vector will be  $5\hat{a}$ .

Let  $\vec{a} = 2\hat{i} - \hat{j}$ . Then,  $|\vec{a}| = \sqrt{2^2 + (-1)^2} = \sqrt{5}$

$\therefore$  Unit vector parallel to  $\vec{a} = \hat{a} = \frac{1}{|\vec{a}|} \cdot \vec{a} = \frac{1}{\sqrt{5}} (2\hat{i} - \hat{j}) = \frac{2}{\sqrt{5}} \hat{i} - \frac{1}{\sqrt{5}} \hat{j}$ .

So, the required vector is  $5\hat{a} = 5 \left( \frac{2}{\sqrt{5}} \hat{i} - \frac{1}{\sqrt{5}} \hat{j} \right) = 2\sqrt{5}\hat{i} - \sqrt{5}\hat{j}$ .

## 2.11 Position Vector

A vector that represents the position of a point P in space with respect to an arbitrary reference origin O is called a position vector (p.v.). It is also known as location vector or radius vector and usually denoted as  $x$ ,  $r$  or  $s$ ; it corresponds to the displacement from O to P.

$r = \overrightarrow{OP}$ .

**Illustration 5:** Show that, the three points A(-2,3,5), B(1,2,3) and C(7,0,-1) are collinear.

**(JEE MAIN)**

**Sol:** By obtaining  $\overrightarrow{AB}$  and  $\overrightarrow{BC}$ , we can conclude that given points are collinear or not.

We have

$$\overrightarrow{AB} = \overrightarrow{OB} - \overrightarrow{OA} = \left( \hat{i} + 2\hat{j} + 3\hat{k} \right) - \left( -2\hat{i} + 3\hat{j} + 5\hat{k} \right) = 3\hat{i} - \hat{j} - 2\hat{k}$$

$$\overrightarrow{BC} = \overrightarrow{OC} - \overrightarrow{OB} = \left( 7\hat{i} + 0\hat{j} - \hat{k} \right) - \left( \hat{i} + 2\hat{j} + 3\hat{k} \right) = 6\hat{i} - 2\hat{j} - 4\hat{k} = 2 \left( 3\hat{i} - \hat{j} - 2\hat{k} \right)$$

Therefore,  $\overrightarrow{BC} = 2\overrightarrow{AB}$ .

This shows that the vectors  $\overrightarrow{AB}$  and  $\overrightarrow{BC}$  are parallel. But, B is a common point. So, the given point A, B and C are collinear.

## 2.12 Equal Vectors

Two vectors having the same corresponding components and direction and represent the same physical quantity are called equal vectors.

**Illustration 6:** Find the values of x, y and z, so that the vectors  $\vec{a} = x\hat{i} + 2\hat{j} + z\hat{k}$  and  $\vec{b} = 2\hat{i} + y\hat{j} + \hat{k}$  are equal.

**(JEE MAIN)**

**Sol:** Two vectors are equal, if their corresponding components are equal.

Note that two vectors are equal, if their corresponding components are equal. Thus, the given vectors  $\vec{a}$  and  $\vec{b}$  will be equal, if and only if  $x = 2$ ,  $y = 2$ ,  $z = 1$ .

**Illustration 7:** Find the vector joining the point P (2, 3, 0) and Q (-1, -2, -4) directed from P to Q.

**(JEE MAIN)**

**Sol:** By subtracting the component of P from Q we will get  $\overrightarrow{PQ}$ .

Since the vector is to be directed from P to Q. Clearly, P is the initial point and Q is the terminal point. So, the required vector joining P and Q is the vector PQ given by

$$\overrightarrow{PQ} = \overrightarrow{OQ} - \overrightarrow{OP} = (-1 - 2)\hat{i} + (-2 - 3)\hat{j} + (-4 - 0)\hat{k} \quad \text{i.e. } \overrightarrow{PQ} = -3\hat{i} - 5\hat{j} - 4\hat{k}$$

**Illustration 8:** Show that, the points  $A(2\hat{i} - \hat{j} + \hat{k})$ ,  $B(\hat{i} - 3\hat{j} - 5\hat{k})$ ,  $C(3\hat{i} - 4\hat{j} - 4\hat{k})$  are the vertices of a right-angled triangle.

**(JEE MAIN)**

**Sol:** Here if  $|\overrightarrow{AB}|^2 = |\overrightarrow{BC}|^2 + |\overrightarrow{CA}|^2$  then only the given points are the vertices of right angled triangle. We have

$$\overrightarrow{AB} = (1 - 2)\hat{i} + (-3 + 1)\hat{j} + (-5 - 1)\hat{k} = -\hat{i} - 2\hat{j} - 6\hat{k}$$

$$\overrightarrow{BC} = (3 - 1)\hat{i} + (-4 + 3)\hat{j} + (-4 + 5)\hat{k} = 2\hat{i} - \hat{j} + \hat{k} \quad \text{and} \quad \overrightarrow{CA} = (2 - 3)\hat{i} + (-1 + 4)\hat{j} + (1 + 4)\hat{k} = -\hat{i} + 3\hat{j} + 5\hat{k}$$

$$\text{Moreover, } |\overrightarrow{AB}|^2 = 41 = 6 + 35 = |\overrightarrow{BC}|^2 + |\overrightarrow{CA}|^2$$

Hence, it is proved that the points form a right-angled triangle.

### 3. RESULTANT OF VECTORS

When two or more vectors are added, they yield the resultant vector. If vectors A and B are added together, the result will be vector R, i.e.  $\vec{R} = \vec{A} + \vec{B}$ . Same technique can also be applied for multiple vectors.

### 4. VECTOR ADDITION

#### 4.1 Triangular Law of Addition

It states that if two vectors can be represented in magnitude and direction by the two sides of a triangle taken in the same order, then their resultant is represented by the third side of the triangle, taken in the opposite direction of the sequence.

#### 4.2 Parallelogram Law of Addition

It states that if two vectors can be represented in magnitude and direction by the two adjacent sides or a parallelogram, then their resultant is represented by the diagonal of the parallelogram.

#### 4.3 Addition in Component Form

Consider two vectors A and B

$$A = \langle a_1, b_1, c_1 \rangle$$

$$B = \langle a_2, b_2, c_2 \rangle$$

$$\text{Then, } A + B = \langle a_1 + a_2, b_1 + b_2, c_1 + c_2 \rangle$$

#### 4.4 Properties of Vector Addition

The properties of vector addition are listed as follows:

- |     |                         |                                     |
|-----|-------------------------|-------------------------------------|
| (a) | $\pi / 2$               | Commutative                         |
| (b) | $\pi / 3$               | Associative                         |
| (c) | $\pi / 4$               | Null vector is an additive identity |
| (d) | $\hat{A}$ and $\hat{B}$ | Additive inverse                    |
| (e) | $\pi$                   |                                     |
| (f) | $ \hat{A} - \hat{B} $   |                                     |
| (g) | $\pi / 2$               |                                     |

#### 4.5 Vector Subtraction

Subtraction is taken as an inverse operation of addition. If  $\vec{u}$  and  $\vec{v}$  are two vectors, the difference  $\vec{u} - \vec{v}$  of two vectors is defined to be the vector added to  $\vec{v}$  to get  $\vec{u}$ . In order to obtain  $\vec{u} - \vec{v}$ , we put the tails of  $\vec{u}$  and  $\vec{v}$  together, the directed segment from the nose of  $\vec{v}$  to the nose of  $\vec{u}$  is a representative of  $\vec{u} - \vec{v}$ .

**Illustration 9:** If  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$  and  $\vec{b} = 2\hat{i} + 4\hat{j} - 5\hat{k}$  represent two adjacent sides of a parallelogram, find the unit vectors parallel to the diagonals of the parallelogram. **(JEE MAIN)**

**Sol:** As mentioned above, if two vector quantities are represented by two adjacent sides or a parallelogram then the diagonal of parallelogram will be equal to the resultant of these two vectors.

Let ABCD be a parallelogram such that,  $\overline{AB} // \vec{b}$  and  $\overline{BC} // \vec{a}$ .

Then,

$$\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC} \Rightarrow \overrightarrow{AC} = \vec{a} + \vec{b} = 3\hat{i} + 6\hat{j} - 2\hat{k} \quad \text{and} \quad \overrightarrow{AB} + \overrightarrow{BD} = \overrightarrow{AD}$$

$$\Rightarrow \overrightarrow{BD} = \overrightarrow{AD} - \overrightarrow{AB} \Rightarrow \overrightarrow{BD} = \vec{b} - \vec{a} = \hat{i} + 2\hat{j} - 8\hat{k}$$

$$\text{Now, } \overrightarrow{AC} = 3\hat{i} + 6\hat{j} - 2\hat{k} \Rightarrow |\overrightarrow{AC}| = \sqrt{9+36+4} = 7$$

$$\text{And } \overrightarrow{BD} = \hat{i} + 2\hat{j} - 8\hat{k}.$$

$$\Rightarrow |\overrightarrow{BD}| = \sqrt{1+4+64} = \sqrt{69}$$

$$\therefore \text{Unit Vector along } \overrightarrow{AC} = \frac{\overrightarrow{AC}}{|\overrightarrow{AC}|} = \frac{1}{7}(3\hat{i} + 6\hat{j} - 2\hat{k})$$

$$\therefore \text{Unit vector along } \overrightarrow{BD} = \frac{\overrightarrow{BD}}{|\overrightarrow{BD}|} = \frac{1}{\sqrt{69}}(\hat{i} + 2\hat{j} - 8\hat{k}).$$




Figure 26.3

**Illustration 10:** ABCDE is a pentagon. Prove that the resultant of the forces  $\overrightarrow{AB}$ ,  $\overrightarrow{AE}$ ,  $\overrightarrow{BC}$ ,  $\overrightarrow{DC}$ ,  $\overrightarrow{ED}$  and  $\overrightarrow{AC}$  is  $3\overrightarrow{AC}$ .  
(JEE MAIN)

**Sol:** By using method of finding resultant of vector we can prove required result.

Let R be the resultant force

$$\therefore R = \overrightarrow{AB} + \overrightarrow{AE} + \overrightarrow{BC} + \overrightarrow{DC} + \overrightarrow{ED} + \overrightarrow{AC}$$

$$\therefore R = (\overrightarrow{AB} + \overrightarrow{BC}) + (\overrightarrow{AE} + \overrightarrow{ED} + \overrightarrow{DC}) + \overrightarrow{AC}$$

$$= \overrightarrow{AC} + \overrightarrow{AC} + \overrightarrow{AC} = 3\overrightarrow{AC}. \text{ Hence proved.}$$




Figure 26.4

**Illustration 11:** ABCD is a parallelogram. If L and M are the middle points of BC and CD, respectively express  $\overrightarrow{AL}$  and  $\overrightarrow{AM}$  in terms of  $\overrightarrow{AB}$  and  $\overrightarrow{AD}$ , also show that  $\overrightarrow{AL} + \overrightarrow{AM} = \frac{3}{2}\overrightarrow{AC}$ .  
(JEE MAIN)

**Sol:** By using mid – point formula and method of finding resultant of vector we can prove given relation.

Let  $\vec{b}$  and  $\vec{d}$  be the position vectors of points B and D, respectively be referred to A as the origin of reference.

$$\text{Then } \overrightarrow{AC} = \overrightarrow{AD} + \overrightarrow{DC} = \overrightarrow{AD} + \overrightarrow{AB} \quad [\because \overrightarrow{DC} = \overrightarrow{AB}]$$

$$= \vec{d} + \vec{b} \quad \therefore \overrightarrow{AB} = \vec{b}, \quad \overrightarrow{AD} = \vec{d}$$

i.e. the position vector of C referred to A is  $\vec{d} + \vec{b}$

$$\overrightarrow{AL} = \text{p.v. of L, the midpoint of } \overrightarrow{BC} \quad \therefore \overrightarrow{AM} = \frac{1}{2}[\vec{a} + \vec{d} + \vec{b}] = \overrightarrow{AD} + \frac{1}{2}\overrightarrow{AB}$$

$$\therefore \overrightarrow{AL} + \overrightarrow{AM} = \vec{b} + \vec{d} + \frac{1}{2}\vec{b} = \frac{3}{2}\vec{b} + \frac{3}{2}\vec{d} = \frac{3}{2}(\vec{b} + \vec{d}) = \frac{3}{2}\overrightarrow{AC}$$




Figure 26.5

## 5. SCALAR MULTIPLE OF A VECTOR

If  $\vec{a}$  is the given vector, then  $k\vec{a}$  is a vector, whose magnitude is  $|k|$  times the magnitude of  $\vec{a}$  and whose direction is the same or opposite as that of  $\vec{a}$  according to whether  $k$  is positive or negative.

## 6. SECTION FORMULA

- (a) If  $\vec{a}$  and  $\vec{b}$  are the position vectors of two points A and B, then the position vector of a point which divides A and B in the ratio  $m:n$  is given by  $\vec{r} = \frac{(n\vec{a} + m\vec{b})}{(m+n)}$ .
- (b) Position vector of the midpoint of  $\overline{AB} = \frac{(\vec{a} + \vec{b})}{2}$ .

### PLANCESS CONCEPTS

- If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are the position vectors of the vertices of any  $\triangle ABC$ . Then the position vector of centroid G will be  $\frac{\vec{a} + \vec{b} + \vec{c}}{3}$ .
- The position vector of incenter of triangle with position vectors of triangle ABC, are A ( $\vec{a}$ ), B ( $\vec{b}$ ), C ( $\vec{c}$ ) is  $\vec{r} = \frac{a\vec{a} + b\vec{b} + c\vec{c}}{a + b + c}$ .

Anurag Saraf (JEE 2011, AIR 226)

**Illustration 12:** If ABCD is a quadrilateral and E and F are the mid points of AC and BD, respectively, prove that  $\overline{AB} + \overline{AD} + \overline{CB} + \overline{CD} = 4\overline{EF}$ . (JEE MAIN)

**Sol:** By using mid-point theorem we can prove given relation.

Since F is the midpoint of BD. Applying the midpoint theorem in triangle ABD,

we have  $\Rightarrow \overline{AB} + \overline{AD} = 2\overline{AF}$

Applying the midpoint theorem in triangle BCD, we have

$\Rightarrow \overline{CB} + \overline{CD} = 2\overline{CF}$

Adding equations (i) and (ii), we obtain

$\Rightarrow \overline{AB} + \overline{AD} + \overline{CB} + \overline{CD} = 2(\overline{AF} + \overline{CF})$

Now applying the midpoint theorem in triangle CFA, we have  $\overline{AF} + \overline{CF} = 2\overline{EF}$

$\Rightarrow \overline{AB} + \overline{AD} + \overline{CB} + \overline{CD} = 2(\overline{AF} + \overline{CF}) = 4\overline{EF}$  Hence proved.




Figure 26.6

**Illustration 13:** If G is the centroid of the triangle ABC, show that  $\vec{GA} + \vec{GB} + \vec{GC} = 0$  and conversely  $\vec{GA} + \vec{GB} + \vec{GC} = 0$ , then G is the centroid of triangle ABC. **(JEE ADVANCED)**

**Sol:** As G is the centroid of triangle ABC, hence  $G = \frac{\vec{a} + \vec{b} + \vec{c}}{3}$ . Therefore

by obtaining  $\vec{GA}$ ,  $\vec{GB}$  and  $\vec{GC}$  we can prove this problem.

Let the position vector of the vertices be  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$ , respectively.

So, the position vector of centroid, G, is  $\frac{\vec{a} + \vec{b} + \vec{c}}{3}$ .

$$\vec{GA} = \vec{OA} - \vec{OG} = \vec{a} - \frac{\vec{a} + \vec{b} + \vec{c}}{3} = \frac{2\vec{a} - \vec{b} - \vec{c}}{3}$$

$$\text{Similarly, } \vec{GB} = \frac{2\vec{b} - \vec{a} - \vec{c}}{3}, \quad \vec{GC} = \frac{2\vec{c} - \vec{a} - \vec{b}}{3}$$

$$\Rightarrow \vec{GA} + \vec{GB} + \vec{GC} = \frac{1}{3}(2\vec{a} - 2\vec{a} + 2\vec{b} - 2\vec{b} + 2\vec{c} - 2\vec{c}) = 0$$

Conversely if  $\vec{GA} + \vec{GB} + \vec{GC} = 0$

$$\Rightarrow (\vec{OA} - \vec{OG}) + (\vec{OB} - \vec{OG}) + (\vec{OC} - \vec{OG}) = 0 \Rightarrow \vec{OA} + \vec{OB} + \vec{OC} = 3\vec{OG} \Rightarrow \vec{OG} = \frac{\vec{OA} + \vec{OB} + \vec{OC}}{3}$$

Hence, G is the centroid of the points A, B and C.




Figure 26.7

**Illustration 14:** Find the values of x and y, for which the vectors  $\vec{a} = (x+2)\hat{i} - (x-y)\hat{j} + \hat{k}$ ,  $\vec{b} = (x-1)\hat{i} + (2x+y)\hat{j} + 2\hat{k}$  are parallel **(JEE MAIN)**

**Sol:** Two vectors are parallel if ratio of their respective components are equal.

$$\vec{a} \text{ and } \vec{b} \text{ are parallel if } \frac{x+2}{x-1} = \frac{y-x}{2x+y} = \frac{1}{2} \Rightarrow x = -5, y = \frac{-20}{3}$$

**Illustration 15:** If ABCD is a parallelogram and E is the midpoint of AB, show by vector method, that DE trisects and is trisected by AC. **(JEE MAIN)**

**Sol:** By using section formula, we can solve this problem.

$$\text{Let } \vec{AB} = \vec{a} \text{ and } \vec{BC} = \vec{b}$$

$$\text{Then } \vec{BC} = \vec{AD} = \vec{b} \text{ and } \vec{AC} = \vec{AB} + \vec{AD} = \vec{a} + \vec{b}$$

Also, let K be a point on AC, such that  $AK:AC = 1:3$

$$\Rightarrow \vec{AK} = \frac{1}{3} \vec{AC} \Rightarrow \vec{AK} = \frac{1}{3} (\vec{a} + \vec{b}) \quad \dots (i)$$

$$\text{Let E be the midpoint of AB, such that } \vec{AE} = \frac{1}{2} \vec{a}$$

Let M be the point on DE such that  $DM:ME = 2:1$

$$\therefore \vec{AM} = \frac{\vec{AD} + 2\vec{AE}}{1+2} = \frac{\vec{b} + \vec{a}}{3} \quad \dots (ii)$$

Comparing equations (i) and (ii), we find that  $\vec{AK} = \frac{(\vec{b} + \vec{a})}{3} = \vec{AM}$ , and thus we conclude that K and M coincide, i.e. DE trisects AC and is trisected by AC. Hence proved.




Figure 26.8



## 7. LINEAR COMBINATION OF VECTORS

### 7.1 Collinear and Non-Collinear Vectors

Let  $\vec{a}$  and  $\vec{b}$  be non-zero vectors. These vectors are said to be collinear if there exists  $\lambda \neq 0$  such that  $\vec{a} + \lambda\vec{b} + \gamma\vec{c} = 0$ .

Given a finite set of vectors  $\vec{a}, \vec{b}, \vec{c}, \dots$ , then the vector  $\vec{r} = x\vec{a} + y\vec{b} + z\vec{c} + \dots$  is called a linear combination of  $\vec{a}, \vec{b}, \vec{c}, \dots$ , for any scalar  $x, y, z, \dots \in \mathbb{R}$ .

### 7.2 Collinearity of Three Points

Let three points with position vectors (non-zero)  $\vec{a}, \vec{b}$ , and  $\vec{c}$  be collinear. Then there exists  $\lambda, \gamma$  both not being 0 such that  $\vec{a} + \lambda\vec{b} + \gamma\vec{c} = 0$

### 7.3 Coplanar Vectors

Let  $\vec{a}$  and  $\vec{b}$ , be non-zero, non-collinear vectors. Then, any vector  $\vec{r}$  coplanar with  $\vec{a}, \vec{b}$  can be uniquely expressed as a linear combination of  $\vec{a}, \vec{b}$ , i.e. there exist some unique  $x, y \in \mathbb{R}$ , such that.

#### PLANCESS CONCEPTS

- If  $\vec{a}, \vec{b}, \vec{c}$  are non-zero, non-coplanar vectors, then  $x\vec{a} + y\vec{b} + z\vec{c} = x'\vec{a} + y'\vec{b} + z'\vec{c} \Rightarrow x = x', y = y', z = z'$
- Let  $\vec{a}, \vec{b}, \vec{c}$  be non-zero, non-coplanar vectors in space. Then any vector  $\vec{r}$  can be uniquely expressed as a linear combination of  $\vec{a}, \vec{b}, \vec{c}$  or there exists some unique  $x, y, z \in \mathbb{R}$ , such that  $x\vec{a} + y\vec{b} + z\vec{c} = \vec{r}$ .

**Vaibhav Krishan (JEE 2009, AIR 22)**

### 7.4 Linear Dependency of Vectors

A set of vectors  $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$  is said to be linearly independent if the vector equation  $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_p \vec{v}_p = 0$  has only a trivial solution. The set  $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$  is said to be linearly dependent if there exists weights  $c_1, \dots, c_p$ , not all 0, such that  $c_1 \vec{v}_1 + c_2 \vec{v}_2 + \dots + c_p \vec{v}_p = 0$

#### PLANCESS CONCEPTS

- Two non-zero, non-collinear vectors are linearly independent.
- Any two collinear vectors are linearly dependent.
- Any three non-coplanar vectors are linearly independent.
- Any three coplanar vectors are linearly dependent.
- Any four vectors in three-dimensional space are linearly dependent.

**Nitish Jhavar (JEE 2009, AIR 7)**

**Illustration 16:** The position vectors of three points  $A = \vec{a} - 2\vec{b} + 3\vec{c}$ ,  $B = 2\vec{a} + 3\vec{b} - 4\vec{c}$  and  $C = -7\vec{b} + 10\vec{c}$ . Prove that the vectors  $\overrightarrow{AB}$  and  $\overrightarrow{AC}$  are linearly dependent. **(JEE MAIN)**

**Sol:** Here obtain  $\overrightarrow{AB}$  and  $\overrightarrow{AC}$  to check its linear dependency.

Let O be the point of reference, then,  $\overrightarrow{OA} = \vec{a} - 2\vec{b} + 3\vec{c}$ ,  $\overrightarrow{OB} = 2\vec{a} + 3\vec{b} - 4\vec{c}$ , and  $\overrightarrow{OC} = -7\vec{b} + 10\vec{c}$

$$\Rightarrow \overrightarrow{AC} = \overrightarrow{OC} - \overrightarrow{OA} = (-7\vec{b} + 10\vec{c}) - (\vec{a} - 2\vec{b} + 3\vec{c}) = -\vec{a} - 5\vec{b} + 7\vec{c}$$

$$\overrightarrow{AB} = \overrightarrow{OB} - \overrightarrow{OA} = (2\vec{a} + 3\vec{b} - 4\vec{c}) - (\vec{a} - 2\vec{b} + 3\vec{c}) = \vec{a} + 5\vec{b} - 7\vec{c}$$

$$\therefore \overrightarrow{AC} = \lambda \overrightarrow{AB}, \text{ where } \lambda = -1.$$

Hence  $\overrightarrow{AB}$  and  $\overrightarrow{AC}$  are linearly dependent.

**Illustration 17:** Prove that the vectors  $5\vec{a} + 6\vec{b} + 7\vec{c}$ ,  $7\vec{a} - 8\vec{b} + 9\vec{c}$  and  $3\vec{a} + 20\vec{b} + 5\vec{c}$  are linearly dependent and  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$ , being linearly independent vectors. **(JEE MAIN)**

**Sol:** We know that if these vectors are linearly dependent, then we can express one of them as a linear combination of the other two.

Now, let us assume that the given vectors are coplanar, and then we can write

$$5\vec{a} + 6\vec{b} + 7\vec{c} = \ell(7\vec{a} - 8\vec{b} + 9\vec{c}) + m(3\vec{a} + 20\vec{b} + 5\vec{c}), \text{ where } \ell \text{ and } m \text{ are scalars.}$$

Comparing the coefficients of  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  on both sides of the equation

$$5 = 7\ell + 3m \quad \dots (i)$$

$$6 = -8\ell + 20m \quad \dots (ii)$$

$$7 = 9\ell + 5m \quad \dots (iii)$$

From equations (i) and (iii), we get

$$4 = 8\ell \Rightarrow \ell = \frac{1}{2} = m, \text{ which evidently satisfies equation (ii) too.}$$

Hence, the given vectors are linearly dependent.

**Illustration 18:** Prove that the four points  $2\vec{a} + 3\vec{b} - \vec{c}$ ,  $\vec{a} - 2\vec{b} + 3\vec{c}$ ,  $3\vec{a} + 4\vec{b} - 2\vec{c}$  and  $\vec{a} - 6\vec{b} + 6\vec{c}$  are coplanar.

**(JEE MAIN)**

**Sol:** Let the given four points be P, Q, R and S respectively. These points are coplanar, if the vectors  $\overrightarrow{PQ}$ ,  $\overrightarrow{PR}$  and  $\overrightarrow{PS}$  are coplanar. These vectors are coplanar if one of them can be expressed as a linear combination of other two.

So, let

$$\overrightarrow{PQ} = x\overrightarrow{PR} + y\overrightarrow{PS}$$

$$\Rightarrow -\vec{a} - 5\vec{b} + 4\vec{c} = x(\vec{a} + \vec{b} - \vec{c}) + y(-\vec{a} - 9\vec{b} + 7\vec{c})$$

$$\Rightarrow -\vec{a} - 5\vec{b} + 4\vec{c} = (x - y)\vec{a} + (x - 9y)\vec{b} + (-x + 7y)\vec{c}$$

$$\Rightarrow x - y = -1, x - 9y = -5, -x + 7y = 4$$

$$\text{Solving the first two of these three equations, we get } x = -\frac{1}{2}, y = \frac{1}{2}$$

On substituting the values of x and y in the third equation, we find that the third equation is satisfied. Hence, the given four points are coplanar.

**Illustration 19:** Show that, the vectors  $2\vec{a} - \vec{b} + 3\vec{c}$ ,  $\vec{a} + \vec{b} - 2\vec{c}$  and  $\vec{a} + \vec{b} - 3\vec{c}$  are non-coplanar vectors.

(JEE MAIN)

**Sol:** If vectors are coplanar then one of them can be expressed as a linear combination of other two otherwise they are non-coplanar. Assume the given vectors are coplanar.

Then one of the given vectors is expressible in terms of the other two.

Let  $2\vec{a} - \vec{b} + 3\vec{c} = x(\vec{a} + \vec{b} - 2\vec{c}) + y(\vec{a} + \vec{b} - 3\vec{c})$  for some scalars  $x$  and  $y$ .

$$\Rightarrow 2\vec{a} - \vec{b} + 3\vec{c} = (x + y)\vec{a} + (x + y)\vec{b} + (-2x - 3y)\vec{c} \Rightarrow 2 = x + y, -1 = x + y \text{ and } 3 = 2x - 3y,$$

Clearly, the first two equations contradict each other. Hence, it is proved that the given vectors are not coplanar.

## 8. SCALAR OR DOT PRODUCT

The scalar product of two vectors  $\vec{a} = (a_1, a_2, a_3)$  and  $\vec{b} = (b_1, b_2, b_3)$  is written using a dot as an operator ( $\cdot$ ) between the two vectors. The component form of the dot product is as follows:

$$\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3 \quad \dots (i)$$

And in geometrical form

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta \quad \dots (ii)$$

where  $\theta$  is the angle between the two vectors and  $0 \leq \theta \leq \pi$ .

From equation (i), it can also be written as

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{a_1 b_1 + a_2 b_2 + a_3 b_3}{|\vec{a}| |\vec{b}|},$$

which can be used to find the angle between two vectors. If  $\vec{a}$  and  $\vec{b}$  are perpendicular then

$$\theta = 90^\circ \Rightarrow \cos \theta = 0 \Rightarrow \vec{a} \cdot \vec{b} = 0$$

### PLANCESS CONCEPTS

- $\vec{a} \cdot \vec{b} \leq |\vec{a}| |\vec{b}|$
- $\vec{a} \cdot \vec{b} > 0 \Rightarrow$  Angle between  $a$  and  $b$  is acute.
- $\vec{a} \cdot \vec{b} < 0 \Rightarrow$  Angle between  $a$  and  $b$  is obtuse.

Shivam Agarwal (JEE 2009, AIR 27)

### Geometrical Interpretation of Dot Product

The scalar product is used to determine the projection of  $\vec{r}$  vector along the given direction.

$\overline{ON}$  is the component of vector  $\overline{OB} (= \vec{b})$  in the direction of vector

$\overline{OA} (= \vec{a})$ ;  $\overline{ON} = b \cos \theta$ . Thus the projection of  $\vec{b}$  along  $\hat{a} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|} \hat{a}$

$$\therefore \overline{ON} = \left( \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|^2} \right) \vec{a}$$




Figure 26.9

$$\text{Projection of } \vec{a} \text{ along } \vec{b} = \left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \right) \hat{b} \quad \therefore \overrightarrow{OM} = \left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \right) \vec{b}$$

## 8.1 Properties of Scalar Product

The properties of scalar product are listed as follows:

- (a)  $\vec{a}, \vec{b}$  are vectors and  $\vec{a} \cdot \vec{b}$  is a number      (b)  $\vec{a} \cdot \vec{a} = |\vec{a}|^2$       (c)  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$   
 (d)  $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$       (e)  $(c\vec{a}) \cdot \vec{b} = c(\vec{a} \cdot \vec{b})$       (f)  $\vec{0} \cdot \vec{a} = 0$   
 (g)  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$       (h)  $\vec{a} \cdot \vec{b} = 0 \Leftrightarrow \vec{a} = \vec{0} \text{ or } \vec{b} = \vec{0} \text{ or } \vec{a} \perp \vec{b}$

**Illustration 20:** Find the angle ' $\theta$ ' between the vectors  $\vec{a} = \hat{i} + \hat{j} - \hat{k}$  and vectors  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$  **(JEE MAIN)**

**Sol:** The angle  $\theta$  between the two vectors  $\vec{a}$  and  $\vec{b}$  is given by  $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$

$$\text{Now } \vec{a} \cdot \vec{b} = (\hat{i} + \hat{j} - \hat{k}) \cdot (\hat{i} - \hat{j} + \hat{k}) = 1 - 1 - 1 = -1$$

Therefore, we have  $\cos \theta = \frac{-1}{3}$ . Hence, the required angle is  $\theta = \cos^{-1} \left( \frac{-1}{3} \right)$

**Illustration 21:** Find the length of the projection of vector  $\vec{a} = 2\hat{i} + 3\hat{j} + 2\hat{k}$  on vector  $\vec{b} = \hat{i} + 2\hat{j} + \hat{k}$ . **(JEE MAIN)**

**Sol:** The projection of vector  $\vec{a}$  on the vector  $\vec{b}$  is given by  $\frac{1}{|\vec{b}|} (\vec{a} \cdot \vec{b})$ .

$$\frac{1}{|\vec{b}|} (\vec{a} \cdot \vec{b}) = \frac{(2 \cdot 1 + 3 \cdot 2 + 2 \cdot 1)}{\sqrt{(1)^2 + (2)^2 + (1)^2}} = \frac{10}{\sqrt{6}} = \frac{5}{3} \sqrt{6}$$

**Illustration 22:** Let  $\vec{a}, \vec{b}, \vec{c}$  be the vectors of lengths 3, 4 and 5, respectively. Let  $\vec{a}$  be perpendicular to  $(\vec{b} + \vec{c})$ ,  $\vec{b}$  to  $(\vec{c} + \vec{a})$  and  $\vec{c}$  to  $(\vec{a} + \vec{b})$ . Then, find the length of the vector  $(\vec{a} + \vec{b} + \vec{c})$ . **(JEE MAIN)**

**Sol:** By using property of scalar product of vector we can solve this illustration.

$$\text{Given } |\vec{a}| = 3, |\vec{b}| = 4, |\vec{c}| = 5$$

$$\begin{aligned} \therefore |\vec{a} + \vec{b} + \vec{c}|^2 &= (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + \vec{a} \cdot (\vec{b} + \vec{c}) + \vec{b} \cdot (\vec{c} + \vec{a}) + \vec{c} \cdot (\vec{a} + \vec{b}) = 9 + 16 + 25 + 0 + 0 + 0 \\ &\Rightarrow |\vec{a} + \vec{b} + \vec{c}| = 5\sqrt{2} \end{aligned}$$

**Illustration 23:** Let  $\vec{a} = 4\hat{i} + 5\hat{j} - \hat{k}$ ,  $\vec{b} = \hat{i} - 4\hat{j} + 5\hat{k}$  and  $\vec{c} = 3\hat{i} + \hat{j} - \hat{k}$ . Find a vector  $\vec{d}$ , which is perpendicular to both  $\vec{a}$  and  $\vec{b}$ , and satisfying  $\vec{d} \cdot \vec{c} = 21$ . **(JEE MAIN)**

**Sol:** If two vector are perpendicular then their product will be zero.

Let  $\vec{d} = x\hat{i} + y\hat{j} + z\hat{k}$ . Since  $\vec{d}$  is perpendicular to both  $\vec{a}$  and  $\vec{b}$ . Therefore,

$$\vec{d} \cdot \vec{a} = 0 \Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (4\hat{i} + 5\hat{j} - \hat{k}) = 0 \Rightarrow 4x + 5y - z = 0 \quad \dots (i)$$

$$\vec{d} \cdot \vec{b} = 0 \Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} - 4\hat{j} + 5\hat{k}) = 0 \Rightarrow x - 4y + 5z = 0 \quad \dots (ii)$$

$$\vec{d} \cdot \vec{c} = 21 \Rightarrow (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (3\hat{i} + \hat{j} - \hat{k}) = 21 \Rightarrow 3x + y - z = 21 \quad \dots (iii)$$

Solving equations (i), (ii) and (iii), we get  $x = 7, y = z = -7$

Hence,  $\vec{d} = 7\hat{i} - 7\hat{j} - 7\hat{k}$

**Illustration 24:** Three vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$  satisfy the condition  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ . Evaluate the quantity

$$\mu = \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}, \text{ if } |\vec{a}| = 1, |\vec{b}| = 4 \text{ and } |\vec{c}| = 2.$$

**(JEE MAIN)**

**Sol:** Simply using property of scalar product we can calculate the value of  $\mu$ .

Since  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ , we have  $\vec{a} \cdot (\vec{a} + \vec{b} + \vec{c}) = \vec{0} \Rightarrow \vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} = 0$ . Therefore,  $\vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} = -|\vec{a}|^2 = -1$

Similarly  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} = -|\vec{b}|^2 = -16$ ,  $\vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{c} = -4$ .

On adding these equations, we have  $2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{a} \cdot \vec{c}) = -21$  or  $2\mu = -21$ , i.e.,  $\mu = \frac{-21}{2}$

**Illustration 25:** Prove, Cauchy-Schwarz inequality,  $(\vec{a} \cdot \vec{b})^2 \leq |\vec{a}|^2 |\vec{b}|^2$ , and hence show that

$$(a_1b_1 + a_2b_2 + a_3b_3)^2 \leq (a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2)$$

**(JEE ADVANCED)**

**Sol:** As we know  $\cos^2 \theta \leq 1$ , solve it by multiplying both side by  $|\vec{a}|^2 |\vec{b}|^2$ .

We have,  $\cos^2 \theta \leq 1$

$$\Rightarrow |\vec{a}|^2 |\vec{b}|^2 \cos^2 \theta \leq |\vec{a}|^2 |\vec{b}|^2 \Rightarrow (\vec{a} \cdot \vec{b})^2 \leq |\vec{a}|^2 |\vec{b}|^2$$

Let  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ . Then,

$$\vec{a} \cdot \vec{b} = a_1b_1 + a_2b_2 + a_3b_3, |\vec{a}|^2 = a_1^2 + a_2^2 + a_3^2 \text{ and } |\vec{b}|^2 = b_1^2 + b_2^2 + b_3^2.$$

$$(\vec{a} \cdot \vec{b})^2 \leq |\vec{a}|^2 |\vec{b}|^2 \Rightarrow (a_1b_1 + a_2b_2 + a_3b_3)^2 \leq (a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2)$$

**Illustration 26:** If  $\vec{a}, \vec{b}, \vec{c}$  are three mutually perpendicular vectors of equal magnitude, prove that  $\vec{a} + \vec{b} + \vec{c}$  is equally inclined with vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$ .

**(JEE ADVANCED)**

**Sol:** Here use formula of dot product to solve the problem. Let  $|\vec{a}| = |\vec{b}| = |\vec{c}| = \lambda$  (say). Since  $\vec{a}, \vec{b}, \vec{c}$  are mutually perpendicular vectors, We have  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$

$$\text{Now, } |\vec{a} + \vec{b} + \vec{c}|^2 = \vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{b} + \vec{c} \cdot \vec{c} + 2\vec{a} \cdot \vec{b} + 2\vec{b} \cdot \vec{c} + 2\vec{c} \cdot \vec{a} = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 = 3\lambda^2 \therefore |\vec{a} + \vec{b} + \vec{c}| = \sqrt{3}\lambda$$

Let  $\vec{a} + \vec{b} + \vec{c}$  makes angles  $\theta_1, \theta_2, \theta_3$  with  $\vec{a}, \vec{b}$  and  $\vec{c}$ , respectively. Then,

$$\cos \theta_1 = \frac{\vec{a} \cdot (\vec{a} + \vec{b} + \vec{c})}{|\vec{a}| |\vec{a} + \vec{b} + \vec{c}|} = \frac{\vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}}{|\vec{a}| |\vec{a} + \vec{b} + \vec{c}|} = \frac{|\vec{a}|^2}{|\vec{a}| |\vec{a} + \vec{b} + \vec{c}|} = \frac{\lambda}{\sqrt{3}\lambda} = \frac{1}{\sqrt{3}} \therefore \theta_1 = \cos^{-1} \left( \frac{1}{\sqrt{3}} \right)$$

$$\text{Similarly, } \theta_2 = \cos^{-1} \left( \frac{1}{\sqrt{3}} \right) \text{ and } \theta_3 = \cos^{-1} \left( \frac{1}{\sqrt{3}} \right) \therefore \theta_1 = \theta_2 = \theta_3$$

Hence,  $\vec{a} + \vec{b} + \vec{c}$  is equally inclined with  $\vec{a}, \vec{b}$  and  $\vec{c}$ .

**Illustration 27:** Using vectors, prove that  $\cos(A+B) = \cos A \cos B - \sin A \sin B$

**(JEE ADVANCED)**

**Sol:** From figure, using vector method we can easily prove that  $\cos(A+B) = \cos A \cos B - \sin A \sin B$ .

Let OX and OY be the coordinate axes and let  $\hat{i}$  and  $\hat{j}$  be unit vectors along OX and OY, respectively.

Let  $\angle XOP = A$  and  $\angle XOQ = B$ . Draw  $PL \perp OX$  and  $QM \perp OY$

Therefore, the angle between  $\overrightarrow{OP}$  and  $\overrightarrow{OL}$  is  $A$  &  $\overrightarrow{OQ}$  and  $\overrightarrow{OM}$  is  $B$ .

In  $\triangle OLP$ ,  $OL = OP \cos A$  and  $LP = OP \sin A$ .

Therefore,  $\overrightarrow{OL} = (OP \cos A)\hat{i}$  and  $\overrightarrow{LP} = (OP \sin A)(-\hat{j})$

Now,  $\overrightarrow{OL} + \overrightarrow{LP} = \overrightarrow{OP}$

$$\Rightarrow \overrightarrow{OP} = OP[(\cos A)\hat{i} - (\sin A)\hat{j}] \quad \dots (i)$$

In  $\triangle OMQ$ ,  $\overrightarrow{OM} = OQ \cos B$  and  $\overrightarrow{MQ} = OQ \sin B$ .

Therefore,  $\overrightarrow{OM} = (OQ \cos B)\hat{i}$ ,  $\overrightarrow{MQ} = (OQ \sin B)\hat{j}$

$$\Rightarrow \overrightarrow{OQ} = OQ[(\cos B)\hat{i} + (\sin B)\hat{j}] \quad \dots (ii)$$

From (i) and (ii), we get

$$\overrightarrow{OP} \cdot \overrightarrow{OQ} = OP[(\cos A)\hat{i} - (\sin A)\hat{j}] \cdot OQ[(\cos B)\hat{i} + (\sin B)\hat{j}] = OP \cdot OQ [\cos A \cos B - \sin A \sin B]$$

But,  $\overrightarrow{OP} \cdot \overrightarrow{OQ} = |\overrightarrow{OP}| |\overrightarrow{OQ}| \cos(A+B) = OP \cdot OQ \cos(A+B)$

$$\therefore OP \cdot OQ \cos(A+B) = OP \cdot OQ [\cos A \cos B - \sin A \sin B]$$

$$\Rightarrow \cos(A+B) = \cos A \cos B - \sin A \sin B$$




Figure 26.10

**Illustration 28:** Find the values of  $c$  for which the vectors  $\vec{a} = (c \log_2 x)\hat{i} - 6\hat{j} + 3\hat{k}$  and  $\vec{b} = (\log_2 x)\hat{i} + 2\hat{j} + (2c \log_2 x)\hat{k}$  made an obtuse angle for any  $x \in (0, \infty)$ . **(JEE ADVANCED)**

**Sol:** For obtuse angle  $\cos \theta < 0$ , therefore by using formula  $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$ , we can solve this problem.

Let  $\theta$  be the angle between the vectors  $\vec{a}$  and  $\vec{b}$ . Then,

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$$

For  $\theta$  to be an obtuse angle, we must have  $\Rightarrow \cos \theta < 0$ , for all  $x \in (0, \infty) \Rightarrow \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} < 0$ , for all  $x \in (0, \infty)$

$$\Rightarrow \vec{a} \cdot \vec{b} < 0, \text{ for all } x \in (0, \infty) \Rightarrow \vec{a} \cdot \vec{b} < 0, \text{ for all } x \in (0, \infty) \Rightarrow c(\log_2 x)^2 - 12 + 6c(\log_2 x) < 0, \text{ for all } x \in (0, \infty)$$

$$\Rightarrow cy^2 + 6cy - 12 < 0, \text{ for all } y \in \mathbb{R}, \text{ where } y = -\log_2 x \quad [\because x > 0 \Rightarrow y = \log_2 x \in \mathbb{R}]$$

$$\Rightarrow c < 0 \text{ and } 36c^2 + 48c < 0 \quad [\because ax^2 + bx^2 + c > 0 \text{ for all } x \Rightarrow a < 0 \text{ and Discriminant} < 0]$$

$$\Rightarrow c < 0 \text{ and } c(3c + 4) < 0$$

$$\Rightarrow c < 0 \text{ and } -\frac{4}{3} < c < 0 \Rightarrow c \in \left(-\frac{4}{3}, 0\right)$$

**Illustration 29:** D is the midpoint of the side  $\overline{BC}$  of a triangle ABC, show that  $AB^2 + AC^2 = 2(AD^2 + BD^2)$  **(JEE MAIN)**

**Sol:** By using the formula of resultant vector we will get the required result.

Given D is midpoint of BC  $\Rightarrow \overrightarrow{BD} = \overrightarrow{DC}$

$$\text{We have } \overrightarrow{AB} = \overrightarrow{AD} + \overrightarrow{DB} \Rightarrow AB^2 = (\overrightarrow{AD} + \overrightarrow{DB})^2$$

$$AB^2 = AD^2 + DB^2 + 2\overrightarrow{AD} \cdot \overrightarrow{DB} \quad \dots (i)$$




Figure 26.11

Also we have  $\overline{AC} = \overline{AD} + \overline{DC} \Rightarrow AC^2 = (\overline{AD} + \overline{DC})^2$

$$AC^2 = AD^2 + DC^2 + 2\overline{AD} \cdot \overline{DC} \quad \dots(ii)$$

Adding equations (i) and (ii), we get

$$AB^2 + AC^2 = 2AD^2 + 2BD^2 + 2\overline{AD} \cdot (\overline{DB} + \overline{DC}) = 2(DA^2 + DB^2)$$

## 9. VECTOR OR CROSS PRODUCT

Let  $\vec{a}$  and  $\vec{b}$  be two vectors. The vector product of these two vectors can be calculated as  $(\vec{a} \times \vec{b}) = |\vec{a}||\vec{b}|\sin\theta\hat{n}$ , where  $\theta$  is the angle between the vectors  $\vec{a}$  and  $\vec{b}$ , ( $0 \leq \theta \leq \pi$ ) and  $\hat{n}$  is the unit vector at right angles to both  $\vec{a}$  and  $\vec{b}$ , i.e.  $\hat{n}$  is vector normal to the plane that contains  $\vec{a}$  and  $\vec{b}$ .  $\vec{a}$ ,  $\vec{b}$  and  $\hat{n}$  are three vectors which form a right-handed set.

The convention is that we choose the direction specified by the right-hand screw rule. Imagine a screw in your right hand. If you turn a right-handed screw from  $\vec{a}$  to  $\vec{b}$ , the screw advances along the unit vector  $\hat{n}$ . It is very important to realize that the result of a vector product is itself a vector.




Figure 26.12

Let us see how the order of multiplication matters from the definition of the right-hand screw rule:

The vector given by  $(\vec{b} \times \vec{a})$  points in the opposite direction to  $(\vec{a} \times \vec{b})$ . So,  $(\vec{a} \times \vec{b}) = -(\vec{b} \times \vec{a})$ .

We can define vector product in terms of matrix notation as  $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$

and in terms of components as  $\vec{a} = \langle a_1, a_2, a_3 \rangle, \vec{b} = \langle b_1, b_2, b_3 \rangle \Rightarrow \vec{a} \times \vec{b} = \langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1 \rangle$

From the definition, the angle can be calculated as  $\sin\theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}||\vec{b}|}$

If  $\vec{a}$  and  $\vec{b}$  are parallel then  $\theta = 0^\circ \Rightarrow \sin\theta = 0$  and  $\vec{a} \times \vec{b} = 0$

### 9.1 Properties of Vector Product

The properties of vector product are listed as follows:

$\vec{a}, \vec{b}$  and  $\vec{a} \times \vec{b}$  are all vectors in three dimensions.

(a)  $\vec{a} \times \vec{b} \perp \vec{a}$  and  $\vec{b}$

- (b)  $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$
- (c)  $\hat{i} \times \hat{j} = \hat{k}, \hat{j} \times \hat{k} = \hat{i}, \hat{k} \times \hat{i} = \hat{j}$
- (d)  $\vec{a} \times \vec{b} = \vec{0} \Leftrightarrow \vec{a} = \vec{0} \text{ or } \vec{b} = \vec{0} \text{ or } \vec{a} \parallel \vec{b}$
- (e)  $\vec{a} \times \vec{b} = -\vec{b} \times \vec{a}$
- (f)  $(c\vec{a}) \times \vec{b} = \vec{a} \times (c\vec{b}) = c(\vec{a} \times \vec{b})$
- (g)  $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$
- (h)  $\vec{a} \cdot (\vec{b} \times \vec{c}) = (\vec{a} \times \vec{b}) \cdot \vec{c}$

**Geometrical interpretation** of  $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$ , denotes the area of parallelogram, in which  $\vec{a}$  and  $\vec{b}$  are the two adjacent sides.

### Vector area of the plane figure

Considering the boundaries of closed, bounded surface, which has been described in a specific manner and that do not cross, it is possible to associate a directed line segment  $\vec{c}$ , such that




Figure 26.13

- (a)  $|\vec{c}|$  is the number of units of area enclosed by the plane figure.
- (b) The support of  $\vec{c}$  is perpendicular to the area and outside the surface.
- (c) The sense of description of the boundaries and the direction of  $\vec{c}$  is in accordance with the R.H.S. screw rule.

### Vector area of a triangle

If  $\vec{a}$  and  $\vec{b}$  are the position vectors, then the vector area of a triangle is given by the formula

$$\vec{\Delta} = \frac{1}{2}(\vec{a} \times \vec{b})$$

If  $\vec{a}, \vec{b}, \vec{c}$  are the position vectors, then the vector area of  $\triangle ABC$  is given by the formula

$$\vec{\Delta} = \frac{1}{2}[(\vec{c} - \vec{b}) \times (\vec{a} - \vec{b})]$$

$$\vec{\Delta} = \frac{1}{2}[(\vec{a} \times \vec{b}) + (\vec{b} \times \vec{c}) + (\vec{c} \times \vec{a})]$$




Figure 26.14




Figure 26.15



## PLANCESS CONCEPTS

- (i) If three points with position vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are collinear, then  $\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} = 0$
- (ii) Unit vector perpendicular to the plane of the  $\triangle ABC$ , when  $\vec{a}, \vec{b}$  and  $\vec{c}$  are the p.v. of its angular point is

$$\hat{n} = \pm \frac{\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}}{2\Delta}$$

Nitish Jhawar (JEE 2009, AIR 7)

## 10. ANGULAR BISECTOR

As discussed earlier, the diagonal of a parallelogram is not necessarily the bisector of the angle formed by two adjacent sides. However, the diagonal of a rhombus bisects the angle formed between two adjacent sides. Consider vectors  $\overrightarrow{AB} = \vec{a}$  and  $\overrightarrow{AD} = \vec{b}$  forming a parallelogram ABCD as shown in the figure.

Consider the two unit vectors along the given vectors, forming a rhombus AB'C'D'.




Figure 26.16

Now,  $\overrightarrow{AB} = \frac{\vec{a}}{|\vec{a}|}$  and  $\overrightarrow{AD} = \frac{\vec{b}}{|\vec{b}|}$ . Therefore  $\overrightarrow{AC'} = \frac{\vec{a}}{|\vec{a}|} + \frac{\vec{b}}{|\vec{b}|}$ . Then, any vector along the internal bisector is  $\lambda \left( \frac{\vec{a}}{|\vec{a}|} + \frac{\vec{b}}{|\vec{b}|} \right)$ .

Similarly, any vector along the external bisector is  $\lambda \left( \frac{\vec{a}}{|\vec{a}|} - \frac{\vec{b}}{|\vec{b}|} \right)$ .

**Illustration 30:** Find a vector of magnitude 9, which is perpendicular to both the vectors  $4\hat{i} - \hat{j} + 3\hat{k}$  and  $-2\hat{i} + \hat{j} - 2\hat{k}$ .  
(JEE MAIN)

**Sol:** By using property of vector product, we can solve this problem. Let  $\vec{a} = 4\hat{i} - \hat{j} + 3\hat{k}$  and  $\vec{b} = -2\hat{i} + \hat{j} - 2\hat{k}$ . Then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & -1 & 3 \\ -2 & 1 & -2 \end{vmatrix} = (2-3)\hat{i} - (-8+6)\hat{j} + (4-2)\hat{k} = -\hat{i} + 2\hat{j} + 2\hat{k} \Rightarrow |\vec{a} \times \vec{b}| = \sqrt{(-1)^2 + 2^2 + 2^2} = 3$$

$$\therefore \text{Required vector} = 9 \left\{ \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} \right\} = \frac{9}{3} (-\hat{i} + 2\hat{j} + 2\hat{k}) = -3\hat{i} + 6\hat{j} + 6\hat{k}$$

**Illustration 31:** Find the area of a parallelogram, whose adjacent sides are given by the vectors  $\vec{a} = 3\hat{i} + \hat{j} + 4\hat{k}$  and  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$ .  
(JEE MAIN)

**Sol:** The area of a parallelogram with  $\vec{a}$  and  $\vec{b}$  as its adjacent sides is given by  $|\vec{a} \times \vec{b}|$ .

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & 4 \\ 1 & -1 & 1 \end{vmatrix} = 5\hat{i} + \hat{j} - 4\hat{k}.$$

Therefore,  $|\vec{a} \times \vec{b}| = \sqrt{25 + 1 + 16} = \sqrt{42}$ ; Hence, the required area is  $\sqrt{42}$ .

**Illustration 32:** Let  $\vec{a}, \vec{b}, \vec{c}$  be the unit vectors such that  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c} = 0$  and the angle between  $\vec{b}$  and  $\vec{c}$  is  $\frac{\pi}{6}$ ,  
Prove that  $\vec{a} = \pm 2(\vec{b} \times \vec{c})$ . (JEE MAIN)

**Sol:** Here  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c} = 0$ , therefore  $\vec{a}$  is perpendicular to the plane of  $\vec{b}$  and  $\vec{c}$  and it is parallel to  $\vec{b} \times \vec{c}$ .

We have  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c} = 0$

$\Rightarrow \vec{a} \perp \vec{b}$  and  $\vec{a} \perp \vec{c} \Rightarrow \vec{a}$  is perpendicular to the plane of  $\vec{b}$  and  $\vec{c}$ .

$\Rightarrow \vec{a}$  is parallel to  $\vec{b} \times \vec{c} \Rightarrow \vec{a} = \lambda(\vec{b} \times \vec{c})$  for some scalar  $\lambda$ .

$$\Rightarrow |\vec{a}| = |\lambda| |\vec{b}| |\vec{c}| \sin \frac{\pi}{6} \Rightarrow 1 = \frac{|\lambda|}{2} \quad [\because |\vec{a}| = |\vec{b}| = |\vec{c}|]$$

$$\Rightarrow |\lambda| = 2 \quad \Rightarrow \lambda = \pm 2$$

$$\therefore \vec{a} = \lambda(\vec{b} \times \vec{c}) \Rightarrow \vec{a} = \pm 2(\vec{b} \times \vec{c}).$$

**Illustration 33:** If  $\vec{a}, \vec{b}, \vec{c}$  are three non-zero vectors, such that  $\vec{a} \times \vec{b} = \vec{c}$  and  $\vec{b} \times \vec{c} = \vec{a}$ , prove that  $\vec{a}, \vec{b}, \vec{c}$  are mutually at right angles and  $|\vec{b}| = 1$  and  $|\vec{c}| = |\vec{a}|$ . (JEE MAIN)

**Sol:** Use property of vector or cross product to prove this illustration.

We have,  $\vec{a} \times \vec{b} = \vec{c}$  and  $\vec{b} \times \vec{c} = \vec{a}$

$$\Rightarrow \vec{c} \perp \vec{a}, \vec{c} \perp \vec{b} \text{ and } \vec{a} \perp \vec{b}, \vec{a} \perp \vec{c} \Rightarrow \vec{a} \perp \vec{b}, \vec{b} \perp \vec{c} \text{ and } \vec{c} \perp \vec{a}.$$

$\Rightarrow \vec{a}, \vec{b}, \vec{c}$  are mutually perpendicular lines.

$$\text{Again } \vec{a} \times \vec{b} = \vec{c} \text{ and } \vec{b} \times \vec{c} = \vec{a} \Rightarrow |\vec{a} \times \vec{b}| = |\vec{c}| \text{ and } |\vec{b} \times \vec{c}| = |\vec{a}|$$

$$\Rightarrow |\vec{a}| |\vec{b}| \sin \frac{\pi}{2} = |\vec{c}| \text{ and } |\vec{b}| |\vec{c}| \sin \frac{\pi}{2} = |\vec{a}| \quad [\because \vec{a} \perp \vec{b} \text{ and } \vec{b} \perp \vec{c}]$$

$$\Rightarrow |\vec{a}| |\vec{b}| = |\vec{c}| \text{ and } |\vec{b}| |\vec{c}| = |\vec{a}|$$

$$\Rightarrow |\vec{b}|^2 |\vec{c}| = |\vec{c}| \quad \left[ \text{Putting } |\vec{a}| = |\vec{b}| |\vec{c}| \text{ in } |\vec{a}| |\vec{b}| = |\vec{c}| \right]$$

$$\Rightarrow |\vec{b}|^2 = 1 \quad [\because |\vec{c}| \neq 0]$$

$$\Rightarrow |\vec{b}| = 1$$

Putting  $|\vec{b}| = 1$  in  $|\vec{a}| |\vec{b}| = |\vec{c}|$ , we obtain  $|\vec{a}| = |\vec{c}|$ .

**Illustration 34:** Prove by vector method, that in a  $\triangle ABC$ ,  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$  (JEE MAIN)

**Sol:** As area of triangle ABC is equal to  $\frac{1}{2} |\vec{AB} \times \vec{AC}| = \frac{1}{2} |\vec{BC} \times \vec{BA}| = \frac{1}{2} |\vec{CA} \times \vec{CB}|$ , therefore by using cross product method we can prove this problem.

Let  $\vec{BC} = \vec{a}$ ,  $\vec{CA} = \vec{b}$ ,  $\vec{AB} = \vec{c}$ . Then

$$\text{The area of } \triangle ABC = \frac{1}{2} |\vec{AB} \times \vec{AC}| = \frac{1}{2} |\vec{BC} \times \vec{BA}| = \frac{1}{2} |\vec{CA} \times \vec{CB}| \Rightarrow bc \sin A = ca \sin B = ab \sin C$$

Dividing the above expression by  $abc$ , we get  $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$

**Illustration 35:** Given the vectors  $\vec{a} = \hat{p} + 2\hat{q}$  and  $\vec{b} = 2\hat{p} + \hat{q}$ , where  $\hat{p}$  and  $\hat{q}$  are unit vectors forming an angle of  $30^\circ$ . Find the area of the parallelogram constructed on these vectors. **(JEE MAIN)**

**Sol:** Simply by applying cross product between  $\vec{a}$  and  $\vec{b}$ , we have  $\vec{a} \times \vec{b} = (\hat{p} + 2\hat{q}) \times (2\hat{p} + \hat{q}) = -3(\hat{p} \times \hat{q})$ .

$$\Rightarrow |\vec{a} \times \vec{b}| = 3|\hat{p} \times \hat{q}| = 3|\hat{p}||\hat{q}|\sin\frac{\pi}{6} = \frac{3}{2}$$

**Illustration 36:** Let  $\vec{OA} = \vec{a}, \vec{OB} = 10\vec{a} + 2\vec{b}$  and  $\vec{OC} = \vec{b}$ , where  $O$  is the origin. Let  $p$  denote the area of the quadrilateral  $OABC$  and  $q$  denote the area of the parallelogram with  $\vec{OA}$  and  $\vec{OC}$  as adjacent sides. Prove that  $p = 6q$ . **(JEE MAIN)**

**Sol:** We have to obtain the area of quadrilateral and parallelogram using cross product method to get the required result.

We have,  $p$  = area of the quadrilateral  $OABC$

$$\begin{aligned} &= \frac{1}{2} |\vec{OB} \times \vec{AC}| = \frac{1}{2} |\vec{OB} \times (\vec{OC} - \vec{OA})| = \frac{1}{2} |(10\vec{a} + 2\vec{b}) \times (\vec{b} - \vec{a})| \\ &= \frac{1}{2} |10(\vec{a} \times \vec{b}) - 10(\vec{a} \times \vec{a}) + 2(\vec{b} \times \vec{b}) - 2(\vec{b} \times \vec{a})| \\ &= \frac{1}{2} |10(\vec{a} \times \vec{b}) - 0 + 0 + 2(\vec{a} \times \vec{b})| = 6(\vec{a} \times \vec{b}) \end{aligned} \quad \dots (i)$$

and  $q$  = area of the parallelogram with adjacent sides  $\vec{OA}$  and  $\vec{OC}$

$$= |\vec{OA} \times \vec{OC}| = (\vec{a} \times \vec{b}) \quad \dots (ii)$$

From equations (i) and (ii), we get  $p = 6q$ .

**Illustration 37:** Given that  $D, E, F$  are the midpoints of the sides of a triangle  $ABC$ , using the vector method, prove that area of  $\triangle DEF = \frac{1}{4}$  (area of  $\triangle ABC$ ) **(JEE MAIN)**

**Sol:** Taking  $A$  as the origin, let the position vectors of  $B$  and  $C$  be  $\vec{b}$  and  $\vec{c}$  respectively.

Then, the position vector of  $D, E$  and  $F$  are  $\frac{1}{2}(\vec{b} + \vec{c})$ ,  $\frac{1}{2}\vec{c}$  and  $\frac{1}{2}\vec{b}$  respectively. Therefore first obtain  $\vec{DE}$  and  $\vec{DF}$ ,

and after that by applying formula of vector area of triangle  $DEF$  we can obtain the required result.

$$\text{Now, } \vec{DE} = \frac{1}{2}\vec{c} - \frac{1}{2}(\vec{b} + \vec{c}) = \frac{-\vec{b}}{2}$$

$$\text{and } \vec{DF} = \frac{1}{2}\vec{b} - \frac{1}{2}(\vec{b} + \vec{c}) = \frac{-\vec{c}}{2}$$

$$\begin{aligned} \therefore \text{Vector area of } \triangle DEF &= \frac{1}{2} (\vec{DE} \times \vec{DF}) = \left( \frac{-\vec{b}}{2} \times \frac{-\vec{c}}{2} \right) \\ &= \frac{1}{8} (\vec{b} \times \vec{c}) = \frac{1}{4} \left\{ \frac{1}{2} (\vec{AB} \times \vec{AC}) \right\} = \frac{1}{4} (\text{vector area of } \triangle ABC) \end{aligned}$$

$$\text{Hence, area of } \triangle DEF = \frac{1}{4} (\text{area of } \triangle ABC)$$




Figure 26.17

**Illustration 38:** Given that P, Q are the midpoints of the non-parallel sides BC and AD of a trapezium ABCD. Show that area of  $\triangle APD = \triangle CQB$ . **(JEE MAIN)**

**Sol:** Use formula of vector area of triangle to solve this problem. Let  $\overrightarrow{AB} = \vec{b}$  and  $\overrightarrow{AD} = \vec{d}$  Now DC is parallel to  $\overrightarrow{AB} \Rightarrow$  there exists a scalar t, such that.  $\overrightarrow{DC} = t\overrightarrow{AB} = t\vec{b}$

$$\therefore \overrightarrow{AC} = \overrightarrow{AD} + \overrightarrow{DC} = \vec{d} + t\vec{b}$$

From geometry we know that  $QP = \frac{AB + DC}{2}$

Now  $\overrightarrow{AP}$  and  $\overrightarrow{AQ}$  are  $\frac{\vec{b} + \vec{d} + t\vec{b}}{2}$  and  $\frac{\vec{d}}{2}$ , respectively.

$$\text{Now, } 2\Delta APD = \overrightarrow{AP} \times \overrightarrow{AD} = \frac{1}{2}(\vec{b} + \vec{d} + t\vec{b}) \times \vec{d} = \frac{1}{2}(1+t)(\vec{b} \times \vec{d})$$

$$\text{Also } 2\Delta CQB = \overrightarrow{BC} \times \overrightarrow{BQ} = [-\vec{b} + \vec{d} + t\vec{b}] \times \left[-\vec{b} + \frac{\vec{d}}{2}\right]$$

$$= -(\vec{d} \times \vec{b}) - \frac{\vec{b} \times \vec{d}}{2} + \frac{t\vec{b} \times \vec{d}}{2} = \frac{1}{2}(1+t)\vec{b} \times \vec{d} = 2\Delta APD \Rightarrow \Delta APD = \Delta CQB$$




Figure 26.18

## 11. TRIPLE PRODUCT OF VECTORS

Two types of triple products are listed below:

Vector triple product  $\Rightarrow (\vec{a} \times \vec{b}) \times \vec{c}$

Scalar triple product  $\Rightarrow (\vec{a} \times \vec{b}) \cdot \vec{c}$

### 11.1 Scalar Triple Product

The scalar triple product has an interesting geometric interpretation:




Figure 26.19

We know that  $(\vec{a} \times \vec{b}) = |\vec{a}||\vec{b}|\sin\theta\hat{n}$  (area of the parallelogram defined by  $\vec{a}$  and  $\vec{b}$ )

Thus,  $(\vec{a} \times \vec{b}) \cdot \vec{c} = (\text{area of the parallelogram})\hat{n} \cdot \vec{c} = (\text{area of the parallelogram})|\hat{n}||\vec{c}|\cos\phi$

But  $|\vec{c}|\cos\phi = h = \text{height of the parallelepiped normal to the plane containing } \vec{a} \text{ and } \vec{b}$ . ( $\phi$  is the angle between  $\vec{c}$  and  $\hat{n}$ ).

So,  $(\vec{a} \times \vec{b}) \cdot \vec{c} = \text{volume of the parallelepiped defined by } \vec{a}, \vec{b} \text{ and } \vec{c}$ . Thus, the following conclusions are arrived:

(a) If any two vectors are parallel, then  $(\vec{a} \times \vec{b}) \cdot \vec{c} = 0$  (zero volume)

(b) If the three vectors are co-planar, then  $(\vec{a} \times \vec{b}) \cdot \vec{c} = 0$  (zero volume)

(c) If  $(\vec{a} \times \vec{b}) \cdot \vec{c} = 0$ , then either

(i)  $\vec{a} = 0$ , or (ii)  $\vec{b} = 0$  or (iii)  $\vec{c} = 0$  or

(iv) two of the vectors are parallel or (v) the three vectors are co-planar

(d)  $(\vec{a} \times \vec{b}) \cdot \vec{c} = \vec{a} \cdot (\vec{b} \times \vec{c}) = \vec{b} \cdot (\vec{c} \times \vec{a}) = \text{The same volume.}$

(e)  $(\vec{a} \times \vec{b}) \cdot \vec{c}$  is also known as box product, which is represented as  $[\vec{a}\vec{b}\vec{c}]$ .

$$\text{Also } [\vec{a} + \vec{b} \quad \vec{c} \quad \vec{d}] = [\vec{a} \quad \vec{c} \quad \vec{d}] + [\vec{b} \quad \vec{c} \quad \vec{d}]$$

- (f) If  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar, then  $[\vec{a}, \vec{b}, \vec{c}] > 0$ , for right-handed system and  $[\vec{a}, \vec{b}, \vec{c}] < 0$ , for left handed system.
- (g) If O is the origin and  $\vec{a}, \vec{b}, \vec{c}$  are the position vectors of A, B and C, respectively, of the tetrahedron OABC, then the volume is given by the formula  $V = \frac{1}{6}[\vec{a} \ \vec{b} \ \vec{c}]$ .

### Reciprocal system of vectors

- (a) If  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{a}', \vec{b}', \vec{c}'$  are the two sets of non-coplanar vectors, such that  $\vec{a} \cdot \vec{a}' = \vec{b} \cdot \vec{b}' = \vec{c} \cdot \vec{c}' = 1$ ,  $\vec{a} \cdot \vec{b}' = \vec{a} \cdot \vec{c}' = 0$ ,  $\vec{b} \cdot \vec{a}' = \vec{b} \cdot \vec{c}' = 0$  and  $\vec{c} \cdot \vec{a}' = \vec{c} \cdot \vec{b}' = 0$ ,  
Then  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{a}', \vec{b}', \vec{c}'$  constitute a reciprocal system of vectors.
- (b) Reciprocal system of vectors exists only in the case of dot product.
- (c)  $\vec{a}', \vec{b}', \vec{c}'$  can be defined in terms of  $\vec{a}, \vec{b}, \vec{c}$  as

$$\vec{a}' = \frac{\vec{b} \times \vec{c}}{[\vec{a} \ \vec{b} \ \vec{c}]}; \vec{b}' = \frac{\vec{c} \times \vec{a}}{[\vec{a} \ \vec{b} \ \vec{c}]}; \vec{c}' = \frac{\vec{a} \times \vec{b}}{[\vec{a} \ \vec{b} \ \vec{c}]} \quad ([\vec{a} \ \vec{b} \ \vec{c}] \neq 0)$$

### Note:

- (i)  $\vec{a} \times \vec{a}' + \vec{b} \times \vec{b}' + \vec{c} \times \vec{c}' = 0 \Rightarrow \vec{a} \times (\vec{b} \times \vec{c}) + \vec{b} \times (\vec{c} \times \vec{a}) + \vec{c} \times (\vec{a} \times \vec{b}) = 0$
- (ii)  $\vec{a} \cdot \vec{a}' = \vec{b} \cdot \vec{b}' = \vec{c} \cdot \vec{c}' = 1$
- (iii)  $(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a}' + \vec{b}' + \vec{c}') = 3$
- (iv) If  $[\vec{a} \ \vec{b} \ \vec{c}] = V$  then  $[\vec{a}' \ \vec{b}' \ \vec{c}'] = \frac{1}{V} \Rightarrow [\vec{a} \ \vec{b} \ \vec{c}][\vec{a}' \ \vec{b}' \ \vec{c}'] = 1$

**Illustration 39:** If  $\vec{\ell}, \vec{m}, \vec{n}$  three non-coplanar vectors, then prove that

$$[\vec{\ell} \ \vec{m} \ \vec{n}](\vec{a} \times \vec{b}) = \begin{vmatrix} \vec{\ell} \cdot \vec{a} & \vec{\ell} \cdot \vec{b} & \vec{\ell} \cdot \vec{c} \\ \vec{m} \cdot \vec{a} & \vec{m} \cdot \vec{b} & \vec{m} \cdot \vec{c} \\ \vec{n} \cdot \vec{a} & \vec{n} \cdot \vec{b} & \vec{n} \cdot \vec{c} \end{vmatrix}.$$

(JEE ADVANCED)

**Sol:** Use scalar triple product method as mentioned above to solve this problem.

$$\text{Let, } \vec{\ell} = \ell_1 \hat{i} + \ell_2 \hat{j} + \ell_3 \hat{k}, \quad \vec{m} = m_1 \hat{i} + m_2 \hat{j} + m_3 \hat{k}, \quad \vec{n} = n_1 \hat{i} + n_2 \hat{j} + n_3 \hat{k},$$

$$\text{and } \vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}, \quad \vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$$

$$\text{Now } [\vec{\ell} \ \vec{m} \ \vec{n}] = \begin{vmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \\ n_1 & n_2 & n_3 \end{vmatrix} \text{ and } (\vec{a} \times \vec{b}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

$$[\vec{\ell} \ \vec{m} \ \vec{n}](\vec{a} \times \vec{b}) = \begin{vmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \\ n_1 & n_2 & n_3 \end{vmatrix} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \begin{vmatrix} \ell_1 \hat{i} + \ell_2 \hat{j} + \ell_3 \hat{k} & \sum \ell_1 a_2 & \sum \ell_1 b_1 \\ m_1 \hat{i} + m_2 \hat{j} + m_3 \hat{k} & \sum m_1 a_1 & \sum m_1 b_1 \\ n_1 \hat{i} + n_2 \hat{j} + n_3 \hat{k} & \sum n_1 a_1 & \sum n_1 b_1 \end{vmatrix}$$

$$\text{Now, } \vec{\ell} \cdot \vec{a} = (\ell_1 \hat{i} + \ell_2 \hat{j} + \ell_3 \hat{k}) \cdot (a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}) = \sum \ell_1 a_2 \text{ etc.}$$

$$\therefore [\vec{\ell} \quad \vec{m} \quad \vec{n}](\vec{a} \times \vec{b}) = \begin{vmatrix} \vec{\ell} & \vec{\ell} \cdot \vec{a} & \vec{\ell} \cdot \vec{b} \\ \vec{m} & \vec{m} \cdot \vec{a} & \vec{m} \cdot \vec{b} \\ \vec{n} & \vec{n} \cdot \vec{a} & \vec{n} \cdot \vec{b} \end{vmatrix} = \begin{vmatrix} \vec{\ell} \cdot \vec{a} & \vec{\ell} \cdot \vec{b} & \vec{\ell} \\ \vec{m} \cdot \vec{a} & \vec{m} \cdot \vec{b} & \vec{m} \\ \vec{n} \cdot \vec{a} & \vec{n} \cdot \vec{b} & \vec{n} \end{vmatrix},$$

Hence proved.

**Illustration 40:** Find the volume of a parallelepiped, whose sides are given by  $-3\hat{i} + 7\hat{j} + 5\hat{k}$ ,  $-5\hat{i} + 7\hat{j} - 3\hat{k}$  and  $7\hat{i} - 5\hat{j} - 3\hat{k}$

(JEE MAIN)

**Sol:** We know that, the volume of a parallelepiped, whose three adjacent edges are  $\vec{a}, \vec{b}, \vec{c}$  is  $[\vec{a} \quad \vec{b} \quad \vec{c}]$ .

Let  $\vec{a} = -3\hat{i} + 7\hat{j} + 5\hat{k}$ ,  $\vec{b} = -5\hat{i} + 7\hat{j} - 3\hat{k}$  and  $\vec{c} = 7\hat{i} - 5\hat{j} - 3\hat{k}$

We know that, the volume of a parallelepiped, whose three adjacent edges are  $\vec{a}, \vec{b}, \vec{c}$  is  $[\vec{a} \quad \vec{b} \quad \vec{c}]$

$$\text{Now, } [\vec{a} \quad \vec{b} \quad \vec{c}] = \begin{vmatrix} -3 & 7 & 5 \\ -5 & 7 & -3 \\ 7 & -5 & -3 \end{vmatrix} = -3(-21 - 15) - 7(15 + 21) + 5(25 - 49) = 108 - 252 - 120 = -264$$

So, the required volume of the parallelepiped  $= [\vec{a} \quad \vec{b} \quad \vec{c}] = |-264| = 264$  cubic units.

**Illustration 41:** Simplify  $[\vec{a} - \vec{b} \quad \vec{b} - \vec{c} \quad \vec{c} - \vec{a}]$

(JEE ADVANCED)

**Sol:** Here by using scalar triple product we can simplify this.

$$\begin{aligned} [\vec{a} - \vec{b} \quad \vec{b} - \vec{c} \quad \vec{c} - \vec{a}] &= \{(\vec{a} - \vec{b}) \times (\vec{b} - \vec{c})\} \cdot (\vec{c} - \vec{a}) \quad [\text{by def.}] \\ &= (\vec{a} \times \vec{b} - \vec{a} \times \vec{c} - \vec{b} \times \vec{b} + \vec{b} \times \vec{c}) \cdot (\vec{c} - \vec{a}) \quad [\text{by dist. law}] \\ &= (\vec{a} \times \vec{b} + \vec{c} \times \vec{a} + \vec{b} \times \vec{c}) \cdot (\vec{c} - \vec{a}) \quad [\because \vec{b} \times \vec{b} = 0] \\ &= (\vec{a} \times \vec{b}) \cdot \vec{c} - (\vec{a} \times \vec{b}) \cdot \vec{a} + (\vec{c} \times \vec{a}) \cdot \vec{c} - (\vec{c} \times \vec{a}) \cdot \vec{a} + (\vec{b} \times \vec{c}) \cdot \vec{c} - (\vec{b} \times \vec{c}) \cdot \vec{a} \quad [\text{by dist. law}] \\ &= [\vec{a} \quad \vec{b} \quad \vec{c}] - 0 + 0 - 0 + 0 - [\vec{b} \quad \vec{c} \quad \vec{a}] \\ &= [\vec{a} \quad \vec{b} \quad \vec{c}] - [\vec{b} \quad \vec{c} \quad \vec{a}] = [\vec{a} \quad \vec{b} \quad \vec{c}] - [\vec{a} \quad \vec{b} \quad \vec{c}] = 0 \quad [\because [\vec{b} \quad \vec{c} \quad \vec{a}] = [\vec{a} \quad \vec{b} \quad \vec{c}]] \end{aligned}$$

**Illustration 42:** Find the volume of the tetrahedron, whose four vertices have position vectors  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{d}$ , respectively.

(JEE MAIN)  
... (i)

**Sol:** Here volume of tetrahedron is equal to  $\frac{1}{6}[\vec{a} - \vec{d} \quad \vec{b} - \vec{d} \quad \vec{c} - \vec{d}]$ .

Let, four vertices be A, B, C, D with p.v.  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{d}$  respectively.

$$\therefore \overrightarrow{DA} = (\vec{a} - \vec{d}), \overrightarrow{DB} = (\vec{b} - \vec{d}), \overrightarrow{DC} = (\vec{c} - \vec{d})$$

$$\begin{aligned} \text{Hence volume} &= \frac{1}{6}[\vec{a} - \vec{d} \quad \vec{b} - \vec{d} \quad \vec{c} - \vec{d}] = \frac{1}{6}(\vec{a} - \vec{d}) \cdot [(\vec{b} - \vec{d}) \times (\vec{c} - \vec{d})] \\ &= \frac{1}{6}(\vec{a} - \vec{d}) \cdot [\vec{b} \times \vec{c} - \vec{b} \times \vec{d} + \vec{c} \times \vec{d}] = \frac{1}{6}\{[\vec{a} \quad \vec{b} \quad \vec{c}] - [\vec{a} \quad \vec{b} \quad \vec{d}] + [\vec{a} \quad \vec{c} \quad \vec{d}] - [\vec{d} \quad \vec{b} \quad \vec{c}]\} \\ &= \frac{1}{6}\{[\vec{a} \quad \vec{b} \quad \vec{c}] - [\vec{d} \quad \vec{b} \quad \vec{c}] - [\vec{a} \quad \vec{d} \quad \vec{c}] - [\vec{a} \quad \vec{b} \quad \vec{d}]\}. \end{aligned}$$

**Illustration 43:** Let  $\vec{u}$  and  $\vec{v}$  be unit vectors and  $\vec{w}$  is a vector, such that  $\vec{u} \times \vec{v} + \vec{u} = \vec{w}$  and  $\vec{w} \times \vec{u} = \vec{v}$ , then find the value of  $[\vec{u} \ \vec{v} \ \vec{w}]$ . **(JEE ADVANCED)**

**Sol:** Here as given  $\vec{u} \times \vec{v} + \vec{u} = \vec{w}$  and  $\vec{w} \times \vec{u} = \vec{v}$ , solve it using scalar triple product.

Given,  $\vec{u} \times \vec{v} + \vec{u} = \vec{w}$  and  $\vec{w} \times \vec{u} = \vec{v}$

$$\Rightarrow (\vec{u} \times \vec{v} + \vec{u}) \times \vec{u} = \vec{w} \times \vec{u}$$

$$\Rightarrow (\vec{u} \times \vec{v}) \times \vec{u} + \vec{u} \times \vec{u} = \vec{v} \quad (\text{as, } \vec{w} \times \vec{u} = \vec{v})$$

$$\Rightarrow (\vec{u} \cdot \vec{u})\vec{v} - (\vec{v} \cdot \vec{u})\vec{u} + \vec{u} \times \vec{u} = \vec{v} \quad (\text{using } \vec{u} \cdot \vec{u} = 1 \text{ and } \vec{u} \times \vec{u} = 0, \text{ since unit vector})$$

$$\Rightarrow \vec{v} - (\vec{v} \cdot \vec{u})\vec{u} = \vec{v} \Rightarrow (\vec{u} \cdot \vec{v})\vec{u} = \vec{0}$$

$$\Rightarrow \vec{u} \cdot \vec{v} = 0 \quad (\text{as; } \vec{u} \neq \vec{0}) \quad \dots\dots(i)$$

$$\therefore [\vec{u} \ \vec{v} \ \vec{w}] = \vec{u} \cdot (\vec{v} \times \vec{w})$$

$$= \vec{u} \cdot (\vec{v} \times (\vec{u} \times \vec{v} + \vec{u})) \quad (\text{given } \vec{w} = \vec{u} \times \vec{v} + \vec{u})$$

$$= \vec{u} \cdot (\vec{v} \times (\vec{u} \times \vec{v}) + \vec{v} \times \vec{u}) = \vec{u} \cdot ((\vec{v} \cdot \vec{v})\vec{u} - (\vec{v} \cdot \vec{u})\vec{v} + \vec{v} \times \vec{u})$$

$$= \vec{u} \cdot (\vec{u} - 0 + \vec{v} \times \vec{u}) \quad (\text{as } \vec{u} \cdot \vec{v} = 0 \text{ from (i)})$$

$$= (\vec{u} \cdot \vec{u}) - \vec{u} \cdot (\vec{v} \times \vec{u}) = 1 - 0 = 1$$

$$\therefore [\vec{u} \ \vec{v} \ \vec{w}] = 1$$

## 11.2 Vector Triple Product

**Definition:**  $(\vec{a} \times \vec{b}) \times \vec{c}$  is a vector, which is coplanar to  $\vec{a}$  and  $\vec{b}$  and perpendicular to  $\vec{c}$ .

$$\text{Hence } (\vec{a} \times \vec{b}) \times \vec{c} = x\vec{a} + y\vec{b} \quad [\text{Linear Combination of } \vec{a} \text{ and } \vec{b}] \quad \dots (i)$$

$$\vec{c} \cdot (\vec{a} \times \vec{b}) \times \vec{c} = x(\vec{a} \cdot \vec{c}) + y(\vec{b} \cdot \vec{c}) \quad \dots (ii)$$

$$0 = x(\vec{a} \cdot \vec{c}) + y(\vec{b} \cdot \vec{c})$$

$$\frac{x}{\vec{b} \cdot \vec{c}} = -\frac{y}{\vec{a} \cdot \vec{c}} = \lambda$$

$$\therefore x = \lambda(\vec{b} \cdot \vec{c}) \text{ and } y = -\lambda(\vec{a} \cdot \vec{c})$$

$$\text{Substituting the values of } x \text{ and } y \text{ we get, } (\vec{a} \times \vec{b}) \times \vec{c} = \lambda(\vec{b} \cdot \vec{c})\vec{a} - \lambda(\vec{a} \cdot \vec{c})\vec{b}$$

This identity must hold true for all values of  $\vec{a}, \vec{b}, \vec{c}$

$$\text{Substitute } \vec{a} = \hat{i}; \vec{b} = \hat{j} \text{ and } \vec{c} = \hat{k}$$

$$(\hat{i} \times \hat{j}) \times \hat{k} = \lambda(\hat{j} \cdot \hat{k})\hat{i} - \lambda(\hat{i} \cdot \hat{k})\hat{j}$$

$$\hat{j} = -\lambda\hat{j} \Rightarrow \lambda = -1$$

$$\Rightarrow (\vec{a} \times \vec{b}) \times \vec{c} = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{b} \cdot \vec{c})\vec{a}$$

**Note:** Unit vector coplanar with  $\vec{a}$  and  $\vec{b}$  perpendicular to  $\vec{a}$  is  $\pm \frac{(\vec{a} \times \vec{b}) \times \vec{a}}{[(\vec{a} \times \vec{b}) \times \vec{a}]}$

**Illustration 44:** Prove that  $\vec{a} \times \{\vec{b} \times (\vec{c} \times \vec{d})\} = (\vec{b} \cdot \vec{d})(\vec{a} \times \vec{c}) - (\vec{b} \cdot \vec{c})(\vec{a} \times \vec{d})$

(JEE MAIN)

**Sol:** By using vector triple product as mention above.

$$\begin{aligned} \text{We have, } \vec{a} \times \{\vec{b} \times (\vec{c} \times \vec{d})\} &= \vec{a} \times \{(\vec{b} \cdot \vec{d})\vec{c} - (\vec{b} \cdot \vec{c})\vec{d}\} \\ &= \vec{a} \times \{(\vec{b} \cdot \vec{d})\vec{c} - \vec{a}(\vec{b} \cdot \vec{c})\vec{d}\} \quad [\text{by distributivelaw}] \\ &= (\vec{b} \cdot \vec{d})(\vec{a} \times \vec{c}) - (\vec{b} \cdot \vec{c})(\vec{a} \times \vec{d}) \end{aligned}$$

**Illustration 45:** Let  $\vec{a} = a\hat{i} + 2\hat{j} - 3\hat{k}$ ,  $\vec{b} = \hat{i} + 2a\hat{j} - 2\hat{k}$ , and  $\vec{c} = 2\hat{i} + a\hat{j} - \hat{k}$ . Find the value (s) of a, if any, such that  $\{(\vec{a} \times \vec{b}) \times (\vec{b} \times \vec{c})\} \times (\vec{c} \times \vec{a}) = 0$ .

(JEE MAIN)

**Sol:** Here use vector triple product to obtain the value of a.

$$\{(\vec{a} \times \vec{b}) \times (\vec{b} \times \vec{c})\} \times (\vec{c} \times \vec{a}) = [\vec{a} \quad \vec{b} \quad \vec{c}] \vec{b} \times (\vec{c} \times \vec{a}) = [\vec{a} \quad \vec{b} \quad \vec{c}] \{(\vec{a} \cdot \vec{b})\vec{c} - (\vec{b} \cdot \vec{c})\vec{a}\},$$

$$\text{Given } \{(\vec{a} \times \vec{b}) \times (\vec{b} \times \vec{c})\} \times (\vec{c} \times \vec{a}) = 0 \Rightarrow (\vec{a} \cdot \vec{b})\vec{c} = (\vec{b} \cdot \vec{c})\vec{a} \text{ or } [\vec{a} \quad \vec{b} \quad \vec{c}] = 0$$

$(\vec{a} \cdot \vec{b})\vec{c} = (\vec{b} \cdot \vec{c})\vec{a}$  leads to three different equations which do not have a common solution.

$$[\vec{a} \quad \vec{b} \quad \vec{c}] = 0 \Rightarrow \begin{vmatrix} a & 2 & -3 \\ 1 & 2a & -2 \\ 2 & a & -1 \end{vmatrix} = 0 \Rightarrow 9a - 6 = 0 \Rightarrow a = \frac{2}{3}$$

**Illustration 46:** Solve for  $\vec{r}$ , from the simultaneous equations  $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$ ,  $\vec{r} \cdot \vec{a} = 0$ , provided  $\vec{a}$  is not perpendicular to  $\vec{b}$ .

(JEE MAIN)

**Sol:** As given  $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$ , solve this using vector triple product to get the result.

$$\text{Given } \vec{r} \times \vec{b} = \vec{c} \times \vec{b}$$

$$\Rightarrow (\vec{r} - \vec{c}) \times \vec{b} = 0 \Rightarrow (\vec{r} - \vec{c}) \text{ and } \vec{b} \text{ are collinear}$$

$$\therefore \vec{r} - \vec{c} = k\vec{b} \Rightarrow \vec{r} = \vec{c} + k\vec{b} \dots (i)$$

$$\vec{r} \cdot \vec{a} = 0 \Rightarrow (\vec{c} + k\vec{b}) \cdot \vec{a} = 0$$

$$\Rightarrow k = -\frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \text{ putting in eq. (i) we get } \vec{r} = \vec{c} - \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \vec{b} = \frac{\vec{a} \times (\vec{c} \times \vec{b})}{\vec{a} \cdot \vec{b}}.$$

**Illustration 47:** If  $\vec{x} \times \vec{a} + k\vec{x} = \vec{b}$ , where k is a scalar and  $\vec{a}, \vec{b}$  are any two vectors, then determine  $\vec{x}$  in terms of  $\vec{a}, \vec{b}$  and k.

(JEE MAIN)

**Sol:** Here as given  $\vec{x} \times \vec{a} + k\vec{x} = \vec{b}$ , Apply cross product of  $\vec{a}$  with both side and solve using vector triple product.

$$\vec{x} \times \vec{a} + k\vec{x} = \vec{b} \quad \dots (i)$$

$$\Rightarrow \vec{a} \times (\vec{x} \times \vec{a}) + k(\vec{a} \times \vec{x}) = (\vec{a} \times \vec{b})$$

$$\Rightarrow (\vec{a} \cdot \vec{a})\vec{x} - (\vec{a} \cdot \vec{x})\vec{a} + k(\vec{a} \times \vec{x}) = \vec{a} \times \vec{b} \quad \dots (ii)$$

$$(i) \Rightarrow \vec{a} \cdot (\vec{x} \times \vec{a}) + k(\vec{a} \cdot \vec{x}) = \vec{a} \cdot \vec{b}$$

$$\Rightarrow k(\vec{a} \cdot \vec{x}) = \vec{a} \cdot \vec{b} \quad \dots (iii)$$



Substituting the values from equations (i) and (iii) in equation (ii), we get,

$$\Rightarrow (\vec{a} \cdot \vec{a})\vec{x} - \frac{1}{k}(\vec{a} \cdot \vec{b})\vec{a} + k(k\vec{x} - \vec{b}) = \vec{a} \times \vec{b}$$

$$\Rightarrow (a^2 + k^2)\vec{x} = (\vec{a} \times \vec{b}) + \frac{1}{k}(\vec{a} \cdot \vec{b})\vec{a} + k\vec{b} \Rightarrow \vec{x} = \frac{1}{a^2 + k^2} \left[ k\vec{b} + (\vec{a} \times \vec{b}) + \frac{\vec{a} \cdot \vec{b}}{k}\vec{a} \right]$$

## 12. APPLICATION OF VECTORS IN 3D GEOMETRY

(a) Direction cosines of  $\vec{r} = a\hat{i} + b\hat{j} + c\hat{k}$  are given by  $\frac{a}{|\vec{r}|}, \frac{b}{|\vec{r}|}, \frac{c}{|\vec{r}|}$ .

(b) **Incentre formula:** The position vector of the incentre of  $\triangle ABC$  is  $\frac{a\vec{a} + b\vec{b} + c\vec{c}}{a + b + c}$

(c) **Orthocentre formula:** The position vector of the orthocenter of

$$\triangle ABC \text{ is } \frac{\vec{a} \tan A + \vec{b} \tan B + \vec{c} \tan C}{\tan A + \tan B + \tan C}$$

(d) The vector equation of a straight line passing through a fixed point with position vector  $\vec{a}$  and parallel to a given vector  $\vec{b}$  is given by  $\vec{r} = \vec{a} + \lambda\vec{b}$ .

(e) The vector equation of a line passing through two points with position vectors  $\vec{a}$  and  $\vec{b}$  is given by  $\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$ .

(f) **Perpendicular distance of a point from a line:** Let L be the foot of perpendicular drawn  $P(\vec{\alpha})$  on the line  $\vec{r} = \vec{a} + \lambda\vec{b}$ , where  $\vec{r}$  is the position vector of any point on the give line. Therefore, let the position vector  $\vec{L}$  be  $\vec{a} + \lambda\vec{b}$ .

$$PL = \frac{|(\vec{a} - \vec{\alpha}) \times \vec{b}|}{|\vec{b}|} \text{ and } \vec{PL} = \vec{a} - \vec{\alpha} + \lambda\vec{b} = (\vec{a} - \vec{\alpha}) - \left( \frac{(\vec{a} - \vec{\alpha}) \cdot \vec{b}}{|\vec{b}|^2} \right) \vec{b}$$

The length PL is the magnitude of  $\vec{PL}$ , and the required length of perpendicular.




Figure 26.20

(g) **Image of a point in a straight line:** If  $Q(\vec{\beta})$  is the image of P in  $\vec{r} = \vec{a} + \lambda\vec{b}$ , then

$$\vec{\beta} = 2\vec{a} - \left( \frac{2(\vec{a} - \vec{\alpha}) \cdot \vec{b}}{|\vec{b}|^2} \right) \vec{b} - \vec{\alpha}$$




Figure 26.21

- (h) **Shortest distance between two skew lines:** Let  $l_1$  and  $l_2$  be two lines whose equations are  $l_1 : \vec{r} = \vec{a}_1 + \lambda \vec{b}_1$  and  $l_2 : \vec{r} = \vec{a}_2 + \lambda \vec{b}_2$ , respectively.

Then, shortest distance is given by  $PQ = \frac{|\left(\vec{b}_1 \times \vec{b}_2\right) \cdot (\vec{a}_2 - \vec{a}_1)|}{|\vec{b}_1 \times \vec{b}_2|} = \frac{|\left[\vec{b}_1 \quad \vec{b}_2 \quad \vec{a}_2 - \vec{a}_1\right]|}{|\vec{b}_1 \times \vec{b}_2|}$

**Shortest distance between two parallel lines:** The shortest distance between the two given parallel lines

$\vec{r} = \vec{a}_1 + \lambda \vec{b}$  and  $\vec{r} = \vec{a}_2 + \mu \vec{b}$  is given by  $d = \frac{|(\vec{a}_2 - \vec{a}_1) \times \vec{b}|}{|\vec{b}|}$ .

If the lines  $\vec{r} = \vec{a}_1 + \lambda \vec{b}_1$  and  $\vec{r} = \vec{a}_2 + \mu \vec{b}_2$  intersect, then the shortest distance between them is zero.

Therefore,  $\left[\vec{b}_1 \quad \vec{b}_2 \quad \vec{a}_2 - \vec{a}_1\right] = 0$

- (i) If the lines  $\vec{r} = \vec{a}_1 + \lambda \vec{b}_1$  and  $\vec{r} = \vec{a}_2 + \lambda \vec{b}_2$  are coplanar, then  $\left[\vec{a}_1 \quad \vec{b}_1 \quad \vec{b}_2\right] = \left[\vec{a}_2 \quad \vec{b}_1 \quad \vec{b}_2\right]$  and the equation of the plane containing them is given by  $\left[\vec{r} \quad \vec{b}_1 \quad \vec{b}_2\right] = \left[\vec{a}_1 \quad \vec{b}_1 \quad \vec{b}_2\right]$ .
- (j) The vector equation of a plane through the point A ( $\vec{a}$ ) and perpendicular to the vector  $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$  is given by  $(\vec{r} - \vec{a}) \cdot \vec{n} = 0$ .
- (k) Vector The vector equation of a plane normal to unit vector  $\vec{n}$  and at a distance  $d$  from the origin is given by  $\vec{r} \cdot \vec{n} = d$ .
- (l) The equation of the plane passing through a point having position vector  $\vec{a}$  and parallel to  $\vec{b}$  and  $\vec{c}$  is given by  $\vec{r} = \vec{a} + \lambda \vec{b} + \mu \vec{c} \Rightarrow \left[\vec{r} \quad \vec{b} \quad \vec{c}\right] = \left[\vec{a} \quad \vec{b} \quad \vec{c}\right]$ , where  $\lambda$  and  $\mu$  are scalars.
- (m) The vector equation of plane passing through a point  $\vec{a}, \vec{b}, \vec{c}$  is given by  $\vec{r} = (1-s-t) \vec{a} + s \vec{b} + t \vec{c}$   
or  $\vec{r} \cdot (\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{a} \times \vec{b}) = \left[\vec{a} \quad \vec{b} \quad \vec{c}\right]$
- (n) The equation of any plane through the intersection of planes  $\vec{r} \cdot \vec{n}_1 = d_1$  and  $\vec{r} \cdot \vec{n}_2 = d_2$  is  $\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = d_1 + \lambda d_2$ , where  $\lambda$  is an arbitrary constant.
- (o) The perpendicular distance of a point having position vector  $\vec{a}$  from the plane  $\vec{r} \cdot \vec{n} = d$  is given by  $p = \frac{|\vec{a} \cdot \vec{n} - d|}{|\vec{n}|}$ .
- (p) The angle  $\theta$  between the planes  $\vec{r}_1 \cdot \vec{n}_1 = d_1$  and  $\vec{r}_2 \cdot \vec{n}_2 = d_2$  is given by  $\cos \theta = \pm \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|}$
- (q) The perpendicular distance of a point P ( $\vec{r}$ ) from a line passing through  $\vec{a}$  and parallel to  $\vec{b}$  is given by
- $$P = \frac{|\left(\vec{r} - \vec{a}\right) \times \vec{b}|}{|\vec{b}|} = \left[ (\vec{r} - \vec{a})^2 - \left\{ \frac{(\vec{r} - \vec{a}) \cdot \vec{b}}{|\vec{b}|} \right\}^2 \right]^{1/2}$$
- (r) The equation of the planes bisecting the angles between the planes  $\vec{r}_1 \cdot \vec{n}_1 = d_1$  and  $\vec{r}_2 \cdot \vec{n}_2 = d_2$  is
- $$\vec{r} \cdot (\vec{n}_1 \pm \vec{n}_2) = \frac{d_1}{|\vec{n}_1|} \pm \frac{d_2}{|\vec{n}_2|}$$
- (s) The perpendicular distance of a point P ( $\vec{r}$ ) from a plane passing through a point  $\vec{a}$  and parallel to points  $\vec{b}$  and  $\vec{c}$  is given by  $PM = \frac{(\vec{r} - \vec{a}) \cdot (\vec{b} \times \vec{c})}{|\vec{b} \times \vec{c}|}$

(t) The perpendicular distance of a point P( $\vec{r}$ ) from a plane passing through the points  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  is given by

$$P = \frac{(\vec{r} - \vec{a}) \cdot (\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{a} \times \vec{b})}{|\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{a} \times \vec{b}|}$$

(u) **Angle between a line and the plane:** If  $\theta$  is the angle between a line  $\vec{r} = (\vec{a} + \lambda \vec{b})$  and the plane  $\vec{r} \cdot \vec{n} = d$ , then

$$\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| |\vec{n}|}.$$

(v) The equation of sphere with center at C( $\vec{c}$ ) and radius 'a' is  $|\vec{r} - \vec{c}| = a$ . If center is the origin then  $|\vec{r}| = a$ .

(w) The plane  $\vec{r} \cdot \vec{n} = d$  touches the sphere  $|\vec{r} - \vec{a}| = R$ , if  $\frac{|\vec{a} \cdot \vec{n} - d|}{|\vec{n}|} = R$ , i.e. the condition of tangency.

(x) If  $\vec{a}$  and  $\vec{b}$  are the position vectors of the extremities of a diameter of a sphere, then its equation is given by  $(\vec{r} - \vec{a}) \cdot (\vec{r} - \vec{b}) = 0$  or  $|\vec{r}|^2 - \vec{r} \cdot (\vec{a} + \vec{b}) + \vec{a} \cdot \vec{b} = 0$  or  $|\vec{r} - \vec{a}|^2 + |\vec{r} - \vec{b}|^2 = |\vec{a} - \vec{b}|^2$ .

## FORMULAE SHEET

(a)  $\vec{OP} = x\hat{i} + y\hat{j}$

(b)  $|\vec{OP}| = \sqrt{x^2 + y^2}$  and direction is  $\tan \theta = \frac{y}{x}$

(c) Unit vector  $\hat{U} = \frac{\text{Vector}}{\text{Its modulus}} = \frac{\vec{a}}{|\vec{a}|}$

(d) Properties of vector addition:

i. $\vec{a} + \vec{b} = \vec{b} + \vec{a}$ commutative	(a) $\vec{a} + (\vec{b} + \vec{c}) = (\vec{a} + \vec{b}) + \vec{c}$ Associative
ii. $\vec{a} + \vec{0} = \vec{a}$ Null vector is an additive identity	(b) $\vec{a} + (-\vec{a}) = \vec{0}$ Additive inverse
iii. $c(\vec{a} + \vec{b}) = c\vec{a} + c\vec{b}$	(c) $(c + d)\vec{a} = c\vec{a} + d\vec{a}$
iv. $(cd)\vec{a} = c(d\vec{a})$	(d) $1 \times \vec{a} = \vec{a}$

(e) **Section formula:**

(i) If  $\vec{a}$  and  $\vec{b}$  are the position vectors of two points A and B, then the position vector of a point which divides

AB in the ratio m:n is given by  $\vec{r} = \frac{(n\vec{a} + m\vec{b})}{(m+n)}$ .

(ii) Position vector of mid-point of  $\vec{AB} = \frac{(\vec{a} + \vec{b})}{2}$ .

(f) **Collinearity of three points:** If  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$  are the position vectors (non-zero) of three points and given they are collinear then there exists  $\lambda, \gamma$  both not being 0 such that  $\vec{a} + \lambda\vec{b} + \gamma\vec{c}$

(g) **Coplanar vectors:** Let  $\vec{a}, \vec{b}$  be non-zero, non-collinear vectors. Then, any vector  $\vec{r}$  coplanar with  $\vec{a}, \vec{b}$  can be expressed uniquely as a linear combination of  $\vec{a}, \vec{b}$  i.e. there exist some unique  $x, y \in \mathbb{R}$ , such that  $x\vec{a} + y\vec{b} = \vec{r}$

(h) **Product of two vectors:**

(i) **Scalar Product (dot product)**

$$\text{If } \vec{a} \cdot \vec{b} = a_1b_1 + a_2b_2 + a_3b_3$$

**Note :** •  $\cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}||\vec{b}|}$

•  $\vec{a}$  and  $\vec{b}$  are perpendicular if  $\theta = 90^\circ$

(ii) **Properties of scalar product:**

i. $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$	ii. $m\vec{a} \cdot n\vec{b} = mn\vec{a} \cdot \vec{b} = \vec{a} \cdot (mn\vec{b})$
iii. $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$	iv. $(\vec{a} + \vec{b})^2 = \vec{a}^2 + 2\vec{a} \cdot \vec{b} + \vec{b}^2$
v. If $\hat{i} = (1, 0, 0), \hat{j} = (0, 1, 0), \hat{k} = (0, 0, 1)$ then $\hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0$	

(iii) **Vector (cross) Product of two vectors:** Let  $\vec{a} = (a_1, a_2, a_3), \vec{b} = (b_1, b_2, b_3)$  be two vectors then the cross product of  $\vec{a} \times \vec{b}$  is devotes by  $\vec{a} \times \vec{b}$  and defined by

$$\vec{a} \times \vec{b} = (a_1, a_2, a_3) \times (b_1, b_2, b_3) = \begin{pmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{pmatrix} = (a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1)$$

OR

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = (a_2b_3 - a_3b_2)\hat{i} + (a_3b_1 - a_1b_3)\hat{j} + (a_1b_2 - a_2b_1)\hat{k}$$

$$|\vec{a} \times \vec{b}| = |\vec{a}| \times |\vec{b}| \sin\theta \hat{n}$$

**Note: (i)**  $\theta$  being angle between  $\vec{a}$  &  $\vec{b}$

(ii) If  $\theta = 0$ , The  $|\vec{a} \times \vec{b}| = 0$  i.e.  $\vec{a} \times \vec{b} = 0$  and  $\vec{a}$  &  $\vec{b}$  are parallel if  $\vec{a} \times \vec{b} = 0$ .

**(iv) Properties of cross product**

i. $\vec{a} \times \vec{b} = 0 \Rightarrow \vec{a} = 0 \text{ or } \vec{b} = 0 \text{ or } \vec{a} \parallel \vec{b}$	ii. $\vec{a} \times \vec{b} = -\vec{b} \times \vec{a}$
iii. $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$	iv. $(n\vec{a}) \times \vec{b} = n(\vec{a} \times \vec{b})$
v. $\vec{a} \times \vec{b}$ is perpendicular to both $\vec{a}$ and $\vec{b}$	vi. $ \vec{a} \times \vec{b} $ is a Area of parallelogram with sides $\vec{a}$ and $\vec{b}$ .

**(v) Scalar Triple Product:** If  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ ,  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ ,  $\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$ .

$$\text{Then } \vec{a} \cdot (\vec{b} \times \vec{c}) = \vec{b} \cdot (\vec{c} \times \vec{a}) = \vec{c} \cdot (\vec{a} \times \vec{b})$$

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

$$\vec{a} \cdot (\vec{b} \times \vec{c}) \text{ is also represented as } [\vec{a} \ \vec{b} \ \vec{c}]$$

$$[\vec{a} \ \vec{b} \ \vec{c}] = [\vec{b} \ \vec{c} \ \vec{a}] = [\vec{c} \ \vec{a} \ \vec{b}]$$

$$[\vec{a} \ \vec{b} \ \vec{c}] = -[\vec{a} \ \vec{c} \ \vec{b}]$$

- If any of the two vectors are parallel, then  $[\vec{a} \ \vec{b} \ \vec{c}] = 0$
- $[\vec{a} \ \vec{b} \ \vec{c}]$  is the volume of the parallelepiped whose coterminous edges are formed by  $\vec{a} \ \vec{b} \ \vec{c}$
- If  $\vec{a} \ \vec{b} \ \vec{c}$  are coplanar,  $[\vec{a} \ \vec{b} \ \vec{c}] = 0$
- $\frac{1}{2}|\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}|$  = area of triangle having  $\vec{a}, \vec{b}, \vec{c}$  as position vectors of vertices of a triangle.

**(vi) Vector Triple Product:**

$$\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}$$

$$(\vec{a} \times \vec{b}) \times \vec{c} = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{b} \cdot \vec{c})\vec{a}$$

$$\text{Unit vector coplanar with } \vec{a} \text{ and } \vec{b} \text{ perpendicular to } \vec{a} \text{ is } \pm \frac{(\vec{a} \times \vec{b}) \times \vec{a}}{|\vec{a} \times \vec{b}| |\vec{a}|}.$$

## Solved Examples

### JEE Main/Boards

**Example 1:** Show that the points A, B & C with position vector  $2\hat{i} - \hat{j} + \hat{k}$ ,  $\hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  respectively, are the vertices of a right angled triangle. Also find the remaining angles of the triangle.

**Sol:** We have,

$$\overrightarrow{AB} = (\hat{i} - 3\hat{j} - 5\hat{k}) - (2\hat{i} - \hat{j} + \hat{k}) = -\hat{i} - 2\hat{j} - 6\hat{k}$$

$$\overrightarrow{BC} = (3\hat{i} - 4\hat{j} - 4\hat{k}) - (\hat{i} - 3\hat{j} - 5\hat{k}) = 2\hat{i} - \hat{j} + \hat{k} \text{ and}$$

$$\overrightarrow{CA} = (2\hat{i} - \hat{j} + \hat{k}) - (3\hat{i} - 4\hat{j} - 4\hat{k}) = -\hat{i} + 3\hat{j} + 5\hat{k}$$

Since  $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA}$

$$= (-\hat{i} - 2\hat{j} - 6\hat{k}) + (2\hat{i} - \hat{j} + \hat{k}) + (-\hat{i} + 3\hat{j} + 5\hat{k}) = \vec{0}$$

So, A, B and C are the vertices of a triangle.

Now,  $\overrightarrow{BC} \cdot \overrightarrow{CA}$

$$= (2\hat{i} - \hat{j} + \hat{k}) \cdot (-\hat{i} + 3\hat{j} + 5\hat{k}) = -2 - 3 + 5 = 0$$

$$\overrightarrow{BC} \perp \overrightarrow{CA} \Rightarrow \angle BCA = \frac{\pi}{2}$$

Hence, ABC is a right angled triangle. Since A is the angle between the vectors  $\overrightarrow{AB}$  and  $\overrightarrow{AC}$ . Therefore,

$$\begin{aligned} \cos A &= \frac{\overrightarrow{AB} \cdot \overrightarrow{AC}}{|\overrightarrow{AB}| |\overrightarrow{AC}|} \\ &= \frac{(-\hat{i} - 2\hat{j} - 6\hat{k}) \cdot (\hat{i} - 3\hat{j} - 5\hat{k})}{\sqrt{(-1)^2 + (-2)^2 + (-6)^2} \sqrt{1^2 + (-3)^2 + (-5)^2}} \\ &= \frac{-1 + 6 + 30}{\sqrt{1+4+36} \sqrt{1+9+25}} = \frac{35}{\sqrt{41} \sqrt{35}} = \sqrt{\frac{35}{41}} \end{aligned}$$

$$A = \cos^{-1} \sqrt{\frac{35}{41}}, \cos B = \frac{\overrightarrow{BA} \cdot \overrightarrow{BC}}{|\overrightarrow{BA}| |\overrightarrow{BC}|}$$

$$= \frac{(\hat{i} + 2\hat{j} + 6\hat{k}) \cdot (2\hat{i} - \hat{j} + \hat{k})}{\sqrt{1^2 + 2^2 + 6^2} \sqrt{2^2 + (-1)^2 + 1^2}}$$


$$\Rightarrow \cos B = \frac{2 - 2 + 6}{\sqrt{41} \sqrt{6}} = \sqrt{\frac{6}{41}} \Rightarrow B = \cos^{-1} \sqrt{\frac{6}{41}}$$

**Example 2:** If ABCDE is a pentagon, prove that the resultant of  $\overrightarrow{AB}, \overrightarrow{AE}, \overrightarrow{BC}, \overrightarrow{DC}, \overrightarrow{ED}$  and  $\overrightarrow{AC}$  is  $3\overrightarrow{AC}$

**Sol:** By using resultant vector formula, we can obtain required result.

If R be the resultant vector then

$$\begin{aligned} R &= \overrightarrow{AB} + \overrightarrow{AE} + \overrightarrow{BC} + \overrightarrow{DC} + \overrightarrow{ED} + \overrightarrow{AC} \\ &= (\overrightarrow{AB} + \overrightarrow{BC}) + (\overrightarrow{AE} + \overrightarrow{ED} + \overrightarrow{DC}) + \overrightarrow{AC} \\ &= \overrightarrow{AC} + \overrightarrow{AC} + \overrightarrow{AC} = 3\overrightarrow{AC} \end{aligned}$$



**Example 3:** Prove that the straight lines joining the mid points of the sides of a quadrilateral ABCD, taken in order, form a parallelogram.

**Sol:** Let the position vectors of A, B, C and D be  $\vec{a}, \vec{b}, \vec{c}$  and  $\vec{d}$ . Hence position vectors of P, Q, R and S (the mid points of AB, BC, CD & DA respectively) are

$$\frac{\vec{a} + \vec{b}}{2}, \frac{\vec{b} + \vec{c}}{2}, \frac{\vec{c} + \vec{d}}{2} \text{ and } \frac{\vec{d} + \vec{a}}{2} \text{ respectively.}$$


$$\overrightarrow{PQ} = \frac{\vec{c} + \vec{b}}{2} - \frac{\vec{a} + \vec{b}}{2}$$

$$\overrightarrow{PQ} = \overrightarrow{PO} + \overrightarrow{OQ} = \frac{\vec{c} - \vec{a}}{2}$$

$$\overrightarrow{RS} = \frac{\vec{a} + \vec{d}}{2} - \frac{\vec{c} + \vec{d}}{2} = \frac{\vec{a} - \vec{c}}{2}$$

$$\Rightarrow \overrightarrow{SR} = \frac{\vec{c} - \vec{a}}{2}$$

$\overrightarrow{PQ}$  is parallel and equal to  $\overrightarrow{SR}$ . Hence PQRS is a parallelogram.



**Example 4:** Write an equation for the plane that contains the points (2, 0, -3), (-4, -5, 2), and (0, 3, -4) in the form  $ax + by + cz = d$ .

**Sol:** Let  $\vec{v} = (-4, -5, 2) - (2, 0, -3) = (-6, -5, 5)$  and  $\vec{w} = (0, 3, -4) - (2, 0, -3) = (-2, 3, -1)$ .

$$\vec{v} \times \vec{w} = \hat{i}(5 - 15) - \hat{j}(6 + 10) +$$

$$\hat{k}(-18 - 10) = (-10, -16, -28)$$

We can choose  $\hat{n}$  to be any vector in the same direction as  $\vec{v} \times \vec{w}$  so let  $\hat{n} = (5, 8, 14)$ . Then the plane has the

form  $5x + 8y + 14z = d$ . Substituting the point  $(2, 0, -3)$  for  $(x, y, z)$  and solving for  $d$  gives  $d = 10 + 0 + 0(-42) = -32$ . So the plane has the equation  $5x + 8y + 14z = -32$ .

**Example 5:** Find a vector that is perpendicular to the vector  $(1, 2, 3)$  with the same length. Also, find a plane perpendicular to  $(1, 2, 3)$  that passes through the point  $(3, 2, 1)$ .

**Sol:** By using formula of perpendicular vector we can obtain the result.

A vector  $\vec{v}$  is perpendicular to  $(1, 2, 3)$  if  $\vec{v} \cdot (1, 2, 3) = 0$ . There are infinite number of possibilities to choose from, but one possible choice for  $\vec{v}$  is  $(2, -1, 0)$ . However, we want this vector to have length  $\|(1, 2, 3)\| = \sqrt{14}$ .

Since  $\|(2, -1, 0)\| = \sqrt{5}$ , we need to rescale our vector to be  $\frac{\sqrt{14}}{\sqrt{5}}(2, -1, 0)$ .

If  $(1, 2, 3)$  is a normal vector to a plane then the plane will have the form  $x + 2y + 3z = d$ . Since the plane passes through the point  $(3, 2, 1)$ , we substitute these values for  $x, y$ , and  $z$  to get  $3 + 4 + 3 = 10 = d$  so our plane equation is  $x + 2y + 3z = 10$ .

**Example 6:** Write an equation for the plane that contains the point  $(1, 0, 3)$  and the line  $(-3, -2, -2) + t(1, 2, -1)$  in the form  $ax + by + cz = d$ .

**Sol:** Since the plane contains the line  $(-3, -2, -2) + t(1, 2, -1)$  we know that one tangent vector to the plane is  $\vec{v} = (1, 2, -1)$ . We can get a second tangent vector by finding the vector between  $(-3, -2, -2)$  and  $(1, 0, 3)$ .

So let  $\vec{w} = (4, 2, 5)$ . Then

$$\vec{v} \times \vec{w} = \vec{i}(10 + 2) - \vec{j}(5 + 4) + \vec{k}(2 - 8) = (12, -9, -6)$$

So we can choose  $\vec{n} = (4, -3, -2)$  and our plane has the form  $4x - 3y - 2z = d$ . Plugging in  $(1, 0, 3)$  for  $(x, y, z)$  and solving for  $d$  yields  $4x - 3y - 2z = -2$ .

**Example 7:** Find the minimum distance between the point  $(3, -3, -3)$  and the plane  $2x + y - z = 3$ .

**Sol:** The point in the plane closest to  $(3, -3, -3)$  lies on a line that is perpendicular to the plane and passes through  $(3, -3, -3)$ . Since  $(2, 1, -1)$  is a normal vector to the plane, we will use it as the direction of this line. Thus a parameterized form of the line is

$$c(t) = (3, -3, -3) + t(2, 1, -1) = (3 + 2t, -3 + t, -3 - t)$$

We substitute this into the plane equation to find its intersection with the plane and get:

$$2(3 + 2t) + 1(-3 + t) - (-3 - t) = 6 + 6t = 3 \Rightarrow t = -\frac{1}{2}.$$

So the point in the plane closest to  $(3, -3, -3)$  is

$$c\left(-\frac{1}{2}\right) = \left(2, -\frac{7}{2}, -\frac{5}{2}\right).$$

The distance between the point and the plane is thus

$$\sqrt{1^2 + \left(\frac{1}{2}\right)^2 + \left(-\frac{1}{2}\right)^2} = \sqrt{\frac{3}{2}}.$$

**Example 8:** Determine if the three vectors  $\vec{a} = (1, 4, -7)$ ,  $\vec{b} = (2, -1, 4)$  and  $\vec{c} = (0, -9, 18)$  lie in the same plane or not.

**Sol:** Three vectors lie in the same plane if volume of the parallelepiped formed by these three vectors is zero.

So, as we noted prior to this example all we need to do is compute the volume of the parallelepiped formed by these three vectors. If the volume is zero, then they lie in the same plane and if the volume isn't zero they don't lie in the same plane.

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = \begin{vmatrix} 1 & 4 & -7 \\ 2 & -1 & 4 \\ 0 & -9 & 18 \end{vmatrix} = -18 + 126 - 144 + 36 = 0$$

So, the volume is zero and so they lie in the same plane.

## JEE Advanced/Boards

**Example 1:** If  $O$  be the circumcenter;  $G$ , the centroid and  $H$ , the orthocenter of triangle  $ABC$ , prove that  $O, G, H$  are collinear and  $G$  divides  $OH$  in the ratio  $1:2$ .

**Sol:** Consider position vector of  $A, B, C$  be taken as  $\vec{a}, \vec{b}, \vec{c}$ . And then use geometry of triangle to solve this problem.

Let  $O$ , the circumcenter of the  $\triangle ABC$  be chosen as origin and position vector of  $A, B, C$  be taken  $\vec{a}, \vec{b}, \vec{c}$ .

Hence position vector of  $G$  the centroid is

$$\vec{OG} = \frac{\vec{a} + \vec{b} + \vec{c}}{3} \quad \dots (i)$$

Since  $O$  is circumcenter


$$\therefore \vec{OA} = \vec{OB} = \vec{OC} \Rightarrow \vec{OA}^2 = \vec{OB}^2 = \vec{OC}^2 \text{ or } a^2 = b^2 = c^2$$

$$a^2 - b^2 = 0, \quad b^2 - c^2 = 0, \quad c^2 - a^2 = 0$$

$$\text{Or } (\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{c}) = 0$$

$$\text{Or } (\vec{a} + \vec{b} + \vec{c} - \vec{c}) \cdot (\vec{a} - \vec{b}) = 0$$

Let P be the point whose position vector is



$$\vec{a} + \vec{b} + \vec{c} \therefore (\vec{OP} - \vec{OC}) \cdot (\vec{OA} - \vec{OB}) = 0$$

$$\text{Or } \vec{CP} \perp \vec{BA}$$

In similar manner we can show that BP is perpendicular to AC and AP is perpendicular to CB.

Hence P is the orthocentre which is H.

$$\vec{OP} = \vec{OH} = \vec{a} + \vec{b} + \vec{c} = 3\vec{OG} \quad \dots (iii)$$

$$\therefore \vec{OH} = 3\vec{OG} \text{ or } \vec{GH} = 2\vec{OG} \text{ or } \frac{\vec{OG}}{\vec{GH}} = \frac{1}{2}$$

Above show that O, G, H are collinear and G divides OH in the ratio 1:2


**Example 2:** Prove using vectors: If two medians of a triangle are equal, then it is isosceles.

**Sol:** Using mid – point formula of vector, we can solve this

Let ABC be a triangle and let BE and CF be two equal medians. Taking A as the origin, let the position vectors of B and C be  $\vec{b}$  and  $\vec{c}$  respectively. Then

$$\text{Position vector of E} = \frac{1}{2}\vec{c} \text{ and}$$

$$\text{Position vector of F} = \frac{1}{2}\vec{b}$$



$$\therefore \vec{BE} = \frac{1}{2}(\vec{c} - 2\vec{b}), \vec{CF} = \frac{1}{2}(\vec{b} - 2\vec{c})$$

$$\text{Now, } BE = CF = |\vec{BE}| = |\vec{CF}|$$

$$\Rightarrow |\vec{BE}|^2 = |\vec{CF}|^2 \Rightarrow \left| \frac{1}{2}(\vec{c} - 2\vec{b}) \right|^2 = \left| \frac{1}{2}(\vec{b} - 2\vec{c}) \right|^2$$

$$\Rightarrow \frac{1}{4}|\vec{c} - 2\vec{b}|^2 = \frac{1}{4}|\vec{b} - 2\vec{c}|^2 \Rightarrow |(\vec{c} - 2\vec{b})|^2 = |(\vec{b} - 2\vec{c})|^2$$

$$\Rightarrow (\vec{c} - 2\vec{b}) \cdot (\vec{c} - 2\vec{b}) = (\vec{b} - 2\vec{c}) \cdot (\vec{b} - 2\vec{c})$$

$$\Rightarrow |\vec{c}|^2 - 4\vec{b} \cdot \vec{c} + 4|\vec{b}|^2 = |\vec{b}|^2 - 4\vec{b} \cdot \vec{c} + 4|\vec{c}|^2$$

$$\Rightarrow 3|\vec{b}|^2 = 3|\vec{c}|^2 \Rightarrow |\vec{b}|^2 = |\vec{c}|^2$$

$$\Rightarrow AB = AC$$

Hence, triangle ABC is an isosceles triangle.

**Example 3:** D, E, F are points dividing side  $\vec{BC}$ ,  $\vec{CA}$ ,  $\vec{AB}$  of a triangle ABC in the ratio 2:3, 1:2 and 3:1 respectively. Show that the lines  $\vec{AD}$ ,  $\vec{BE}$ ,  $\vec{CF}$  are concurrent and hence find the position vector of their point of intersection.


**Sol:** By using section formula we can obtain required result.

If  $\vec{d}$ ,  $\vec{e}$ ,  $\vec{f}$  are position vector of points D, E & F respectively then, by section formula

$$\vec{d} = \frac{2\vec{c} + 3\vec{b}}{5} \quad \dots (i)$$

$$\vec{e} = \frac{2\vec{c} + \vec{a}}{3} \quad \dots (ii)$$

$$\vec{f} = \frac{3\vec{b} + \vec{a}}{4} \quad \dots (iii)$$



$$\text{Equation of line AD is } \vec{r} = \vec{a} + t(\vec{d} - \vec{a})$$

$$\text{Equation of line BE is } \vec{r} = \vec{b} + m(\vec{e} - \vec{b})$$

For intersection of  $\vec{AD}$  and  $\vec{BE}$  we need that

" $\vec{a} + t(\vec{d} - \vec{a}) = \vec{b} + m(\vec{e} - \vec{b})$ " be true for some  $0 < t, m < 1$ .

$$\vec{a} + t \frac{(2\vec{c} + 3\vec{b} - 5\vec{a})}{5} = \vec{b} + m \frac{(2\vec{c} + \vec{a} - 3\vec{b})}{3}$$



$$\therefore 1-t = \frac{m}{3}; \frac{3t}{5} = 1-m; \frac{2t}{5} = \frac{2m}{3}$$

$$\therefore t = \frac{5}{6}, m = \frac{1}{2}$$

The existence of  $t$  and  $m$  assures the intersection of  $\overrightarrow{AD}$  and  $\overrightarrow{BE}$ .

The point of intersection is

$$\vec{r} = \vec{a} + \frac{5}{6}(\vec{d} - \vec{a}) = \frac{(\vec{a} + 3\vec{b} + 2\vec{c})}{6}$$

**Example 4:** Find a parametric form for the line passing through the point  $(1,2)$  in the direction  $(3,4)$ , which we will call  $c_1(t)$ . Set  $c_1(t)$  equal to  $(x,y)$  and eliminate  $t$  to get the line into  $y = mx + b$  form. Now find a different parametrization  $c_2(t)$  of the same line such that  $c_2(0) = (-2,-2)$  and  $c_2(2) = (-5,-6)$ .

**Sol:**  $c_1(t) = (1,2) + t(3,4) = (1+3t, 2+4t)$ . Setting  $(x,y) = (1+3t, 2+4t)$  yields  $x = 1+3t$  and  $y = 2+4t$ .

Solving the former equation for  $t$  yields  $t = (x-1)/3$ . Substituting this into the second equation then gives

$$\text{us } y = \frac{4}{3}x + \frac{2}{3}.$$

Let  $c_2(t) = p + t\vec{v}$ .  $c_2$  will then be a parameterization of the same line given by  $c_1$  if  $p$  is a point on the same line and  $\vec{v}$  is in the same direction as  $(3,4)$  (i.e. some scalar multiple of  $(3,4)$ ). Since  $c_2(0) = (-2,-2)$  we will choose  $p = (-2,-2)$  (you can check that this point indeed lies on the line parameterized by  $c_1$ ). Then

$c_2(2) = (-2,-2) + 2\vec{v} = (-5,-6)$ , so we get that

$\vec{v} = (-3/2, -2)$ , which is indeed a scalar multiple of  $(3,4)$ . So

$c_2(t) = (-2,-2) + t\left(\frac{-3}{2}, -2\right)$  is a different

parameterization of the line parameterized by  $c_1$ .

**Example 5:** Find the vector projection of  $(3,2)$  onto  $(-1,-1)$ . Then find the area of the triangle with one side vector  $(3,2)$  and another side the result of this projection.

**Sol:** Use projection method to obtain vector projection of  $(3,2)$  and area of triangle will be half of the area of parallelogram.

$$\text{proj}_{(-1,-1)}(3,2) = \frac{-5}{2}(-1,-1) = \left(\frac{5}{2}, \frac{5}{2}\right).$$

Then the area of the triangle with sides  $(3,2)$  and  $\left(\frac{5}{2}, \frac{5}{2}\right)$  is one half the area of the parallelogram with sides  $(3,2)$  and  $\left(\frac{5}{2}, \frac{5}{2}\right)$ . So, the area of the triangle is

$$\frac{1}{2} \left\| (3,2,0) \times \left(\frac{5}{2}, \frac{5}{2}, 0\right) \right\| = \frac{1}{2} \left\| \left(0, 0, \frac{5}{2}\right) \right\| = \left[\frac{5}{4}\right].$$

**Example 6:** Find the minimum distance between the point  $(4, 2, -3)$  and the line  $(1, 0, 2) + t(-1, -1, 2)$ .

**Sol:** Let  $\vec{v}(t)$  represents the vector from the point  $(4, 2, -3)$  and line  $(1, 0, 2) + t(-1, -1, 2) = (1-t, -t, 2+2t)$  at any  $t \in \mathbb{R}$ .

So,  $\vec{v}(t) = (4, 2, -3) - (1-t, -t, 2+2t) = (3+t, 2+t, -5-2t)$ . We want to find the  $t$  such that  $\vec{v}(t)$  is perpendicular to the line, which is when  $\vec{v}(t) \cdot (-1, -1, 2) = 0$ .

$$(3+t, 2+t, -5-2t) \cdot (-1, -1, 2) = -15 - 6t = 0$$

$\Rightarrow t = -\frac{5}{2}$ . So the length of  $\vec{v}\left(-\frac{5}{2}\right)$  should represent

the minimum distance from  $(4, 2, -3)$  and the line.

$$\left\| \vec{v}\left(-\frac{5}{2}\right) \right\| = \left\| (1/2, -1/2, 0) \right\| = \frac{1}{\sqrt{2}}.$$

**Example 7:** Prove that, in any triangle ABC

$$(i) c^2 = a^2 + b^2 - 2ab \cos C \quad (ii) c = b \cos A + a \cos B$$

**Sol:** By using simple scalar product method we can prove given relation.

$$(i) \text{ In } \triangle ABC, \overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = 0$$

$$\text{or, } \overrightarrow{BC} + \overrightarrow{CA} = -\overrightarrow{AB}$$

....(i)

Squaring both sides

$$(\overrightarrow{BC})^2 + (\overrightarrow{CA})^2 + 2(\overrightarrow{BC}) \cdot (\overrightarrow{CA}) = (\overrightarrow{AB})^2$$

$$\Rightarrow a^2 + b^2 + 2(\overrightarrow{BC} \cdot \overrightarrow{CA}) = c^2$$


$$\Rightarrow c^2 = a^2 + b^2 = 2ab \cos(\pi - C)$$

$$\Rightarrow c^2 = a^2 + b^2 - 2ab \cos C$$

$$(ii) (\overrightarrow{BC} + \overrightarrow{CA}) \cdot \overrightarrow{AB} = -\overrightarrow{AB} \cdot \overrightarrow{AB} \Rightarrow \overrightarrow{BC} \cdot \overrightarrow{AB} + \overrightarrow{CA} \cdot \overrightarrow{AB}$$

$$= -c^2 - a \cos B - b \cos A$$


$$\Rightarrow a \cos B + b \cos A = c$$



**Example 8:** In any triangle, show that the perpendicular bisectors of the sides are concurrent.

**Sol:** By using formula of Dot product and Mid – point we can solve this problem.

Let ABC be the triangle and D, E and F are respectively middle points of sides  $\overline{BC}$ ,  $\overline{CA}$  and  $\overline{AB}$ . Let the perpendicular through D and E meet at O join  $\overline{OF}$ . We are required to prove that  $\overline{OF}$  is  $\perp$  to  $\overline{AB}$ . Let the position vectors of A, B, C with O as origin of reference be  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  respectively.



$$\therefore \overline{OD} = \frac{1}{2}(\vec{b} + \vec{c}),$$

$$\overline{OE} = \frac{1}{2}(\vec{c} + \vec{a}),$$

$$\text{and } \overline{OF} = \frac{1}{2}(\vec{a} + \vec{b})$$

Also

$$\overline{BC} = \vec{c} - \vec{b}, \quad \overline{CA} = \vec{a} - \vec{c} \quad \text{and} \quad \overline{AB} = \vec{b} - \vec{a}$$

$$\text{Since, } \overline{OD} \perp \overline{BC}, \quad \frac{1}{2}(\vec{b} + \vec{c}) \cdot (\vec{c} - \vec{b}) = 0$$

$$\Rightarrow b^2 = c^2 \quad \dots (i)$$

$$\text{Similarly, } \overline{OE} \perp \overline{CA}, \quad \frac{1}{2}(\vec{a} + \vec{c}) \cdot (\vec{a} - \vec{c}) = 0$$

$$\Rightarrow a^2 = c^2 \quad \dots (ii)$$

From (i) and (ii) we have  $b^2 - a^2 = 0$

$$(\vec{a} + \vec{b}) \cdot (\vec{b} + \vec{a}) = 0 \Rightarrow \frac{1}{2}(\vec{b} + \vec{a}) \cdot (\vec{b} - \vec{a}) = 0$$

$$\Rightarrow \overline{OF} \perp \overline{AB}$$

Hence proved.

## JEE Main/Boards

### JEE Main/Boards

#### Exercise 1

**Q.1** The line  $L_1$  passes through the points (2, -3, 1) and (-1, -2, -4). The line  $L_2$  passes through the point (3, 2, -9) and is parallel to the vector  $4\hat{i} - 4\hat{j} + 5\hat{k}$ .

(i) Find an equation for  $L_1$  in the form  $\vec{r} = \vec{a} + t\vec{b}$

(ii) Prove that  $L_1$  and  $L_2$  are skew.

**Q.2** Two lines have vector equations

$$\vec{r} = \begin{pmatrix} 4 \\ 2 \\ -6 \end{pmatrix} + t \begin{pmatrix} -8 \\ 1 \\ -2 \end{pmatrix} \quad \text{and} \quad \vec{r} = \begin{pmatrix} -2 \\ a \\ -2 \end{pmatrix} + s \begin{pmatrix} -9 \\ 2 \\ -5 \end{pmatrix},$$

Where 'a' is a constant.

(i) Calculate the acute angle between the lines.

(ii) Given that these two lines intersect, find the point of intersection.

**Q.3** The points A and B have position vectors  $\vec{a}$  and  $\vec{b}$  relative to an origin O, where  $\vec{a} = 4\hat{i} + 3\hat{j} - 2\hat{k}$  and  $\vec{b} = -7\hat{i} + 5\hat{j} + 4\hat{k}$

(i) Find the length of AB.

(ii) Use a scalar product to find angle OAB.

**Q.4** The position vectors of the points P and Q with respect to an origin O are  $5\hat{i} + 2\hat{j} - 9\hat{k}$  and  $4\hat{i} + 4\hat{j} - 6\hat{k}$  respectively.

(i) Find the vector equation for the line PQ

The position vector of the point T is  $\hat{i} + 2\hat{j} - \hat{k}$

(ii) Write down a vector equation for the line OT and show that OT is perpendicular to PQ.

It is given that OT intersects PQ.

(iii) Find the position vector of the point of intersection of OT and PQ.

(iv) Hence find the perpendicular distance from O to PQ, giving your answer in an exact form.

**Q.5** ABCD is a parallelogram. The position vectors of A, B and C are given respectively by

$$\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}, \vec{b} = 3\hat{i} - 2\hat{j}, \vec{c} = \hat{i} - \hat{j} - 2\hat{k}$$

- (i) Find the position vector of D.  
(ii) Determine, to the nearest degree, the angle ABC.

**Q.6** The position vectors of three points A, B and C relative to an origin O are given respectively by  $\vec{OA} = 7\hat{i} + 3\hat{j} - 3\hat{k}$ ,  $\vec{OB} = 4\hat{i} + 2\hat{j} - 4\hat{k}$ ,  $\vec{OC} = 5\hat{i} + 4\hat{j} - 5\hat{k}$ .

- (i) Find the angle between AB and AC.  
(ii) Find the area of triangle ABC.

**Q.7** Two lines have vector equations

$$\vec{r} = \hat{i} - 2\hat{j} + 4\hat{k} + \lambda(3\hat{i} + \hat{j} + a\hat{k}) \text{ and}$$

$$\vec{r} = -8\hat{i} + 2\hat{j} + 3\hat{k} + \mu(\hat{i} - 2\hat{j} - \hat{k}),$$

Where 'a' is a constant

- (i) Given that the lines are skew, find the value that a cannot take.  
(ii) Given instead that the lines intersect, find the point of intersection.

**Q.8** Lines,  $L_1$ ,  $L_2$  and  $L_3$  have vector equations

$$L_1 : \vec{r} = (5\hat{i} - \hat{j} - 2\hat{k}) + s(-6\hat{i} + 8\hat{j} - 2\hat{k}),$$

$$L_2 : \vec{r} = (3\hat{i} - 8\hat{j}) + t(\hat{i} + 3\hat{j} + 2\hat{k})$$

$$L_3 : \vec{r} = (2\hat{i} + \hat{j} + 3\hat{k}) + u(3\hat{i} + \hat{j} + \hat{k}).$$

- (i) Calculate the acute angle between  $L_1$  and  $L_2$ .  
(ii) Given that  $L_1$  and  $L_3$  are parallel, find the value of c.  
(iii) Given that  $L_2$  and  $L_3$  intersect, find the value of c

**Q.9** Given that  $\vec{u} = \hat{i} - 2\hat{j} + 3\hat{k}$ ;

$$\vec{v} = 2\hat{i} + \hat{j} + 4\hat{k}; \vec{w} = \hat{i} + 3\hat{j} + 3\hat{k} \text{ and}$$

$$(\vec{u} \cdot \vec{R} - 10)\hat{i} + (\vec{v} \cdot \vec{R} - 20)\hat{j} + (\vec{w} \cdot \vec{R} - 20)\hat{k} = \vec{0}.$$

Find the unknown vector  $\vec{R}$ .

**Q.10** The base vectors  $\vec{a}_1$ ,  $\vec{a}_2$ ,  $\vec{a}_3$  are given in terms of base vectors  $\vec{b}_1$ ,  $\vec{b}_2$ ,  $\vec{b}_3$  as,

$$\vec{a}_1 = 2\vec{b}_1 + 3\vec{b}_2 - \vec{b}_3;$$

$$\vec{a}_2 = \vec{b}_1 - 2\vec{b}_2 + 2\vec{b}_3 \text{ \& } \vec{a}_3 = -2\vec{b}_1 + \vec{b}_2 - 2\vec{b}_3.$$

If  $\vec{F} = 3\vec{b}_1 - \vec{b}_2 + 2\vec{b}_3$ , then express  $\vec{F}$  in terms of  $\vec{a}_1$ ,  $\vec{a}_2$ ,  $\vec{a}_3$ .

**Q.11** If  $\vec{r}$  and  $\vec{s}$  are non zero constant vectors and the scalar b is chosen such that  $|\vec{r} + b\vec{s}|$  is minimum, then show that the value of  $|\vec{b}\vec{s}|^2 + |\vec{r} + b\vec{s}|^2$  is equal to  $|\vec{r}|^2$ .

## Exercise 2

### Single Correct Choice Type

**Q.1** Let  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  be three vectors such that  $|\vec{a}| = |\vec{c}| = 1$ ;  $|\vec{b}| = 4$  and  $|\vec{b} \times \vec{c}| = \sqrt{15}$ . If  $\vec{b} - 2\vec{c} = \lambda\vec{a}$  then a value of  $\lambda$  is

- (A) 1 (B) -1 (C) 2 (D) -4

**Q.2** Vector  $\vec{r}$  which is equally inclined to coordinate axes such that  $|\vec{r}| = 15\sqrt{3}$  is

- (A)  $\hat{i} + \hat{j} + \hat{k}$  (B)  $15(\hat{i} + \hat{j} + \hat{k})$   
(C)  $7(\hat{i} + \hat{j} + \hat{k})$  (D) None of these

**Q.3** For 3 vectors  $\vec{u}$ ,  $\vec{v}$ ,  $\vec{w}$ , which of the following expressions is  $\neq$  to any remaining three.

- (A)  $\vec{u}(\vec{v} \times \vec{w})$  (B)  $(\vec{v} \times \vec{w}) \cdot \vec{u}$   
(C)  $\vec{v}(\vec{u} \times \vec{w})$  (D)  $(\vec{w} \times \vec{u}) \cdot \vec{v}$

**Q.4** If  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ ,  $|\vec{a}| = 3$ ,  $|\vec{b}| = 5$  &  $|\vec{c}| = 7$  then  $\angle\theta$  between  $\vec{a}$  and  $\vec{b}$  is

- (A)  $40^\circ$  (B)  $30^\circ$  (C)  $150^\circ$  (D) None of these

**Q.5** If 2 out of 3 vectors  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  are unit vectors,

$\vec{a} + \vec{b} + \vec{c} = \vec{0}$  and  $2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) + 3 = 0$ , then third vector is length-

- (A) 3 (B) 1 (C) 2 (D) None of these

**Q.6** Let  $\vec{a} + \vec{b}$  is orthogonal to  $\vec{b}$  and  $\vec{a} + 2\vec{b}$  is orthogonal to  $\vec{a}$ , then

- (A)  $|\vec{a}| = \sqrt{2}|\vec{b}|$  (B)  $|\vec{a}| = 2|\vec{b}|$   
(C)  $|\vec{a}| = |\vec{b}|$  (D)  $2|\vec{a}| = |\vec{b}|$

**Q.7** Magnitude of projection of vector  $\hat{i} + 2\hat{j} + \hat{k}$  on vector  $4\hat{i} + 4\hat{j} + 7\hat{k}$  is

- (A) 3 (B)  $3\sqrt{6}$  (C)  $\sqrt{6}/3$  (D) None of these

**Q.8** Magnitude of moment of force  $-2\hat{i} + 6\hat{j} - 8\hat{k}$  acting at point  $2\hat{i} - \hat{j} + 3\hat{k}$  about point  $\hat{i} + 2\hat{j} - \hat{k}$

- (A)  $\sqrt{211}$  (B) 0 (C)  $\sqrt{54}$  (D) None of these

**Q.9** If  $\hat{a}$  &  $\hat{b}$  are unit vectors represented by  $\vec{OA}$  and  $\vec{OB}$ , then unit vector along bisector of  $\angle AOB$  is scalar multiple of

- (A)  $\hat{a} - \hat{b}$  (B)  $\hat{a} \times \hat{b}$  (C)  $\hat{b} \times \hat{a}$  (D) None of these

**Q.10** If  $[2\vec{a} + 4\vec{b} \quad \vec{c} \quad \vec{d}] = \lambda[\vec{a} \quad \vec{c} \quad \vec{d}] + \mu[\vec{b} \quad \vec{c} \quad \vec{d}]$  then  $\lambda + \mu =$

- (A) 6 (B) -6 (C) 10 (D) None of these

## Previous Years' Questions

**Q.1** The volume of the parallelepiped whose sides are given by  $\vec{OA} = 2\hat{i} - 3\hat{j}$ ,  $\vec{OB} = \hat{i} + \hat{j} - \hat{k}$ ,  $\vec{OC} = 3\hat{i} - \hat{k}$ , is

(1983)

- (A)  $\frac{4}{13}$  (B) 4 (C)  $\frac{2}{7}$  (D) None of these

**Q.2** A vector  $\vec{a}$  has components  $2p$  and  $1$  with respect to a rectangular Cartesian system. This system is rotated through a certain angle about the origin in the counter clockwise sense. If, with respect to the new system,  $\vec{a}$  has components  $p+1$  and  $l$ , then

(1986)

- (A)  $p = 0$  (B)  $p = 1$  or  $p = -\frac{1}{3}$   
(C)  $p = -1$  or  $p = \frac{1}{3}$  (D)  $p = 1$  or  $p = -1$

**Q.3** Let  $a, b, c$  be distinct non-negative numbers. If the vectors  $a\hat{i} + a\hat{j} + c\hat{k}$ ,  $\hat{i} + \hat{k}$  and  $\hat{c}\hat{i} + \hat{c}\hat{j} + b\hat{k}$  lie in a plane, then  $c$  is

(1993)

- (A) The Arithmetic Mean of  $a$  and  $b$ .  
(B) The Geometric Mean of  $a$  and  $b$ .  
(C) The Harmonic Mean of  $a$  and  $b$ .  
(D) Equal to zero.

**Q.4** If  $\vec{a}, \vec{b}, \vec{c}$  are non-coplanar unit vectors such that  $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{(\vec{b} + \vec{c})}{\sqrt{2}}$ , then the angle between  $\vec{a}$  and  $\vec{b}$  is

(1995)

- (A)  $\frac{3\pi}{4}$  (B)  $\frac{\pi}{4}$  (C)  $\frac{\pi}{2}$  (D)  $\pi$

**Q.5** If  $\vec{a}$  and  $\vec{b}$  are two unit vectors such that  $\vec{a} + 2\vec{b}$  and  $5\vec{a} - 4\vec{b}$  are perpendicular to each other, then the angle between  $\vec{a}$  and  $\vec{b}$  is

(2002)

- (A)  $45^\circ$  (B)  $60^\circ$  (C)  $\cos^{-1}\left(\frac{1}{3}\right)$  (D)  $\cos^{-1}\left(\frac{2}{7}\right)$

**Q.6** Let  $\vec{V} = 2\vec{i} + \vec{j} - \vec{k}$  and  $\vec{W} = \vec{i} + 3\vec{k}$ . If  $\vec{U}$  is a unit vector, then the maximum value of the scalar triple product  $[\vec{U} \vec{V} \vec{W}]$  is

(2002)

- (A) -1 (B)  $\sqrt{10} + \sqrt{6}$  (C)  $\sqrt{59}$  (D)  $\sqrt{60}$

**Q.7** The unit vector which is orthogonal to the vector  $3\hat{i} + 2\hat{j} + 6\hat{k}$  and is coplanar with the vectors  $2\hat{i} + \hat{j} + \hat{k}$  and  $\hat{i} - \hat{j} + \hat{k}$  is

(2004)

- (A)  $\frac{2\hat{i} - 6\hat{j} + \hat{k}}{\sqrt{41}}$  (B)  $\frac{2\hat{i} - 3\hat{j}}{\sqrt{13}}$   
(C)  $\frac{3\hat{j} - \hat{k}}{\sqrt{10}}$  (D)  $\frac{4\hat{i} + 3\hat{j} - 3\hat{k}}{\sqrt{34}}$

**Q.8** Two adjacent sides of a parallelogram ABCD are given by  $\vec{AB} = 2\hat{i} + 10\hat{j} + 11\hat{k}$  and  $\vec{AD} = -\hat{i} + 2\hat{j} + 2\hat{k}$ . The side AD is rotated by an acute angle  $\alpha$  in the plane of the parallelogram so that AD becomes AD'. If AD' makes a right angle with the side AB, then the cosine of the angle  $\alpha$  is given by

(2010)

- (A)  $\frac{8}{9}$  (B)  $\frac{\sqrt{17}}{9}$  (C)  $\frac{1}{9}$  (D)  $\frac{4\sqrt{5}}{9}$

**Q.9** Let  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$  and  $\vec{c} = \hat{i} - \hat{j} - \hat{k}$  be three vectors. A vector  $\vec{v}$  in the plane of  $\vec{a}$  and  $\vec{b}$ , whose projection on  $\vec{c}$  is  $\frac{1}{\sqrt{3}}$ , is given by

(2011)

**Q.10** Let  $\vec{a}, \vec{b}$  and  $\vec{c}$  be three unit vectors such that  $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{\sqrt{3}}{2}(\vec{b} + \vec{c})$ . If  $\vec{b}$  is not parallel to  $\vec{c}$  then the angle between  $\vec{a}$  and  $\vec{b}$  is:

(2016)

- (A)  $\frac{\pi}{2}$  (B)  $\frac{2\pi}{3}$  (C)  $\frac{5\pi}{6}$  (D)  $\frac{3\pi}{4}$

**Q.11** Let  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  be three non-zero vectors such that no two of them are collinear and  $(\vec{a} \times \vec{b}) \times \vec{c} = \frac{1}{3} |\vec{b}| |\vec{c}| \vec{a}$ . If  $\theta$  is the angle between vectors  $\vec{b}$  and  $|\vec{c}|$  then a value of  $\sin \theta$  is **(2015)**

- (A)  $\frac{-\sqrt{2}}{3}$  (B)  $\frac{2}{3}$  (C)  $-\frac{2\sqrt{3}}{3}$  (D)  $\frac{2\sqrt{3}}{3}$

**Q.12** If  $[\vec{a} \times \vec{b} \vec{b} \times \vec{c} \times \vec{a}] = \lambda [\vec{a} \vec{b} \vec{c}]^2$  then  $\lambda$  is equal to **(2014)**

- (A) 1 (B) 3 (C) 0 (D) 1

**Q.13** If the vectors  $\vec{AB} = 3\hat{j} + 4\hat{k}$  and  $\vec{AC} = 2\hat{j} + 4\hat{k}$  are the sides of a triangle ABC, then the length of the median through A is **(2013)**

- (A)  $\sqrt{72}$  (B)  $\sqrt{33}$  (C)  $\sqrt{45}$  (D)  $\sqrt{18}$

**Q.14** Let  $\hat{a}$  and  $\hat{b}$  be two unit vectors. If the vectors  $\vec{c} = \vec{a} + 2\hat{b}$  and  $\vec{d} = 5\hat{a} - 4\hat{b}$  are perpendicular to each other then the angle between  $\hat{a}$  and  $\hat{b}$  is **(2012)**

- (A)  $\frac{\pi}{6}$  (B)  $\frac{\pi}{2}$  (C)  $\frac{\pi}{3}$  (D)  $\frac{\pi}{4}$

**Q.15** If the vectors  $\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$  and  $\vec{c} = \lambda\hat{i} + \hat{j} + \mu\hat{k}$  are mutually orthogonal then  $(\lambda, \mu) =$  **(2010)**

- (A) (2, -3) (B) (-2, 3) (C) (3, -2) (D) (-3, 2)

**Q.16** Let  $\vec{a} = \hat{j} - \hat{k}$ . Then vector  $\vec{b}$  satisfying  $\vec{a} \times \vec{b} + \vec{c} = \vec{0}$  and  $\vec{a} \cdot \vec{b}$  is **(2010)**

- (A)  $2\hat{i} - \hat{j} + 2\hat{k}$  (B)  $\hat{i} - \hat{j} + 2\hat{k}$   
(C)  $\hat{i} + \hat{j} - 2\hat{k}$  (D)  $-\hat{i} + \hat{j} - 2\hat{k}$

**Q.17** If  $\vec{a} = \frac{1}{\sqrt{10}}(3\hat{i} + \hat{k})$  and  $\vec{b} = \frac{1}{7}(2\hat{i} + 3\hat{j} - 6\hat{k})$ , then the value of  $(2\vec{a} - \vec{b}) \cdot [(\vec{a} \times \vec{b}) \times (\vec{a} + 2\vec{b})]$  is **(2011)**

- (A) -3 (B) 5 (C) 3 (D) -5

**Q.18** The vector  $\vec{a}$  and  $\vec{b}$  are not perpendicular and  $\vec{a}$  and  $\vec{b}$  are two vectors satisfying:  $\vec{b} \times \vec{c} = \vec{b} \times \vec{d}$  and  $\vec{a} \cdot \vec{d} = 0$ . Then the vector is equal to **(2011)**

- (A)  $\vec{c} + \left( \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \right) \vec{b}$  (B)  $\vec{b} + \left( \frac{\vec{b} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \right) \vec{c}$   
(C)  $\vec{c} - \left( \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \right) \vec{b}$  (D)  $\vec{b} + \left( \frac{\vec{b} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \right) \vec{c}$

## JEE Advanced/Boards

### Exercise 1

**Q.1** What will be the value of  $\frac{(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2}{2a^2 b^2}$ ?

**Q.2** What will be the area of the triangle determined by the vectors  $3\hat{i} + 4\hat{j}$  and  $-5\hat{i} + 7\hat{j}$ ?

**Q.3** What will be the value of  $a$  if points whose position vectors are  $60\hat{i} + 3\hat{j}$ ,  $40\hat{i} - 8\hat{j}$ ,  $a\hat{i} - 52\hat{j}$  are collinear?

**Q.4** What will be the angle between diagonals which adjacent sides of  $\square$  are along  $\vec{a} = \hat{i} + 2\hat{j}$  &  $\vec{b} = 2\hat{i} + \hat{j}$ ?

**Q.5** What will be the angle between  $\vec{a}$  and  $\vec{b}$  if  $\vec{a}$  &  $\vec{b}$  are unit vectors such that  $\vec{a} + 3\vec{b}$  is  $\perp$  to  $7\vec{a} - 5\vec{b}$ ?

**Q.6** If the unit vectors  $\hat{A}$  and  $\hat{B}$  are inclined at  $\pi$  then what will be the value of  $|\hat{A} - \hat{B}|/2$ ?

**Q.7** A particle acted upon by forces  $3\hat{i} + 2\hat{j} + 5\hat{k}$  and  $2\hat{i} + \hat{j} + 3\hat{k}$  is displaced from a point P to a point Q whose respective position vectors are  $2\hat{i} + \hat{j} + 3\hat{k}$  and  $4\hat{i} + 3\hat{j} + 7\hat{k}$ . What will be the work done by the force?

**Q.8** A force  $F = 6\hat{i} + \lambda\hat{j} + 4\hat{k}$  acting on a particle displaces it from A (3,4,5) to B(1,1,1). If the work done is 2 units, then What will be the value of  $\lambda$ ?

**Q.9** What will be the length of longer diagonal of  $\Delta$  constructed on  $5\vec{a} + 2\vec{b}$  &  $\vec{a} - 3\vec{b}$ . Given  $|\vec{b}| = 3$  &  $|\vec{a}| = 2\sqrt{2}$  and angle between  $\vec{a}$  &  $\vec{b}$  is  $\pi/4$

**Q.10** The vectors  $\hat{i} + \hat{j} + 3\hat{k}$  is rotated through an angle  $\theta$  and doubled in magnitude, then it becomes  $4\hat{i} + (4x - 2)\hat{j} + 2\hat{k}$ . Find  $x$ ?

## Exercise 2

### Single Correct Choice Type

**Q.1** Moment of couple formed by forces  $5\hat{i} + \hat{j}$  &  $-5\hat{i} + \hat{j}$  acting at  $[9, -1, 2]$  and  $[3, -2, 1]$

- (A)  $-\hat{i} + 5\hat{j} + \hat{k}$  (B)  $\hat{i} - \hat{j} - 5\hat{k}$   
(C)  $2\hat{i} - 2\hat{j} - 10\hat{k}$  (D)  $-2\hat{i} - 2\hat{j} + 10\hat{k}$

**Q.2** Let  $\vec{a}, \vec{b}, \vec{c}$  be vectors such that

$\vec{a} \cdot (\vec{b} + \vec{c}) + \vec{b} \cdot (\vec{c} + \vec{a}) + \vec{c} \cdot (\vec{a} + \vec{b}) = 0$  and  
 $|\vec{a}| = 1, |\vec{b}| = 4, |\vec{c}| = 8$  then  $|\vec{a} + \vec{b} + \vec{c}|$  equals

- (A) 13 (B) 81 (C) 9 (D) None of these

**Q.3** Position vectors of A and B are  $2\hat{i} + 2\hat{j} + \hat{k}$  and  $2\hat{i} + 4\hat{j} + 4\hat{k}$ . Length of internal bisector of  $\angle BOA$  of  $\Delta AOB$  is

- (A)  $\sqrt{\frac{136}{9}}$  (B)  $\sqrt{\frac{136}{9}}$  (C)  $\frac{20}{3}$  (D) None of these

**Q.4** If  $\vec{a} + 2\vec{b} + 3\vec{c} = 0$ , then  $\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}$  is equal to

- (A)  $6(\vec{b} \times \vec{c})$  (B)  $6(\vec{a} \times \vec{b})$   
(C)  $6(\vec{c} \times \vec{a})$  (D) None of these

**Q.5** Value of  $|\vec{a} - \vec{b}, \vec{b} - \vec{c}, \vec{c} - \vec{a}|$  where  $|\vec{a}| = 1, |\vec{b}| = 2$ , and  $|\vec{c}| = 3$  is

- (A) 1 (B) -6 (C) 0 (D) None of these

**Q.6** If P and Q be two given points on the curve  $y = x + 1/x$  such that  $OP \cdot I = 1$  and  $OQ \cdot I = -1$  where I is a unit vector along the x-axis, then the length of vector  $2OP + 3OQ$  is

- (A)  $5\sqrt{5}$  (B)  $3\sqrt{5}$  (C)  $2\sqrt{5}$  (D)  $\sqrt{5}$

**Q.7** Let A, B, C be three vectors such that  $A \cdot (B + C) + B \cdot C = 0$  and  $|A| = 1, |B| = 4, |C| = 8$ , then  $|A + B + C|$  equals

- (A) 13 (B) 81 (C) 9 (D) 5

**Q.8** If the unit vectors  $\hat{A}$  and  $\hat{B}$  are inclined at an angle  $2\theta$  and  $|\hat{A} - \hat{B}| \leq 1$ , then for  $\theta \in [0, \pi]$ ,  $\theta$  may lie in the interval

- (A)  $[\pi/6, \pi/3]$  (B)  $[\pi/6, \pi/2]$   
(C)  $[5\pi/6, \pi]$  (D)  $[\pi/2, 5\pi/6]$

**Q.9** If unit vectors  $\hat{A}$  and  $\hat{B}$  such that  $STP$   $|\hat{A} \cdot \hat{B} \cdot \hat{A} \times \hat{B}| = 1/4$  then  $\hat{A}$  and  $\hat{B}$  are inclined

- (A)  $\pi/6$  (B)  $\pi/2$  (C)  $\pi/3$  (D)  $\pi/4$

**Q.10** If  $\hat{A}$  and  $\hat{B}$  unit vectors then greatest value of  $|\hat{A} - \hat{B}| + |\hat{A} + \hat{B}|$  is

- (A) 2 (B) 4 (C)  $2\sqrt{2}$  (D)  $\sqrt{2}$

## Previous Years' Questions

**Q.1** (i) If C be a given non zero scalar and  $\vec{A}$  and  $\vec{B}$  be given non-zero vectors such that  $\vec{A} \perp \vec{B}$ , find the vector  $\vec{X}$  which satisfies the equation  $\vec{A} \cdot \vec{X} = c$  and  $\vec{A} \times \vec{X} = \vec{B}$ .

(ii)  $\vec{A}$  vector A has components  $A_1, A_2, A_3$  in a right-handed rectangular Cartesian coordinate system  $oxyz$ . The coordinate system is rotated about the x-axis through an angle  $\frac{\pi}{2}$ . Find the components of A in the new coordinate system, in terms of  $A_1, A_2, A_3$ . (1983)

**Q.2** If vectors  $\vec{a}, \vec{b}, \vec{c}$  are coplanar, show that

$$\begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \\ \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} \cdot \vec{a} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \end{vmatrix} = 0$$

(1989)

**Q.3** Let  $\vec{A} = 2\hat{i} + \hat{k}$ ,  $\vec{B} = \hat{i} + \hat{j} + \hat{k}$ , and  $\vec{C} = 4\hat{i} - 3\hat{j} + 7\hat{k}$ . Determine a vector  $\vec{R}$  satisfying  $\vec{R} \times \vec{B} = \vec{C} \times \vec{B}$  and  $\vec{R} \cdot \vec{A} = 0$ . (1990)

**Q.4** If  $\vec{a}, \vec{b}, \vec{c}, \vec{d}$  are four distinct vectors satisfying the conditions  $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$  and  $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$  then prove that  $\vec{a} \cdot \vec{b} + \vec{c} \cdot \vec{d} \neq \vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{d}$ . (2004)

**Q.5** Incident ray is along the unit vector  $\hat{v}$  and the reflected ray is along the unit vector  $\hat{w}$ . The normal is along unit vector  $\hat{a}$  outwards. Express  $\hat{w}$  in terms of  $\hat{a}$  and  $\hat{v}$ . (2005)

**Q.6** Let  $\vec{A}$  be vector parallel to line of intersection of planes  $P_1$  and  $P_2$  through origin.  $P_1$  is parallel to the vectors  $2\hat{j} + 3\hat{k}$  and  $4\hat{j} - 3\hat{k}$  and  $P_2$  is parallel to  $\hat{j} - \hat{k}$  and  $3\hat{i} + 3\hat{j}$ , then the angle between vector  $\vec{A}$  and  $2\hat{i} + \hat{j} - 2\hat{k}$  is (2006)

- (A)  $\frac{\pi}{2}$  (B)  $\frac{\pi}{4}$  (C)  $\frac{\pi}{4}$  (D)  $\frac{3\pi}{4}$

**Q.7** The vector(s) which is /are coplanar with vectors  $\hat{i} + \hat{j} + 2\hat{k}$  and  $\hat{i} + 2\hat{j} + \hat{k}$ , are perpendicular to the vector  $\hat{i} + \hat{j} + \hat{k}$  is/are (2011)

- (A)  $\hat{j} - \hat{k}$  (B)  $-\hat{i} + \hat{j}$  (C)  $\hat{i} - \hat{j}$  (D)  $-\hat{j} + \hat{k}$

**Q.8** If  $\vec{a}$  and  $\vec{b}$  are vectors in space by  $\vec{a} = \frac{\hat{i} - 2\hat{j}}{\sqrt{5}}$  and  $\vec{b} = \frac{2\hat{i} + \hat{j} + 3\hat{k}}{\sqrt{14}}$ , then the value of  $(2\vec{a} + \vec{b}) \cdot [(\vec{a} \times \vec{b}) \times (\vec{a} - 2\vec{b})]$  is (2010)

**Q.9** Let  $\vec{a} = -\hat{i} - \hat{k}$ ,  $\vec{b} = -\hat{i} + \hat{j}$ , and  $\vec{c} = \hat{i} + 2\hat{j} + 3\hat{k}$  be three given vectors. If  $\vec{r}$  is a vector such that  $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$ ,  $\vec{r} \cdot \vec{a} = 0$ , then the value of  $\vec{r} \cdot \vec{b}$  is..... (2011)

**Q.10** Let  $\hat{u} = u_1\hat{i} + u_2\hat{j} + u_3\hat{k}$  be a unit vector in  $R^2$  and  $\hat{w} = \frac{1}{\sqrt{6}}(\hat{i} + \hat{j} + 2\hat{k})$ . Given that there exists a vector  $\vec{v}$  in  $R^2$  such that  $|\hat{u} \times \vec{v}| = 1$  and  $\hat{w} \cdot (\hat{u} \times \vec{v}) = 1$ . Which of the following statements (s) is (are) correct? (2016)

- (A) There is exactly one choice for such  $\vec{v}$   
 (B) There are infinitely many choices for such  $\vec{v}$   
 (C) If  $\hat{u}$  lies in the xy-plane then  $|u_1| = |u_2|$   
 (D) If  $\hat{u}$  lies in the xz-plane then  $2|u_1| = |u_3|$

**Q.11** Let  $\Delta PQR$  be a triangle. Let  $\vec{a} = \overrightarrow{QR}$ ,  $\vec{b} = \overrightarrow{RP}$  and  $\vec{c} = \overrightarrow{PQ}$ . If  $|\vec{a}| = 12$ ,  $|\vec{b}| = 4\sqrt{3}$  and  $\vec{b} \cdot \vec{c} = 24$ , then which of the following is (are) true (2015)

- (A)  $\frac{|\vec{c}|^2}{2} - |\vec{a}| = 12$   
 (B)  $\frac{|\vec{c}|^2}{2} - |\vec{a}| = 30$   
 (C)  $|\vec{a} \times \vec{b} + \vec{c} \times \vec{a}| = 48\sqrt{3}$   
 (D)  $\vec{a} \cdot \vec{b} = -72$

**Q.12** Let  $\vec{a} = -\hat{i} - \hat{k}$ ,  $\vec{b} = -\hat{i} + \hat{j}$  and  $\vec{c} = \hat{i} + 2\hat{j} + 3\hat{k}$  be three given vectors. If  $\vec{r}$  is a vector such that  $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$  and  $\vec{r} \cdot \vec{a} = 0$  then the value of  $\vec{r} \cdot \vec{b}$  is (2011)

**Q.13** If  $\vec{a}$  and  $\vec{b}$  are vectors in space given by  $\vec{a} = \frac{\hat{i} - 2\hat{j}}{\sqrt{5}}$  and  $\vec{b} = \frac{2\hat{i} + \hat{j} + 3\hat{k}}{\sqrt{14}}$  then the value of  $(2\vec{a} + \vec{b}) \cdot [(\vec{a} \times \vec{b}) \times (\vec{a} - 2\vec{b})]$  is (2010)

**Q.14** Let  $\vec{a}, \vec{b}$  and  $\vec{c}$  be three non-coplanar unit vectors such that the angle between every pair of them is  $\frac{\pi}{3}$ . If  $\vec{a} \times \vec{b} + \vec{b} \times \vec{c} = p\vec{a} + q\vec{b} + r\vec{c}$  where  $p, q$  and  $r$  are scalars, then the value of  $\frac{p^2 + 2q^2 + r^2}{q^2}$  is (2014)

**Q.15** Let  $\overrightarrow{PR} = 3\hat{i} + \hat{j} - 2\hat{k}$  and  $\overrightarrow{SQ} = \hat{i} - 3\hat{j} - 4\hat{k}$  determine diagonals of a parallelogram PQRS and  $\overrightarrow{PT} = \hat{i} + 2\hat{j} - 3\hat{k}$  be another vector. Then the volume of the parallelepiped determined by the vectors  $\overrightarrow{PT}, \overrightarrow{PQ}$  and  $\overrightarrow{PS}$  is (2013)

(A) 5 (B) 20 (C) 10 (D) 30



**Q.16** A line  $\ell$  passing through the origin is perpendicular to the lines

$$\ell_1 : (3+t)\hat{i} + (-1+2t)\hat{j} + (4+2t)\hat{k}, -\infty < t < \infty$$

$$\ell_2 : (3+2s)\hat{j} + (3+2s)\hat{j} + (2+s)\hat{k}, -\infty < s < \infty$$

Then, the coordinate (s) of the point(s) on  $\ell_2$  at a distance of  $\sqrt{17}$  from the point of intersection of  $\ell$  and  $\ell_2$  is (are) **(2013)**

(A)  $\left(\frac{7}{3}, \frac{7}{3}, \frac{5}{3}\right)$  (B)  $(-1, -1, 0)$

(C)  $(1, 1, 1)$  (D)  $\left(\frac{7}{9}, \frac{7}{9}, \frac{8}{9}\right)$

**Q.17** Consider the set of eight vectors  $V = \{a\hat{i} + b\hat{j} + c\hat{k} : a, b, c \in \{-1, 1\}\}$ . Three non-coplanar vectors can be chosen from  $V$  in  $2^n$  ways. Then  $p$  is **(2013)**

**Q.18** Match list I with list II and select the correct answer using the code given below the lists : **(2013)**

List I	List II
(i) Volume of parallelepiped by vectors $\vec{a}, \vec{b}$ and $\vec{c}$ is 2. Then the volume of the parallelepiped determined by vectors $2(\vec{a} \times \vec{b}), 3(\vec{b} \times \vec{c})$ and $(\vec{c} \times \vec{a})$ is	(p) 100
(ii) Volume of parallelepiped determined by vectors $\vec{a}, \vec{b}$ and $\vec{c}$ is 5. Then the volume of the parallelepiped determined by vectors $3(\vec{a} + \vec{b}), (\vec{b} + \vec{c})$ and $2(\vec{c} + \vec{a})$ is	(q) 30
(iii) Area of a triangle with adjacent sides determined by vectors $\vec{a}$ and $\vec{b}$ is 20. Then the area of the triangle with adjacent sides determined by vectors $(2\vec{a} + 3\vec{b})$ and $(\vec{a} - \vec{b})$ is	(r) 24
(iv) Area of a parallelogram with adjacent sides determined by vectors $\vec{a}$ and $\vec{b}$ is 30. Then the area of the parallelogram with adjacent sides determined by vectors $(\vec{a} + \vec{b})$ and $\vec{a}$ is	(s) 60

(A)  $i \rightarrow s, ii \rightarrow q, iii \rightarrow r, iv \rightarrow p$

(B)  $i \rightarrow q, ii \rightarrow r, iii \rightarrow p, iv \rightarrow s$

(C)  $i \rightarrow r, ii \rightarrow s, iii \rightarrow p, iv \rightarrow q$

(D)  $i \rightarrow p, ii \rightarrow s, iii \rightarrow r, iv \rightarrow q$

**Q.19** If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are unit vectors satisfying

$$|\vec{a} - \vec{b}|^2 + |\vec{b} - \vec{c}|^2 + |\vec{c} - \vec{a}|^2 = 9 \text{ then } |2\vec{a} + 5\vec{b} + 5\vec{c}| \text{ is } \textbf{(2012)}$$

**Q.20** If  $\vec{a}$  and  $\vec{b}$  are vectors such that

$$|\vec{a} + \vec{b}| = \sqrt{29} \text{ and } \vec{a} \times (2\hat{i} + 3\hat{j} + 4\hat{k}) = (2\hat{i} + 3\hat{j} + 4\hat{k}) \times \vec{b}$$

then a possible value of  $(\vec{a} + \vec{b}) \cdot (-7\hat{i} + 2\hat{j} + 3\hat{k})$  is **(2012)**

(A) 0 (B) 3 (C) 4 (D) 8

**Q.21** Let  $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} - \hat{j} + \hat{k}$  and  $\vec{c} = \hat{i} - \hat{j} + \hat{k}$  be three vectors. A vector  $\vec{v}$  in the plane of  $\vec{a}$  and  $\vec{b}$  whose projection on  $\vec{c}$  is  $\frac{1}{\sqrt{3}}$ , is given by **(2011)**

(A)  $\hat{i} + 3\hat{j} + 3\hat{k}$  (B)  $-3\hat{i} - 3\hat{j} + \hat{k}$

(C)  $3\hat{i} - \hat{j} + 3\hat{k}$  (D)  $\hat{i} - 3\hat{j} - 3\hat{k}$

**Q.22** The vector(s) which is/are coplanar with vectors  $\hat{i} + \hat{j} + 2\hat{k}$  and  $\hat{i} + 2\hat{j} + \hat{k}$ , and perpendicular to the vector  $\hat{i} + \hat{j} + \hat{k}$  is/are to **(2011)**

(A)  $\hat{j} - \hat{k}$  (B)  $-\hat{i} + \hat{j}$  (C)  $\hat{i} - \hat{j}$  (D)  $-\hat{j} + \hat{k}$



# PlancEssential Questions

## JEE Main/Boards

### Exercise 1

Q.2      Q.4      Q.7      Q.8  
Q.10

### Exercise 2

Q.3      Q.8      Q.10

### Previous Years' Questions

Q.2      Q.8      Q.9

## JEE Advanced/Boards

### Exercise 1

Q.2      Q.9      Q.12      Q.15

### Exercise 2

Q.1      Q.3

### Previous Years' Questions

Q.1      Q.5      Q.6      Q.8  
Q.10

## Answer Key

## JEE Main/Boards

### Exercise 1

**Q.1** (i)  $r = (2\hat{i} - 3\hat{j} + \hat{k})$  or  $-\hat{i} - 2\hat{j} - 4\hat{k}$  +  $t(3\hat{i} - \hat{j} + 5\hat{k})$

**Q.2** (i)  $15^\circ$  (15.38.....), 0.268 rad (ii)  $a = 1$  and intersection is  $(-20, 5, -12)$

**Q.3** (i)  $\sqrt{161}$  (ii)  $43^\circ$

**Q.4** (i)  $r = (\text{either point}) + t(\hat{i} - 2\hat{j} - 3\hat{k})$  or  $-\hat{i} + 2\hat{j} + 3\hat{k}$ , (ii)  $s(\hat{i} + 2\hat{j} - \hat{k})$  (iii)  $3\hat{i} + 6\hat{j} - 3\hat{k}$  (iv)  $\sqrt{54}$

**Q.5** (i)  $2\hat{j} + \hat{k}$  (ii)  $86^\circ$

**Q.6** (i)  $45.3^\circ$  (ii) 3.54

**Q.7** (i) A cannot be 2. (ii)  $-5\hat{i} - 4\hat{j}$

**Q.8** (i)  $68.5^\circ$  (ii)  $c = -4$  (iii)  $c = -3$

**Q.9**  $-\hat{i} + 2\hat{j} + 5\hat{k}$

**Q.10**  $\vec{F} = 2\vec{a}_1 + 5\vec{a}_2 + 3\vec{a}_3$

### Exercise 2

#### Single Correct Choice Type

**Q.1** D

**Q.2** B

**Q.3** C

**Q.4** B

**Q.5** B

**Q.6** A

**Q.7** D

**Q.8** B

**Q.9** A

**Q.10** A

## Previous Years' Questions

<b>Q.1</b> B	<b>Q.2</b> B	<b>Q.3</b> B	<b>Q.4</b> A	<b>Q.5</b> B	<b>Q.6</b> C
<b>Q.7</b> C	<b>Q.8</b> A	<b>Q.9</b> $3\hat{i} - \hat{j} + 3\hat{k}$	<b>Q.10</b> C	<b>Q.11</b> D	<b>Q.12</b> A
<b>Q.13</b> B	<b>Q.14</b> C	<b>Q.15</b> D	<b>Q.16</b> D	<b>Q.17</b> D	<b>Q.18</b> C

## JEE Advanced/Boards

### Exercise 1

<b>Q.1</b> $\frac{1}{2}$	<b>Q.2</b> $\frac{41}{2}$	<b>Q.3</b> -40	<b>Q.4</b> $90^\circ$ and $90^\circ$	<b>Q.5</b> $\frac{\pi}{3}$	<b>Q.6</b> 1
<b>Q.7</b> 48 units	<b>Q.8</b> -10	<b>Q.9</b> $\sqrt{593}$	<b>Q.10</b> 2 or $-\frac{2}{3}$		

### Exercise 2

#### Single Correct Choice Type

<b>Q.1</b> A	<b>Q.2</b> C	<b>Q.3</b> A	<b>Q.4</b> A	<b>Q.5</b> C	<b>Q.6</b> D
<b>Q.7</b> C	<b>Q.8</b> C	<b>Q.9</b> C	<b>Q.10</b> C		

## Previous Years' Questions

<b>Q.1</b> (i) $\vec{X} = \left( \frac{c}{ \vec{A} ^2} \right) \vec{A} - \left( \frac{1}{ \vec{A} ^2} \right) (\vec{A} \times \vec{B})$	(ii) $(A_2\hat{i} - A_1\hat{j} + A_3\hat{k})$	<b>Q.3</b> $-\hat{i} - 8\hat{j} + 2\hat{k}$			
<b>Q.5</b> $\hat{w} = \hat{v} - 2(\hat{a} \cdot \hat{v})\hat{a}$	<b>Q.6</b> B, D	<b>Q.7</b> A, D	<b>Q.8</b> 5	<b>Q.9</b> 9	
<b>Q.10</b> B, C	<b>Q.11</b> A, B, C	<b>Q.12</b> 9	<b>Q.13</b> 5	<b>Q.14</b> 4	<b>Q.15</b> C
<b>Q.16</b> B, D	<b>Q.17</b> 5	<b>Q.18</b> C	<b>Q.19</b> 3	<b>Q.20</b> C	<b>Q.21</b> C
<b>Q.22</b> A, D					

## Solutions

### JEE Main/Boards

#### Exercise 1

**Sol 1:** (i) For (either point) + t(diff b/w vectors)

$$r = (2i - 3j + k \text{ or } -i - 2j - 4k) + t(3i - j + 5k)$$

$$(ii) L(2)(r) = 3i + 2j - 9k + s(4i - 4j + 5k)$$

L(1) and L(2) must be of form  $r = a + bt$

$$2 + 3t = 3 + 4s, -3 - t = 2 - 4s, 1 + 5t = -9 + 5s$$

$$(t, s) = (+/-3, 2) \text{ or } (-/+1, 1) \text{ or } (-/+9, -7)$$

$$\text{Or } (+/-4, 2) \text{ or } (0, 1) \text{ or } (-/+8, -7)$$

**Sol 2:** (i) Angle between the lines

$$\cos \theta = \frac{-8 \times 9 + 1 \times 2 + (-2) \times (-5)}{\sqrt{64 + 1 + 4} \sqrt{81 + 4 + 25}} = \frac{84}{\sqrt{69} \sqrt{110}} = 0.9641$$

$$\Rightarrow \theta = \cos^{-1}(0.9641) = 15.38 \text{ degree}$$

(ii) Let P be the point of intersection

$$\text{Equation of lines are } \frac{x-4}{-8} = \frac{y-2}{1} = \frac{z+6}{-2} = r_1$$

$$\text{Point P be } (-8r_1 + 4, r_1 + 2, -2r_1 - 6) \quad \dots(i)$$

$$\text{Similarly for second line } \frac{x+2}{-9} = \frac{y-a}{2} = \frac{z+2}{-5} = r_2$$

$$\text{The point P be } (-9r_2 - 2, 2r_2 - a, -5r_2 - 2) \quad \dots(ii)$$

From (i) and (ii), we get

$$-8r_1 + 4 = -9r_2 - 2$$

$$r_1 + 2 = 2r_2 + a$$

$$-2r_1 - 6 = -5r_2 - 2$$

On solving, we get  $r_1 = 3, r_2 = 2$  and  $a = 1$

The points of intersection is  $(-20, 5, -12)$

**Sol 3:** (i) Find  $\vec{a} - \vec{b}$  or  $\vec{b} - \vec{a}$  irrespective of label  
(expect  $11\hat{i} - 2\hat{j} - 6\hat{k}$  or  $-11\hat{i} + 2\hat{j} + 6\hat{k}$ )

$$\text{Magnitude of vector} = \sqrt{161}$$

(ii) Using  $(\vec{AO} \text{ or } \vec{OA})$  and  $(\vec{AB} \text{ or } \vec{BA})$

$$\cos \theta = \frac{\text{Scalar product of any two vector}}{\text{Product of their moduli}}$$

$$\begin{aligned} & \frac{(4\hat{i} + 3\hat{j} - 2\hat{k}) \cdot (11\hat{i} - 2\hat{j} - 6\hat{k})}{\sqrt{4^2 + 3^2 + 2^2} \sqrt{11^2 + 2^2 + 6^2}} \\ &= \frac{44 - 6 + 12}{\sqrt{29} \sqrt{161}} = \frac{50}{\sqrt{29} \sqrt{161}} = 43^\circ \end{aligned}$$

**Sol 4:** (i) For (either point) + t(diff between position vectors)

$$(ii) r = s(i + 2j - k) \text{ or } (i + 2j - k) + s(i + 2j - k)$$

Evaluate scalar product of  $i + 2j - k$  and their dir vect in (i)

$$\text{Show as } (1 \times 1 \text{ or } 1) + (2 \times 2 \text{ or } -4) + (-1 \times -3 \text{ or } 3)$$

$$= 0 \text{ and}$$

$$(iii) \text{ Obtain } t = -2 \text{ or } s = 3 \text{ (possibly } -3 \text{ or } 2 \text{ or } -2)$$

Check if  $t = 2, 1$  or  $-1$

Subst. into eqn AB or OT and to produce  $3i + 6j - 3k$

(iv)  $|\vec{OC}|$  is to be found, where C is their point of intersection

$$|\vec{OC}| = \sqrt{54}$$

**Sol 5:** (i)  $\vec{OD} = \vec{OA} + \vec{AD}$  or  $\vec{OB} + \vec{BC} + \vec{CD}$  AEF

$$\vec{AD} = \vec{BC} \text{ or } \vec{CD} = \vec{BA}$$

$$\vec{OD} = 2\hat{i} + \hat{k}$$

$$(ii) \vec{AB} \cdot \vec{CB} = |\vec{AB}| |\vec{CB}| \cos \theta \Rightarrow \cos \theta = 86^\circ$$

**Sol 6:** (i) Work out  $\vec{b} - \vec{a}$  or  $\vec{a} - \vec{b}$  or  $\vec{c} - \vec{a}$  or  $\vec{a} - \vec{c}$

$$= \pm(-3\hat{i} - \hat{j} - \hat{k}) \text{ or } \pm(-2\hat{i} + \hat{j} - 2\hat{k})$$

Use cosine rule and find angle as  $45.3^\circ$

$$(ii) \text{ Use of } \frac{1}{2} |\vec{AB}| \times |\vec{AC}| \sin \theta$$

$$= \frac{1}{2} (\sqrt{11}) (3) \sin 45.3^\circ = 3.54$$

**Sol 7:** (i) Produce at least 2 of the 3 relevant eqns in  $\lambda$  and  $\mu$

Solving we get

1<sup>st</sup> solution:  $\lambda = -2$  or  $\mu = 3$

2<sup>nd</sup> solution:  $\mu = 3$  or  $\lambda = -2$

Substitute their  $\lambda$  and  $\mu$  into 3<sup>rd</sup> eqn and find 'a'

We get  $a=2$  but  $a$  cannot be 2

(ii) Subst their  $\lambda$  or  $\mu$  (& pass a) into either line eqn

Point of intersection is  $-5\hat{i} - 4\hat{j}$

$$\begin{aligned}\text{Sol 8: (i) } \cos\theta &= \frac{-6 \times 1 + 8 \times 3 - 2 \times 2}{\sqrt{36 + 64 + 4} \sqrt{1 + 9 + 4}} \\ &= \frac{14}{\sqrt{104} \sqrt{14}} = 68.47^\circ\end{aligned}$$

(ii) Since, and are parallel

$$\cos\phi = \frac{-6 \times 3 + 8 \times C - 2 \times 1}{\sqrt{104} \sqrt{9 + C^2 + 1}} = 1$$

$$\Rightarrow 8C - 20 = \sqrt{104} \sqrt{10 + C^2}$$

$$\Rightarrow (8C - 20)^2 = 104(10 + C^2)$$

$$\Rightarrow 64C^2 + 400 - 320C = 1040 + 104C^2$$

$$\Rightarrow 40C^2 + 320C + 640 = 0$$

$$\Rightarrow C^2 + 8C + 16 = 0$$

$$\Rightarrow (C + 4)^2 = 0 \Rightarrow C = -4$$

$$\text{(iii) } L_2 \equiv \frac{x-3}{1} = \frac{y+3}{3} = \frac{z-0}{2} = m$$

Any point  $(m+3, 3m-8, 2m)$

$$L_3 \equiv \frac{x-2}{3} = \frac{y-1}{C} = \frac{z-3}{1} = n$$

Any point  $(3n+2, Cn+1, n+3)$

If  $L_2$  and  $L_3$  intersect, then

$$m+3 = 3n+2 \quad \dots \text{(i)}$$

$$3m-8 = Cn+1 \quad \dots \text{(ii)}$$

$$2m = n+3 \quad \dots \text{(iii)}$$

On solution, we get  $C = -3$

**Sol 9:** Let  $\vec{R} = R_1\hat{i} + R_2\hat{j} + R_3\hat{k}$

$$(\vec{u} \cdot \vec{R} - 10)\hat{i} + (\vec{v} \cdot \vec{R} - 20)\hat{j} + (\vec{w} \cdot \vec{R} - 20)\hat{k} = 0$$

$$\begin{aligned}(R_1 - 2R_2 + 3R_3 - 10)\hat{i} + (2R_1 + R_2 + 4R_3 - 20)\hat{j} \\ + (R_1 + 3R_2 + 3R_3 - 20)\hat{k} = 0\end{aligned}$$

$$\Rightarrow R_1 - 2R_2 + 3R_3 = 10 \quad \dots \text{(i)}$$

$$2R_1 + R_2 + 4R_3 = 20 \quad \dots \text{(ii)}$$

$$R_1 + 3R_2 + 3R_3 = 20 \quad \dots \text{(iii)}$$

On solving, we get  $R_1 = -1, R_2 = 2, R_3 = 5$

$$\vec{R} = -\hat{i} + 2\hat{j} + 5\hat{k}$$

$$\text{Sol 10: } \vec{a}_1 = 2\vec{b}_1 + 3\vec{b}_2 - \vec{b}_3$$

$$\vec{a}_2 = \vec{b}_1 - 2\vec{b}_2 + 2\vec{b}_3$$

$$\vec{a}_3 = 3\vec{b}_1 - \vec{b}_2 + 2\vec{b}_3$$

On solving, we get  $\vec{F} = 2\vec{a}_1 + 5\vec{a}_2 + 3\vec{a}_3$

$$\text{Sol 11: } |\vec{r} + b\vec{s}|^2 = |\vec{r}|^2 + b^2|\vec{s}|^2 + 2b\vec{r} \cdot \vec{s}$$

$|\vec{r} + b\vec{s}|$  is minimum when  $\vec{r} \cdot \vec{s} = -|\vec{r}||\vec{s}|$  and  $\vec{r} = -b\vec{s}$

$$\begin{aligned}\text{Then, } |\vec{b}\vec{s}|^2 + |\vec{r} + b\vec{s}|^2 &= b^2|\vec{s}|^2 + |\vec{r}|^2 + b^2|\vec{s}|^2 - 2b|\vec{r}||\vec{s}| \\ &= 2b^2|\vec{s}|^2 + |\vec{r}|^2 - 2b^2|\vec{s}|^2 = |\vec{r}|^2\end{aligned}$$

## Exercise 2

### Single Correct Choice Type

$$\text{Sol 1: (D) } (\vec{b} - 2\vec{c}) = \lambda\vec{a} \text{ or } \frac{1}{\lambda}|\vec{b} - 2\vec{c}| = 1$$

$$\text{or } (\vec{b} - 2\vec{c})^2 = (\lambda)^2 \text{ or } (\vec{b} - 2\vec{c}) \cdot (\vec{b} - 2\vec{c}) = \lambda^2$$

$$\text{or } \vec{b} \cdot \vec{b} - 2\vec{b} \cdot \vec{c} - 2\vec{c} \cdot \vec{b} + 4\vec{c} \cdot \vec{c}$$

$$\Rightarrow 16 - 4|\vec{b}| \cdot |\vec{c}| \cdot \cos\theta + 4 = \lambda^2$$

$$\sin\theta = \frac{\sqrt{15}}{4} \Rightarrow \cos\theta = \frac{1}{4}$$

$$\Rightarrow 16 - 4 \times \frac{1}{4} \times 4 \times 1 + 4 = \lambda^2$$

$$\Rightarrow \lambda^2 = 16 \Rightarrow \lambda = \pm 4$$

$$\text{Sol 2: (B) } l^2 + m^2 + n^2 = 1$$

$$\text{or } \cos^2\alpha + \cos^2\beta + \cos^2\gamma = 1$$

$$\text{or } 3\cos^2\theta = 1 \text{ or } \cos\theta = \frac{1}{\sqrt{3}}$$

$\therefore$  The desired vector is

$$15\sqrt{3}\left(\frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} + \frac{1}{\sqrt{3}}\hat{k}\right) = 15(\hat{i} + \hat{j} + \hat{k})$$


**Sol 3: (C)** Hint: Scalar Triple product

**Sol 4: (B)**  $\vec{a} + \vec{b} + \vec{c} = 0$

$$|\vec{a}| = 3, |\vec{b}| = 5, |\vec{c}| = 7$$

$$\cos \theta = \frac{(5)^2 + (3)^2 - (7)^2}{2 \times 5 \times 3} = \frac{25 + 9 - 49}{30}$$

$$= \frac{-15}{30} = -\frac{1}{2} \Rightarrow \theta = 150^\circ \text{ or } -30^\circ$$



**Sol 5: (B)** Let  $|\vec{a}| = 1 = |\vec{b}|$

$$2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) + 3 = 0$$

$$(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = 0$$

$$\Rightarrow 1 + 1 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow |\vec{c}| + 2 - 3 = 0 \Rightarrow |\vec{c}| = 1$$

**Sol 6: (A)**  $(\vec{a} + \vec{b}) \cdot \vec{b} = 0 \Rightarrow 2\vec{a} \cdot \vec{b} + 2\vec{b} \cdot \vec{b} = 0$

$$\text{Similarly } (\vec{a} + 2\vec{b}) \cdot \vec{a} = 0 \Rightarrow \vec{a} \cdot \vec{a} + 2\vec{a} \cdot \vec{b} = 0$$

$$\Rightarrow |\vec{a}|^2 = -2|\vec{b}|^2 \Rightarrow |\vec{a}| = \sqrt{2}|\vec{b}|$$

**Sol 7: (D)** Projection of  $\vec{a}$  on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$

$$\text{Let } \vec{a} = \hat{i} + 2\hat{j} + \hat{k} \text{ and } \vec{b} = 4\hat{i} + 4\hat{j} + 7\hat{k}$$

$$\therefore \text{Projection} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$$

$$= \frac{(\hat{i} + 2\hat{j} + \hat{k}) \cdot (4\hat{i} + 4\hat{j} + 7\hat{k})}{\sqrt{(4)^2 + (4)^2 + (7)^2}} = \frac{4 + 8 + 7}{\sqrt{16 + 16 + 49}} = \frac{19}{9}$$


Hence, the correct option is d.

**Sol 8: (B)**

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -2 & 6 & -8 \\ 1 & -3 & 4 \end{vmatrix} = \hat{i}(24 - 24) - \hat{j}(-8 + 8) + \hat{k}(6 - 6) = 0$$

$\therefore$  Magnitude = 0

**Sol 9: (A)**



**Sol 10: (A)**  $[2\vec{a} + 4\vec{b} \times \vec{c} \times \vec{d}] = \lambda [\vec{a} \times \vec{c} \times \vec{d}] + \mu [\vec{b} \times \vec{c} \times \vec{d}]$

$$[2\vec{a} + 4\vec{b} \times \vec{c} \times \vec{d}] = 2\vec{a} + 4\vec{b} \cdot (\vec{c} \times \vec{d}) = 2\vec{a} \cdot (\vec{c} \times \vec{d}) + 4\vec{b} \cdot (\vec{c} \times \vec{d})$$

$$= 2[\vec{a} \times \vec{c} \times \vec{d}] + 4[\vec{b} \times \vec{c} \times \vec{d}] \Rightarrow \lambda = 2, \mu = 4 \Rightarrow \lambda + \mu = 6$$


## Previous Years' Questions

**Sol 1: (B)** The volume of parallelepiped

$$= \begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \end{vmatrix} = \begin{vmatrix} 2 & -3 & 0 \\ 1 & 1 & -1 \\ 3 & 0 & -1 \end{vmatrix}$$

$$= 2(-1) + 3(-1+3) = -2+6 = 4$$

**Sol 2: (B)**  $\vec{a} = 2p\hat{i} + \hat{j}$



New vector

$$\vec{a} = \sqrt{4p^2 + 1} \cos \alpha \hat{i} + \sqrt{4p^2 + 1} \sin \alpha \hat{j}$$

$$\Rightarrow \sqrt{4p^2 + 1} \cos \alpha = p + 1$$

$$\Rightarrow \cos \alpha = \frac{p+1}{\sqrt{4p^2 + 1}}$$

$$\text{And } \sqrt{4p^2 + 1} \sin \alpha = 1$$

$$\Rightarrow \sin \alpha = \frac{1}{\sqrt{4p^2 + 1}}$$

$$\Rightarrow \sin^2 \alpha + \cos^2 \alpha = 1 = \frac{(p+1)^2}{4p^2 + 1} + \frac{1}{4p^2 + 1}$$

$$\Rightarrow 4p^2 + 1 = p^2 + 1 + 2p + 1$$

$$\Rightarrow 3p^2 - 2p - 1 = 0 \Rightarrow 3p^2 - 3p + p - 1 = 0$$

$$\Rightarrow 3p(p-1) + 1(p-1) = 0$$

$$\Rightarrow (p-1)(3p+1) = 0$$

$$\Rightarrow p = 1, \frac{-1}{3}$$

**Sol 3: (B)** Since, three vectors are coplanar.

$$\begin{vmatrix} a & a & c \\ 1 & 0 & 1 \\ c & c & b \end{vmatrix} = 0$$

Applying  $C_1 \rightarrow C_1 - C_2$

$$\Rightarrow \begin{vmatrix} 0 & a & c \\ 1 & 0 & 1 \\ 0 & c & b \end{vmatrix} = 0$$

$$\Rightarrow -1(ab - c^2) = 0 \Rightarrow ab = c^2$$

**Sol 4: (A)** Since,  $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{\vec{b} + \vec{c}}{\sqrt{2}}$

$$\Rightarrow (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c} = \frac{1}{\sqrt{2}}\vec{b} + \frac{1}{\sqrt{2}}\vec{c}$$

On equating the coefficient of  $\vec{c}$ , we get

$$\vec{a} \cdot \vec{b} = -\frac{1}{\sqrt{2}}$$

$$\Rightarrow |\vec{a}| |\vec{b}| \cos \theta = -\frac{1}{\sqrt{2}}$$

$$\therefore \cos \theta = -\frac{1}{\sqrt{2}} \Rightarrow \theta = \frac{3\pi}{4}$$

**Sol 5: (B)** Since,  $(\vec{a} + 2\vec{b}) \cdot (5\vec{a} - 4\vec{b}) = 0$

$$\Rightarrow 5|\vec{a}|^2 + 6\vec{a} \cdot \vec{b} - 8|\vec{b}|^2 = 0$$

$$\Rightarrow 6\vec{a} \cdot \vec{b} = 3[\because |\vec{a}| = |\vec{b}| = 1]$$

$$\Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = 60^\circ$$

**Sol 6: (C)** Given,  $\vec{v} = 2\hat{i} + \hat{j} - \hat{k}$  and  $\vec{w} = \hat{i} + 3\hat{k}$

$$\therefore [\vec{u} \ \vec{v} \ \vec{w}] = \vec{u} \cdot [(\vec{v} \times \vec{w})]$$

$$= \vec{u} \cdot (3\hat{i} - 7\hat{j} - \hat{k}) = |\vec{u}| |3\hat{i} - 7\hat{j} - \hat{k}| \cos \theta$$

which is maximum, if angle between  $\vec{u}$  and  $3\hat{i} - 7\hat{j} - \hat{k}$

is 0 and maximum value =  $|3\hat{i} - 7\hat{j} - \hat{k}| = \sqrt{59}$

**Sol 7: (C)** As we know, a vector coplanar to  $\vec{a}, \vec{b}$  and orthogonal to  $\vec{c}$  is  $\lambda \{(\vec{a} \times \vec{b}) \times \vec{c}\}$

$\therefore$  A vector coplanar to  $(2\hat{i} + \hat{j} + \hat{k}), (\hat{i} - \hat{j} + \hat{k})$  and

Orthogonal to  $3\hat{i} + 2\hat{j} + 6\hat{k}$

$$= \lambda \{[(2\hat{i} + \hat{j} + \hat{k}) \times (\hat{i} - \hat{j} + \hat{k})] \times (3\hat{i} + 2\hat{j} + 6\hat{k})\}$$

$$= \lambda [(2\hat{i} - \hat{j} - 3\hat{k}) \times (3\hat{i} + 2\hat{j} + 6\hat{k})]$$

$$= \lambda (21\hat{j} - 7\hat{k})$$

$$\therefore \text{Unit vector} = +\frac{(21\hat{j} - 7\hat{k})}{\sqrt{(21)^2 + (7)^2}} = +\frac{(3\hat{j} - \hat{k})}{\sqrt{10}}$$

**Sol 8: (A)**  $\vec{AB} = 2\hat{i} + 10\hat{j} + 11\hat{k}$

$$\vec{AD} = -\hat{i} + 2\hat{j} + 2\hat{k}$$

Angle ' $\theta$ ' between  $\vec{AB}$  and  $\vec{AD}$  is

$$\cos(\theta) = \frac{|\vec{AB} \cdot \vec{AD}|}{|\vec{AB}| |\vec{AD}|} = \frac{|-2 + 20 + 22|}{(15)(3)} = \frac{8}{9}$$

**Sol 9:**  $\vec{V} = \vec{i} + \vec{j} + \vec{k} + \lambda(\vec{i} - \vec{j} + \vec{k})$

$$= (1+\lambda)\vec{i} + (1-\lambda)\vec{j} + (1+\lambda)\vec{k}$$

Projection on  $\vec{C}$  is  $\frac{1}{\sqrt{3}}$

$$\frac{\vec{V} \cdot \vec{C}}{|\vec{C}|} = \frac{1}{\sqrt{3}}$$

$$\frac{(1+\lambda) - (1-\lambda) - (1+\lambda)}{\sqrt{3}} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow 1 + \lambda - 1 + \lambda - 1 - \lambda = 1 \Rightarrow \lambda = 2$$

$$\vec{V} = 3\hat{i} - \hat{j} + 3\hat{k}$$

**Sol 10: (C)**  $\vec{a} \times (\vec{b} \times \vec{c}) - \frac{\sqrt{3}}{2}(\vec{b} + \vec{c})$

$$\Rightarrow (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c} = \frac{\sqrt{3}}{2}(\vec{b} + \vec{c})$$

$$\Rightarrow \vec{a} \cdot \vec{c} = \frac{\sqrt{3}}{2} \text{ and } \vec{a} \cdot \vec{b} = -\frac{\sqrt{3}}{2}$$

$$\Rightarrow |\vec{a}| |\vec{b}| \cos \theta = -\frac{\sqrt{3}}{2}$$

$$\Rightarrow \cos \theta = \frac{-\sqrt{3}}{2}$$

$$\Rightarrow \theta = \frac{5\pi}{6}$$

**Sol 11: (D)**  $(\vec{a} \times \vec{b}) \times \vec{c} = \frac{1}{3} |\vec{b}| |\vec{c}| \vec{a}$

$$\Rightarrow (\vec{c} \cdot \vec{a}) \vec{b} - (\vec{c} \cdot \vec{b}) \vec{a} = \frac{1}{3} |\vec{b}| |\vec{c}| \vec{a}$$

$$\Rightarrow \vec{c} \cdot \vec{b} = -\frac{1}{3} |\vec{b}| |\vec{c}| \Rightarrow \cos \theta = -\frac{1}{3}$$

$$\Rightarrow \sin^2 \theta = 1 - \cos^2 \theta = 1 - \frac{1}{9} = \frac{8}{9}$$

$$\Rightarrow \sin \theta = \pm \frac{2\sqrt{3}}{3}$$

But  $\sin \theta = \frac{2\sqrt{3}}{3}$

**Sol 12: (A)**  $[\vec{a} \times \vec{b} \cdot \vec{b} \times \vec{c} \cdot \vec{c} \times \vec{a}] = \lambda [\vec{a} \cdot \vec{b} \cdot \vec{c}]^2$

We know that

$$[\vec{a} \times \vec{b} \cdot \vec{b} \times \vec{c} \cdot \vec{c} \times \vec{a}] = [\vec{a} \cdot \vec{b} \cdot \vec{c}]^2$$

$$\Rightarrow \lambda = 1$$

**Sol 13: (B)** The length of median through A

$$= \frac{|\vec{AB} + \vec{AC}|}{2} = \frac{3\hat{i} + 4\hat{k} + 5\hat{i} - 2\hat{j} + 4\hat{k}}{2} = \frac{8\hat{i} - 2\hat{j} + 8\hat{k}}{2}$$

$$= 4\hat{i} - \hat{j} + 4\hat{k}$$

$$\text{Length} = \sqrt{16 + 1 + 16} = \sqrt{33}$$

**Sol 14: (C)**  $\vec{c} = \vec{a} + 2\vec{b}$  and  $\vec{d} = 5\vec{a} - 4\vec{b}$

$$\vec{c} \perp \vec{d}$$

$$\vec{c} \cdot \vec{d} = 0$$

$$\Rightarrow 5|\vec{a}|^2 - 4\vec{a} \cdot \vec{b} + 10\vec{a} \cdot \vec{b} - 8|\vec{b}|^2 = 0$$

$$\Rightarrow 5 + 6\vec{a} \cdot \vec{b} - 8 = 0$$

$$\Rightarrow 6\vec{a} \cdot \vec{b} = 3$$

$$\Rightarrow \vec{a} \cdot \vec{b} = \frac{1}{2}$$

$$\Rightarrow \cos \phi = \frac{1}{2}$$

$$\Rightarrow \phi = \frac{\pi}{3}$$

**Sol 15: (D)**  $\vec{a} \cdot \vec{b} = 0$        $\vec{b} \cdot \vec{c} = 0$        $\vec{c} \cdot \vec{a} = 0$

$$\Rightarrow 2\lambda + 4 + \mu = 0 \quad \lambda - 1 + 2\mu = 0$$

Solving we get :  $\lambda = -3, \mu = 2$

**Sol 16: (D)**  $\vec{a} = \hat{j} - \hat{k}$  and  $\vec{c} = \hat{i} - \hat{j} - \hat{k}$

Let  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

$$\therefore \vec{a} \times \vec{b} + \vec{c} = 0, \vec{a} \times \vec{b} = -\vec{c}$$

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & -1 \\ b_1 & b_2 & b_3 \end{vmatrix} = -\hat{i} + \hat{j} + \hat{k}$$

$$\hat{i}(b_3 + b_2) - \hat{j}(b_1) + \hat{k}(-b_1) = -\hat{i} + \hat{j} + \hat{k}$$

$$b_3 + b_2 = -1 \quad \dots (i)$$

$$b_1 = -1 \quad \dots (ii)$$

$$\vec{a} \cdot \vec{b} = 3$$

$$b_2 - b_3 = 3 \quad \dots (iii)$$

Solve (i) and (iii)

$$2b_2 = 2 \quad b_1 = 2 \quad b_3 = -2$$

$$\therefore b_1 = -1 \quad b_2 = 1 \quad b_3 = -2$$

$$\text{Hence } \vec{b} = -\hat{i} + \hat{j} - 2\hat{k}$$

**Sol 17: (D)**  $(2\vec{a} - \vec{b}) \cdot \{(\vec{a} \times \vec{b}) \times (\vec{a} + 2\vec{b})\}$   
 $= (2\vec{a} - \vec{b}) \cdot \{[\vec{a} \cdot (\vec{a} + 2\vec{b})]\vec{b} - [\vec{b} \cdot (\vec{a} + 2\vec{b})\vec{a}]\}$   
 $= -5(\vec{a})^2 (\vec{b})^2 + 5(\vec{a} \cdot \vec{b})^2 = -5$

**Sol 18: (C)**  $\vec{b} \times \vec{c} = \vec{b} \times \vec{d}$

$$\Rightarrow \vec{a} \times (\vec{b} \times \vec{c}) = \vec{a} \times (\vec{b} \times \vec{d})$$

$$\Rightarrow (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c} = (\vec{a} \cdot \vec{d})\vec{b} - (\vec{a} \cdot \vec{b})\vec{d}$$

$$\Rightarrow (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c} = -(\vec{a} \cdot \vec{b})\vec{d}$$

$$\therefore \vec{d} = \vec{c} - \left( \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \right) \vec{b}$$

## JEE Advanced/Boards

### Exercise 1

**Sol 1:**  $\frac{(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2}{2a^2b^2}$

$$\begin{aligned}
 &= \frac{(\vec{a} \times \vec{b}) \cdot (\vec{a} \times \vec{b}) + (\vec{a} \cdot \vec{b}) \cdot (\vec{a} \cdot \vec{b})}{2|\vec{a}|^2 \cdot |\vec{b}|^2} = \frac{|\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2}{2|\vec{a}|^2 \cdot |\vec{b}|^2} \\
 &= \frac{|\vec{a}|^2 \cdot |\vec{b}|^2 \cdot \sin^2 \theta + |\vec{a}|^2 \cdot |\vec{b}|^2 \cdot \cos^2 \theta}{2|\vec{a}|^2 \cdot |\vec{b}|^2}
 \end{aligned}$$

**Sol 2:** Area =  $\frac{1}{2} |(\hat{3}\hat{i} + 4\hat{j}) \times (-5\hat{i} + 7\hat{j})| = \frac{1}{2} |21\hat{k} + 20\hat{k}| = \frac{41}{2}$

**Sol 3:**  $20\hat{i} + 11\hat{j} = \lambda \{(40 - a)\hat{i} + 44\hat{j}\}$

$\therefore \lambda(40 - a) = 20$  and  $\lambda(4) = 11$

$\Rightarrow a = -40$

**Sol 4:** Note that since  $|\vec{a}| = |\vec{b}|$  hence the parallelogram will be a rhombus.

**Sol 5:**  $\therefore (\vec{a} + 3\vec{b})$  is perpendicular to  $(7\vec{a} - 5\vec{b})$

$\therefore (\vec{a} + 3\vec{b}) \cdot (7\vec{a} - 5\vec{b}) = 0$

$\Rightarrow 7|\vec{a}|^2 - 5\vec{a} \cdot \vec{b} + 21\vec{a} \cdot \vec{b} - 15|\vec{b}|^2 = 0 \Rightarrow 16\vec{a} \cdot \vec{b} = 8$

$\therefore \vec{a} \cdot \vec{b} = \frac{1}{2} \Rightarrow |\vec{a}| \cdot |\vec{b}| \cdot \cos \theta = \frac{1}{2} \Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$

**Sol 6:**  $|\hat{A} - \hat{B}|^2 = (\hat{A} - \hat{B}) \cdot (\hat{A} - \hat{B})$

$= |\hat{A}|^2 + |\hat{B}|^2 - 2\hat{A} \cdot \hat{B} = 1 + 1 - 2|1||1| \cos \pi = 2 + 2 = 4$

$\Rightarrow |\hat{A} - \hat{B}| = 2 \Rightarrow \frac{|\hat{A} - \hat{B}|}{2} = 1$

**Sol 7:**  $\vec{F}_1 = 3\hat{i} + 2\hat{j} + 5\hat{k}$  and  $\vec{F}_2 = 2\hat{j} + \hat{j} + 3\hat{k}$

$\therefore \vec{F} = \vec{F}_1 + \vec{F}_2 = 5\hat{i} + 3\hat{j} + 8\hat{k}$

$\Delta \vec{x} = (4\hat{i} + 3\hat{j} + 7\hat{k}) - (2\hat{i} + \hat{j} + 3\hat{k}) = 2\hat{i} + 2\hat{j} + 4\hat{k}$

$\therefore$  Work done =  $\vec{F} \cdot \Delta \vec{x} = 10 + 6 + 32 = 48$  units

**Sol 8:**  $\Delta \vec{x} = -2\hat{i} - 3\hat{j} - 4\hat{k}$

Work done =  $\vec{F} \cdot \Delta \vec{x}$

$\Rightarrow 2 = -12 - 3\lambda - 16 \Rightarrow 30 = -3\lambda \Rightarrow \lambda = -10$

**Sol 9:** Let  $\vec{b} = 3\hat{i}$  and  $\vec{a} = 2(\hat{i} + \hat{j})$

The two diagonals will be  $6\vec{a} - \vec{b}$  and  $4\vec{a} + 5\vec{b}$

Length of  $6\vec{a} - \vec{b} = |9\hat{i} + 12\hat{j}| = 15$

Length of  $4\vec{a} + 5\vec{b}$

$= |8\hat{j} + 23\hat{i}| = \sqrt{(23)^2 + (8)^2} = \sqrt{593}$

**Sol 10:**  $2 \times |\hat{i} + x\hat{j} + 3\hat{k}| = |4\hat{i} + (4x - 2)\hat{j} + 2\hat{k}|$

or,  $2 \cdot \sqrt{1^2 + x^2 + 9} = \sqrt{16 + (4x - 2)^2 + 4}$

$\therefore 4(10 + x^2) = 20 + 16x^2 + 4 - 16x$

or,  $40 + 4x^2 = 24 + 16x^2 - 16x$

or,  $12x^2 - 16x - 16 = 0$  or,  $3x^2 - 4x - 4 = 0$

$\therefore x = \frac{4 \pm \sqrt{16 + 4 \cdot 4 \cdot 3}}{6} = \frac{4 \pm 8}{6} = 2$  or  $\frac{-2}{3}$

## Exercise 2

**Sol 1: (A)**  $\vec{r} = (9\hat{i} - \hat{j} + 2\hat{k}) - (3\hat{i} - 2\hat{j} + \hat{k}) = (6\hat{i} + \hat{j} + \hat{k})$

$\therefore$  Moment of couple =  $\vec{r} \times \vec{F} = (6\hat{i} + \hat{j} + \hat{k}) \times (5\hat{i} + \hat{j})$

$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 6 & 1 & 1 \\ 5 & 1 & 0 \end{vmatrix} = \hat{i}(0 - 1) - \hat{j}(0 - 5) + \hat{k}(6 - 5)$

$= -\hat{i} + 5\hat{j} + \hat{k}$

**Sol 2: (C)**  $(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c})$

$= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + \vec{a} \cdot (\vec{b} + \vec{c}) + \vec{b} \cdot (\vec{a} + \vec{c}) + \vec{c} \cdot (\vec{a} + \vec{b})$

$= (1)^2 + (4)^2 + (8)^2 + 0 = 81$

$\therefore |\vec{a} + \vec{b} + \vec{c}| = 9$

**Sol 3: (A)** Let M be the point of intersection of internal bisector with AB.

$\therefore \frac{AM}{MB} = \frac{1}{2}$

$\therefore \vec{OM} = \frac{1(2\hat{i} + 4\hat{j} + 4\hat{k}) + 2(2\hat{i} + 2\hat{j} + \hat{k})}{3} = \frac{6\hat{i} + 8\hat{j} + 6\hat{k}}{3}$

$\therefore |\vec{OM}| = \sqrt{4 + \left(\frac{8}{3}\right)^2 + 4} = \sqrt{\frac{72 + 64}{9}} = \sqrt{\frac{136}{9}}$



**Sol 4: (A)**  $\vec{a} + 2\vec{b} + 3\vec{c} = 0$

$$\Rightarrow \vec{a} \times \vec{b} + 2\vec{b} \times \vec{b} + 3\vec{c} \times \vec{b} = 0$$

$$\Rightarrow \vec{a} \times \vec{b} + 3\vec{c} \times \vec{b} = 0 \Rightarrow \vec{a} \times \vec{b} = 3\vec{b} \times \vec{c} \quad \dots (i)$$

Similarly,  $\vec{a} + 2\vec{b} + 3\vec{c} = 0$

$$\vec{a} \times \vec{a} + 2\vec{b} \times \vec{a} + 3\vec{c} \times \vec{a} = 0 \Rightarrow \vec{c} \times \vec{a} = \frac{2}{3}\vec{a} \times \vec{b}$$

$$= \frac{2}{3} \times 3(\vec{b} \times \vec{c}) = 2(\vec{b} \times \vec{c}) \quad \dots (ii)$$

$$\Rightarrow \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} = 6(\vec{b} \times \vec{c})$$

**Sol 5: (C)**  $[\vec{a} - \vec{b} \quad \vec{b} - \vec{c} \quad \vec{c} - \vec{a}] = (\vec{a} - \vec{b}) \cdot (\vec{b} - \vec{c}) \times (\vec{c} - \vec{a})$

$$= (\vec{a} - \vec{b}) \cdot \{\vec{b} \times \vec{c} - \vec{b} \times \vec{a} + \vec{c} \times \vec{a}\} = \vec{a} \cdot (\vec{b} \times \vec{c}) - \vec{b} \cdot (\vec{c} \times \vec{a}) = 0$$

**Sol 6: (D)** A general point on the curve will have vector

$$x\hat{i} + y\hat{j} = x\hat{i} + \left(x + \frac{1}{x}\right)\hat{j}$$

$$\therefore \overrightarrow{OP} \cdot \vec{I} = 1$$

$$\therefore \left\{x\hat{i} + \left(x + \frac{1}{x}\right)\hat{j}\right\} \cdot \hat{i} = 1 \Rightarrow x = 1$$

$$\therefore \overrightarrow{OP} = \hat{i} + 2\hat{j}$$

Again,  $\overrightarrow{OQ} \cdot \vec{I} = -1 \Rightarrow x = -1$

$$\therefore \overrightarrow{OQ} = -\hat{i} - 2\hat{j}$$

$$\therefore 2\overrightarrow{OP} + 3\overrightarrow{OQ} = -\hat{i} - 2\hat{j}$$

$$\therefore |2\overrightarrow{OP} + 3\overrightarrow{OQ}| = \sqrt{(-1)^2 + (2)^2} = \sqrt{5}$$

**Sol 7: (C)**  $\vec{A} \cdot (\vec{B} \cdot \vec{C}) + \vec{B} \cdot \vec{C}(\vec{A} + \vec{B}) = 0$

$$\Rightarrow A \cdot B + A \cdot C + B \cdot C = 0$$

$$|A + B + C|^2 = |A|^2 + |B|^2 + |C|^2 + 2(A \cdot B + B \cdot C + C \cdot A)$$

$$= 1 + 16 + 64 + 0 = 81$$

$$\Rightarrow |A + B + C| = 9$$

**Sol 8: (C)** Given

$$|A - B| \leq 1$$

$$\therefore \sqrt{1^2 + 1^2 - 2\cos 2\theta} \leq 1$$

$$\Rightarrow 2 - 2\cos 2\theta \leq 1 \Rightarrow 2\cos 2\theta \geq 1 \Rightarrow \cos 2\theta \geq \frac{1}{2}$$

$$\therefore \frac{5\pi}{3} \leq 2\theta \leq 2\pi \Rightarrow \frac{5\pi}{6} \leq \theta \leq \pi$$

**Sol 9: (C)**  $|\vec{A} \cdot \vec{B} \cdot \vec{A} \times \vec{B}| = \frac{1}{4}$

$$\Rightarrow \vec{A} \cdot (\vec{B} \times \vec{A} \times \vec{B}) = \frac{1}{4} \Rightarrow \vec{A} \cdot [(\vec{B} \cdot \vec{B})\vec{A} - (\vec{A} \cdot \vec{B})\vec{B}] = \frac{1}{4}$$

$$\Rightarrow \vec{A} \cdot [\vec{A} - (\vec{A} \cdot \vec{B})\vec{B}] = \frac{1}{4} \Rightarrow \vec{A} \cdot \vec{A} - (\vec{A} \cdot \vec{B})^2 = \frac{1}{4}$$

$$\Rightarrow (\vec{A} \cdot \vec{B})^2 = 1 - \frac{1}{4} = \frac{3}{4} \Rightarrow \vec{A} \cdot \vec{B} = \frac{\sqrt{3}}{2}$$

$$\Rightarrow \cos \theta = \frac{\sqrt{3}}{2} \Rightarrow \theta = \frac{\pi}{3}$$

**Sol 10: (C)**  $|\vec{A} - \vec{B}| + |\vec{A} + \vec{B}| = \sqrt{2 - 2\cos \theta} + \sqrt{2 + 2\cos \theta}$

$$= \sqrt{2}(\sqrt{1 - \cos \theta} + \sqrt{1 + \cos \theta})$$

$$= \sqrt{2} \cdot \left\{ \sqrt{2} \sin \frac{\theta}{2} + 2 \cos \frac{\theta}{2} \right\} = 2 \left( \sin \frac{\theta}{2} + \cos \frac{\theta}{2} \right)$$

Greatest value is  $2\sqrt{2}$

## Previous Years' Questions

**Sol 1:**  $\vec{A} \cdot \vec{X} = C$  and  $\vec{A} \times \vec{X} = \vec{B}$

Let  $\vec{A} = A_1\hat{i} + A_2\hat{j} + A_3\hat{k}$

(i)  $\vec{A} \times \vec{X} = \vec{B}$

$$\vec{A} \times (\vec{A} \times \vec{X}) = \vec{A} \times \vec{B} \Rightarrow (\vec{A} \cdot \vec{X})\vec{A} - (\vec{A} \cdot \vec{A})\vec{X} = \vec{A} \times \vec{B}$$

$$\Rightarrow |\vec{A}|^2 \vec{X} = C\vec{A} - \vec{A} \times \vec{B} \Rightarrow \vec{X} = \frac{C\vec{A}}{|\vec{A}|^2} - \frac{\vec{A} \times \vec{B}}{|\vec{A}|^2}$$

(ii) If coordinate system is rotated about the x-axis through an angle  $\frac{\pi}{2}$ , then

x- component =  $A_2$

y- component =  $A_1$

z- component =  $A_3$

$$\vec{A} = A_2\hat{i} - A_1\hat{j} + A_3\hat{k}$$

[New coordinates system]

**Sol 2:** 
$$\begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \\ \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} \cdot \vec{a} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \end{vmatrix} = \begin{vmatrix} \vec{a} + \vec{b} + \vec{c} & \vec{b} & \vec{c} \\ \vec{a} \cdot (\vec{a} + \vec{b} + \vec{c}) & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} \cdot (\vec{a} + \vec{b} + \vec{c}) & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \end{vmatrix}$$

$$\begin{aligned}
&= (\vec{a} + \vec{b} + \vec{c}) \cdot \begin{vmatrix} 1 & \vec{b} & \vec{c} \\ \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \end{vmatrix} \\
&= (\vec{a} + \vec{b} + \vec{c}) \cdot \left[ (\vec{a} \cdot \vec{b})(\vec{b} \cdot \vec{c}) - (\vec{b} \cdot \vec{b})(\vec{a} \cdot \vec{c}) - (\vec{a} \cdot \vec{b})(\vec{b} \cdot \vec{c}) + \right. \\
&\quad \left. (\vec{b} \cdot \vec{b})(\vec{a} \cdot \vec{c}) + (\vec{a} \cdot \vec{c})(\vec{b} \cdot \vec{b}) - (\vec{b} \cdot \vec{c})(\vec{a} \cdot \vec{b}) \right] \\
&= (\vec{a} + \vec{b} + \vec{c}) \cdot 0 \\
&\Rightarrow \text{if } \vec{a} + \vec{b} + \vec{c} = 0 \text{ [coplanar condition]}
\end{aligned}$$

**Sol 3:**  $\vec{R} \times \vec{B} = \vec{C} \times \vec{B} \Rightarrow \vec{A} \times (\vec{R} \times \vec{B}) = \vec{A} \times (\vec{C} \times \vec{B})$


$$\begin{aligned}
&\Rightarrow (\vec{A} \cdot \vec{B})\vec{R} - (\vec{A} \cdot \vec{R})\vec{B} = (\vec{A} \cdot \vec{B})\vec{C} - (\vec{A} \cdot \vec{C})\vec{B} \\
&(2+0+1)\vec{R} - 0 = (2+0+1)\vec{C} - (8+0+7)\vec{B} \\
&\Rightarrow 3\vec{R} = 3\vec{C} - 15\vec{B} \Rightarrow \vec{R} = \vec{C} - 5\vec{B} \\
&= 4\hat{i} - 3\hat{j} + 7\hat{k} - 5\hat{i} - 5\hat{j} - 5\hat{k} = -\hat{i} - 8\hat{j} + 2\hat{k}
\end{aligned}$$

**Sol 4:** Given,  $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$

and  $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$

$$\begin{aligned}
&\Rightarrow \vec{a} \times \vec{b} - \vec{a} \times \vec{c} = \vec{c} \times \vec{d} - \vec{b} \times \vec{d} \\
&\Rightarrow \vec{a} \times (\vec{b} - \vec{c}) = (\vec{c} - \vec{b}) \times \vec{d} \\
&\Rightarrow \vec{a} \times (\vec{b} - \vec{c}) - (\vec{c} - \vec{b}) \times \vec{d} = 0 \\
&\Rightarrow \vec{a} \times (\vec{b} - \vec{c}) - \vec{d} \times (\vec{b} - \vec{c}) = 0 \\
&\Rightarrow (\vec{a} - \vec{d}) \times (\vec{b} - \vec{c}) = 0 \Rightarrow (\vec{a} - \vec{d}) \parallel (\vec{b} - \vec{c}) \\
&\therefore (\vec{a} - \vec{d}) \cdot (\vec{b} - \vec{c}) \neq 0 \Rightarrow \vec{a} \cdot \vec{b} + \vec{d} \cdot \vec{c} \neq \vec{d} \cdot \vec{b} + \vec{a} \cdot \vec{c}
\end{aligned}$$

**Sol 5:** Since,  $\hat{v}$  is unit vector along the incident ray and  $\hat{w}$  is the unit vector along the reflected ray.



Hence,  $\hat{a}$  is a unit vector along the external bisector of  $\hat{v}$  and  $\hat{w}$ .

$$\therefore \hat{w} - \hat{v} = \lambda \hat{a}$$

On squaring both sides, we get

$$\Rightarrow 1 + 1 - \hat{w} \cdot \hat{v} = \lambda^2 \Rightarrow 2 - 2\cos 2\theta = \lambda^2 \Rightarrow \lambda = 2\sin\theta$$

where  $2\theta$  is the angle between  $\hat{v}$  and  $\hat{w}$ .

Hence,  $\hat{w} - \hat{v} = 2\sin\theta \cdot \hat{a}$

$$= 2\cos(90^\circ - \theta) \hat{a} = -(2\hat{a} \cdot \hat{v}) \hat{a}$$

$$\hat{w} = \hat{v} - 2(\hat{a} \cdot \hat{v}) \hat{a}$$

**Sol 6: (B, D)** Let vector  $\vec{AO}$  be parallel to line of intersection of planes  $P_1$  and  $P_2$  through origin

Normal to plane  $P_1$  is

$$\vec{n}_1 = [(2\hat{j} + 3\hat{k}) \times (4\hat{j} - 3\hat{k})] = -18\hat{i}$$

Normal to plane  $P_2$  is

$$\vec{n}_2 = (\hat{j} - \hat{k}) \times (3\hat{i} + 3\hat{j}) = 3\hat{i} - 3\hat{j} - \hat{k}$$

$$\therefore \vec{OA} \text{ is parallel to } \pm (\vec{n}_1 \times \vec{n}_2) = 54\hat{j} - 54\hat{k}$$

$\therefore$  Angle between  $54(\hat{j} - \hat{k})$  and  $(2\hat{i} + \hat{j} - 2\hat{k})$  is

$$\cos\theta = \pm \left( \frac{54 + 108}{3 \cdot 54 \cdot \sqrt{2}} \right) = \pm \frac{1}{\sqrt{2}} \quad \therefore \theta = \frac{\pi}{4}, \frac{3\pi}{4}$$

**Sol 7: (A, D)** Let,  $\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$ ,  $\vec{b} = \hat{i} + 2\hat{j} + \hat{k}$  and  $\vec{c} = \hat{i} + \hat{j} + \hat{k}$

$\therefore$  A vector coplanar to  $\vec{a}$  and  $\vec{b}$ , and perpendicular to  $\vec{c}$

$$\begin{aligned}
\vec{r} &= \lambda(\vec{a} \times \vec{b}) \times \vec{c} = \lambda\{(\vec{a} \cdot \vec{c})\vec{b} - (\vec{b} \cdot \vec{c})\vec{a}\} \\
&= \lambda\{(1+1+4)(\hat{i} + 2\hat{j} + \hat{k}) - (1+2+1)(\hat{i} + \hat{j} + 2\hat{k})\} \\
&= \lambda\{6\hat{i} + 12\hat{j} + 6\hat{k} - 6\hat{i} - 6\hat{j} - 12\hat{k}\} = \lambda\{6\hat{j} - 6\hat{k}\} = 6\lambda(\hat{j} - \hat{k})
\end{aligned}$$

For  $\lambda = \frac{1}{6} \Rightarrow$  (a) is correct.

and  $\lambda = -\frac{1}{6} \Rightarrow$  (d) is correct.

**Sol 8:** From the given information, it is clear that

$$\vec{a} = \frac{\hat{i} - 2\hat{j}}{\sqrt{5}}$$

$$\Rightarrow |\vec{a}| = 1, |\vec{b}| = 1, |\vec{a} \cdot \vec{b}| = 0$$

Now,  $(2\vec{a} + \vec{b}) \cdot [(\vec{a} \times \vec{b}) \times (\vec{a} - 2\vec{b})]$

$$= (2\vec{a} + \vec{b}) \cdot [\vec{a}^2 \vec{b} - (\vec{a} \cdot \vec{b}) \cdot \vec{a} + 2\vec{b}^2 \cdot \vec{a} - 2(\vec{b} \cdot \vec{a}) \cdot \vec{a}]$$

$$= [2\vec{a} + \vec{b}] \cdot [\vec{b} + 2\vec{a}] = 4\vec{a}^2 + \vec{b}^2 = 4 \cdot 1 + 1 = 5 \quad [\text{as } \vec{a} \cdot \vec{b} = 0]$$

**Sol 9:**  $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$

$$\Rightarrow (\vec{r} - \vec{c}) \times \vec{b} = 0 \Rightarrow \vec{r} - \vec{c} = \lambda \vec{b}$$

or  $\vec{r} = \vec{c} + \lambda \vec{b}$  ... (i)

Given,  $\vec{r} \cdot \vec{a} = 0$ , taking dot product with  $\vec{a}$  for Eq. (i)

$$\Rightarrow \vec{r} \cdot \vec{a} = \vec{a} \cdot \vec{c} + \lambda \vec{a} \cdot \vec{b}$$

$$\therefore \lambda = \frac{-\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \quad (\because \vec{r} \cdot \vec{a} = 0) \quad \dots (ii)$$

From Eqs. (i) and (ii), we get

$$\vec{r} = \vec{c} - \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \vec{b}, \text{ taking dot with } \vec{b}, \text{ we get}$$

$$\vec{r} \cdot \vec{b} = \vec{c} \cdot \vec{b} - \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} (\vec{b} \cdot \vec{b})$$

$$= (-1+2) - \frac{(-1-3)}{(1)}(1+1) \text{ where,}$$

$$\begin{bmatrix} \vec{a} = -\hat{i} - \hat{k} \\ \vec{b} = -\hat{i} + \hat{j} \\ \vec{c} = \hat{i} + 2\hat{j} + 3\hat{k} \end{bmatrix} = 1+8 = 9$$

**Sol 10: (B, C)** Given:  $w \cdot (\hat{u} \times \hat{v}) = 1$

$$\Rightarrow |w| |(\hat{u} \times \hat{v})| \cos \theta = 1 \Rightarrow \cos \theta = 1$$

$$w \perp \hat{u} \times \hat{v} \Rightarrow w \perp \hat{u} \text{ and } w \perp \hat{v} \text{ and } |\hat{u} \times \hat{v}| = 1$$

Angle between  $\hat{u}$  and  $\hat{v}$  can change to have initially many of vectors  $\hat{v}$  as  $\hat{w} \perp \hat{v}$

$$\text{If } \hat{u} \text{ lies in } xy \text{ plane then } \hat{u} = u_1 \hat{i} + u_2 \hat{j}$$

$$\Rightarrow \hat{w} \cdot \hat{u} = 0 \Rightarrow u_1 + u_2 = 0 \Rightarrow |u_1| = |u_2|$$

**Sol 11: (A, C, D)** In  $\Delta PQR$

$$-\vec{a} = \vec{b} + \vec{c}$$

$$\Rightarrow \vec{a} \cdot \vec{a} = (\vec{b} + \vec{c}) \cdot (\vec{b} + \vec{c})$$


$$\Rightarrow |\vec{a}|^2 = |\vec{b}|^2 + |\vec{c}|^2 + 2\vec{b} \cdot \vec{c}$$

$$\Rightarrow |\vec{a}|^2 = (4\sqrt{3})^2 + |\vec{c}|^2 + 2 \times 24$$

$$\Rightarrow (12)^2 = (4\sqrt{3})^2 + |\vec{c}|^2 + 2 \times 24$$

$$\Rightarrow |\vec{c}|^2 = 144 - 96$$

$$\Rightarrow |\vec{c}| = 4\sqrt{3}$$



$$\Rightarrow \frac{|\vec{c}|^2}{2} - |\vec{a}|^2 = \frac{48}{2} - 12 = 24 - 12 = 12$$

Given  $\vec{b} \cdot \vec{c} = 24$

$$-|\vec{b}| |\vec{c}| \cos \theta = 24$$

$$-4\sqrt{3} \times 4\sqrt{3} \cos \theta = 24$$

$$\cos \theta = \frac{-1}{2}$$

Since  $|\vec{b}| = |\vec{c}|$

$$\angle PQR = \angle PRQ \Rightarrow \angle QPR = 120^\circ$$

$$\text{and } \angle PQR = \angle PRQ = 30^\circ \Rightarrow |\vec{a} \times \vec{b} + \vec{c} \times \vec{a}| = 48\sqrt{3}$$

$$\text{And } \vec{a} \times \vec{b} = -72$$

**Sol 12:**  $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$

Taking cross with  $\vec{a}$

$$\Rightarrow \vec{a} \times (\vec{r} \times \vec{b}) = \vec{a} \times (\vec{c} \times \vec{b})$$

$$\Rightarrow (\vec{a} \cdot \vec{b}) \vec{r} - (\vec{a} \cdot \vec{r}) \vec{b} = \vec{a} \times (\vec{c} \times \vec{b})$$

$$\Rightarrow \vec{r} = -3\hat{i} + 6\hat{j} + 3\hat{k}$$

$$\vec{r} \cdot \vec{b} = 3 + 6 = 9$$

**Sol 13:**  $(2\vec{a} + \vec{b}) \cdot [(\vec{a} \times \vec{b}) \times (\vec{a} - 2\vec{b})]$

$$\vec{a} = \frac{\hat{i} - 2\hat{j}}{\sqrt{5}} \quad \vec{b} = \frac{2\hat{i} + \hat{j} + 3\hat{k}}{\sqrt{14}}$$

$$\Rightarrow (2\vec{a} + \vec{b}) \cdot [\vec{a} \times (\vec{a} \times \vec{b}) - (\vec{a} \times \vec{b}) \times \vec{b}]$$

$$= (2\vec{a} \times \vec{b}) \cdot [(\vec{a} \cdot \vec{a}) \vec{b} - (\vec{a} \cdot \vec{b}) \vec{a} + 2(\vec{a} \cdot \vec{b}) \vec{b} + 2(\vec{b} \cdot \vec{b}) \vec{a}]$$

$$= (2\vec{a} + \vec{b}) \cdot [\vec{b} + 2\vec{a}] \quad \{\vec{a} \cdot \vec{b} = 0\}$$

$$= 2\vec{a} \cdot \vec{b} + 4|\vec{a}|^2 + |\vec{b}|^2 - 2\vec{b} \cdot \vec{a}$$

$$= 4|\vec{a}|^2 + |\vec{b}|^2$$

$$= 4 + 1 = 5$$

**Sol 14:** Given:  $\vec{a} \times \vec{b} + \vec{b} \times \vec{c} = p\vec{a} + q\vec{b} + r\vec{c}$

$$\Rightarrow \vec{a} \cdot (\vec{a} \times \vec{b}) + \vec{a} \cdot (\vec{b} \times \vec{c}) = p + q(\vec{a} \cdot \vec{b}) + r(\vec{a} \cdot \vec{c})$$

$$\Rightarrow \vec{a} \cdot (\vec{b} \times \vec{c}) = p + \frac{q}{2} + \frac{r}{2} \quad \dots (i)$$

$$\text{Similarly, } \vec{b} \cdot (\vec{a} \times \vec{b}) + \vec{b} \cdot (\vec{b} \times \vec{c}) = \frac{p}{2} + q + \frac{r}{2}$$

$$\Rightarrow \frac{p}{2} + q + \frac{r}{2} = 0 \quad \dots (ii)$$

$$\text{and } \frac{p}{2} + \frac{q}{2} + r = a(b \times c) \quad \dots(iii)$$

From (i), (ii) and (iii)

$$P = -q = r \Rightarrow \frac{p^2 + 2q^2 + r^2}{q^2} = 4$$

$$\text{Sol 15: (C)} \Rightarrow \vec{x} + \vec{y} = 3\hat{i} + \hat{j} - 2\hat{k}$$

$$\text{and } \vec{x} - \vec{y} = \hat{i} - 3\hat{j} - 4\hat{k}$$

On solving we get

$$\vec{x} = 2\hat{i} - \hat{j} - 3\hat{k}$$

$$\vec{y} = \hat{i} + 2\hat{j} + \hat{k}$$

Volume of parallelepiped

$$\begin{aligned} &= \begin{vmatrix} 2 & -1 & -3 \\ 1 & 2 & 1 \\ 1 & 2 & 3 \end{vmatrix} \\ &= 2(6-2) + 1(3-1) - 3(2-0) \\ &= 8 + 2 = 10 \end{aligned}$$

**Sol 16: (B, D)** Vector perpendicular to  $\ell_1$  and  $\ell_2$  is given by

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 2 & 1 \\ 1 & 2 & 2 \end{vmatrix} = \hat{j}(4-2) - \hat{j}(4-1) + \hat{k}(4-2) = 2\hat{i} - 3\hat{j} + 2\hat{k}$$

The eq. of line  $\perp$  to  $\ell_1$  and  $\ell_2$

$$\frac{x-0}{2} = \frac{y-0}{-3} = \frac{z-0}{2} = \gamma$$

$$\Rightarrow Q \equiv (2\gamma, -3\gamma, 2\gamma)$$

$$\text{The point Q lies on } \ell_2, \text{ then } = \frac{2\gamma-3}{1} = \frac{-3\gamma+1}{2} = \frac{2\gamma-4}{2}$$

$$\Rightarrow \gamma = 1$$


$$\Rightarrow Q \equiv (2, -3, 2)$$

Distance of P from Q in  $\sqrt{17}$

$$PQ^2 = 17 = (2-3-2)^2 + (-3-3-25)^2 + (2-5-2)^2$$

$$\Rightarrow S = -2, \frac{-10}{9}$$

$$\Rightarrow P \equiv (-1, -1, 0) \text{ and } \left(\frac{7}{9}, \frac{7}{9}, \frac{8}{9}\right)$$



$$\text{Sol 17: } V = a\hat{i} + b\hat{j} + c\hat{k} : a, b, c \in \{-1, 1\}$$

$$\text{Total number of selection} = {}^8C_3$$

$$\text{No. of coplanar vectors} = 6 \times 4 = 24$$

$$\text{Total number of non co-planar vet}$$

$$= {}^8C_3 - 24 = 32 = 2^5$$

$$= P = 5$$

$$\text{Sol 18: (C)} \quad (i) \left[ 2\vec{a} \times \vec{b} \quad 3\vec{b} \times \vec{c} \quad \vec{c} \times \vec{a} \right] = 6 \left[ \vec{a} \quad \vec{b} \quad \vec{c} \right]^2 = 6(2)^2 = 24 \left[ \vec{a} \cdot \vec{b} \cdot \vec{c} = 2 \right]$$

$$\begin{aligned} (ii) \left[ 3(\vec{a} + \vec{b}) \vec{b} + \vec{c} \quad 2\vec{c} + \vec{a} \right] &= 6 \left[ \vec{a} + \vec{b} \quad \vec{b} + \vec{c} \quad \vec{c} + \vec{a} \right] \\ &= 6(\vec{a} + \vec{b}) \cdot [(\vec{b} + \vec{c}) \times (\vec{c} + \vec{a})] \\ &= 6(\vec{a} \times \vec{b}) \cdot [\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{c} + \vec{c} \times \vec{a}] = 12\vec{a} \cdot (\vec{b} \times \vec{c}) \\ &= 12 \left[ \vec{a} \quad \vec{b} \quad \vec{c} \right] = 12 \times 5 = 60 \quad \left( \left[ \vec{a} \quad \vec{b} \quad \vec{c} \right] = 5 \right) \end{aligned}$$

$$\begin{aligned} (iii) \frac{1}{2} \left| (2\vec{a} + 3\vec{b}) \times (\vec{a} - \vec{b}) \right| &= \frac{1}{2} \left| (2\vec{a} \times \vec{a} - 2\vec{a} \times \vec{b} + 3\vec{b} \times \vec{a} - 3\vec{b} \times \vec{b}) \right| \\ &= \frac{1}{2} |5\vec{a} \times \vec{b}| = \frac{1}{2} \times 5 \times 40 = 100 \left[ \frac{1}{2} |\vec{a} \times \vec{b}| = 20 \right] \end{aligned}$$

$$\begin{aligned} (iii) \left| (\vec{a} \times \vec{b}) \times \vec{a} \right| &= \left| \vec{a} \times \vec{a} + \vec{b} \times \vec{a} \right| = \left| \vec{b} \times \vec{a} \right| \\ &= 30 \left[ |\vec{a} \times \vec{b}| = 30 \right] \end{aligned}$$

$$\text{Sol 19: } |\vec{a} - \vec{b}|^2 + |\vec{b} - \vec{c}|^2 + |\vec{c} - \vec{a}|^2 = 9$$

$$\Rightarrow 3 \left( |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 \right) - |\vec{a} + \vec{b} + \vec{c}|^2 = 9$$

$$\Rightarrow 3(1+1+1) - |\vec{a} + \vec{b} + \vec{c}|^2 = 9$$

$$\Rightarrow |\vec{a} + \vec{b} + \vec{c}|^2 = 0 \Rightarrow \vec{a} + \vec{b} + \vec{c} = 0$$

$$\Rightarrow \vec{b} + \vec{c} = -\vec{a}$$

$$\text{Now, } |2\vec{a} + 5\vec{b} + 5\vec{c}| = |2\vec{a} + 5(\vec{b} + \vec{c})|$$

$$= |2\vec{a} - 5\vec{a}| = 3|\vec{a}| = 3$$

**Sol 20: (C)** Let  $\vec{c} = 2\hat{i} + 3\hat{j} + 4\hat{k}$

$$\vec{a} \times \vec{c} = \vec{c} \times \vec{b} \Rightarrow (\vec{a} + \vec{b}) \times \vec{c} = 0$$

$$\Rightarrow (\vec{a} + \vec{b}) \parallel \vec{c}$$

$$\text{Let } (\vec{a} + \vec{b}) = \lambda \vec{c}$$

$$\Rightarrow |\vec{a} + \vec{b}| = |\lambda| |\vec{c}|$$

$$\Rightarrow \sqrt{29} = |\lambda| \cdot \sqrt{29} \Rightarrow \lambda = \pm 1$$

$$\therefore \vec{a} + \vec{b} = \pm (2\hat{i} + 3\hat{j} + 4\hat{k})$$

$$\text{Now, } (\vec{a} + \vec{b}) \cdot (-7\hat{i} + 2\hat{j} + 3\hat{k}) = \pm (-14 + 6 + 12) = \pm 4$$

**Sol 21: (C)** Any vectors  $\vec{v}$  coplanar with  $\vec{a}$  and  $\vec{b}$  is given by

$$\vec{v} = m\vec{a} + n\vec{b}$$

$$= m(\hat{i} + \hat{j} + \hat{k}) + n(\hat{i} + \hat{j} - \hat{k})$$

$$= (m+n)\hat{i} + (m+n)\hat{j} + (m-n)\hat{k} \quad \dots (i)$$

Projection to  $\vec{v}$  on  $\vec{c}$  is given by  $\frac{\vec{v} \cdot \vec{c}}{|\vec{c}|} = \frac{1}{\sqrt{3}}$

$$\Rightarrow (m+n) - (m+n) - (m-n) = 1$$

$$\Rightarrow m+n-m+n-m-n=1$$

$$\Rightarrow m+1=n$$

$$\Rightarrow m=n-1$$

Substituting in (i)

$$(2n-1)\hat{i} - \hat{j} + 2n-1\hat{k}$$

$$\text{for } n=2$$

$$3\hat{i} - \hat{j} + 3\hat{k}$$

**Sol 22: (A, D)** Let  $\vec{r}$  the vector coplanar with  $\vec{i} + \vec{j} + 2\vec{k}$  and  $\vec{i} + 2\vec{j} + \vec{k}$  then

$$\vec{r} = m(\vec{i} + \vec{j} + 2\vec{k}) + n(\vec{i} + 2\vec{j} + \vec{k})$$

$$= (m+n)\vec{i} + (m+2n)\vec{j} + (2m+n)\vec{k}$$

$$\vec{r} \perp \vec{c}, \text{ then}$$

$$m+n+m+2n+2m+n=0$$

$$\Rightarrow m+n=0$$

$$\Rightarrow \vec{r} = (0)\vec{i} + (0+n)\vec{j} + (n+0)\vec{k}$$

$$= n\vec{j} + m\vec{k}$$