

# IONIC EQUILIBRIUM

## 1. INTRODUCTION

Ionic equilibrium deals with the equilibrium of any substance with its ions in solution. The substance producing ions are called electrolytes.

According to conductivity, substances are of two types :

(i) **Non-Conductor :**

Those substances which do not show the flow of current or electricity.

**Ex.** Non - metals, plastic rubber, wood, etc.

Exception – Graphite is a non-metal but shows conductivity due to motion of free electrons.

(ii) **Conductors :**

Those substances which show conductivity or flow of current are called conductors. These are of 2 types :

(a) **Metallic or electronic conductors :**

Those conductor which show conductivity due to motion of free electrons. Resistance increases with temperature.

**Ex.** All metals, Graphite

(b) **Ionic or electrolytic conductors :**

Those conductors which show conductivity due to movement of free ions. Ions are in free state in the solutions of ionic compounds. On passing electric current through the solution, ions move towards oppositely charged electrodes, i.e., the cation moves towards cathode (negative electrode) and the anion moves towards anode (positive electrode).

The current flows through the solution due to the movement of the ions. Resistance decrease with temperature.

According to strength, ionic conductors are of two types :

(i) **Strong electrolyte :** Those ionic conductors which are completely ionized in aqueous solution are called as strong electrolyte.


For strong electrolyte the value of degree of dissociation is 100%.

**i.e. :  $\alpha = 1$**

- Ex.** (a) Strong acid  $\rightarrow$   $\text{H}_2\text{SO}_4$ ,  $\text{HCl}$ ,  $\text{HNO}_3$ ,  $\text{HClO}_4$ ,  $\text{H}_2\text{SO}_5$ ,  $\text{HBr}$ ,  $\text{HI}$ ,  $\text{HBrO}_4$ ,  $\text{HIO}_4$ ,  $\text{RSO}_3\text{H}$   
 (b) Strong base  $\rightarrow$   $\text{KOH}$ ,  $\text{NaOH}$ ,  $\text{Ba}(\text{OH})_2$ ,  $\text{CsOH}$ ,  $\text{RbOH}$   
 (c) All soluble salts  $\rightarrow$   $\text{NaCl}$ ,  $\text{KCl}$ ,  $\text{CuSO}_4$ , .....

(ii) **Weak electrolytes :** Those electrolytes which are partially ionized in aqueous solution are called as weak electrolytes. For weak electrolytes the value of  $\alpha$  is less than one.

- Ex.** (a) **Weak acid :**  $\text{HCN}$ ,  $\text{CH}_3\text{COOH}$ ,  $\text{HCOOH}$ ,  $\text{H}_2\text{CO}_3$ ,  $\text{H}_3\text{PO}_3$ ,  $\text{H}_3\text{PO}_2$ , etc.  
 (b) **Weak base :**  $\text{NH}_4\text{OH}$ ,  $\text{Cu}(\text{OH})_2$ ,  $\text{Zn}(\text{OH})_2$ ,  $\text{Fe}(\text{OH})_3$ ,  $\text{Al}(\text{OH})_3$ , etc.



## 1.2 DEGREE OF DISSOCIATION / IONISATION





- When an electrolyte is dissolved in a solvent ( $H_2O$ ), it spontaneously dissociates into ions.
- It may dissociate partially ( $\alpha < 1$ ) or sometimes completely ( $\alpha \cong 1$ )
- The degree of dissociation ( $\alpha$ ) of an electrolyte is the fraction of mole of the electrolyte that has dissociated under the given conditions.

$$\alpha = \frac{\text{No. of moles dissociated}}{\text{No. of moles taken initially}}$$

## 1.3 FACTORS AFFECTING THE VALUE OF DEGREE OF DISSOCIATION:

- (i) **Dilution** :  $\alpha \propto \sqrt{V}$ . So on dilution,  $\alpha$  increases
- (ii) **Temperature** : On increasing temperature, ionization increases so,  $\alpha$  increases
- (iii) **Nature of electrolyte** :
- |                        |                      |
|------------------------|----------------------|
| (a) Strong electrolyte | (b) Weak electrolyte |
| $\alpha = 100\%$       | $\alpha \ll 100\%$   |
- (iv) **Nature of solvent** :

If dielectric constant,  $\mu$ , of solvent increases, then the value of  $\alpha$  increases.



**Ex.1** Which one has greater  $\alpha_1$  or  $\alpha_2$  for the following equation :




**Sol.** Dielectric constant of  $H_2O$  is more than that of  $D_2O$ , so  $\alpha_1 > \alpha_2$

### (v) Mixing of Ions :

Common ion Effect	Odd ion Effect
$NH_4OH \rightleftharpoons NH_4^+ + OH^-$ On mixing $NH_4Cl$ $NH_4Cl \rightarrow NH_4^+ + Cl^-$ Due to mixing of common ion, concentration of ammonium ion will increase therefore equilibrium will shift in backward direction means $\alpha$ decreases.	$NH_4OH \rightleftharpoons NH_4^+ + OH^-$ On mixing $HCl$ $HCl \rightarrow H^+ + Cl^-$ Due to reaction of $OH^-$ ions with $H^+$ ion, concentration of $OH^-$ will decrease $\therefore$ Equilibrium will shift in forward direction means $\alpha$ increases.

## 1.4 OSTWALD'S DILUTION LAW (FOR WEAK ELECTROLYTE)

- For a weak electrolyte  $A^+B^-$  dissolved in water, if  $\alpha$  is the degree of dissociation then



Then according to law of mass action,


$$K_{\text{diss}} = \frac{[A^+][B^-]}{[AB]} = \frac{C\alpha \cdot C\alpha}{C(1-\alpha)} = \frac{C\alpha^2}{(1-\alpha)} = \text{dissociation constant of the weak electrolyte.}$$

$$\left[ C = \frac{1}{V}, \text{ then } V = 1/C \text{ (volume of solution in which 1 mole is present) is called dilution, so } K_{\text{diss}} = \frac{\alpha^2}{(1-\alpha)V} \right]$$

If  $\alpha$  is negligible in comparison to unity,  $1 - \alpha \approx 1$ . so  $K_{\text{diss}} = \alpha^2 C \Rightarrow \alpha = \sqrt{\frac{K_{\text{diss}}}{C}} = \sqrt{K_{\text{diss}} V}$

$$\alpha \propto \frac{1}{\sqrt{\text{concentration}}}$$

- As concentration increases  $\Rightarrow \alpha$  decreases
- At infinite dilution  $\alpha$  reaches its maximum value, unity.




## 2. ACIDS BASES AND SALTS

### 2.1 Arrhenius concept :

- (i) **Arrhenius Acid :** Substance which gives  $H^+$  ion on dissolving in water ( $H^+$  donor)


**Ex.**  $HNO_3$ ,  $HClO_4$ ,  $HCl$ ,  $HI$ ,  $HBr$ ,  $H_2SO_4$ ,  $H_3PO_4$  etc.

#### Types of acids




- (ii) **Arrhenius base :** Any substance which releases  $OH^-$  (hydroxyl) ion in water ( $OH^-$  ion donor)

#### Types of bases




**(iii) Strength of Acid or Base :**

- (a) Strength of acids or bases depends on the extent of its ionisation. Hence equilibrium constant  $K_a$  or  $K_b$  respectively of the following equilibria give a quantitative measurement of the strength of the acid or base.



$$K_a = \frac{[H^+][A^-]}{[HA]} = \text{dissociation or ionisation constant of acid.}$$

- (c) Similarly




$$K_b = \frac{[B^+][OH^-]}{[BOH]} = \text{dissociation or ionisation constant of base}$$

- (d) Larger the value of  $K_a$  or  $K_b$ , stronger is the acid or base respectively.

**2.2 Bronsted - Lowry concept : (Conjugate acid - base concept) (Protonic concept)**

- (i) **Acid** : substances which donate  $H^+$  are Bronsted Lowry acids ( $H^+$  donor)
- (ii) **Base** : substances which accept  $H^+$  are Bronsted Lowry bases ( $H^+$  acceptor)
- (iii) **Conjugate acid - base pairs** : In a typical acid base reaction,  $HX + B \rightleftharpoons X^- + HB^+$




- Forward reaction - Here HX being a proton donor is an acid  
B being a proton acceptor is a base.
- Backward reaction - Here  $HB^+$  being a proton donor is an acid  
 $X^-$  being a proton acceptor is a base.


Acid	Base		Conjugate Acid		Conjugate Base
• HCl	+ H <sub>2</sub> O	$\rightleftharpoons$	H <sub>3</sub> O <sup>+</sup>	+	Cl <sup>-</sup>
• HSO <sub>4</sub> <sup>-</sup>	+ NH <sub>3</sub>	$\rightleftharpoons$	NH <sub>4</sub> <sup>+</sup>	+	SO <sub>4</sub> <sup>-2</sup>
• [Fe(H <sub>2</sub> O) <sub>6</sub> ] <sup>3+</sup>	+ H <sub>2</sub> O	$\rightleftharpoons$	H <sub>3</sub> O <sup>+</sup>	+	[Fe(H <sub>2</sub> O) <sub>5</sub> (OH)] <sup>2+</sup>
• Conjugate acid - base pair differ by only one proton.					
• Strong acid will have weak conjugate base and vice versa.					

Acid	Conjugate base	Base	Conjugate acid
HCl	$\text{Cl}^-$	$\text{NH}_3$	$\text{NH}_4^+$
$\text{H}_2\text{SO}_4$	$\text{HSO}_4^-$	$\text{H}_2\text{O}$	$\text{H}_3\text{O}^+$
$\text{HSO}_4^-$	$\text{SO}_4^{2-}$	$\text{RNH}_2$	$\text{RNH}_3^+$
$\text{H}_2\text{O}$	$\text{OH}^-$		

(iv) **Amphoteric (amphiprotic)** : Substances which can act as acid as well as base are known as amphoteric



base



acid

(v) **Classification of Bronsted - Lowry Acids and Bases :**

Bronsted - Lowry acids and bases can be

(i) Molecular      (ii) Cationic      and      (iii) Anionic

**Table - 1**

Type	Acid	Base
Molecular	$\text{HCl}$ , $\text{HNO}_3$ , $\text{HClO}_4$ , $\text{H}_2\text{SO}_4$ , $\text{H}_3\text{PO}_4$ , $\text{H}_2\text{O}$ etc.	$\text{NH}_3$ , $\text{N}_2\text{H}_4$ , Amines, $\text{H}_2\text{O}$ , Alcohol, Ethers, etc.
Cationic	$\text{NH}_4^+$ , $\text{N}_2\text{H}_5^+$ , $\text{PH}_4^+$ , $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ , $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$ etc.	$[\text{Fe}(\text{H}_2\text{O})_5\text{OH}]^{2+}$ $[\text{Al}(\text{H}_2\text{O})_5\text{OH}]^{2+}$ etc.
Anionic	$\text{HS}^-$ , $\text{HSO}_3^-$ , $\text{H}_2\text{PO}_4^-$ , $\text{HSO}_4^-$ $\text{HCO}_3^-$ , $\text{HPO}_4^{2-}$ , etc. all amphiprotic anions	$\text{Cl}^-$ , $\text{Br}^-$ , $\text{OH}^-$ , $\text{HSO}_4^-$ , $\text{CN}^-$ , $\text{CO}_3^{2-}$ , $\text{SO}_4^{2-}$ , $\text{NH}_2^-$ , $\text{CH}_3\text{COO}^-$ , etc. all amphiprotic anions

### 2.3 Lewis concept (electronic concept) :

(i) **Acid** : An acid is a molecule/ion which can accept an electron pair with the formation of a coordinate bond.

**Ex.** Electron deficient molecules :  $\text{BF}_3$ ,  $\text{AlCl}_3$ , etc.

Cations :  $\text{H}^+$ ,  $\text{Fe}^{2+}$ ,  $\text{Na}^+$ , etc.

Molecules with vacant orbitals :  $\text{SF}_4$ ,  $\text{PF}_3$

(ii) **Base** : A base is any molecule/ion which has a pair of electrons which can be donated.

**Ex.** Molecules with lone pairs :  $\text{NH}_3$ ,  $\text{PH}_3$ ,  $\text{H}_2\text{O}$ ,  $\text{CH}_3\text{OH}$

Anions :  $\text{OH}^-$ ,  $\text{H}^-$ ,  $\text{NH}_2^-$ , etc.

### 3. PROPERTIES OF WATER

#### (i) Molar concentration / Molarity of water :

$$\text{Molarity} = \text{No. of moles/litre} = \frac{1000 \text{ g/litre}}{18 \text{ g/mole}} = 55.55 \text{ mole/litre} = \mathbf{55.55 \text{ M}} \text{ (density} = 1 \text{ g/cc)}$$

#### (ii) Ionic product of water :

According to arrhenius concept,  $\text{H}_2\text{O} \rightleftharpoons \text{H}^+ + \text{OH}^-$

So, ionic product of water,  $K_w = [\text{H}^+][\text{OH}^-] = 10^{-14}$  at  $25^\circ$  (experimental)

Dissociation of water is endothermic, so on increasing temperature  $K_w$  increases.

#### (iii) Degree of dissociation of water :

$$\begin{aligned} \text{H}_2\text{O} &\rightleftharpoons \text{H}^+ + \text{OH}^- \Rightarrow \alpha = \frac{\text{decrease in concentration}}{\text{initially concentration}} \\ &= \frac{10^{-7}}{55.55} = 18 \times 10^{-10} \text{ or } 1.8 \times 10^{-7} \% \quad [\text{at } 25^\circ\text{C}] \end{aligned}$$

#### (iv) Dissociation or ionisation constant of water :

$$\text{H}_2\text{O} \rightleftharpoons \text{H}^+ + \text{OH}^- \quad K_a = K_b = \frac{[\text{H}^+][\text{OH}^-]}{[\text{H}_2\text{O}]} = \frac{10^{-7} \times 10^{-7}}{55.55} = 1.8 \times 10^{-16}$$

$$\text{So, } pK_a = pK_b = -\log(1.8 \times 10^{-16}) = 16 - \log 1.8 = 15.74$$

**Ex.2.** At dissociation constant of heavy water is  $4 \times 10^{-15}$  at  $35^\circ\text{C}$ . If its density is  $1.04 \text{ g/mL}$ . Calculate its ionic product & degree of dissociation.

$$\text{Sol. } K_w = K_d[D_2O] = \left(4 \times 10^{-15} \times \frac{1040}{20}\right) = 2.08 \times 10^{-13}$$

$$d = \sqrt{\frac{K_w}{C}} = \sqrt{\frac{2.08 \times 10^{-13}}{52}} = 12.64 \times 10^{-8}$$

**Ex.3** Calculate ionic product of  $\text{H}_2\text{O}$  at  $50^\circ\text{C}$ .

$$\text{Sol. } \Delta H = 13.7 \times 10^3 \text{ cal}$$

$$\log \frac{K_2}{10^{-14}} = \frac{13.7 \times 10^3}{2} \left( \frac{1}{298} - \frac{1}{323} \right)$$

**Ex.4** The hydronium ion conc. in an aq.  $\text{H}_2\text{CO}_3$  solution is  $4 \times 10^{-4} \text{ M}$  at  $25^\circ\text{C}$   $\text{OH}^-$  ion conc. in the solution is :

- (A) 0                      (B)  $2.5 \times 10^{-10}$                       (C)  $2.5 \times 10^3$                       (D)  $2.5 \times 10^{-11} \text{ M}$

Answer : (D)

**Ex.5** Select the correct option from the following ?


- (A)  $pK_w$  increases with increase of temperature  
 (B)  $pK_w$  decreases with increase of temperature  
 (C)  $pK_w = 14$  at all temperatures  
 (D)  $pK_w = pH$  at all temperatures

Sol. (B)

#### 4. Acidity and pH scale :

- (i) Acidic strength means the tendency of an acid to give  $\text{H}_3\text{O}^+$  or  $\text{H}^+$  ions in water.  
So greater then tendency to give  $\text{H}^+$ , more will be the acidic strength of the substance.
- (ii) Basic strength means the tendency of a base to give  $\text{OH}^-$  ions in water.  
So greater the tendency to give  $\text{OH}^-$  ions, more will be basic strength of the substance.
- (iii) The concentration of  $\text{H}^+$  ions is written in a simplified form introduced by **Sorenson** known as pH scale. pH is defined as negative logarithm of activity of  $\text{H}^+$  ions.  
 $\therefore \text{pH} = -\log a_{\text{H}^+}$  (where  $a_{\text{H}^+}$  is the activity of  $\text{H}^+$  ions)
- (iv) Activity of  $\text{H}^+$  ions is the molar concentration of free  $\text{H}^+$  ions or  $\text{H}_3\text{O}^+$  ions in a dilute solution, but unitless.
- (v) Now  $\text{pH} = -\log[\text{H}^+] = 7$  and  $\text{pOH} = -\log[\text{OH}^-] = 7$  for water at  $25^\circ\text{C}$  (experimental)
 

$\text{pH} = 7 = \text{pOH} \Rightarrow \text{neutral}$   
 $\text{pH} < 7 \text{ or } \text{pOH} > 7 \Rightarrow \text{acidic}$   
 $\text{pH} > 7 \text{ or } \text{pOH} < 7 \Rightarrow \text{Basic}$




#### 4.1 pH Calculation of different Types of solutions :

##### (a) Strong acid solution :

- (i) If concentration of  $\text{H}^+$  ions is greater than  $10^{-6} \text{ M}$ ,  $\text{H}^+$  ions coming from water can be neglected,  
So  $[\text{H}^+] = \text{normality of strong acid solution}$
- (ii) If concentration is less than  $10^{-6} \text{ M}$ ,  $\text{H}^+$  ions coming from water cannot be neglected. So  
 $[\text{H}^+] = \text{normality of strong acid} + \text{H}^+ \text{ ions coming from water in presence of this strong acid}$

**Ex.6 Calculate pH of  $10^{-8} \text{ M HCl}$  solution.**



$$K_w = [\text{H}^+][\text{OH}^-]$$

$$10^{-14} = x(x + 10^{-8})$$

$$\Rightarrow x^2 + x \times 10^{-8} - 10^{-14} = 0$$

$$x = \frac{-10^{-8} \pm \sqrt{10^{-16} + 4 \times 10^{-14}}}{2} = \frac{-10^{-8} + 10^{-7} \sqrt{4 + \frac{1}{100}}}{2} = \frac{(\sqrt{401} - 1)10^{-8}}{2} = 0.95 \times 10^{-7}$$

$$[\text{H}^+] = 10.5 \times 10^{-8} = 1.05 \times 10^{-7}$$

$$\text{pH} = -\log [\text{H}^+]$$

$$\text{pH} = 7 - \log 1.05 \approx 6.98$$

### 4.2 Strong base solution :


Calculate the  $[\text{OH}^-]$  which will be equal to normality of the strong base solution and then use

$$K_w = [\text{H}^+] \times [\text{OH}^-] = 10^{-14}, \text{ to calculate } [\text{H}^+]$$

#### Ex.7 Calculate pH of $10^{-7} \text{ M}$ of NaOH solution

**Sol.**  $[\text{OH}^-]$  from NaOH =  $10^{-7}$

$[\text{OH}^-]$  from water =  $x < 10^{-7} \text{ M}$  (due to common ion effect)



$$K_w = [\text{H}^+] [\text{OH}^-] = 10^{-14} = x (x + 10^{-7})$$

$$x^2 + 10^{-7}x - 10^{-14} = 0$$

$$\Rightarrow x = \frac{\sqrt{5}-1}{2} \times 10^{-7} = 0.618 \times 10^{-7} \quad (\sqrt{5} = 2.236)$$

$$[\text{OH}^-] = 10^{-7} + 0.618 \times 10^{-7} = 1.618 \times 10^{-7}$$

$$\text{pOH} = 7 - \log(1.618) = 6.79$$

$$\text{pH} = 14 - 6.79 = 7.21$$

### 4.3 pH of mixture of two strong acids : If $V_1$ volume of a strong acid solution of normality $N_1$ is mixed with $V_2$ volume of another strong acid solution of normality $N_2$ , then

$$\text{Number of } \text{H}^+ \text{ ions from I-solution} = N_1 V_1$$

$$\text{Number of } \text{H}^+ \text{ ions from II-solution} = N_2 V_2$$

If final normality is  $N$  and final volume is  $V$ , then

$$NV = N_1 V_1 + N_2 V_2$$

[dissociation equilibrium of none of these acids will be disturbed as both are strong acid]

$$[\text{H}^+] = N = \frac{N_1 V_1 + N_2 V_2}{V_1 + V_2} \quad \left[ \begin{array}{l} \text{where } N = M \times n \\ n = \text{Basicity of acid} \end{array} \right]$$

### 4.4 pH of mixture of two strong bases :

Similar to above calculation

$$[\text{OH}^-] = N = \frac{N_1 V_1 + N_2 V_2}{V_1 + V_2} \quad [\text{H}^+] = \frac{10^{-14}}{[\text{OH}^-]}$$

$$\left[ \begin{array}{l} \text{where } N = M \times n \\ n = \text{Acidity of base} \end{array} \right]$$

#### Ex.8 Calculate pH of mixture of $(400 \text{ mL}, \frac{1}{200} \text{ M H}_2\text{SO}_4) + (400 \text{ mL}, \frac{1}{100} \text{ M HCl}) + (200 \text{ mL of water})$

**Sol.**  $N_1 V_1 = \frac{1}{200} \times \frac{400}{1000} \times 2 = \frac{4}{1000}$ ,  $N_2 V_2 = \frac{4}{1000}$ ,  $\text{H}^+$  ions from water will be neglected

$$N_1 V_1 + N_2 V_2 = 8 \times 10^{-3} \quad [\text{H}^+] = \frac{8 \times 10^{-3}}{1} = 8 \times 10^{-3}$$

$$\text{pH} = 3 - \log 8 = 2.1$$



**Ex.9** 500 mL of  $10^{-5}$  M NaOH is mixed with 500 mL of  $2.5 \times 10^{-5}$  M of  $\text{Ba}(\text{OH})_2$ . To the resulting solution 99 L water is added. Calculate pH.

**Sol.**  $[\text{OH}^-] = \frac{500 \times 10^{-5} + 500 \times 2 \times 2.5 \times 10^{-5}}{1000} = 3 \times 10^{-5} \text{ M}$

$$M_1 = 3 \times 10^{-5} \text{ M}$$


$$V_2 + V_1 = 1 \text{ L}$$

$$V_F = 100 \text{ L}$$

no. of moles of  $[\text{OH}^-]$  initially = no. of moles of  $[\text{OH}^-]$

$$3 \times 10^{-5} = M_2 \times 100$$

$$\therefore M_2 = 3 \times 10^{-7} < 10^{-6}$$



$$x \quad (x + 3 \times 10^{-7})$$

$$K_w = x(x + 3 \times 10^{-7}) = 10^{-14}$$

$$\therefore x = \left( \frac{\sqrt{13} - 3}{2} \right) \times 10^{-7}$$

$$x = 0.302 \times 10^{-7}$$

$$[\text{OH}^-]_{\text{Net}} = 3.302 \times 10^{-7}$$

#### 4.5 pH of mixture of a strong acid and a strong base :

- Acid Base neutralisation reaction will take place.
- The solution will be acidic or basic depending on which component has been taken in excess.
- If  $V_1$  volume of a strong acid solution of normality  $N_1$  is mixed with  $V_2$  volume of a strong base solution of normality  $N_2$ , then

$$\text{Number of } \text{H}^+ \text{ ions from I-solution} = N_1 V_1$$

$$\text{Number of } \text{OH}^- \text{ ions from II-solution} = N_2 V_2$$

$\downarrow$ If $N_1 V_1 > N_2 V_2$ $[\text{H}^+] = N = \frac{N_1 V_1 - N_2 V_2}{V_1 + V_2}$	$\downarrow$ If $N_2 V_2 > N_1 V_1$ $[\text{OH}^-] = N = \frac{N_2 V_2 - N_1 V_1}{V_1 + V_2}$
--	---

Solution will  
be acidic in nature

Solution will  
be basic in nature

$$[\text{H}^+] = \frac{10^{-14}}{[\text{OH}^-]}$$

**Ex.10** Calculate pH of mixture of (400 mL,  $\frac{1}{200}$  M Ba(OH)<sub>2</sub>) + (400 mL,  $\frac{1}{50}$  M HCl) + (200 mL of water)

**Sol.**  $[H^+] = \frac{400 \times \frac{1}{50} - 400 \times \frac{1}{200} \times 2}{1000} = 4 \times 10^{-3}$ , so  $pH = 3 - 2 \log 2 = 2.4$

**Ex.11** What will be the resultant pH when 150 mL of an aqueous solution of HCl (pH = 2.0) is mixed with 350 mL of an aqueous solution of NaOH (pH = 12.0) ?

**Sol.** pH of HCl = 2

$\therefore [HCl] = 10^{-2}$  M

pH of NaOH = 12, pOH = 2  $\therefore [NaOH] = 10^{-2}$  M

	HCl	+	NaOH	→	NaCl	+	H <sub>2</sub> O
Meq. initial	$150 \times 10^{-2}$		$350 \times 10^{-2}$		0		0
	= 1.5		= 3.5				
Meq. final	0		2		1.5		1.5

$\therefore [OH^-] \text{ from NaOH} = \frac{2}{500} = 4 \times 10^{-3}$  M


pOH =  $-\log[OH^-] = -\log(4 \times 10^{-3})$

$\therefore$  pOH = 2.3979

$\therefore$  pH = 14 - pOH = 14 - 2.3979 = 11.6021

#### 4.6 pH of a weak acid or weak base (monoprotic) Solution :

- Weak acid does not dissociated 100 % therefore we have to calculate the percentage dissociation using  $K_a$  dissociation constant of the acid.
- We have to use Ostwald's Dilution law (as have been derived earlier)




If  $\alpha \ll 1 \Rightarrow (1 - \alpha) \approx 1 \Rightarrow K_a \approx C\alpha^2 \Rightarrow \alpha = \sqrt{\frac{K_a}{C}}$  (is valid if  $\alpha < 0.1$  or 10%)

$[H^+] = C\alpha = C\sqrt{\frac{K_a}{C}} = \sqrt{K_a \times C}$  So  $pH = \frac{1}{2}(pK_a - \log C)$

On increasing the dilution  $\Rightarrow C \downarrow = \alpha \uparrow$  and  $[H^+] \downarrow \Rightarrow pH \uparrow$

**Ex.12 Calculate pH of : (a)  $10^{-1} M CH_3COOH$  (b)  $10^{-3} M CH_3COOH$  (c)  $10^{-6} M CH_3COOH$**

Take  $K_a = 2 \times 10^{-5}$



$$\begin{array}{ccc} C & 0 & 0 \\ C(1-\alpha) & C\alpha & C\alpha \end{array}$$

$$K_a = \frac{C\alpha^2}{1-\alpha} \Rightarrow \alpha = \sqrt{\frac{K_a}{C}} = \sqrt{\frac{2 \times 10^{-5}}{10^{-1}}} = \sqrt{2 \times 10^{-4}} \quad (\alpha \ll 0.1)$$

$$\text{So, } [H^+] = 10^{-1} \times \sqrt{2} \times 10^{-2} \Rightarrow \text{pH} = 3 - \frac{1}{2} \log 2 = 2.85$$

**(b)**  $\alpha = \sqrt{\frac{K_a}{C}} \Rightarrow \alpha = \sqrt{\frac{K_a}{C}} = \sqrt{\frac{2 \times 10^{-5}}{10^{-3}}} = \sqrt{2 \times 10^{-2}} \quad (\alpha > 0.1)$

So we have to do the exact calculations

$$K_a = \frac{C\alpha^2}{1-\alpha} \Rightarrow 2 \times 10^{-5} = \frac{10^{-3} \times \alpha^2}{1-\alpha} \Rightarrow \alpha = 13.14 \%$$

$$[H^+] = 10^{-3} \times 0.1314 = 1.314 \times 10^{-4} \Rightarrow \text{pH} = 4 - \log(1.314) \approx 3.8$$

**(c)** If approximation is used the,  $\alpha = \sqrt{\frac{2 \times 10^{-5}}{10^{-6}}} = \sqrt{20} > 1$ ,

So we have to do the exact calculation,  $2 \times 10^{-5} = 10^{-6} \frac{\alpha^2}{1-\alpha} \Rightarrow \alpha \approx 0.95$  or 95%


$$[H^+] = 0.95 \times 10^{-6} = 9.5 \times 10^{-7} \Rightarrow \text{pH} = 7 - \log(9.5) = 6.022$$

- At very low concentration (at infinite dilution) weak electrolyte will be almost 100% dissociate, so behave as strong electrolyte.

$$(\text{pH}) \text{ of } 10^{-6} M HCl \approx \text{pH of } 10^{-6} M CH_3COOH \approx 6)$$

**Ex.13  $K_a$  for acid HA is  $2.5 \times 10^{-8}$  calculate for its decimolar solution at  $25^\circ C$ .**

**(i) % dissociation (ii) pH (iii)  $OH^-$  ion concentration**



$$\begin{array}{ccc} C & 0 & 0 \\ C(1-\alpha) & C\alpha & C\alpha \end{array}$$

$$K_a = \frac{[H^+][A^-]}{[HA]} \Rightarrow K_a = \frac{C\alpha \cdot C\alpha}{C(1-\alpha)} = \frac{C\alpha^2}{(1-\alpha)} \approx C\alpha^2$$

**(i)**  $\therefore \alpha = \sqrt{\frac{K_a}{C}} = \sqrt{\frac{2.5 \times 10^{-8}}{1/10}} \quad (C = 1/10 M)$   
 $= 5 \times 10^{-4} = 0.05\%$


**(ii)**  $[H^+] = C\alpha = \frac{1}{10} \times 5 \times 10^{-4} = 5 \times 10^{-5} \text{ mol/L}$

So  $\text{pH} = 5 - \log 5 = 4.30$

**(iii)**  $[H^+][OH^-] = 1 \times 10^{-14}$

$$\therefore [OH^-] = \frac{10^{-14}}{5 \times 10^{-5}} = 2 \times 10^{-10} \text{ mol/L}$$

**Ex.14** Determine the degree of dissociation of 0.05 M  $\text{NH}_4\text{OH}$  at  $25^\circ\text{C}$  in a solution of  $\text{pH} = 10$ .



Given,  $\text{pH} = 10$

$$[\text{H}^+] = 10^{-10}$$

$$[\text{H}^+][\text{OH}^-] = 1 \times 10^{-14}$$

$$\therefore [\text{OH}^-] = \frac{1 \times 10^{-14}}{10^{-10}} = 10^{-4} = C\alpha$$

$$\therefore \alpha = \frac{[\text{OH}^-]}{C} = \frac{10^{-4}}{0.05} = 2 \times 10^{-3} \text{ or } 0.2 \%$$

**Ex.15** The concentration of  $[\text{H}^+]$  and  $[\text{OH}^-]$  of the  $10^{-1}$  M aqueous solution of 2% ionised weak acid is :

(A)  $2 \times 10^{-3}$  M and  $5 \times 10^{-12}$  M

(B)  $1 \times 10^{-3}$  M and  $3 \times 10^{-11}$  M

(C)  $2 \times 10^{-4}$  M and  $5 \times 10^{-11}$  M

(D)  $3 \times 10^{-2}$  M and  $4 \times 10^{-13}$  M

**Sol.** (A)

$$[\text{H}^+] = C\alpha = 2 \times 10^{-3} \text{ M or } [\text{OH}^-] = \frac{10^{-14}}{[\text{H}^+]} = 5 \times 10^{-12} \text{ M}$$

**Ex.16** When a 0.1 N solution of an acid at  $25^\circ\text{C}$  has a degree of ionisation of 4%, the concentration of  $\text{OH}^-$  present is :

(A)  $2.5 \times 10^{-3}$

(B)  $2.5 \times 10^{-11}$


(C)  $2.5 \times 10^{-12}$

(D)  $2.5 \times 10^{-13}$

**Sol.** (C)

$$[\text{H}^+] = C\alpha = 0.1 \times 4 \times 10^{-2} = 4 \times 10^{-3} \text{ M or } [\text{OH}^-] = \frac{10^{-14}}{[\text{H}^+]} = 2.5 \times 10^{-12} \text{ N}$$

**Ex.17** The degree of dissociation of acetic acid in a 0.1 M solution is  $1.32 \times 10^{-2}$ . Calculate dissociation constant of acid and its  $\text{pK}_a$  value :



$$K_a = \frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{[\text{CH}_3\text{COOH}]} = \frac{0.1 \times 0.0132 \times 0.1 \times 0.0132}{0.1(1-0.0132)} = 1.76 \times 10^{-5}$$

$$\text{pK}_a = -\log K_a = -\log (1.76 \times 10^{-5}) = 4.75$$

**4.7 pH of a mixture of weak acid (monoprotic) and a strong acid solution :**

- Weak acid and Strong acid both will contribute  $H^+$  ion.
- For the first approximation we can neglect the  $H^+$  ions coming from the weak acid solution and calculate the pH of the solution from the concentration of the strong acid only.
- To calculate exact pH, we have to take the effect of presence of strong acid on the dissociation equilibrium of the weak acid.
- If  $[SA] = C_1$  and  $[WA] = C_2$ , then  $[H^+]$  from  $SA = C_1$  the weak acid will dissociate as follows.



(The weak acids dissociation will be further suppressed because of presence of strong acid, common ion effect)


$$K_a = (C_2\alpha + C_1)\alpha$$

Total  $H^+$  ion concentration =  $C_1 + C_2\alpha$

- If the total  $[H^+]$  from the acid is more than  $10^{-6}$  M, then contribution from the water can be neglected, if not then we have to take  $[H^+]$  from the water also.

**4.8 pH of a mixture of two weak acid (both monoprotic) solution :**

- Both acids will dissociate partially.
- Let the acid are  $HA_1$  &  $HA_2$  and their final concentrations are  $C_1$  &  $C_2$  respectively, then



(Since  $\alpha_1, \alpha_2$  both are small in comparison to unity)

$$K_{a_1} = (C_1\alpha_1 + C_2\alpha_2)\alpha_1 ; K_{a_2} = (C_1\alpha_1 + C_2\alpha_2)\alpha_2 \Rightarrow \frac{K_{a_1}}{K_{a_2}} = \frac{\alpha_1}{\alpha_2}$$

$$\begin{aligned}
 [H^+] = C_1\alpha_1 + C_2\alpha_2 &= \frac{C_1K_{a_1}}{\sqrt{C_1K_{a_1} + C_2K_{a_2}}} + \frac{C_2K_{a_2}}{\sqrt{C_1K_{a_1} + C_2K_{a_2}}} \Rightarrow [H^+] \\
 &= \sqrt{C_1K_{a_1} + C_2K_{a_2}}
 \end{aligned}$$

- If the dissociation constant of one of the acid is very much greater than that of the second acid then contribution from the second acid can be neglected.

$$\text{So, } [H^+] = C_1\alpha_1 + C_2\alpha_2 \approx C_1\alpha_1$$

**Ex.18** Calculate pH of solution obtained by mixing equal vol. of 0.02 M HOCl & 0.2 M CH<sub>3</sub>COOH solution given that  $K_{a_1}(\text{HOCl}) = 2 \times 10^{-4}$ ,  $K_{a_2}(\text{CH}_3\text{COOH}) = 2 \times 10^{-5}$

Also calculate OH<sup>-</sup>, OCl<sup>-</sup>, CH<sub>3</sub>COO<sup>-</sup>

**Sol.** Final solution volume become double



$$C_1 = 0.01, \quad C_2 = 0.1$$

$$[\text{H}^+] = \sqrt{K_{a_1}C_1 + K_{a_2}C_2} = \sqrt{2 \times 10^{-4} \times 0.01 + 2 \times 10^{-5} \times 0.1}$$

$$= \sqrt{2 \times 10^{-6} + 2 \times 10^{-6}} = 2 \times 10^{-3}$$

$$\text{pH} = 3 - \log 2 = 3 - 0.3010 = 2.69$$

$$\alpha_1 = \frac{2 \times 10^{-4}}{2 \times 10^{-3}} = 10^{-1} \quad \alpha_2 = \frac{2 \times 10^{-5}}{2 \times 10^{-3}} = 10^{-2}$$



$$[\text{OH}^-] = \frac{K_w}{[\text{H}^+]} = \frac{10^{-14}}{2 \times 10^{-3}} = 0.5 \times 10^{-11} = 5 \times 10^{-12} \text{ M}$$

$$[\text{HOCl}] = 10^{-2}(1 - 0.1) = 9 \times 10^{-3} \text{ M}$$

$$[\text{CH}_3\text{COOH}] = 10^{-1}(1 - 0.01) \approx 10^{-1}$$

#### 4.9 pH of a solution of a polyprotic weak acid :

- Diprotic acid is the one, which is capable of giving 2 protons per molecule in water. Let us take a weak diprotic acid (H<sub>2</sub>A) in water whose concentration is c M.


In an aqueous solution, following equilibria exist.

If

$\alpha_1$  = degree of ionization of H<sub>2</sub>A in presence of HA<sup>-</sup>       $K_{a_1}$  = first ionisation constant of H<sub>2</sub>A

$\alpha_2$  = degree of ionisation of HA<sup>-</sup> in presence of H<sub>2</sub>A       $K_{a_2}$  = second ionisation constant of H<sub>2</sub>A


##### I step



$$\text{at eq. } c(1 - \alpha_1) \quad c\alpha_1(1 - \alpha_2) \quad (c\alpha_1 + c\alpha_1\alpha_2)$$

$$(K_{\text{eq}})_1 [\text{H}_2\text{O}] = \frac{[\text{H}_3\text{O}^+][\text{HA}^-]}{[\text{H}_2\text{A}]} = K_{a_1}$$

##### II step



$$\text{at eq. } c\alpha_1(1 - \alpha_2) \quad c\alpha_1\alpha_2 \quad (c\alpha_1 + c\alpha_1\alpha_2)$$

$$(K_{\text{eq}})_2 [\text{H}_2\text{O}] = \frac{[\text{H}_3\text{O}^+][\text{A}^{2-}]}{[\text{HA}^-]} = K_{a_2}$$

$$\therefore K_{a_1} = \frac{(c\alpha_1 + c\alpha_1\alpha_2)[c\alpha_1(1-\alpha_2)]}{c(1-\alpha_1)} \quad K_{a_2} = \frac{(c\alpha_1 + c\alpha_1\alpha_2)[c\alpha_1\alpha_2]}{c\alpha_1(1-\alpha_2)}$$

$$= \frac{[c\alpha_1(1+\alpha_2)][\alpha_1(1-\alpha_2)]}{1-\alpha_1} \dots\dots (i) \quad = \frac{[c\alpha_1(1+\alpha_2)]\alpha_2}{1-\alpha_2} \dots\dots (ii)$$

Knowing the values of  $K_{a_1}$ ,  $K_{a_2}$  and  $c$ , the values of  $\alpha_1$  and  $\alpha_2$  can be calculated using equations (i) and (ii) After getting the values of  $\alpha_1$  and  $\alpha_2$ ,  $[H_3O^+]$  can be calculated as

$$[H_3O^+]_T = c\alpha_1 + c\alpha_1\alpha_2$$

Finally, for calculation of pH

- If the total  $[H_3O^+] < 10^{-6} M$ , the contribution of  $H_3O^+$  from water should be added.
- If the total  $[H_3O^+] > 10^{-6} M$ , then  $[H_3O^+]$  contribution from water can be ignored.

Using this  $[H_3O^+]$ , pH of the solution can be calculated.

### ❖ Approximation :

For diprotic acids,  $K_{a_2} \ll K_{a_1}$  and  $\alpha_2$  would be even smaller than  $\alpha_1$

$$\therefore 1 - \alpha_2 \approx 1 \text{ and } 1 + \alpha_2 \approx 1$$

Thus, equation (i) can be reduced to  $K_{a_1} = \frac{C\alpha_1 \times \alpha_1}{1 - \alpha_1}$

This is expression similar to the expression for a weak monoprotic acid.

- Hence, for a diprotic acid (or a polyprotic acid) the  $[H_3O^+]$  can be calculated from its first equilibrium constant expression alone provided  $K_{a_2} \ll K_{a_1}$ .


**Ex.19 Calculate pH of  $[HS^-]$ ,  $[S^{2-}]$ ,  $[Cl^-]$  in a solution which is 0.1 M HCl & 0.1 M  $H_2S$  given that  $K_{a_1}(H_2S) = 10^{-7}$ ,  $K_{a_2}(H_2S) = 10^{-14}$  also calculate  $\alpha_1$  &  $\alpha_2$ .**

**Sol.**  $HCl + H_2S$

$$0.1 \quad 0.1$$


$$C_1 = C_2 = 0.1$$

$$\therefore pH = 1 \quad (\text{most of } [H^+] \text{ comes from HCl})$$



$$K_{a_1} = \frac{C\alpha_1 \times 10^{-1}}{C(1 - \alpha_1)} = \frac{10^{-7}}{10^{-1}} = \alpha_1 \quad (\because 1 - \alpha_1 = 1)$$

$$\Rightarrow \alpha_1 = 10^{-6}$$



$$\Rightarrow \alpha_2 = 10^{-13}$$

$$[S^{2-}] = C\alpha_1\alpha_2$$

$$= 10^{-6} \times 10^{-1} \times 10^{-13} = 10^{-20} M$$

#### 4.10 ISOHYDRIC SOLUTIONS

- (i) Solutions of electrolytes are said to be isohydric if the concentration of the common ion present in them is the same and on mixing such solutions, there occurs no change in the degree of dissociation of either of the electrolyte.
- (ii) Let the isohydric solution is made by  $HA_1$  and  $HA_2$  acids, then  $[H^+]$  of both acids should be equal i.e.

$$\sqrt{K_{a_1} C_1} = \sqrt{K_{a_2} C_2} \quad \text{or} \quad \frac{K_{a_1}}{K_{a_2}} = \frac{C_2}{C_1}$$

#### 4.11 RELATIVE STRENGTH OF WEAK ACIDS AND BASES

For two acids of equimolar concentrations.

$$\frac{\text{Strength of acid (I)}}{\text{Strength of acid (II)}} = \sqrt{\frac{K_{a_1}}{K_{a_2}}}$$

Similarly for bases, 
$$\frac{\text{Strength of base (I)}}{\text{Strength of base (II)}} = \sqrt{\frac{K_{b_1}}{K_{b_2}}}$$

The modern method is to convert  $K_a$  as a power of 10 and express acid strength by power of 10 with sign changed and call this new unit  $pK_a$ . Thus, if  $K_a$  for acid is equal to  $10^{-4}$ ,  $pK_a = 4$ . So higher  $pK_a$  value means lower acid strength, that is,  $pK_a = -\log K_a$

Also,  $pK_b = -\log K_b$

### 5. SALTS

- (i) Salts are the ionic compounds formed when its positive part (Cation) come from a base and its negative part (Anion) come from an acid.
- (ii) Salts may taste salty, bitter or sweet or tasteless.
- (iii) Solution of salts may be acidic, basic or neutral.
- (iv) Fused salts and their aqueous solutions conduct electricity and undergo electrolysis.
- (v) The salts are generally crystalline solids.

#### 5.1 Classification of salts :

The salts may be classified into four categories.

##### (a) Normal salt :

- (i) The salt formed by the loss of all possible protons (replaceable  $H^+$  ions)

**Ex.**  $NaCl$ ,  $NaNO_3$ ,  $K_2SO_4$ ,  $Ca_3(PO_4)_2$ ,  $Na_3BO_3$ ,  $Na_2HPO_3$ ,  $NaH_2PO_2$  etc.

##### (b) Acid salts :

- (i) Salts formed by incomplete neutralisation of polybasic acids. Such salts contain one or more replaceable H atom.

**Ex.**  $NaHCO_3$ ,  $NaHSO_4$ ,  $NaH_2PO_4$ ,  $Na_2HPO_4$  etc.

- (ii) Above salts when neutralized by base form normal salts.

##### (c) Basic salts :



- (i) Salts formed by in complete neutralisation of poly acidic bases are called basic salts. These salt contain one or more hydroxyl groups.


**Ex.**  $\text{Zn(OH)Cl}$ ,  $\text{Mg(OH)Cl}$ ,  $\text{Fe(OH)}_2\text{Cl}$ ,  $\text{Bi(OH)}_2\text{Cl}$  etc.

- (ii) Above salts when neutralised by acids form normal salts.

## 5.2 HYDROLYSIS OF SALTS



Salt hydrolysis is defined as the process in which water reacts with cation or anion or both of a salt to change the concentration of  $\text{H}^+$  and  $\text{OH}^-$  ions of water.

Salt hydrolysis is reverse process of neutralization.




### 5.2.1 Hydrolysis of strong acid - weak base [SA - WB] type salt –

**Ex.**  $\text{CaSO}_4$ ,  $\text{NH}_4\text{Cl}$ ,  $(\text{NH}_4)_2\text{SO}_4$ ,  $\text{Ca(NO}_3)_2$ ,  $\text{ZnCl}_2$ ,  $\text{CuCl}_2$ ,  $\text{CaCl}_2$




- (i) In this type of salt hydrolysis, cation reacts with  $\text{H}_2\text{O}$ , therefore called as **cationic hydrolysis**.  
 (ii) Solution is acidic in nature (SAWB) as  $[\text{H}^+]$  is increased.  
 (iii) pH of the solution is less than 7.  
 (iv) **Relation between  $K_h$ ,  $K_w$  &  $K_b$**



$$\text{Hydrolysis constant } K_h = \frac{[\text{NH}_4\text{OH}][\text{H}^+]}{[\text{NH}_4^+]} \quad \dots(i)$$



$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]} \quad \dots(ii)$$



$$K_w = [\text{OH}^-][\text{H}^+] \quad \dots(iii)$$


Now multiplying Eq. (1) & (2) = Eq. (3)

$$\frac{[\text{NH}_4\text{OH}][\text{H}^+]}{[\text{NH}_4^+]} \times \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]} = [\text{H}^+][\text{OH}^-]$$

$$\text{i.e. } K_h \times K_b = K_w$$

$$\boxed{K_h = \frac{K_w}{K_b}}$$

- (v) **Degree of hydrolysis – (Represented by h)**



$$K_h = \frac{[\text{NH}_4\text{OH}][\text{H}^+]}{[\text{NH}_4^+]} = \frac{\text{Ch}^2}{(1-h)}$$

Since  $h \ll \ll \ll 1$

then  $(1-h) \approx 1$

$$\therefore K_h = \text{Ch}^2$$

$$\Rightarrow h = \sqrt{\frac{K_h}{C}}$$

$$\therefore \Rightarrow h = \sqrt{\frac{K_w}{K_b C}} \Rightarrow \boxed{h = \sqrt{\frac{K_w}{K_b \times C}}}$$

(vi) **pH of the solution :**

$$\text{pH} = -\log [\text{H}^+]$$

$$\Rightarrow [\text{H}^+] = \sqrt{\frac{K_w \times C}{K_b}}$$

On taking  $-\log$  on both sides

$$\text{pH} = -\log \left( \frac{K_w \times C}{K_b} \right)^{1/2}$$

$$\text{pH} = -\frac{1}{2} \log K_w - \frac{1}{2} \log C - \frac{1}{2} (-\log K_b)$$

$$\text{pH} = 7 - \frac{1}{2} \text{p}K_b - \frac{1}{2} \log C$$

**Ex.20** Find out the  $K_h$  of centi normal [ $10^{-2}$  N] solution of  $\text{NH}_4\text{Cl}$  (SA - WB) if dissociation constant of  $\text{NH}_4\text{OH}$  is  $10^{-6}$  and  $K_w = 10^{-14}$ . Find out degree of hydrolysis and also find  $[\text{H}^+]$  and pH of solution?

(Given :  $K_w = 10^{-14}$  ;  $K_b = 10^{-6}$ )

**Sol.** (1)  $K_h = \frac{K_w}{K_b} = \frac{10^{-14}}{10^{-6}} = 10^{-8}$

(2)  $h = \sqrt{\frac{K_h}{C}} = \sqrt{\frac{10^{-8}}{10^{-2}}} = \sqrt{10^{-6}} = 10^{-3}$

(3)  $[\text{H}^+] = \text{Ch}$   
 $= 10^{-2} \times 10^{-3} = 10^{-5}$

(4)  $\text{pH} = -\log [\text{H}^+] = -\log [10^{-5}] = +5 \log 10 = +5 \times 1 = 5$

**Ex.21** How many grams of  $\text{NH}_4\text{Cl}$  should be dissolved per litre of solution to have a pH of 5.13 ?  $K_b$  for  $\text{NH}_3$  is  $1.8 \times 10^{-5}$ .

**Sol.**  $\text{NH}_4\text{Cl}$  is a salt of strong acid and weak base for solutions of such salts.

$$\text{pH} = \frac{1}{2} [\text{p}K_w - \log C - \text{p}K_b]$$

$$\Rightarrow 10.26 = 14 - \log C - 4.74$$

$$\Rightarrow \log C = 9.26 - 10.26 = -1.0$$

$$\therefore C = 10^{-1} \text{ M}$$




$$[\text{NH}_4\text{Cl}] = 10^{-1} \text{ M}$$

$$W_{\text{NH}_4\text{NO}_3} = 10^{-1} \times 53.5 \text{ gL}^{-1}$$

$$= 5.35 \text{ gL}^{-1}$$

### 5.2.2 Hydrolysis of [WA – SB] type salt :

**Ex.** KCN, NaCN,  $\text{K}_2\text{CO}_3$ ,  $\text{BaCO}_3$ ,  $\text{K}_3\text{PO}_4$




(i) In this type of salt hydrolysis anion reacts with water therefore called as anionic hydrolysis.

(ii) Solution is basic in nature as  $[\text{OH}^-]$  increases.

(iii) pH of the solution is greater than 7.

(iv) **Relation between  $K_h$ ,  $K_w$ ,  $K_a$**




$$K_h = \frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]} \quad \dots (i)$$

$$\frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]} \times \frac{[\text{CN}^-][\text{H}^+]}{[\text{HCN}]} = [\text{H}^+][\text{OH}^-]$$

$$\boxed{K_h = \frac{K_w}{K_a}}$$

(v) **Degree of hydrolysis :**



Initial concentration at equilibrium

$$K_h = \frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]}$$

$$K_h = \frac{\text{Ch}^2}{(1-h)}$$

Since  $h \ll 1$ , therefore  $(1-h) \approx 1$

$$\therefore K_h = \text{Ch}^2$$

$$h^2 = \frac{K_h}{C} \quad \Rightarrow \quad \boxed{h = \sqrt{\frac{K_h}{C}}}$$

$$\boxed{h = \sqrt{\frac{K_w}{K_a \times C}}}$$

## (vi) pH of the solution

$$[\text{OH}^-] = Ch$$

$$[\text{OH}^-] = \sqrt{\frac{K_w \times C}{K_a}}$$

$$[\text{H}^+] = \frac{K_w}{\sqrt{\frac{K_w \times C}{K_a}}} \Rightarrow [\text{H}^+] = \sqrt{\frac{K_w \times K_a}{C}}$$


On taking  $-\log$  on both sides

$$\text{pH} = -\frac{1}{2} [\log K_w + \log K_a - \log C]$$

$$\text{pH} = 7 + \frac{1}{2} \text{p}K_a + \frac{1}{2} \log C$$

**Ex.22** Calculate the pH and degree of hydrolysis of 0.01 M solution of NaCN,  $K_a$  for HCN is  $6.2 \times 10^{-12}$ .

**Sol.** NaCN is a salt of strong base NaOH and weak acid HCN.  $\text{Na}^+$  does not react with water whereas  $\text{CN}^-$  reacts with water as here under



$$K_h = \frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]} = \frac{K_w}{K_a} = \frac{10^{-14}}{6.2 \times 10^{-12}} = 1.6 \times 10^{-3}$$

Let, x moles of salt undergo hydrolysis then concentrations of various species would be

$$[\text{CN}^-] = (0.01 - x) \approx 0.01, [\text{HCN}] = x$$

$$[\text{OH}^-] = x$$

$$\therefore K_h = \frac{x \cdot x}{0.01} = 1.6 \times 10^{-3}$$

$$\therefore x^2 = 1.6 \times 10^{-5}$$

$$\therefore x = 4 \times 10^{-3}$$

$$[\text{OH}^-] = x = 4 \times 10^{-3} \text{ M}$$

$$[\text{H}_3\text{O}^+] = \frac{K_w}{[\text{OH}^-]} = \frac{10^{-14}}{4 \times 10^{-3}} = 0.25 \times 10^{-11}$$

$$\text{pH} = -\log(0.25 \times 10^{-11}) = 11.6020$$

$$\text{Degree of hydrolysis} = \frac{x}{0.01} = \frac{4 \times 10^{-3}}{0.01} = 4 \times 10^{-1}$$

**Ex.23. Calculate for 0.01 N solution of sodium acetate -**

(i) **Hydrolysis constant**

(ii) **Degree of hydrolysis**

(iii) **pH**

**Given  $K_a$  of  $\text{CH}_3\text{COOH} = 1.9 \times 10^{-5}$ .**

**Sol.** For  $\text{CH}_3\text{COONa} + \text{H}_2\text{O} \rightleftharpoons \text{CH}_3\text{COOH} + \text{NaOH}$

Initial    C                    0                    0

After    C(1-h)                Ch                    Ch

$$(i) \quad K_h = \frac{K_w}{K_a} = \frac{10^{-14}}{1.9 \times 10^{-5}} = 5.26 \times 10^{-10}$$

$$(ii) \quad h = \sqrt{\frac{K_h}{C}} = \sqrt{\frac{5.26 \times 10^{-10}}{0.01}} = 2.29 \times 10^{-4} \text{ M}$$



$$(iii) \quad [\text{OH}^-] \text{ from NaOH, a strong base} = \text{Ch} = 0.01 \times 2.29 \times 10^{-4} = 2.29 \times 10^{-6} \text{ M}$$

$$\text{pOH} = 5.64$$

$$\therefore \text{pH} = 14 - 5.64 = 8.36$$

### 5.2.3 Hydrolysis of (WA - WB) type salt :


**Ex.**  $\text{NH}_4\text{CN}$ ,  $\text{CaCO}_3$ ,  $(\text{NH}_4)_2\text{CO}_3$ ,  $\text{ZnHPO}_3$



Solution is almost neutral but it may be acidic or basic depending upon the nature of acid & base & pH of the solution is near to 7.

**For WA - WB types of salt :**

	$K_a > K_b$	$K_b > K_a$	$K_a = K_b$
1. Hydrolysis	Cationic-anionic	Anionic-cationic	Neutral hydrolysis
2. Nature	Acidic	Basic	Neutral
3. pH	$\text{pH} < 7$	$\text{pH} > 7$	$\text{pH} = 7$


**(i) Relation between  $K_h$ ,  $K_w$ ,  $K_a$  &  $K_b$** 

$$K_h = \frac{[\text{NH}_4\text{OH}][\text{HCN}]}{[\text{NH}_4^+][\text{CN}^-]} \quad \dots (i)$$

$$\frac{[\text{NH}_4\text{OH}][\text{HCN}]}{[\text{NH}_4^+][\text{CN}^-]} \times \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]} \times \frac{[\text{H}^+][\text{CN}^-]}{[\text{HCN}]} = [\text{H}^+][\text{OH}^-]$$

$$K_h \times K_b \times K_a = K_w$$

$$\boxed{K_h = \frac{K_w}{K_a \times K_b}}$$

**(ii) Degree of Hydrolysis :**

Initial concentration  
at equilibrium

$$K_h = \frac{[\text{NH}_4\text{OH}][\text{HCN}]}{[\text{NH}_4^+][\text{CN}^-]}$$

Since  $h \ll \ll \ll 1$

Then  $(1 - h) \approx 1$

$$\therefore \boxed{K_h = h^2} \quad \text{or} \quad h^2 = \frac{K_w}{K_a \times K_b}$$

$$h = \sqrt{\frac{K_w}{K_a \times K_b}} \quad \dots (v)$$

**(iii) pH of the solution**

From eq. (iii)

$$K_a = \frac{[\text{H}^+][\text{CN}^-]}{[\text{HCN}]}$$

$$[\text{H}^+] = \frac{K_a \times [\text{HCN}]}{[\text{CN}^-]}$$

$$[\text{H}^+] = \frac{K_a \times \text{Ch}}{\text{C} - \text{Ch}} = \frac{K_a \times h}{1 - h}$$

Since  $h \ll \ll \ll 1$ ,  $(1 - h) \approx 1$

$$[\text{H}^+] = K_a \times h \quad [\text{Now put the value of } h \text{ from eq. (5)}]$$

$$= K_a \times \sqrt{\frac{K_w}{K_a \times K_b}}$$

$$\boxed{[\text{H}^+] = \sqrt{\frac{K_w \times K_a}{K_b}}}$$

On taking  $-\log$  on both sides

$$-\log [H^+] = -\log \left( \frac{K_w \times K_a}{K_b} \right)^{1/2}$$

$$pH = -\frac{1}{2} [\log K_w + \log K_a - \log K_b]$$

$$pH = 7 + \frac{1}{2} pK_a - \frac{1}{2} pK_b$$

**Note :** Degree of hydrolysis of [WA – WB] type salt does not depend on the concentration of salt.

**Ex.24 Salt of weak acid and weak base**

(i) Calculate pH of the mixture (25 mL of 0.1 M  $NH_4OH$  + 25 mL of 0.1 M  $CH_3COOH$ ).

Given that  $K_a : 1.8 \times 10^{-5}$ , and  $K_b = 1.8 \times 10^{-5}$

<b>Sol.</b>	$NH_4OH$	+	$CH_3COOH$	$\rightarrow$	$CH_3COONH_4$	+	$H_2O$
Initial milli moles	$25 \times 0.1$		$25 \times 0.1$		0		0
	= 2.5		= 2.5		–		
Final milli moles	0		0		2.5		2.5

As salt is formed (salt of weak acid and weak base) and pH will be decided by salt hydrolysis

$$pH = \frac{pK_w + pK_a - pK_b}{2} = \frac{1}{2} (-\log 10^{-14} - \log 1.8 \times 10^{-5} + \log 1.8 \times 10^{-5}) = 7$$

**Ex.25 In the following which one has highest / maximum degree of hydrolysis.**

(1) 0.01 M –  $NH_4Cl$

(2) 0.1 M –  $NH_4Cl$

(3) 0.001 M –  $NH_4Cl$

(4) Same

**Sol.** [3]

$$\left( h = \sqrt{\frac{K_h}{C}} \quad \text{if } C \text{ decreases, } h \text{ increases} \right)$$

**Ex.26 In the following which one has lowest value of degree of hydrolysis.**

(1) 0.01 M –  $CH_3COONH_4$

(2) 0.1 M –  $CH_3COONH_4$

(3) 0.001 M –  $CH_3COONH_4$

(4) Same

**Sol.** [4]

**Ex.27 Find out the concentration of  $[H^+]$  in 0.1M  $CH_3COONa$  solution ( $K_a = 10^{-5}$ )**

**Sol.** Salt is [WA – SB] type

$$\therefore [H^+] = \sqrt{\frac{K_w \times K_a}{C}} = \sqrt{\frac{10^{-14} \times 10^{-5}}{10^{-1}}} = \sqrt{10^{-19} \times 10^{+1}} = \sqrt{10^{-18}} = 10^{-9}$$

**Ex.28** Calculate the degree of hydrolysis of a mixture containing 0.1N  $\text{NH}_4\text{OH}$  and 0.1N  $\text{HCN}$




$$K_a = 10^{-5} \quad \& \quad K_b = 10^{-5}$$

**Sol.** Salt is [WA – WB]

$$h = \sqrt{\frac{K_w}{K_a \times K_b}} = \sqrt{\frac{10^{-14}}{10^{-5} \times 10^{-5}}} \\ = \sqrt{10^{-14} \times 10^{+10}} = \sqrt{10^{-4}} = 10^{-2}$$

### 5.2.4 Hydrolysis of [SA – SB] type salt :



**Ex.**  $\text{NaCl}$ ,  $\text{BaCl}_2$ ,  $\text{Na}_2\text{SO}_4$ ,  $\text{KClO}_4$  etc.



- (1) Hydrolysis of salt of [SA – SB] is not possible
- (2) Solution is neutral in nature ( $\text{pH} = \text{pOH} = 7$ )
- (3) pH of the solution is 7

### 5.2.5 Hydrolysis of Amphiprotic Anion :




$\text{NaHCO}_3$ ,  $\text{NaHS}$ , etc., can undergo ionisation to form  $\text{H}^+$  ion and can undergo hydrolysis to form  $\text{OH}^-$  ( $\text{Na}^+$  ion is not hydrolysed)



$$\text{pH}(\text{HCO}_3^-) = \left( \frac{\text{pK}_{a_1} + \text{pK}_{a_2}}{2} \right)$$

(b) Similarly for  $\text{H}_2\text{PO}_4^-$  and  $\text{HPO}_4^{2-}$  amphiprotic anions.

$$\text{pH}_{(\text{H}_2\text{PO}_4^-)} = \left( \frac{\text{pK}_{a_1} + \text{pK}_{a_2}}{2} \right) \quad \text{and} \quad \text{pH}_{(\text{HPO}_4^{2-})} = \left( \frac{\text{pK}_{a_2} + \text{pK}_{a_3}}{2} \right)$$



$\therefore \text{H}^+$  and  $\text{OH}^-$  also react

$\therefore$  We can safely assume that both reactions have nearly same degree of dissociation

$$\therefore [\text{H}_2\text{CO}_3] \approx [\text{CO}_3^{2-}] \quad \dots(1)$$

$$\frac{K_w}{K_{a_1}} = \frac{[\text{H}_2\text{CO}_3][\text{OH}^-]}{[\text{HCO}_3^-]} \Rightarrow \frac{1}{K_{a_1}} = \frac{[\text{H}_2\text{CO}_3]}{[\text{H}^+][\text{HCO}_3^-]} \quad \dots(2)$$




$$\frac{[\text{CO}_3^{2-}][\text{H}^+]}{[\text{HCO}_3^-]} = K_{a_2} \quad \dots(3)$$

Divide (2) by (3)

$$[\text{H}^+] = \sqrt{K_{a_1}K_{a_2}} \Rightarrow \text{pH} = \frac{\text{p}K_{a_1} + \text{p}K_{a_2}}{2}$$

**Ex.29 Calculate the pH of 0.5 M  $\text{Na}_3\text{PO}_4$  in aqueous solution ?**



**Sol.**  $\text{HPO}_4^{2-}$  and  $\text{PO}_4^{3-}$  are conjugate acid and base so  $K_a \times K_b = 10^{-14}$

$$K_a(\text{HPO}_4^{2-}) = \frac{10^{-14}}{2.4 \times 10^{-2}} = 4.17 \times 10^{-13}$$

$$\text{p}K_a = -\log K_a = 12.38$$

$$\text{or } \text{pH} = 7 + \frac{1}{2} \text{p}K_a + \frac{1}{2} \log C$$

$$\text{pH} = 13.04$$

## 6 BUFFER SOLUTIONS

A solution that resists change in pH value upon addition of small amount of strong acid or base or when solution is diluted is called buffer solution.

The capacity of a solution to resist alteration in its pH value is known as buffer capacity and the mechanism of buffer solution is called buffer action.

### 6.1 Types of buffer solutions

(A) Simple buffer solution


(B) Mixed buffer solution

### 6.2 SIMPLE BUFFER SOLUTION :


A salt of weak acid and weak base in water e.g.  $\text{CH}_3\text{COONH}_4$ ,  $\text{HCOONH}_4$ ,  $\text{AgCN}$ ,  $\text{NH}_4\text{CN}$ .

#### Buffer action of simple buffer solution

Consider a simple buffer solution of  $\text{CH}_3\text{COONH}_4$ , since it is a salt will dissociated completely.



If a strong acid such as HCl is added then




The  $\text{H}^+$  ions from the added acid (HCl) combine with  $\text{CH}_3\text{COO}^-$  ions to form  $\text{CH}_3\text{COOH}$ , which is a weak acid so will not further ionized.

Thus there is no rise in  $\text{H}^+$  ion concentration and the pH remains constant.



- If a strong base is added as NaOH



Thus change in  $\text{OH}^-$  ion concentration is resisted by  $\text{NH}_4^+$  ions by forming  $\text{NH}_4\text{OH}$  which is a weak base. So it will not further ionized and pH remains constant.




pH of a simple buffer solution :-

$$\text{pH} = 7 + \frac{1}{2} \text{p}K_a - \frac{1}{2} \text{p}K_b$$


### 6.3 MIXED BUFFER SOLUTIONS :

#### 6.3.1 Acidic buffer solution :

An acidic buffer solution consists of solution of a weak acid and its salt with strong base. The best known example is a mixture of solution of acetic acid and its salt with strong base ( $\text{CH}_3\text{COONa}$ ). Other example :



When a few drops of an acid ( $\text{HCl}$ ) are added to it, the  $\text{H}^+$  ions from the added acid ( $\text{HCl}$ ) combine with the  $\text{CH}_3\text{COO}^-$  ions to form  $\text{CH}_3\text{COOH}$ . Thus there is no rise in  $\text{H}^+$  ion concentration and the pH of solution remains constant. On the other hand, when a few drops of base ( $\text{NaOH}$ ) are added, the  $\text{OH}^-$  of the added base reacts with acetic acid to form unionise water and acetate ions.





Thus there is no increase in  $\text{OH}^-$  ion concentration and hence the pH of the solution remains constant.

#### pH of a acidic buffer solution (Henderson equation) :

Consider a buffer mixture (acidic buffer)



where  $\text{A} = \text{CH}_3\text{COO}$ ,  $\text{A}^- = \text{CH}_3\text{COO}^-$



Applying law of mass action to dissociation equilibrium of HA

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]} ; \text{ so } [\text{H}^+] = \frac{K_a[\text{HA}]}{[\text{A}^-]}$$

taking log,  $\log [\text{H}^+] = \log K_a + \log \frac{[\text{HA}]}{[\text{A}^-]}$

$$-\log [\text{H}^+] = -\log K_a - \log \frac{[\text{HA}]}{[\text{A}^-]}$$


$$\text{pH} = \text{p}K_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

$[\text{A}^-]$  = Initial concentration of salt as it is mainly comes from salt.

$[\text{HA}]$  = Initial concentration of the acid.

$$\text{pH} = \text{p}K_a + \log \frac{[\text{Salt}]}{[\text{Acid}]} \quad (\text{it is known as Henderson-Hasselbalch equation.})$$

**Note :** A solution can act as buffer only if ratio of concentration of salt to acid is between 0.1 to 10.



Thus pH range of an acidic buffer solution is  $(\text{pK}_a + 1)$  to  $(\text{pK}_a - 1)$

$$\text{pH range} = \text{pK}_a \pm 1$$

Maximum buffer action will be only when ratio of concentration of acid and salt is 1. So for maximum buffer action,  $\text{pH} = \text{pK}_a$

**Ex.30** How much volume of 0.2 M solution of acetic acid should be added to 100 mL of 0.2 M solution of sodium acetate to prepare a buffer solution of  $\text{pH} = 6.00$  ? ( $\text{pK}_a$  for acetic acid is 4.74)

**Sol.**  $\text{pH} = \text{pK}_a + \log \frac{[\text{Salt}]}{[\text{Acid}]}$

$$\log \frac{[\text{Salt}]}{[\text{Acid}]} = \text{pH} - \text{pK}_a = 6.00 - 4.74 = 1.26 \quad \therefore \quad \frac{[\text{Salt}]}{[\text{Acid}]} = 18.2$$

$$\text{Moles of } \text{CH}_3\text{COONa} \text{ in solution} = \frac{100 \times 0.2}{1000} = 0.02$$

Let, volume of 0.2 acetic acid added = V mL


$$\therefore \quad \text{Moles of acetic acid} = \frac{V \times 0.2}{1000}$$

$$\therefore \quad \frac{0.02}{V \times \frac{0.2}{1000}} = 18.2$$

$$\therefore \quad V = 5.49 \text{ mL}$$

**Ex.31** Calculate the pH after the addition of 80 mL and 100 mL respectively of 0.1 N NaOH to 100 mL, 0.1 N  $\text{CH}_3\text{COOH}$ . (Given  $\text{pK}_a$  for  $\text{CH}_3\text{COOH} = 4.74$ )

**Sol.** If 80 mL of 0.1 N NaOH is added to 100 mL of 0.1 N  $\text{CH}_3\text{COOH}$ , acidic buffer will form as



$$\text{pH} = \text{pK}_a + \log \frac{[\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]} = 4.74 + \log \frac{0.008}{0.002} = 5.342$$

If 100 mL of 0.1 N NaOH is added to 100 mL of 0.1 N  $\text{CH}_3\text{COOH}$ , complete neutralization takes place and the concentration of  $\text{H}_3\text{CCOONa} = \frac{0.1}{2} \text{ M} = 0.05 \text{ M}$

$$\text{Now, } \text{pH} = 7 + \frac{1}{2} \text{pK}_a + \frac{1}{2} \log C = 8.72$$

**Ex.32** Calculate the pH of a solution when 0.20 moles of HCl is added to one litre solution containing -

(a) 1 M each of acetic acid and acetate ion ?


(b) 0.1 M each of acetic acid and acetate ion ?

Given  $K_a$  for acetic acid is  $1.8 \times 10^{-5}$ .

**Sol. (a)** Initially [Acetic acid] = 1 M

[Acetate] = 1 M

Now 0.2 moles of HCl are added to it.




$$\therefore \text{pH} = \text{p}K_a + \log \frac{[\text{conjugate}]}{[\text{acid}]}$$

$$\therefore \text{pH} = -\log 1.8 \times 10^{-5} + \log \frac{0.8}{1.2} = 4.5686$$

**(b)** In II case initially [Acetic acid] = 0.1 M

[Acetate] = 0.1 M

Now 0.2 mole of HCl are added to it




$$\therefore \text{pH} = 1$$


**Note:** CH<sub>3</sub>COOH no doubt gives H<sup>+</sup> but being weak acid as well as in presence of HCl does not dissociate appreciably and thus, H<sup>+</sup> from CH<sub>3</sub>COOH may be neglected.

### 6.3.2 Basic buffer solution :


A basic buffer solution consists of a mixture of a weak base and its salt with strong acid. The best known example is a mixture of NH<sub>4</sub>OH and NH<sub>4</sub>Cl.



When a few drops of a base (NaOH) are added, the OH<sup>-</sup> ions from NaOH combine with NH<sub>4</sub><sup>+</sup> ions to form feebly ionised NH<sub>4</sub>OH thus there is no rise in the concentration of OH<sup>-</sup> ions and hence the pH value remains constant.




If a few drops of an acid (HCl) are added the H<sup>+</sup> from acid combine with NH<sub>4</sub>OH to form H<sub>2</sub>O and NH<sub>4</sub><sup>+</sup> ions.



Thus the addition of acid does not increase the H<sup>+</sup> ion concentration and hence pH remains unchanged.

#### • pH of basic buffer solution :



$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]}$$

$$[\text{OH}^-] = \frac{K_b [\text{NH}_4\text{OH}]}{[\text{NH}_4^+]} = \frac{K_b [\text{Base}]}{[\text{Salt}]}$$

( $\text{NH}_4^+$  mainly comes from salt)

taking  $-\log$  on both side

$$-\log \text{OH}^- = -\log \frac{K_b [\text{Base}]}{[\text{Salt}]} \Rightarrow \text{pOH} = -\log K_b - \log \frac{[\text{Base}]}{[\text{Salt}]}$$

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{Salt}]}{[\text{Base}]} \Rightarrow \text{pH} = 14 - \text{pOH}$$

❖ **pOH range :**

A solution can act as buffer solution only if ratio of concentration of salt to base is from 0.1 to 10.

$\text{NH}_4\text{OH}$	:	$\text{NH}_4\text{Cl}$	
1		10	$\text{pOH} = \text{p}K_b + 1$
10		1	$\text{pOH} = \text{p}K_b - 1$

So pOH range is  $\text{p}K_b \pm 1$

❖ **Condition for maximum buffer action :**

$[\text{NH}_4\text{OH}]$	:	$[\text{NH}_4\text{Cl}]$
1		1

$$\text{pOH} = \text{p}K_b + \log \frac{1}{1}$$

$$\text{pOH} = \text{p}K_b \quad \text{and} \quad \text{pH} = 14 - \text{p}K_b$$

Maximum buffer action because pH remains constant.

**Ex.33** An organic base  $B$  has  $K_b$  value equal to  $1 \times 10^{-8}$ . In what amounts should 0.01 M HCl and 0.01 M solution of  $B$  be mixed to prepare 1 L of a buffer solution having  $\text{pH} = 7.0$  ?

**Sol.**  $B + \text{H}_2\text{O} \rightleftharpoons \text{BH}^+ + \text{OH}^-$

$$K_b = \frac{[\text{BH}^+][\text{OH}^-]}{[B]} = 1 \times 10^{-8}$$

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{BH}^+]}{[B]}$$

$$\Rightarrow 7 = -\log(10^{-8}) + \log \frac{[\text{BH}^+]}{[B]} \Rightarrow 7 = 8 + \log \frac{[\text{BH}^+]}{[B]}$$

$$\log \frac{[\text{BH}^+]}{[B]} = -1$$

$$\therefore \frac{[\text{BH}^+]}{[\text{B}]} = 10^{-1} = 0.1$$

Let, volume of HCl taken = xL

$$\therefore \text{Volume of base taken} = (1 - x) \text{ L}$$

After the reaction, millimole of  $\text{BH}^+$  formed =  $0.01 \times (x)$

Millimoles of base left =  $0.01 (1 - 2x)$

$$\therefore \frac{[\text{BH}^+]}{[\text{B}]} = \frac{x}{[1 - 2x]} = 0.1$$

$$\therefore x = 0.083 \text{ L} = \text{Volume of HCl}$$

$$\therefore \text{Volume of base} = 0.917 \text{ L}$$

**Ex.34** Which of the following buffers containing  $\text{NH}_4\text{OH}$  and  $\text{NH}_4\text{Cl}$  show the lowest pH value?

	conc. of $\text{NH}_4\text{OH} (\text{mol L}^{-1})$	conc. of $\text{NH}_4\text{Cl} (\text{mol L}^{-1})$
(A)	0.50	0.50
(B)	0.10	0.50
(C)	0.50	1.50
(D)	0.50	0.10

**Sol. (B)**  $\text{pOH} = \text{p}K_b + \log \frac{[\text{salt}]}{[\text{base}]}$  for  $\text{NH}_4\text{Cl} = 0.5$  and  $\text{NH}_4\text{OH} = 0.1$

pOH will be maximum and so pH will be minimum.

**Ex.35** A solution of weak base BOH was titrated with 0.1 N HCl. The pH of the solution was found to be 10.04 and 9.14 after the addition of 5 mL and 20 mL of the acid respectively. Find the dissociation constant of the base.

**Sol. Case I :**

	BOH	+	HCl	→	BCl	+	H <sub>2</sub> O
Millimole before reac.	a		$0.1 \times 5 = 0.5$		0		0
Millimole after reac.	$(a - 0.5)$		0		0.5		0.5

$$\therefore \text{pOH} = -\log K_b + \log \frac{[\text{BCl}]}{[\text{BOH}]} \dots\dots\dots(\text{i})$$

$$\therefore \text{pH} = 10.04 \quad \text{so} \quad \text{pOH} = 3.96$$

$$\therefore 3.96 = -\log K_b + \log \frac{0.5}{(a - 0.5)} \dots\dots\dots(\text{ii})$$

**Case II :**

	BOH	+	HCl	→	BCl	+	H <sub>2</sub> O
Millimole before reac.	a		$0.1 \times 20 = 2$		0		0
Millimole after reac.	$(a - 2)$		0		2		2

$$\therefore \text{pOH} = -\log K_b + \log \frac{[\text{BCl}]}{[\text{BOH}]} \dots\dots\dots(\text{iii})$$

$$\because \text{pH} = 9.14 \quad \therefore \text{pOH} = 4.86$$


$$\therefore 4.86 = -\log K_b + \log \frac{2}{(a-2)} \dots\dots\dots(\text{iv})$$

Solving Eqs. (ii) and (iv),  $K_b = 1.81 \times 10^{-5}$

## 7. INDICATORS

The stage of titration when complete reaction occur between the solution is called **equivalent point**. The stage of titration when sudden change in colour of solution is observed is called **end point**. A perfect indicator response sudden colour change exactly on completion of reaction. An **indicator** is a substance which response sudden change in colour of solution at the end point or neutral point of the acid-base titration. **At end point**  $N_1 V_1 = N_2 V_2$

- The indicators in acid-base titration changes colour on changing the pH of solution.
- All the acid-base indicators are either weak organic acid or base and having different colour for unionized and ionised form.
- A mixture of two colour is recognized in a single colour if the conc. of one is 10 times or more than that of others. (This 10 time is flexible)



$$\text{Diss. const. or (Ionisation const)} = K_a = K_m = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

$$\text{pH} = \text{pK}_m + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

$$(a) \quad \text{The solution will appear only of colour Y, if } \frac{[\text{A}^-]}{[\text{HA}]} \geq 10 \Rightarrow \text{pH} \geq (\text{pK} + 1)$$

$$(b) \quad \text{The solution will appear only of colour X, if } \frac{[\text{A}^-]}{[\text{HA}]} \leq \frac{1}{10} \Rightarrow \text{pH} \leq (\text{pK} - 1)$$

pH of solution below and above which solution appears in a single colour is called pH range of indicator.

Indicator	pH range	Colour change	pK <sub>a</sub>
Methyl orange	3.2 – 4.5	Pink to yellow	3.7
Methyl red	4.4 – 6.5	Red to yellow	5.1
Litmus	5.5 – 7.5	Red to blue	7.0
Phenol red	6.8 – 8.4	Yellow to red	7.8
Phenolphthalein	8.3 – 10.5	Colourless to pink	9.6

**Ex.36** The disso. const. of a basic indicator is  $2 \times 10^{-7}$ . Calculate its pH range.

**Sol.**  $5.7 - 7.7 = \text{pOH} \quad \therefore \quad \text{pH} = 6.3 - 8.3$


**Ex.37** The pH range of an acidic indicator  $\text{HIn}$  is 4.0 - 5.2. Calculate dissociation constant. Also

calculate  $\frac{\text{In}^-}{\text{HIn}}$  for the appearance of solution in single colour.





**Sol.** Diss. constant =  $2.5 \times 10^{-5}$ , 4

## 9.1 TITRATION OF STRONG ACID AGAINST STRONG ALKALI :

The graph (A) shows how pH changes during the titration of  $50 \text{ cm}^3$  of  $0.1 \text{ M HCl}$  with  $0.1 \text{ M NaOH}$ .



The pH of  $0.1 \text{ M}$  solution of  $\text{HCl}$  in the beginning would be 1. As alkali is added, the pH changes slowly in the beginning. However, at the equivalence point pH changes rapidly from about 3.5 to 10. It can be shown by simple calculations that pH of the solution is 3.7 when  $49.8 \text{ cm}^3$  of  $0.1 \text{ M NaOH}$  solution have been added. The pH suddenly changes to 10 after addition of  $50.1 \text{ cm}^3$  of the  $\text{NaOH}$  solution. Thus, any indicator having pH range between 3.5 to 10 will identify the equivalence point. This means that any one of phenolphthalein, methyl orange or bromothymol blue could be used as an indicator.




**Titration curves :** (A) strong base with strong acid ; (B) weak base with strong acid ; (C) strong base with weak acid ; (D) weak base with weak acid.



## 7.2 TITRATION OF STRONG ACID AGAINST WEAK ALKALI :


The graph (B) shows how pH changes during titration of 50 cm<sup>3</sup> of 0.1 M HCl with 0.1 M NH<sub>3</sub>.



In this case, the pH changes rapidly from 3.5 to 7.0 at the equivalence point. Methyl orange, methyl red and bromocresol green are suitable indicators for this type of titration. Phenolphthalein is unsuitable because its pH range lies outside the vertical portion of the curve.

## 7.3 TITRATION OF WEAK ACID AGAINST STRONG BASE :


The graph (C) shows how pH changes during titration of 50 cm<sup>3</sup> of 0.1 M CH<sub>3</sub>COOH with 0.1 M NaOH.



The vertical portion of this titration curve lies between pH range 7 to 10.6. Phenolphthalein is suitable indicator for this titration. Methyl orange is not suitable for this titration because its pH range lies on the flat portion of the curve.

## 7.4 TITRATION OF WEAK ACID AGAINST WEAK BASE :

The graph (D) represents the titration curve obtained for titration of 50 cm<sup>3</sup> of 0.1 M CH<sub>3</sub>COOH with 0.1 M NH<sub>3</sub>.



For this type of titration there is no sharp increase in pH at the equivalence point. No indicator is suitable for this type of titration.

**Ex.38** Bromophenol blue is an indicator with a value of  $K_a = 6.84 \times 10^{-6}$ . At what pH it will work as an indicator? Also report the % of this indicator in its basic form at a pH of 5.84.

**Sol.**  $\text{HBPh} \rightleftharpoons \text{H}^+ + \text{BPh}^-$

$$K_a = \frac{[\text{H}^+][\text{BPh}^-]}{[\text{HBPh}]}, \text{ when } \text{BPh}^- = \text{HBPh}, \text{ indicator will work. Thus}$$

$$[\text{H}^+] = 6.84 \times 10^{-6}$$

$$\therefore \text{pH} = 5.165$$

Also if pH = 5.84

$$\text{or } [\text{H}^+] = 1.44 \times 10^{-6}, \text{ then}$$

$$K_a = \frac{[\text{H}^+][\text{BPh}^-]}{[\text{HBPh}]} \quad \text{or} \quad 6.84 \times 10^{-6} = \frac{1.44 \times 10^{-6} \cdot C\alpha}{C(1-\alpha)} \quad \text{or} \quad \alpha = 0.83 \text{ or } 83 \%$$

### 7.5 Titration of Poly protic acid v/s S.B. :

50 mL of 0.1 M  $\text{H}_3\text{PO}_4$  agent 0.1 M NaOH. Calculate pH when vol. of NaOH added is

- (a) 0 mL                      (b) 25 mL                      (c) 50 mL                      (d) 75 mL                      (e) 100 mL  
 (f) 125 mL                      (g) 150 mL                      (h) 200 mL                      (i) 90 mL

$$pK_{a1} = 3 \quad pK_{a2} = 7, pK_{a3} = 11$$

(a)  $\text{pH} = 1/2 (pK_{a1} - \log C) = 1/2 (3 + 1) = 2$



t = 0    5 Mmol        2.5 Mmol                      0                      –

          2.5 M mol        0                      2.5 M mol

$$\text{pH} = pK_{a1} + \log 1/1 = pK_{a1} = 3$$

(c) Solution of  $\text{H}_2\text{PO}_4^-$  amphoteric species

$$\text{pH} = \frac{pK_{a1} + pK_{a2}}{2} = \frac{3 + 7}{2} = 5$$



t = 0    5 M mol                      2.5 M mol                      0                      –

          2.5 M mol                                      2.5 M mol                      –

$$\text{pH} = \log 1/1 + pK_{a2} = 7$$

(e)  $\text{HPO}_4^{2-}$  solution (amphoteric species)

$$\text{pH} = \frac{pK_{a2} + pK_{a3}}{2} = 9$$



          5                      2.5                      0                      –

          2.5                      0                      2.5                      –

$$\text{pH} = pK_{a3} + \log 1/1 = 11$$

(g) 3<sup>rd</sup> eq. pt  $\text{Na}_3\text{PO}_4$  solution

$$[\text{Na}_3\text{PO}_4] = 5/200 = 1/40$$

$$\text{pH} = 1/2 \{pK_w + pK_{a3} + \log C\} = 1/2 (14 + 11 - 2 + 0.4) = 11.7$$

(h) 200 mL


hydrolysis of  $\text{Na}_3\text{PO}_4$  can be neglected in presence of NaOH

$$[\text{NaOH}] = 5/250 = 1/50$$

$$\text{pOH} = 1.7$$

$$\text{pH} = 12.3$$


(i) 90 mL



          5                      4                      0

          1                      0                      4

$$\text{pH} = pK_{a2} + \log 4 = 7.6$$



## 10. SOLUBILITY (s) & SOLUBILITY PRODUCT ( $K_{sp}$ )

### 10.1 SOLUBILITY :

At constant temperature, the maximum number of moles of solute which can be dissolved in a solvent to obtain 1 litre of saturated solution is called solubility.


Solubility depends on the following –

- (i) Temperature
- (ii) Presence of common ion
- (iii) Nature of solvent

### 10.2 SOLUBILITY PRODUCT( $K_{sp}$ ) :

When a sparingly soluble salt such as AgCl is put into water, a very small amount of AgCl dissolves in water and most of the salt remains undissolved in its saturated solution.

- A solution which remains in contact with undissolved solute is said to be saturated.
- The salt AgCl is an electrolyte, its dissociation occurs in solution. Hence, the quantity of AgCl that dissolves in water dissociates into  $Ag^+$  and  $Cl^-$  ions. Thus, in the saturated solution of AgCl an equilibrium exists between undissolved solid AgCl and its ions,  $Ag^+$  and  $Cl^-$  ions.



according to law of mass action

$$K = \frac{[Ag^+].[Cl^-]}{[AgCl]}$$

Since, the concentration of undissolved solid AgCl is constant. Thus, the product  $K.[AgCl]$  gives another constant which is designated as  $K_{sp}$

So,  $K.[AgCl] = [Ag^+].[Cl^-]$

$\therefore K_{sp} = [Ag^+] \cdot [Cl^-]$

- **$K_{sp}$  for  $CaCl_2$**   $CaCl_2(s) \rightleftharpoons Ca^{+2}(aq) + 2Cl^-(aq)$

Solubility product in terms of concentration of ions

$$K_{sp} = [Ca^{+2}] [Cl^-]^2$$

- **$K_{sp}$  for  $AlCl_3$**   $AlCl_3(s) \rightleftharpoons Al^{+3}(aq) + 3Cl^-(aq)$

Solubility product in terms of concentration of ions  $K_{sp} = [Al^{+3}] [Cl^-]^3$

- **General form**  $A_xB_y(s) \rightleftharpoons xA^{+y}(aq) + yB^{-x}(aq)$

$$K_{sp} = [A^{+y}]^x [B^{-x}]^y$$


Thus, solubility product is defined as the product of concentrations of the ions raised to a power equal to the number of ions given by the dissociation of electrolyte at a given temperature when the solution is saturated.

### 10.3 APPLICATION OF SOLUBILITY PRODUCT ( $K_{sp}$ ) :

#### 10.3.1 To find out the solubility (S) :

- (i)  **$K_{sp}$  of AB (Mono-mono, di-di, tri-tri valency) type salt –**

**Ex.** NaCl,  $BaSO_4$ ,  $CH_3COONa$ ,  $CaCO_3$ , NaCN, KCN,  $NH_4CN$ ,  $NH_4Cl$  etc.




$$K_{sp} = [A^+] [B^-]$$

$$K_{sp} = s^2 \quad \text{or} \quad s = \sqrt{K_{sp}}$$

(ii)  **$K_{sp}$  of  $AB_2$  or  $A_2B$  (Mono-di or di-mono valency) type salt –**

**Ex.**  $CaCl_2$ ,  $CaBr_2$ ,  $K_2S$ ,  $(NH_4)_2SO_4$ ,  $K_2SO_4$ ,  $K_2CO_3$  etc.



$$K_{sp} = [A^{+2}] [B^-]^2$$

$$K_{sp} = s \times (2s)^2 = s \times 4s^2 = 4s^3$$

$$s = \left( \frac{K_{sp}}{4} \right)^{1/3}$$

(iii) **General form :**



$$K_{sp} = [A^{+y}]^x \cdot [B^{-x}]^y$$

$$K_{sp} = (xs)^x \cdot (ys)^y$$

$$K_{sp} = x^x \cdot y^y \cdot s^{(x+y)}$$

## 10.4 COMMON ION EFFECT ON SOLUBILITY :

Solubility of substances always decreases in the presence of common ion. According to Le-Chatelier's principle, on increasing common ion concentration equilibrium shifts in backward direction until the equilibrium is reestablished so, the solubility of substances decreases.


**Ex.** Find out the solubility of  $AgCl$  in water and in the presence of CM –  $NaCl$  solution?




$$K_{sp} = [Ag^+] [Cl^-]$$

$$K_{sp} = S^2$$

In  $NaCl$  solution



Let solubility of AgCl in the presence of NaCl solution is  $S'$  mol  $L^{-1}$



$$K_{sp} = [\text{Ag}^+] [\text{Cl}^-]$$

$$K_{sp} = S' (S' + C) = S'^2 + S'C \quad (\text{Neglecting the higher power terms of } S')$$

$$K_{sp} = S' C$$

$$\boxed{S' = \frac{K_{sp}}{C}}$$

### 10.5 SIMULTANEOUS SOLUBILITY :



When two sparingly soluble salts are added in water simultaneously, there will be two simultaneous equilibria in the solution.

### 10.6 SOLUBILITY IN APPROPRIATE BUFFER SOLUTIONS :

Appropriate buffer means that the components of buffer should not interfere with the salt or only  $H^+$  or  $OH^-$  ions should be interacting with the ions of the salt.

### 10.7 EFFECT ON SOLUBILITY BECAUSE OF COMPLEX FORMATION :

Solubility of AgCl in aqueous  $NH_3$  is roughly 10,000 times as its solubility in water, due to complex formation.



$$\text{and } \frac{1}{K_{stability}} = K_{dissociation} = K_{instability}$$

### 10.8 CONDITION OF PRECIPITATION / IONIC PRODUCT (IP OR $Q_{sp}$ ) :

- Ionic product (IP) of an electrolyte is defined in the same way as  $K_{sp}$ . The only difference is that ionic product expression contains the initial concentration of ions or the concentration at any time whereas the expression of  $K_{sp}$  contains only equilibrium concentration. Thus, for AgCl.

$$IP = [\text{Ag}^+]_i [\text{Cl}^-]_i \quad \text{and} \quad K_{sp} = [\text{Ag}^+]_{eq} [\text{Cl}^-]_{eq}$$

Ionic product changes with concentration but  $K_{sp}$  does not.

To decide whether an ionic compound will precipitate, its  $K_{sp}$  is compared with the value of ionic product.


The following three cases arise :

- $IP < K_{sp}$  : The solution is unsaturated and precipitation will not occur.
- $IP = K_{sp}$  : The solution is saturated and solubility equilibrium exists.
- $IP > K_{sp}$  : The solution is supersaturated and hence precipitation of the compound will occur.

Thus, a salt is precipitated when its ionic product exceeds the solubility product of the salt.

## 10.9 SELECTIVE PRECIPITATION :


When the  $K_{sp}$  values differ then one of the salt can be selectively precipitated.



$$K_{sp} = 2^2 \times 3^3 \times (S)^{2+3} = 4 \times 27 \times S^5 = 108 S^5$$



$$K_{sp} = 2^2 \times 1^1 \times 1^1 (S)^{2+1+1} = 4S^4$$



$$K_{sp} = 1^1 \times 1^1 \times 1^1 \times 1^1 \times (S)^{1+1+1+1} = S^4$$

**Ex. 40.** If solubility product of the base  $M(OH)_3$  is  $2.7 \times 10^{-11}$ , the concentration of  $OH^-$  will be

(1)  $3 \times 10^{-3}$

(2)  $3 \times 10^{-4}$


(3)  $10^{-3}$

(4)  $10^{-11}$

Answer : (3)

**Ex. 41.** The solubility of  $BaSO_4$  in water is  $1.07 \times 10^{-5} \text{ mol dm}^{-3}$ . Estimate its solubility product.

**Sol.** Solubility equilibrium for  $BaSO_4$  is




Now,  $S = 1.07 \times 10^{-5} \text{ M}$

Hence,  $K_{sp} = (1.07 \times 10^{-5})^2 = 1.145 \times 10^{-10}$

**Ex. 42.** The solubility product of  $AgBr$  is  $5.2 \times 10^{-13}$ . Calculate its solubility in  $\text{mol dm}^{-3}$  and  $\text{g dm}^{-3}$ . (Molar mass of  $AgBr$  =  $187.8 \text{ g mol}^{-1}$ )

**Sol.** The solubility equilibrium of  $AgBr$  is



The molar solubility  $S$  of  $AgBr$  is given by

$$S = \sqrt{K_{sp}} = \sqrt{5.2 \times 10^{-13}} = 7.2 \times 10^{-7} \text{ mol dm}^{-3}$$

The solubility in  $\text{g dm}^{-3}$  = molar solubility ( $\text{mol dm}^{-3}$ )  $\times$  molar mass ( $\text{g mol}^{-1}$ )

$$= 7.2 \times 10^{-7} \times 187.8 = 1.35 \times 10^{-4} \text{ g dm}^{-3}$$

**Ex. 43.** What is the maximum volume of water required to dissolve 1 g of calcium sulphate at 25°C. For calcium sulphate,  $K_{sp} = 9.0 \times 10^{-6}$ .

**Sol.**  $\text{CaSO}_4 (aq) \rightleftharpoons \text{Ca}^{2+} (aq) + \text{SO}_4^{2-} (aq)$

If  $S$  is the solubility of  $\text{CaSO}_4$  in moles  $\text{L}^{-1}$

$$K_{sp} = [\text{Ca}^{2+}] \times [\text{SO}_4^{2-}] = S^2$$

$$\therefore S = \sqrt{K_{sp}} = \sqrt{9.0 \times 10^{-6}}$$

$$= 3 \times 10^{-3} \text{ mol L}^{-1}$$


$$= 3 \times 10^{-3} \times 136 \text{ g L}^{-1} = 0.408 \text{ g L}^{-1}$$

For dissolving 0.408 g of  $\text{CaSO}_4$  water required = 1 L

$$\therefore \text{For dissolving 1 g } \text{CaSO}_4 \text{ water required} = \frac{1}{0.408} \text{ L} = 2.45 \text{ L}$$

**Ex. 44.** Equal volumes of 0.04 M  $\text{CaCl}_2$  and 0.0008 M  $\text{Na}_2\text{SO}_4$  are mixed. Will a precipitate form?  $K_{sp}$  for  $\text{CaSO}_4 = 2.4 \times 10^{-5}$

**Sol.**



Suppose  $V$  mL of both are mixed

$$\therefore [\text{Ca}^{2+}] = \frac{0.04 V}{2V}$$

$$[\text{SO}_4^{2-}] = \frac{0.0008 V}{2V}$$

$$\therefore [\text{Ca}^{2+}] [\text{SO}_4^{2-}] = \frac{0.04 V}{2V} \times \frac{0.0008 V}{2V} = 8 \times 10^{-6}$$

Thus,  $[\text{Ca}^{2+}] [\text{SO}_4^{2-}]$  in solution  $< K_{sp}$

$$8 \times 10^{-6} < 2.4 \times 10^{-5}$$

$\therefore \text{CaSO}_4$  will not precipitate.

**Ex. 45.** Calculate simultaneous solubility of silverthiocyanate and silver bromide in water given that  $k_{sp}$  of silver thiocyanate =  $10^{-12}$  and  $k_{sp}$  of silver bromide =  $5 \times 10^{-13}$  respectively.

**Sol.** Let the solubility of  $\text{AgSCN}$  be  $x$  and that of  $\text{AgBr}$  is  $y$ , then



$$10^{-12} = x(x + y) \quad \dots\dots (i)$$

$$5 \times 10^{-13} = y(x + y) \quad \dots\dots (ii)$$

On solving we get,  $x = 2y$

$$\text{So} \quad y = 4.08 \times 10^{-7} \quad \text{and} \quad x = 8.16 \times 10^{-7}$$

**Ex. 46.** What  $[H^+]$  must be maintained in saturated  $H_2S(0.1\text{ M})$  to precipitate  $CdS$  but not  $ZnS$ , if  $[Cd^{2+}] = [Zn^{2+}] = 0.1$  initially ?

$$K_{sp} = (CdS) = 8 \times 10^{-27}$$

$$K_{sp} = (ZnS) = 1 \times 10^{-21}$$

$$K_a = (H_2S) = 1.1 \times 10^{-21}$$

**Sol.** In order to prevent precipitation of  $ZnS$

$$[Zn^{2+}][S^{2-}] < K_{sp}(ZnS) = 1 \times 10^{-21}$$

(ionic product)

$$\text{or } (0.1)[S^{2-}] < 1 \times 10^{-21}$$

$$\text{or } [S^{2-}] < 1 \times 10^{-20}$$

This is the maximum value of  $[S^{2-}]$  before  $ZnS$  will precipitate. Let  $[H^+]$  to maintain this  $[S^{2-}]$  be  $x$ .

Thus for  $H_2S \rightleftharpoons 2H^+ + S^{2-}$

$$K_a = \frac{[H^+]^2[S^{2-}]}{[H_2S]} = \frac{x^2(1 \times 10^{-20})}{0.1} = 1.1 \times 10^{-21}$$

$$\text{or } x = [H^+] = 0.1\text{ M}$$

$\therefore$  No  $ZnS$  will precipitate at a concentration of  $H^+$  greater than  $0.1\text{ M}$

**Ex. 47.** What must be the concentration of aq.  $NH_3(\text{eq.})$  which must be added to a solution containing  $4 \times 10^{-3}\text{ M Ag}^+$  and  $0.001\text{ M NaCl}$ , to prevent the precipitation of  $AgCl$ .

Given that  $K_{sp}(AgCl) = 1.8 \times 10^{-10}$  and the formation constant of  $[Ag(NH_3)_2]^+$  is


$$K_{\text{formation}} = \frac{10^8}{6}.$$

**Sol.** Calculate silver ion concentration which can be allowed to remain in the solution,

$$1.8 \times 10^{-10} = [Ag^+][Cl^-]$$

$$[Ag^+] = \frac{1.8 \times 10^{-10}}{0.001} = 1.8 \times 10^{-7}\text{ M},$$

This quantity is so small that almost all the  $Ag^+$  ion will be consumed.



$$4 \times 10^{-3} \quad b \quad 0$$

$$1.8 \times 10^{-7} \quad (b - 8 \times 10^{-3}) \quad 4 \times 10^{-3} \quad K = \frac{10^8}{6} = \frac{4 \times 10^{-3}}{1.8 \times 10^{-7} \times (b - 8 \times 10^{-3})^2}$$


$$\Rightarrow b = 0.0445$$



**Ex. 48.** 0.10 mol sample of  $\text{AgNO}_3$  is dissolved in one litre of 2.00 M  $\text{NH}_3$ . Is it possible to form  $\text{AgCl}(s)$  in the solution by adding 0.010 mol of  $\text{NaCl}$ ?

$$(K_{\text{sp}}(\text{AgCl}) = 1.8 \times 10^{-10}, K_{\text{f}}[\text{Ag}(\text{NH}_3)_2^+] = 1.6 \times 10^7)$$

**Sol.**



It is assumed that all  $\text{Ag}^+$  ions have been complexed and only  $x$  amount is left

$$K_{\text{f}} = \frac{[\text{Ag}(\text{NH}_3)_2^+]}{[\text{Ag}^+][\text{NH}_3]^2} \Rightarrow 1.6 \times 10^7 = \frac{0.10}{x(1.80)^2}$$

$$\therefore x = 1.93 \times 10^{-9} \text{ M} = [\text{Ag}^+] \text{ undissolved}$$

$$[\text{Cl}^-] = 1.0 \times 10^{-2} \text{ M}$$


$$\therefore [\text{Ag}^+][\text{Cl}^-] = 1.93 \times 10^{-9} \times 1.0 \times 10^{-2} = 1.93 \times 10^{-11} < 1.8 \times 10^{-10} [K_{\text{sp}}(\text{AgCl})]$$

Hence,  $\text{AgCl}(s)$  will not precipitate.

**Ex. 49.** What is the concentration of  $\text{Ag}^+$  ions in  $0.01 \text{ M AgNO}_3$  that is also  $1.0 \text{ M NH}_3$ ? Will  $\text{AgCl}$  precipitate from a solution that is  $0.01 \text{ M AgNO}_3$ ,  $0.01 \text{ M NaCl}$  and  $1 \text{ M NH}_3$ ?

$$K_d(\text{Ag}[\text{NH}_3]_2^+) = 5.88 \times 10^{-8}; K_{sp}(\text{AgCl}) = 1.8 \times 10^{-10}.$$

**Sol.** Let us first assume that  $0.01 \text{ M AgNO}_3$  shall combine with  $0.02 \text{ M NH}_3$  to form  $0.01 \text{ M Ag}(\text{NH}_3)_2^+$  and then consider its dissociation.



Since  $x \ll 1$

$$K_d = \frac{[\text{Ag}^+][\text{NH}_3]^2}{[\text{Ag}(\text{NH}_3)_2^+]} = 5.88 \times 10^{-8}$$

$$\therefore [\text{Ag}^+] = \frac{5.88 \times 10^{-8} \times 0.01}{(0.98)^2} = 6.12 \times 10^{-10} \text{ M}$$

Further, ionic product of  $\text{AgCl} = [\text{Ag}^+][\text{Cl}^-] = (6.12 \times 10^{-10})(0.01) = 6.12 \times 10^{-12}$

Because the ionic product is smaller than  $K_{sp} = 1.8 \times 10^{-10}$ , no precipitate should form.

**EXERCISE # S-I****IONIZATION CONSTANTS AND pH**

- Q.1 Calculate the number of  $H^+$  present in one ml of solution whose pH is 13. **IE0001**
- Q.2 (i)  $K_w$  for  $H_2O$  is  $9 \times 10^{-14}$  at  $60^\circ C$ . What is pH of water at  $60^\circ C$ . ( $\log 3 = 0.47$ ) **IE0002**
- (ii) What is the nature of solution at  $60^\circ C$  whose **IE0003**
- (a) pH = 6.7 (b) pH = 6.35
- Q.3 The value of  $K_w$  at the physiological temperature ( $37^\circ C$ ) is  $2.56 \times 10^{-14}$ . What is the pH at the neutral point of water at this temperature? ( $\log 2 = 0.3$ ) **IE0004**
- Q.4 Calculate pH of following solutions :
- (a) 0.1 M HCl **IE0005**
- (b) 0.1 M  $CH_3COOH$  ( $K_a = 1.8 \times 10^{-5}$ ) ( $\log \sqrt{1.8} = 0.13$ ) **IE0006**
- (c) 0.1 M  $NH_4OH$  ( $K_b = 1.8 \times 10^{-5}$ ) **IE0007**
- (d)  $10^{-8}$  M HCl [ $\sqrt{401} = (20.02)$ ] [ $\log 1.051 = 0.03$ ] **IE0008**
- (e)  $10^{-10}$  M NaOH **IE0009**
- (f)  $10^{-6}$  M  $CH_3COOH$  ( $K_a = 1.8 \times 10^{-5}$ ) **IE0010**
- (g)  $10^{-8}$  M  $CH_3COOH$  ( $K_a = 1.8 \times 10^{-5}$ ) [ $\sqrt{401} = (20.02)$ ] [ $\log 1.051 = 0.03$ ] **IE0011**
- (h) Decimolar solution of Baryta ( $Ba(OH)_2$ ), diluted 100 times. ( $\log 2 = 0.3$ ) **IE0012**
- (i)  $10^{-3}$  mole of KOH dissolved in 100 L of water. **IE0013**
- (j) Equal volume of HCl solution (PH = 4) + 0.0019 N HCl solution **IE0014**
- Q.5 Calculate :
- (a)  $K_a$  for a monobasic acid whose 0.10 M solution has pH of 4.50.
- (b)  $K_b$  for a monoacidic base whose 0.10 M solution has a pH of 10.50. **IE0015**
- Q.6 Calculate the ratio of degree of dissociation ( $\alpha_2/\alpha_1$ ) when an acetic acid solution is diluted 100 times. Assume  $\alpha < 1$ , even on dilution. [Given  $K_a = 10^{-5}$  M] **IE0016**
- Q.7 Calculate the ratio of degree of dissociation of acetic acid and hydrocyanic acid (HCN) in 1 M their respective solution of acids. [Given  $K_{a(CH_3COOH)} = 1.8 \times 10^{-5}$ ;  $K_{a(HCN)} = 6 \times 10^{-10}$ ] **IE0017**
- Q.8 How many moles of HCl must be removed from 1 litre of aqueous HCl solution to change its pH from 2 to 3 ? **IE0018**
- Q.9 pH of a dilute solution of HCl is 6.95. Calculate molarity of HCl solution.
- $$\left[ \begin{array}{l} 10^{-6.95} = 11.22 \times 10^{-8} \\ 10^{-7.05} = 8.90 \times 10^{-8} \end{array} \right]$$
- IE0205**
- Q.10 The pH of aqueous solution of ammonia is 10. Find molarity of solution.  $K_b(NH_4OH) = 10^{-5}$ . **IE0019**

Q.11 The solution of weak monoprotic acid which is 0.01 M, has pH = 3. Calculate  $K_a$  of weak acid.

IE0020

Q.12 Boric acid is a weak monobasic acid. It ionizes in water as



Calculate pH of 0.5 M boric acid.

IE0021

### MIXTURE OF TWO OR MORE ACIDS / BASES

Q.13 The pH of the solution produced when an aqueous solution of strong acid pH 5 is mixed with equal volume of an aqueous solution of strong acid of pH 3 is :-

IE0022

Q.14 Calculate pH of following solutions :  $[\log 0.3 = -0.522]$

(a) 0.1 M  $\text{H}_2\text{SO}_4$  (50 ml) + 0.4 M HCl 50 (ml)

(b) 0.1 M HA + 0.1 M HB  $[K_a(\text{HA}) = 5 \times 10^{-5} ; K_a(\text{HB}) = 4 \times 10^{-5}]$

IE0023

Q.15 Calculate pH of a solution containing 0.1M HA ( $K_a = 10^{-5}$ ) & 0.1 M HCl.

IE0024

Q.16 Calculate  $[\text{H}^+]$  and  $[\text{CHCl}_2\text{COO}^-]$  in a solution that is 0.01 M in HCl and 0.01 M in  $\text{CHCl}_2\text{COOH}$ .

Take ( $K_a = 3 \times 10^{-2}$ ) ( $\sqrt{28} = 5.3$ )

IE0206

Q.17 Calculate  $[\text{H}^+]$ ,  $[\text{CH}_3\text{COO}^-]$  and  $[\text{C}_7\text{H}_5\text{O}_2^-]$  in a solution that is 0.2 M in acetic acid and 0.1M in benzoic acid.  $K_a(\text{acetic}) = 1.8 \times 10^{-5}$ ,  $K_a(\text{benzoic}) = 5.4 \times 10^{-5}$ .

IE0207

### POLYPROTIC ACIDS & BASES

Q.18 What are the concentration of  $\text{H}^+$ ,  $\text{H}_2\text{C}_2\text{O}_4$ ,  $\text{HC}_2\text{O}_4^-$  and  $\text{C}_2\text{O}_4^{2-}$  in a 0.1 M solution of oxalic acid ?

$[K_1 = 10^{-2} \text{ M and } K_2 = 10^{-5} \text{ M}]$  ( $\sqrt{41} = 6.4$ )

IE0025

Q.19 Calculate  $[\text{H}^+]$ ,  $[\text{H}_2\text{PO}_4^-]$ ,  $[\text{HPO}_4^{2-}]$  and  $[\text{PO}_4^{3-}]$  in a 0.01M solution of  $\text{H}_3\text{PO}_4$ .

Take  $K_1 = 10^{-3}$ ,  $K_2 = 10^{-8}$ ,  $K_3 = 10^{-13}$ ,  $\sqrt{41} = 6.4$

IE0208

Q.20 Calculate pH of 0.2 M –  $\text{B(OH)}_2$  solution.

( $K_{b1} = 2 \times 10^{-5}$ ;  $K_{b2} = 4 \times 10^{-11}$ ,  $\log 2 = 0.3$ )

IE0026

### HYDROLYSIS

Q.21 What is the  $\text{OH}^-$  concentration of a 0.18 M solution of  $\text{CH}_3\text{COONa}$ .  $[K_a(\text{CH}_3\text{COOH}) = 1.8 \times 10^{-5}]$

IE0027

Q.22 Calculate the pH of a 2.0 M solution of  $\text{NH}_4\text{Cl}$ .  $[K_b(\text{NH}_3) = 2 \times 10^{-5}]$

IE0028

Q.23 0.25 M solution of pyridinium chloride  $\text{C}_5\text{H}_6\text{N}^+\text{Cl}^-$  was found to have a pH of 2.699. What is  $K_b$  for pyridine,  $\text{C}_5\text{H}_5\text{N}$ ? ( $\log 2 = 0.3010$ )

IE0029

Q.24 Calculate the extent of hydrolysis & the pH of 0.02 M  $\text{CH}_3\text{COONH}_4$ .

$[K_b(\text{NH}_3) = 1.8 \times 10^{-5}$ ,  $K_a(\text{CH}_3\text{COOH}) = 1.8 \times 10^{-5}]$

IE0030

- Q.25 Calculate the percent hydrolysis in a 0.06 M solution of KCN. [ $K_a(\text{HCN}) = 6 \times 10^{-10}$ ] **IE0031**
- Q.26 Calculate the extent of hydrolysis of 0.005 M  $\text{K}_2\text{CrO}_4$ . [ $K_2 = 3.2 \times 10^{-7}$  for  $\text{H}_2\text{CrO}_4$ ]  
(It is essentially strong for first ionization). **IE0209**
- Q.27 A 0.010 M solution of  $\text{PuO}_2(\text{NO}_3)_2$  was found to have a pH of 4.0. What is the hydrolysis constant,  $K_h$ , for  $\text{PuO}_2^{2+}$ , and what is  $K_b$  for  $\text{PuO}_2\text{OH}^+$ ? **IE0210**
- Q.28 What is the pH of 0.1M  $\text{NaHCO}_3$ ?  $K_1 = 5 \times 10^{-7}$ ,  $K_2 = 5 \times 10^{-11}$  for carbonic acids. **IE0032**
- Q.29 Calculate pH of 0.05M potassium hydrogen phthalate,  $\text{KHC}_8\text{H}_4\text{O}_4$ .
- $$\text{H}_2\text{C}_8\text{H}_4\text{O}_4 + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HC}_8\text{H}_4\text{O}_4^- \quad \text{p}K_1 = 2.94$$
- $$\text{HC}_8\text{H}_4\text{O}_4^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{C}_8\text{H}_4\text{O}_4^{2-} \quad \text{p}K_2 = 5.44$$
- IE0211**
- Q.30 The acid ionization (hydrolysis) constant of  $\text{Zn}^{2+}$  is  $1.0 \times 10^{-9}$   
(a) Calculate the pH of a 0.001 M solution of  $\text{ZnCl}_2$   
(b) What is the basic dissociation constant of  $\text{Zn}(\text{OH})^+$ ? **IE0033**

### BUFFER SOLUTION

- Q.31 Calculate the pH of solution containing 0.1M - HCN and 0.1M - NaCN.  $K_a$  of HCN =  $10^{-9}$  **IE0034**
- Q.32 Calculate the pH of solution containing 0.2 M -  $\text{NH}_4\text{OH}$  and 0.1 M -  $\text{NH}_4\text{Cl}$ .  $K_b$  of  $\text{NH}_4\text{OH}$  =  $1.8 \times 10^{-5}$ . ( $\log 2 = 0.3$ ,  $\log 1.8 = 0.26$ ) **IE0035**
- Q.33 0.4 mole  $\text{CH}_3\text{COONa}$  is added in 500 ml 0.4 M -  $\text{CH}_3\text{COOH}$  solutions. What is the pH of final solution ?  $K_a$  of  $\text{CH}_3\text{COOH} = 1.8 \times 10^{-5}$ . ( $\log 2 = 0.3$ ,  $\log 1.8 = 0.26$ ). **IE0036**
- Q.34 A buffer of pH 9.26 is made by dissolving x moles of ammonium sulphate and 0.1 mole of ammonia into 100 mL solution. If  $\text{p}K_b$  of ammonia is 4.74, calculate value of x. **IE0037**
- Q.35 Determine  $[\text{OH}^-]$  of a 0.050 M solution of ammonia to which sufficient  $\text{NH}_4\text{Cl}$  has been added to make the total  $[\text{NH}_4^+]$  equal to 0.100. [ $K_{b(\text{NH}_3)} = 1.8 \times 10^{-5}$ ,  $\text{p}K_b = 4.74$ ] **IE0038**
- Q.36 Calculate the pH of a solution containing 0.2 M  $\text{HCO}_3^-$  and 0.1 M  $\text{CO}_3^{2-}$   
[ $K_1(\text{H}_2\text{CO}_3) = 4 \times 10^{-7}$ ;  $K_2(\text{HCO}_3^-) = 4 \times 10^{-11}$ ] **IE0039**
- Q.37 Calculate the pH of a solution prepared by mixing 50.0 mL of 0.200 M  $\text{HC}_2\text{H}_3\text{O}_2$  and 50.0 mL of 0.100 M NaOH. [ $K_a(\text{CH}_3\text{COOH}) = 1.8 \times 10^{-5}$ ,  $\text{p}K_a = 4.74$ ] **IE0040**
- Q.38 50 mL of 0.1 M NaOH is added to 75 mL of 0.1 M  $\text{NH}_4\text{Cl}$  to make a basic buffer. If  $\text{p}K_a$  of  $\text{NH}_4^+$  is 9.26, calculate pH. **IE0041**

Q.39 Calculate the pH of a solution which results from the mixing of 50.0 ml of 0.3 M HCl with 50.0 ml of 0.4 M  $\text{NH}_3$ . [ $K_b(\text{NH}_3) = 1.8 \times 10^{-5}$ ,  $\text{p}K_b = 4.74$ ]

IE0042

Q.40 In 100 ml buffer solution of 0.1M  $\text{CH}_3\text{COOH}$  & 0.1M  $\text{CH}_3\text{COONa}$ , how many millimoles of NaOH should be added to increase its pH by 0.3.

Given ( $\log 2 = 0.3$ )

IE0043

### ACID BASE REACTIONS & TITRATIONS

Q.41 Calculate  $\text{OH}^-$  concentration at the equivalent point when a solution of 0.2 M acetic acid is titrated with a solution of 0.2 M NaOH.  $K_a$  for the acid =  $10^{-5}$ .

IE0044

Q.42 Calculate the hydronium ion concentration and pH at the equivalence point in the reaction of 22.0 mL of 0.10M acetic acid,  $\text{CH}_3\text{COOH}$ , with 22.0 mL of 0.10 M NaOH. [ $K_a = 2 \times 10^{-5}$ ]

IE0045

Q.43 Calculate the hydronium ion concentration and the pH at the equivalence point in a titration of 50.0 mL of 0.40 M  $\text{NH}_3$  with 0.40M HCl. [ $K_b = 2 \times 10^{-5}$ ]

IE0046

Q.44  $\text{CH}_3\text{COOH}$  (50 ml, 0.1 M) is titrated against 0.1 M NaOH solution. Calculate the pH at the addition of 0 ml, 10 ml, 20 ml, 25 ml, 40 ml, 50 ml of NaOH.  $K_a$  of  $\text{CH}_3\text{COOH}$  is  $2 \times 10^{-5}$ .  
[ $\log 2 = 0.3010$ ,  $\log 3 = 0.4771$ ]

IE0047

### INDICATORS

Q.45 For the acid indicator thymol blue, pH is 3 when half the indicator is in unionised form. Find the % of indicator in unionised form in the solution with  $[\text{H}^+] = 4 \times 10^{-3}$  M.

IE0048

Q.46 Bromophenol blue is an acid indicator with a  $K_a$  value of  $6 \times 10^{-5}$ . What % of this indicator is in its basic form at a pH of 5 ?

IE0049

Q.47 At what pH does an indicator change colour if the indicator is a weak acid with  $K_{\text{ind}} = 4 \times 10^{-4}$ . For which one(s) of the following neutralizations would the indicator be useful ? Explain.

(a)  $\text{NaOH} + \text{CH}_3\text{COOH}$                       (b)  $\text{HCl} + \text{NH}_3$                       (c)  $\text{HCl} + \text{NaOH}$

IE0050

Q.48 An acid indicator has a  $K_a$  of  $3 \times 10^{-5}$ . The acid form of the indicator is red & the basic form is blue. By how much must the pH change in order to change the indicator from 75% red to 75% blue?  
[ $\log 3 = 0.4771$ ]

IE0051

### SOLUBILITY & SOLUBILITY PRODUCT'S

Q.49 The values of  $K_{\text{sp}}$  for the slightly soluble salts MX and  $\text{QX}_2$  are each equal to  $4.0 \times 10^{-18}$ . Which salt is more soluble? Explain your answer fully.

IE0052

Q.50 The solubility of  $\text{PbSO}_4$  in water is 0.0608 g/L. Calculate the solubility product constant of  $\text{PbSO}_4$ .  
Molar mass  $\text{PbSO}_4 = 304$  g/mole

IE0053

Q.51 How many mole CuI ( $K_{\text{sp}} = 5 \times 10^{-12}$ ) will dissolve in 1.0 L of 0.10 M NaI solution ?

IE0054

Q.52 A solution of saturated  $\text{CaF}_2$  is found to contain  $4 \times 10^{-4}$  M fluoride ion. Calculate the  $K_{\text{sp}}$  of  $\text{CaF}_2$ . Neglect hydrolysis.

IE0055

Q.53 The solubility of  $\text{ML}_2$  (formula weight = 60 g/mol) in water is  $2.4 \times 10^{-5}$  g/100 mL solution. Calculate the solubility product constant for  $\text{ML}_2$ .

IE0056

Q.54 Calculate the solubility of  $\text{A}_2\text{X}_3$  in pure water, assuming that neither kind of ion reacts with water. For  $\text{A}_2\text{X}_3$ ,  $K_{\text{sp}} = 1.08 \times 10^{-23}$

IE0057

Q.55 Determine the solubility of  $\text{AgCl}$  in 0.1 M  $\text{BaCl}_2$ . [ $K_{\text{sp}}$  for  $\text{AgCl} = 1 \times 10^{-10}$ ]

IE0058

Q.56 Calculate solubility of  $\text{Ca}_3(\text{PO}_4)_2$  ( $K_{\text{sp}} = 10^{-15}$ ) in presence of 0.1 M  $\text{CaCl}_2$  solution.

IE0059

### SIMULTANEOUS SOLUBILITY

Q.57 Calculate the Simultaneous solubility of  $\text{AgSCN}$  and  $\text{AgBr}$ .  $K_{\text{sp}}$  ( $\text{AgSCN}$ ) =  $3.2 \times 10^{-12}$ ,  $K_{\text{sp}}$  ( $\text{AgBr}$ ) =  $8 \times 10^{-13}$ .

IE0060

Q.58 Calculate  $\text{F}^-$  in a solution saturated with respect of both  $\text{MgF}_2$  and  $\text{SrF}_2$ .  $K_{\text{sp}}$  ( $\text{MgF}_2$ ) =  $9.5 \times 10^{-9}$ ,  $K_{\text{sp}}$  ( $\text{SrF}_2$ ) =  $4 \times 10^{-9}$ .

IE0212

### COMPLEX FORMATION

Q.59 Calculate the solubility of  $\text{AgCl}$  in 0.2 M -  $\text{NH}_3$  solution.

Given :  $K_{\text{sp}}$  of  $\text{AgCl} = 2 \times 10^{-10}$ ,  $K_{\text{f}}$  of  $\text{Ag}(\text{NH}_3)_2^+ = 8 \times 10^6$ .

IE0061

Q.60 Calculate the solubility of  $\text{AgCN}$  in 0.4 M -  $\text{KCN}$  solution

(i) neglecting complex formation

(ii) considering complex formation. Given :  $K_{\text{sp}}$  of  $\text{AgCN} = 8 \times 10^{-10}$ ,  $K_{\text{d}}$  of  $\text{Ag}(\text{CN})_2^- = 4 \times 10^{-8}$ .

IE0213

### SOLUBILITY, CONSIDERING HYDROLYSIS

Q.61 Calculating the solubility of  $\text{MX}$  in water. Also calculate pH of solution.

Given :  $K_{\text{sp}}$  of  $\text{MX} = 4 \times 10^{-8}$ ;  $K_{\text{a}}$  of  $\text{HX} = 2 \times 10^{-6}$  and  $\text{MOH}$  is strong base.

IE0214

Q.62 Calculate the solubility of  $\text{AgCN}$  in a buffer solution at pH = 3.0. .

Given :  $K_{\text{sp}}$  of  $\text{AgCN} = 8 \times 10^{-10}$ ,  $K_{\text{a}}$  of  $\text{HCN} = 5 \times 10^{-10}$ .

IE0062

### PRECIPITATION

Q.63 A solution has a  $\text{Mg}^{2+}$  concentration of 0.0010 mol/L. Will  $\text{Mg}(\text{OH})_2$  precipitate if the  $\text{OH}^-$  concentration of the solution is [ $K_{\text{sp}} = 1.2 \times 10^{-11}$ ]

(a)  $10^{-5}$  mol/L

(b)  $10^{-3}$  mol/L ?

IE0063

Q.64 200 ml of  $2 \times 10^{-4}$  M -  $\text{AgNO}_3$  solution is mixed with 400 ml of  $1.2 \times 10^{-6}$  M -  $\text{NaCl}$  solution. Predict whether precipitation of  $\text{AgCl}$  will occur or not.  $K_{\text{sp}}$  of  $\text{AgCl} = 2 \times 10^{-10}$ .

IE0064

Q.65 Calculate the minimum mass of  $\text{Na}_2\text{SO}_4$  needed to just start precipitation of  $\text{BaSO}_4$  from 500 ml of  $2 \times 10^{-5}$  M -  $\text{BaCl}_2$  solution.  $K_{\text{sp}}$  of  $\text{BaSO}_4 = 8 \times 10^{-8}$ .

IE0065

## EXERCISE # S-II

- Q.1 What are the concentrations of  $\text{H}^+$ ,  $\text{HSO}_4^-$ ,  $\text{SO}_4^{2-}$  and  $\text{H}_2\text{SO}_4$  in a 0.20 M solution of sulphuric acid?

Given:  $\text{H}_2\text{SO}_4 \longrightarrow \text{H}^+ + \text{HSO}_4^-$ ; strong



IE0066

- Q.2 Calculate the pH of a 0.1M solution of  $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$ ; ethylenediamine (en). Determine the en  $\text{H}_2^{2+}$  concentration in the solution.  $K_{b1}$  and  $K_{b2}$  values of ethylenediamine are  $9 \times 10^{-5}$  and  $7.1 \times 10^{-8}$  respectively.

IE0215

- Q.3 Nicotine,  $\text{C}_{10}\text{H}_{14}\text{N}_2$ , has two basic nitrogen atoms and both can react with water to give a basic solution  
 $\text{Nic (aq)} + \text{H}_2\text{O (l)} \rightleftharpoons \text{NicH}^+ \text{ (aq)} + \text{OH}^- \text{ (aq)}$   
 $\text{NicH}^+ \text{ (aq)} + \text{H}_2\text{O (l)} \rightleftharpoons \text{NicH}_2^{2+} \text{ (aq)} + \text{OH}^- \text{ (aq)}$   
 $K_{b1}$  is  $8 \times 10^{-7}$  and  $K_{b2}$  is  $10^{-10}$ . Calculate the approximate pH of a 0.20 M solution.

IE0067

- Q.4 Determine the  $[\text{S}^{2-}]$  in a saturated (0.1M)  $\text{H}_2\text{S}$  solution to which enough HCl has been added to produce a  $[\text{H}^+]$  of  $2 \times 10^{-4}$ .  $K_1 = 10^{-7}$ ,  $K_2 = 10^{-14}$ .

IE0216

- Q.5 An aqueous solution contains 0.01 M  $\text{RNH}_2$  ( $K_b = 2 \times 10^{-6}$ ) &  $10^{-4}$  M NaOH. The concentration of  $\text{OH}^-$  is nearly :

IE0068



- Q.6 Calculate the pH of  $1.0 \times 10^{-3}$  M sodium phenoxide,  $\text{NaOC}_6\text{H}_5$ .  $K_a$  for  $\text{HOC}_6\text{H}_5$  is  $0.6 \times 10^{-10}$ .

IE0069

- Q.7 Calculate the  $\text{OH}^-$  concentration and the  $\text{H}_3\text{PO}_4$  concentration of a solution prepared by dissolving 0.1 mol of  $\text{Na}_3\text{PO}_4$  in sufficient water to make 1L of solution.  $K_1 = 7.1 \times 10^{-3}$ ,  $K_2 = 6.3 \times 10^{-8}$ ,  $K_3 = 4.5 \times 10^{-13}$ .

IE0217

- Q.8 Calculate the pH of 0.1 M solution of (i)  $\text{NaHCO}_3$ , (ii)  $\text{Na}_2\text{HPO}_4$  and (iii)  $\text{NaH}_2\text{PO}_4$ . Given that:



( $\log 4.2 = 0.62$ ,  $\log 4.8 = 6.8$ ,  $\log 6.2 = 0.80$ ,  $\log 7.5 = 0.88$ )

IE0070

- Q.9 An ammonia-ammonium chloride buffer has a pH value of 9 with  $[\text{NH}_3] = 0.25$ . What will be the new pH if 500 ml 0.1 M KOH is added to 200 ml buffer solution ( $K_b = 2 \times 10^{-5}$ )  
 $[\log 2 = 0.3]$

IE0071



- Q.10 A weak base (50.0mL) was titrated with 0.1 M HCl. The pH of the solution after the addition of 10.0 mL and 25.0 mL were found to be 9.84 and 9.24, respectively. Calculate  $K_b$  of the base and pH at the equivalence point.  $[\log 2 = 0.3]$
- IE0218**
- Q.11 A weak acid (50.0mL) was titrated with 0.1 M NaOH. The pH values when 10.0 mL and 25.0 mL of base have been added are found to be 4.16 and 4.76, respectively. Calculate  $K_a$  of the acid and pH at the equivalence point.  $[\log 2 = 0.3]$
- IE0219**
- Q.12 10 ml of 0.1M weak acid  $HA (K_a = 10^{-5})$  is mixed with 10 ml 0.2M HCl and 10 ml 0.1M NaOH. Find the value of  $[A^-]$  in the resulting solution.
- IE0072**
- 
- Q.13 150 ml of 0.5 M HCN ( $K_a = 3.75 \times 10^{-9}$ ) was reacted with 1.5 M KOH for complete neutralisation. What will be molarity of HCN at equilibrium.
- IE0073**
- Q.14 The indicator phenol red is half in the ionic form when pH is 7.2. If the ratio of the undissociated form to the ionic form is 1 : 5, find the pH of the solution. With the same pH for solution, if indicator is altered such that the ratio of undissociated form to dissociated form becomes 1 : 4, find the pH when 50 % of the new indicator is in ionic form.  $[\log 2 = 0.3]$
- IE0220**
- Q.15 How much AgBr could dissolve in 1.0 L of 0.40 M  $NH_3$  ? Assume that  $Ag(NH_3)_2^+$  is the only complex formed.  $[K_f (Ag(NH_3)_2^+) = 1 \times 10^8 ; K_{sp} (AgBr) = 5 \times 10^{-13}]$
- IE0074**
- $[\sqrt{50} \approx 7]$
- Q.16 Calculate solubility of  $PbI_2 (K_{sp} = 1.4 \times 10^{-8})$  in water at  $25^\circ$ , which is 90% dissociated.
- IE0075**
- $$\left( \frac{1.4}{(0.81)(3.6)} \right)^{1/3} = 0.78$$
- Q.17 A recent investigation of the complexation of  $SCN^-$  with  $Fe^{3+}$  led to 130, 16, and 1.0 for  $K_1$ ,  $K_2$ , and  $K_3$ , respectively. What is the overall formation constant of  $Fe(SCN)_3$  from its component ions, and what is the dissociation constant of  $Fe(SCN)_3$  into its simplest ions on the basis of these data ?
- IE0221**

## EXERCISE # O-I

## Single correct

- Q.1 The conjugate acid of  $\text{NH}_2^-$  is  
 (A)  $\text{NH}_3$  (B)  $\text{NH}_2\text{OH}$  (C)  $\text{NH}_4^+$  (D)  $\text{N}_2\text{H}_4$  IE0076
- Q.2 Which of the following is not a Bronsted acid:-  
 (A)  $\text{CH}_3\text{NH}_4^+$  (B)  $\text{CH}_3\text{COO}^-$  (C)  $\text{H}_2\text{O}$  (D)  $\text{HSO}_4^-$  IE0077
- Q.3 In the reaction  
 $\text{HNO}_3 + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{NO}_3^-$ , the conjugate base of  $\text{HNO}_3$  is :-  
 (A)  $\text{H}_2\text{O}$  (B)  $\text{H}_3\text{O}^+$  (C)  $\text{NO}_3^-$  (D)  $\text{H}_3\text{O}^+$  and  $\text{NO}_3^-$  IE0078
- Q.4 Out of the following, amphoteric species in aqueous medium are  
 I:  $\text{HPO}_3^{2-}$  II  $\text{OH}^-$  III  $\text{H}_2\text{PO}_4^-$  IV  $\text{HCO}_3^-$   
 (A) I, III, IV (B) I and III (C) III and IV (D) All IE0079
- Q.5 When ammonia is added to water, it decreases the concentration of which of the following ion  
 (A)  $\text{OH}^-$  (B)  $\text{H}_3\text{O}^+$  (C)  $\text{NH}_4^+$  (D)  $\text{NH}_4^+$  &  $\text{OH}^-$  IE0080
- Q.6 Which of the following pair is Lewis acid & Lewis base & Product of these is also Lewis base  
 (A)  $\text{BF}_3$ ,  $\text{NH}_3$  (B)  $\text{SiCl}_4$ ,  $2\text{Cl}^-$  (C)  $\text{CH}_3^+$ ,  $^-\text{OC}_2\text{H}_5$  (D) All of these IE0081
- Q.7 Ionic product of water will increase, if :-  
 (A) Pressure is decreased (B)  $\text{H}^+$  is added  
 (C)  $\text{OH}^-$  is increased (D) Temperature is increased IE0082
- Q.8 At  $60^\circ\text{C}$ , pure water has  $[\text{H}_3\text{O}^+] = 10^{-6.7} \text{ mol/lit.}$  what is the value of  $K_w$  at  $60^\circ\text{C}$  :-  
 (A)  $10^{-6}$  (B)  $10^{-12}$  (C)  $10^{-67}$  (D)  $10^{-13.4}$  IE0083
- Q.9 Liquid  $\text{NH}_3$  ionises to a slight extent. At a certain temperature its self ionization constant  $K_{\text{SIC}(\text{NH}_3)} = 10^{-30}$ . The number of  $\text{NH}_4^+$  ions present per  $100 \text{ cm}^3$  of pure liquid are  
 (A)  $10^{-15}$  (B)  $6.022 \times 10^8$  (C)  $6.022 \times 10^7$  (D) None IE0084
- Q.10 The pH of solution is increased from 3 to 6. Its  $\text{H}^+$  ion conc. will be :-  
 (A) Reduced to half (B) Doubled  
 (C) Reduced by 1000 times (D) Increased by 1000 times IE0085
- Q.11. pOH of  $[1/200] \text{ mol/m}^3 \text{ H}_2\text{SO}_4 \text{ (aq.)}$  solution at  $25^\circ\text{C}$  is-  
 (A) 2 (B) 5 (C) 9 (D) 12 IE0086

- Q.12 Degree of dissociation of 0.1 N  $\text{CH}_3\text{COOH}$  is :- (Dissociation constant =  $1 \times 10^{-5}$ )  
(A)  $10^{-5}$  (B)  $10^{-4}$  (C)  $10^{-3}$  (D)  $10^{-2}$  **IE0087**
- Q.13 The pH of a 0.02 M ammonia solution which is 5% ionised will be :-  
(A) 2 (B) 11 (C) 5 (D) 7 **IE0088**
- Q.14 The pH of an aqueous solution of 1.0 M solution of a weak monoprotic acid which is 1% ionised is  
(A) 1 (B) 2 (C) 3 (D) 11 **IE0089**
- Q.15 The concentration of  $[\text{H}^+]$  and concentration of  $[\text{OH}]^-$  of a 0.1 M aqueous solution of 2% ionised weak acid is [ionic product of water =  $1 \times 10^{-14}$ ]  
(A)  $0.02 \times 10^{-3}$  M and  $5 \times 10^{-11}$  M (B)  $1 \times 10^{-3}$  M and  $3 \times 10^{-11}$  M  
(C)  $2 \times 10^{-3}$  M and  $5 \times 10^{-12}$  M (D)  $3 \times 10^{-2}$  M and  $4 \times 10^{-13}$  M **IE0090**
- Q.16 What is the quantity of NaOH present in 250 cc of the solution, so that it gives a pH = 13 :-  
(A)  $10^{-13}$  g (B)  $10^{-1}$  g (C) 1.0 g (D) 4.0 g **IE0091**
- Q.17 An aqueous solution of HCl is  $10^{-9}$  M HCl. The pH of the solution should be:-  
(A) 9 (B) Between 6 and 7 (C) 7 (D) Unpredictable **IE0092**
- Q.18. The moles of  $\text{H}^+$  from  $\text{H}_2\text{O}$  in a 1l,  $\sqrt{5} \times 10^{-7}$  M HCl solution at  $25^\circ\text{C}$ , is  
( $\sqrt{5} = 2.23$ )  
(A)  $10^{-7}$  (B)  $6.85 \times 10^{-8}$  (C)  $3.85 \times 10^{-8}$  (D)  $10^{-8}$  **IE0093**
- Q.19 Which one of the following has highest pH:-  
(A) Distilled water (B) 1 M  $\text{NH}_3$   
(C) 1 M NaOH (D) Water saturated with chlorine **IE0094**
- Q.20 8 gm NaOH and 4.9 gm  $\text{H}_2\text{SO}_4$  are present in one litre of the solution. What is its pH  
(A) 1 (B) 13 (C) 12 (D) 2 **IE0095**
- Q.21 10 ml of  $\frac{\text{M}}{200} \text{H}_2\text{SO}_4$  is mixed with 40 ml of  $\frac{\text{M}}{200} \text{H}_2\text{SO}_4$ . The pH of the resulting solution is  
(A) 1 (B) 2 (C) 2.3 (D) none of these **IE0096**
- Q.22 Which of the following solution will have pH close to 1.0?  
(A) 100 ml of M/100 HCl + 100 ml of M/10 NaOH  
(B) 55 ml of M/10 HCl + 45 ml of M/10 NaOH  
(C) 10 ml of M/10 HCl + 90 ml of M/10 NaOH  
(D) 75 ml of M/5 HCl + 25 ml of M/5 NaOH **IE0097**

- Q.23 A solution with pH 2.0 is more acidic than the one with pH 6.0 by a factor of:  
 (A) 3 (B) 4 (C) 3000 (D) 10000 **IE0098**
- Q.24 The first and second dissociation constants of an acid  $H_2A$  are  $1.0 \times 10^{-5}$  and  $5.0 \times 10^{-10}$  respectively. The overall dissociation constant of the acid will be :  
 (A)  $5.0 \times 10^{-5}$  (B)  $5.0 \times 10^{15}$  (C)  $5.0 \times 10^{-15}$  (D)  $0.2 \times 10^5$  **IE0099**
- Q.25 If  $pK_b$  for fluoride ion at  $25^\circ C$  is 10.4, the ionisation constant of hydrofluoric acid in water at this temperature is :  
 (A)  $4 \times 10^{-11}$  (B)  $3 \times 10^{-3}$  (C)  $2.5 \times 10^{-4}$  (D)  $2 \times 10^{-2}$  **IE0100**
- Q.26 pH of an aqueous solution of NaCl at  $85^\circ C$  should be  
 (A) 7 (B)  $> 7$  (C)  $< 7$  (D) 0 **IE0101**
- Q.27 1 cc of 0.1 N HCl is added to 99 cc solution of NaCl. The pH of the resulting solution will be  
 (A) 7 (B) 3 (C) 4 (D) 1 **IE0102**
- Q.28 The degree of hydrolysis of a salt of weak acid and weak base in its 0.1 M solution is found to be 50%. If the molarity of the solution is 0.2 M, the percentage hydrolysis of the salt should be  
 (A) 100% (B) 50% (C) 25% (D) none of these **IE0103**
- Q.29 What is the percentage hydrolysis of NaCN in N/80 solution when the dissociation constant for HCN is  $2 \times 10^{-9}$  and  $K_w = 1.0 \times 10^{-14}$   
 (A) 2 (B) 5.26 (C) 8.2 (D) 9.6 **IE0104**
- Q.30 The compound whose 0.1 M solution is basic is  
 (A) Ammonium acetate (B) Ammonium chloride  
 (C) Ammonium sulphate (D) Sodium acetate **IE0105**
- Q.31 If equilibrium constant of  
 $CH_3COOH + H_2O \rightleftharpoons CH_3COO^- + H_3O^+$   
 is  $1.8 \times 10^{-5}$ , equilibrium constant for  
 $CH_3COOH + OH^- \rightleftharpoons CH_3COO^- + H_2O$  is  
 (A)  $1.8 \times 10^{-9}$  (B)  $1.8 \times 10^9$  (C)  $5.55 \times 10^{-9}$  (D)  $5.55 \times 10^{10}$  **IE0106**
- Q.32 The  $pK_a$  of a weak acid, HA, is 4.80. The  $pK_b$  of a weak base, BOH, is 4.78. The pH of an aqueous solution of the corresponding salt, BA, will be :  
 (A) 8.58 (B) 4.79 (C) 7.01 (D) 9.22 **IE0107**
- Q.33 The highest pH value is of :-  
 (A) 0.1 M NaCl (B) 0.1 M  $NH_4Cl$   
 (C) 0.1 M  $CH_3COONa$  (D) 0.1 M  $CH_3COONH_4$  **IE0108**

- Q.34 pH of  $K_2S$  solution is:-  
(A) 7 (B) Less than 7 (C) More than 7 (D) 0  
**IE0109**
- Q.35 Degree of Hydrolysis of  $\frac{N}{100}$  solution of KCN is (Given  $K_a = 1.6 \times 10^{-9}$ )  
(A)  $2.5 \times 10^{-3}$  (B)  $2.5 \times 10^{-2}$  (C)  $2.5 \times 10^{-4}$  (D)  $2.5 \times 10^{-5}$   
**IE0110**
- Q.36 A solution of  $FeCl_3$  in water acts as acidic due to :-  
(A) Acidic impurities (B) Ionisation (C) Hydrolysis of  $Fe^{3+}$  (D) Dissociation  
**IE0111**
- Q.37 If 40 ml of 0.2 M KOH is added to 160 ml of 0.1 M  $HCOOH$  [ $K_a = 2 \times 10^{-4}$ ], the pOH of the resulting solution is  
(A) 3.4 (B) 3.7 (C) 7 (D) 10.3  
**IE0112**
- Q.38 1 M NaCl and 1M HCl are present in an aqueous solution. The solution is  
(A) not a buffer solution and with  $pH < 7$   
(B) not a buffer solution with  $pH > 7$   
(C) a buffer solution with  $pH < 7$   
**IE0113**
- Q.39 The  $pK_a$  of a weak acid (HA) is 4.5. The pOH of an aqueous buffered solution of HA in which 50% of the acid is ionized is :  
(A) 4.5 (B) 2.5 (C) 9.5 (D) 7.0  
**IE0114**
- Q.40 To a 50 ml. of 0.05M formic acid, how much volume of 0.10M sodium formate must be added to get a buffer solution of  $pH = 4.0$  ?  
( $pK_a$  of the acid is 3.7) ( $\log 2 = 0.3$ )  
(A) 40 ml. (B) 4 ml. (C) 50 ml. (D) 100 ml.  
**IE0115**
- Q.41 Which can act as buffer :-  
(A)  $NH_4OH + NaOH$   
(B)  $HCOOH + HCl$   
(C) 40 ml. of 0.1 M NaCN + 20 ml. of 0.1 M HCl  
(D) All of them  
**IE0116**
- Q.42 If equal volume of 0.05 M ammonium hydroxide solution is dissolved in 0.001 M ammonium chloride solution. What will be the  $OH^-$  ion concentration of this solution :  
 $K_b(NH_4OH) = 1.8 \times 10^{-5}$   
(A)  $3.0 \times 10^{-3}$  (B)  $4.6 \times 10^{-4}$  (C)  $9.0 \times 10^{-3}$  (D)  $9.0 \times 10^{-4}$   
**IE0117**
- Q.43 Calculate the pH of a buffer prepared by mixing 600 cc of 0.6 M  $NH_3$  and 400 cc of 0.5 M  $NH_4Cl$ .  
 $K_b$  for  $NH_3 = 1.8 \times 10^{-5}$ , ( $\log 1.8 = 0.26$ )  
(A) 11.3 (B) 9.0 (C) 9.52 (D) 5  
**IE0118**

- Q.44  $pK_b$  for  $NH_4OH$  at certain temperature is 4.74. The pH of basic buffer containing equimolar concentration of  $NH_4OH$  and  $NH_4Cl$  will be:-  
 (A) 7.74 (B) 4.74 (C) 2.37 (D) 9.26  
**IE0119**
- Q.45 On addition of NaOH to  $CH_3COOH$  solution, 60% of the acid is neutralised. If  $pK_a$  of  $CH_3COOH$  is 4.7 then the pH of the resulting solution is :-  
 (A) More than 4.7 but less than 5.0 (B) Less than 4.7 but more than 4.0  
 (C) More than 5.0 (D) Remains unchanged  
**IE0120**
- Q.46 Henderson equation  $pH - pK_a = 5$  will be applicable to an acidic buffer when :-  
 (A)  $[Acid] = [Conjugate\ base]$  (B)  $[Acid] \times 10^5 = [Conjugate\ base]$   
 (C)  $[Acid] = [Conjugate\ base] \times 10^5$  (D)  $[acid] = 2 [conjugate\ base]$   
**IE0121**
- Q.47 What amount of sodium propanoate should be added to one litre of an aqueous solution containing 0.02 mole of propanoic acid ( $K_a = 3 \times 10^{-5}$  at  $25^\circ C$ ) to obtain a buffer solution of pH 4.7  
 (A)  $4.52 \times 10^{-2}$  mol (B)  $3.52 \times 10^{-2}$  mol (C)  $2.52 \times 10^{-2}$  mol (D)  $3 \times 10^{-2}$  mol  
**IE0122**
- Q.48 In a buffer solution the ratio of concentration of  $NH_4Cl$  and  $NH_4OH$  is 1 : 1. When it changes in 2 : 1, what will be the value of pH of buffer ?  
 (A) Increase (B) Decrease (C) No effect (D) None  
**IE0123**
- Q.49 The buffer solution play an important role in :-  
 (A) Increasing the pH value (B) Decreasing the pH value  
 (C) Keeping the pH constant (D) Solution will be neutral  
**IE0124**
- Q.50 The total number of different kind of acidic buffers obtained during the titration of  $H_3PO_4$  with NaOH are :  
 (A) 3 (B) 1 (C) 2 (D) 0  
**IE0125**
- Q.51 Which of the following solutions does not act as buffer :  
 (A)  $H_3PO_4 + NaH_2PO_4$  (B)  $NaHCO_3 + H_2CO_3$   
 (C)  $NH_4Cl + HCl$  (D)  $CH_3COOH + CH_3COONa$   
**IE0126**
- Q.52 Half of the formic acid solution is neutralised on addition of a KOH solution to it. If  $K_a (HCOOH) = 2 \times 10^{-4}$  then pH of the solution is : - ( $\log 2 = 0.3010$ )  
 (A) 3.6990 (B) 10.3010 (C) 3.85 (D) 4.3010  
**IE0127**
- Q.53 When 0.02 moles of NaOH are added to a litre of buffer solution, its pH changes from 5.75 to 5.80. What is its buffer capacity :-  
 (A) 0.4 (B) 0.05 (C) - 0.05 (D) 2.5  
**IE0128**
- Q.54 Calculate pH when 100 ml of 0.2M NaOH is reacted with 100 ml of 0.2 M  $CH_3COOH$  ( $K_a = 10^{-5}$ )  
 (A) 9 (B) 7 (C) 5 (D) 2  
**IE0129**
- Q.55 At  $90^\circ C$ , pure water has  $[H^+] = 10^{-6} M$ , if 100 ml of 0.2 M  $HNO_3$  is added to 20 ml of 1 M NaOH at  $90^\circ C$  then pH of the resulting solution will be  
 (A) 5 (B) 6 (C) 7 (D) None of these  
**IE0130**

Q.56 When 20 ml of  $\frac{M}{20}$  NaOH are added to 10 ml of  $\frac{M}{10}$  HCl, the resulting solution will:-

- (A) Turn blue litmus red  
 (B) Turn phenolphthalein solution pink colour  
 (C) Turn methyl orange red  
 (D) Will have no effect on either red or blue litmus

**IE0131**

Q.57 The rapid change of pH near the stoichiometric point of an acid-base titration is the basis of indicator detection. pH of the solution is related to ratio of the concentrations of the conjugate acid (HIn) and base ( $\text{In}^-$ ) forms of the indicator by the expression :-

- (A)  $\log \frac{[\text{HIn}]}{[\text{In}^-]} = \text{pK}_{\text{In}} - \text{pH}$  (B)  $\log \frac{[\text{HIn}]}{[\text{In}^-]} = \text{pH} - \text{pK}_{\text{In}}$   
 (C)  $\log \frac{[\text{In}^-]}{[\text{HIn}]} = \text{pH} + \text{pK}_{\text{In}}$  (D)  $\log \frac{[\text{In}^-]}{[\text{HIn}]} = \text{pK}_{\text{In}} - \text{pH}$

**IE0132**

Q.58 Calculate the pH range in which an acid indicator with  $K_{\text{acid}}(\text{indicator}) = 1.0 \times 10^{-5}$  changes colour when the concentration of the indicator is  $1 \times 10^{-3} \text{M}$ .

- (A)  $5 \pm 1$  (B)  $11 \pm 1$  (C)  $3 \pm 1$  (D)  $8 \pm 1$

**IE0133**

Q.59 In what pH range will a  $1 \times 10^{-4} \text{M}$  solution of an indicator will  $K_{\text{b}}(\text{indicator}) = 1 \times 10^{-11}$  change colour?

- (A)  $7.0 \pm 1$  (B)  $3.0 \pm 1$  (C)  $5.5 \pm 1$  (D)  $11.0 \pm 1$

**IE0134**

Q.60 Indicator which is used in the titration of  $\text{CH}_3\text{COOH}$  & NaOH :-

- (A) Methyl orange (B) Methyl red (C) Phenolphthalein (D) Litmus

**IE0135**

Q.61 Phenolphthalein is a :-

- (A) Strong acid (B) Strong base (C) Weak base (D) Weak acid

**IE0136**

Q.62 pH-range of Methyl red indicator is :-

- (A) 4.2–6.2 (B) 6.8–10.8 (C) 8 – 9.6 (D) 6.8 – 8.2

**IE0137**

Q.63 In the volumetric estimation of HCl, if we make use of phenolphthalein as an indicator, which base is unsuitable for the titration :-

- (A) NaOH (B) RbOH (C) KOH (D)  $\text{NH}_4\text{OH}$

**IE0138**

Q.64 Phenolphthalein does not act as an indicator for the titration between :-

- (A) KOH and  $\text{H}_2\text{SO}_4$  (B) NaOH and  $\text{CH}_3\text{COOH}$   
 (C) Oxalic acid and  $\text{KMnO}_4$  (D)  $\text{Ba}(\text{OH})_2$  and HCl

**IE0139**

Q.65 For weak acid and strong base titration, the indicator used is :-

- (A) Potassium di-chromate (B) Methyl orange  
 (C) Litmus (D) Phenolphthalein

**IE0140**

Q.66 The solubility of  $\text{A}_2\text{X}_3$  is  $y \text{ mol dm}^{-3}$ . Its solubility product is

- (A)  $6y^2$  (B)  $64y^4$  (C)  $36y^5$  (D)  $108y^5$

**IE0141**

- Q.67 If  $K_{sp}$  for  $HgSO_4$  is  $6.4 \times 10^{-5}$ , then solubility of this substance in mole per  $m^3$  is  
 (A)  $8 \times 10^{-3}$  (B)  $6.4 \times 10^{-5}$  (C)  $8 \times 10^{-6}$  (D) 8 **IE0142**
- Q.68 If the solubility of  $AgCl$  (formula mass=143) in water at  $25^\circ C$  is  $1.43 \times 10^{-4}$  gm/100 ml of solution then the value of  $K_{sp}$  will be :-  
 (A)  $1 \times 10^{-5}$  (B)  $2 \times 10^{-5}$  (C)  $1 \times 10^{-10}$  (D)  $2 \times 10^{-10}$  **IE0143**
- Q.69 One litre of saturated solution of  $CaCO_3$  is evaporated to dryness, 7.0 g of residue is left. The solubility product for  $CaCO_3$  is:-  
 (A)  $4.9 \times 10^{-3}$  (B)  $4.9 \times 10^{-5}$  (C)  $4.9 \times 10^{-9}$  (D)  $4.9 \times 10^{-7}$  **IE0144**
- Q.70  $A_3B_2$  is a sparingly soluble salt of molar mass  $M$  ( $g\ mol^{-1}$ ) and solubility  $x\ g\ lit^{-1}$ . The ratio of the molar concentration of  $B^{3-}$  to the solubility product of the salt is  
 (A)  $108 \frac{x^5}{M^5}$  (B)  $\frac{1}{108} \frac{M^4}{x^4}$  (C)  $\frac{1}{54} \frac{M^4}{x^4}$  (D) None **IE0145**
- Q.71. Solubility of  $Ag_2CrO_4$  ( $K_{sp} = 4 \times 10^{-13}$ ) in  $0.1\ M\ K_2CrO_4$  solution will be :-  
 (A)  $10^{-3}\ M$  (B)  $10^{-6}\ M$  (C)  $4 \times 10^{-6}\ M$  (D)  $5 \times 10^{-7}\ M$  **IE0146**
- Q.72. How many times solubility of  $CaF_2$  is decreased in  $4 \times 10^{-3}\ M\ KF$  (aq.) solution as compare to pure water at  $25^\circ C$ . Given  $K_{sp}(CaF_2) = 3.2 \times 10^{-11}$   
 (A) 50 (B) 100 (C) 500 (D) 1000 **IE0147**
- Q.73 At  $30^\circ C$ , In which of the one litre solution, the solubility of  $Ag_2CO_3$  (solubility product =  $8 \times 10^{-12}$ ) will be maximum :-  
 (A)  $0.05\ M\ Na_2CO_3$  (B) Pure water  
 (C)  $0.05\ M\ AgNO_3$  (D)  $0.05\ M\ NH_3$  **IE0148**
- Q.74 What will happen if the pH of the solution of  $0.001\ M\ Mg(NO_3)_2$  solution is adjusted to  $pH = 9$  ( $K_{sp}$  of  $Mg(OH)_2 = 8.9 \times 10^{-12}$ )  
 (A) ppt will take place  
 (B) ppt will not take place  
 (C) Solution will be saturated  
 (D) None of these **IE0149**
- Q.75  $Na_3PO_4$  which should be added in  $10\ L$  of  $1.0 \times 10^{-5}\ M$  -  $BaCl_2$  solution without any precipitation of  $Ba_3(PO_4)_2$  is [ $K_{sp}$  of  $Ba_3(PO_4)_2$ ] =  $4 \times 10^{-23}$   
 (A)  $2 \times 10^{-4}\ gm$  (B)  $0.328\ gm$  (C)  $0.164\ gm$  (D)  $0.82\ gm$  **IE0150**



**EXERCISE # O-II****Single correct :**

- Q.1 The pH of the solution produced when an aqueous solution of strong acid pH 5 is mixed with equal volume of an aqueous solution of strong acid of pH 3 is :-  
 (A) 3.3 (B) 3.5 (C) 4.5 (D) 4.0

**IE0151**

- Q.2 How many moles of HCl must be removed from 1 litre of aqueous HCl solution to change its pH from 2 to 3 :-  
 (A) 1 (B) 0.02 (C) 0.009 (D) 0.01

**IE0152**

- Q.3 Which of the following is most soluble in water?  
 (A)  $\text{MnS}$  ( $K_{\text{sp}} = 8 \times 10^{-37}$ ) (B)  $\text{ZnS}$  ( $K_{\text{sp}} = 7 \times 10^{-16}$ )  
 (C)  $\text{Bi}_2\text{S}_3$  ( $K_{\text{sp}} = 1 \times 10^{-72}$ ) (D)  $\text{Ag}_3(\text{PO}_4)$  ( $K_{\text{sp}} = 1.8 \times 10^{-18}$ )

**IE0153**

- Q.4 Solubility of AgBr will be minimum in :-  
 (A) Pure water (B) 0.1 M  $\text{CaBr}_2$  (C) 0.1 M NaBr (D) 0.1 M  $\text{AgNO}_3$

**IE0154**

- Q.5 pH of solution at first  $1/4^{\text{th}}$  equivalence point of  $\text{Na}_2\text{CO}_3$  when titrated with HCl will be (for  $\text{H}_2\text{CO}_3$   $K_{\text{a}1} = 10^{-7}$ ;  $K_{\text{a}2} = 10^{-11}$ )  
 (A)  $7 + \log 3$  (B)  $7 - \log 3$  (C)  $11 + \log 3$  (D)  $11 - \log 3$

**IE0155**

- Q.6 An acid-base indicator has a  $K_{\text{a}} = 1.0 \times 10^{-5}$ . The acid form of the indicator is red and the basic form is blue. Calculate the pH change required to change the colour of the indicator from 80% red to 80% blue.  
 (A) 1.20 (B) 0.80 (C) 0.20 (D) 1.40

**IE0156****Assertion /Reason :**

- Q.7 **Statement-1** pH of  $10^{-7}$  M NaOH solution is exist between 7 to 7.3 at  $25^\circ\text{C}$ .

**Statement-2** Due to common ion effect ionization of water is reduced.

- (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.  
 (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.  
 (C) Statement-1 is true, statement-2 is false.  
 (D) Statement-1 is false, statement-2 is true.

**IE0157**

- Q.8 **Statement-1** In general phenolphthalein is used as an indicator for the titration of weak acid (HA) against strong base (NaOH)

**Statement-2** At equivalent point solution is basic.

- (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.  
 (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.  
 (C) Statement-1 is true, statement-2 is false.  
 (D) Statement-1 is false, statement-2 is true.

**IE0222**

- Q.9 **Statement-1** : Moles of  $\text{Sr}^{2+}$  furnished by sparingly soluble substance  $\text{Sr}(\text{OH})_2$  decreases due to dilution.  
**Statement-2** : Solubility product constant of  $\text{Sr}(\text{OH})_2$  is not affected by dilution.  
 (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.  
 (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.  
 (C) Statement-1 is true, statement-2 is false.  
 (D) Statement-1 is false, statement-2 is true.

IE0223

- Q.10 **Statement-1** : On dilution of a concentrated solution of  $\text{CH}_3\text{COOH}$ , the concentration of  $[\text{H}^+]$  decreases.  
**Statement-2** : Increase in volume is more than the increase in degree of ionisation.  
 (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.  
 (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.  
 (C) Statement-1 is true, statement-2 is false.  
 (D) Statement-1 is false, statement-2 is true.

IE0158

**Multiple correct :**

- Q.11 Which of the following is correct for 0.1 M BOH solution ( $K_b = 10^{-5}$ )  
 (A) pH of solution is 11  
 (B)  $\text{OH}^-$  concentration is  $10^{-3}$  mol/L  
 (C) it's salt with HCl (i.e. BCl) form the acidic solution in water  
 (D) Phenolphthalein indicator can be used during the titration of BOH with HCl

IE0159

- Q.12 For weak monobasic acid, HA, the dissociation constant is  $2 \times 10^{-6}$ , at  $25^\circ\text{C}$ . Which of the following is/are correct regarding this acid? [ $\log 2 = 0.3$ ]  
 (A)  $\text{A}^- + \text{H}_2\text{O} \rightleftharpoons \text{HA} + \text{OH}^-$ ;  $K_{\text{eq}} = 5 \times 10^{-9}$   
 (B) The equilibrium constant for the reaction of HA with aq. NaOH is  $2 \times 10^8$   
 (C) The pH of 0.1 M, HA solution is 3.35  
 (D) solution of  $\text{A}^-$  is basic

IE0160

- Q.13 Select correct statement for 50ml 0.1M  $\text{H}_2\text{A}(\text{aq.})$  solution ;  $K_{a_1} = 10^{-5}$  ;  $K_{a_2} = 10^{-8}$   
 (A)  $[\text{H}^+] = 2[\text{A}^{2-}]$   
 (B) pH of 0.1M  $\text{H}_2\text{A}$  solution is 3  
 (C) In above  $\text{H}_2\text{A}$  solution when 5 milimoles of NaHA are added then pH increases by 2 units  
 (D) 50 ml of 0.1M NaOH required to neutralised completely 50 ml of 0.1M  $\text{H}_2\text{A}$  solution

IE0224

- Q.14 A solution containing 0.01M each of  $\text{Pb}^{2+}$ ,  $\text{Ag}^+$ ,  $\text{Zn}^{2+}$  &  $\text{Cr}^{3+}$  ion. If solid  $\text{Na}_2\text{S}$  is added slowly to the solution then correct statement is based on given data -

Precipitate	PbS	Ag <sub>2</sub> S	ZnS	Cr <sub>2</sub> S <sub>3</sub>
K <sub>sp</sub>	10 <sup>-11</sup>	10 <sup>-12</sup>	10 <sup>-8</sup>	10 <sup>-10</sup>

- (A)  $\text{Pb}^{2+}$  will start precipitating first  
 (B)  $\text{Cr}^{3+}$  will start precipitating last  
 (C)  $\text{Zn}^{2+}$  will start precipitating before  $\text{Ag}^+$   
 (D) When  $\text{Zn}^{2+}$  just starts precipitating then  $\text{Pb}^{2+}$  ion gets 99.9% precipitated

IE0225

- Q.15 If  $K_1$  &  $K_2$  be first and second ionisation constant of  $\text{H}_3\text{PO}_4$  and  $K_1 \gg K_2$  which is/are incorrect.

- (A)  $[\text{H}^+] = [\text{H}_2\text{PO}_4^-]$  (B)  $[\text{H}^+] = \sqrt{K_1[\text{H}_3\text{PO}_4]}$   
 (C)  $K_2 = [\text{HPO}_4^{2-}]$  (D)  $[\text{H}^+] = 3[\text{PO}_4^{3-}]$

IE0161

- Q.16 10 ml. of a solution contains 0.1 M  $\text{NH}_4\text{Cl}$  + 0.01M  $\text{NH}_4\text{OH}$ . Which addition would not change the pH of solution :-

- (A) Adding 1 ml. water  
 (B) Adding 5 ml. of 0.1 M  $\text{NH}_4\text{Cl}$   
 (C) Adding 5 ml. of 0.1 M  $\text{NH}_4\text{OH}$   
 (D) Adding 10 ml. of 0.1 M  $\text{NH}_4\text{Cl}$

IE0162

- Q.17 When equal volumes of the following solutions are mixed, precipitation of  $\text{AgCl}$  ( $K_{sp} = 1.8 \times 10^{-10}$ ) will occur only with:

- (A)  $10^{-4}$  M ( $\text{Ag}^+$ ) and  $10^{-4}$  M ( $\text{Cl}^-$ )  
 (B)  $10^{-5}$  M ( $\text{Ag}^+$ ) and  $10^{-5}$  M ( $\text{Cl}^-$ )  
 (C)  $10^{-6}$  M ( $\text{Ag}^+$ ) and  $10^{-6}$  M ( $\text{Cl}^-$ )  
 (D)  $10^{-10}$  M ( $\text{Ag}^+$ ) and  $10^{-10}$  M ( $\text{Cl}^-$ )

IE0163

### Paragraph for Q.18 to Q.20

8 gm weak acid HX (molecular mass = 80) is dissolved in 100 ml water. ( $K_a = 10^{-4}$ )

- Q.18 Find pH of solution-

- (A) 3.3 (B) 2 (C) 2.3 (D) 3

IE0164

- Q.19 If it is titrated with 0.25 M NaOH find pH at equivalence point ( $\log 5 = 0.7$ )

- (A) 9.15 (B) 8.65 (C) 4.65 (D) 4.85

IE0165

- Q.20 Find  $[\text{H}^+]$  if  $10^{-3}$  mol HCl is added to 100 ml original solution ( $\sqrt{41} = 6.4$  ;  $\sqrt{5} = 2.24$ )

- (A)  $0.62 \times 10^{-2}$  (B)  $1.62 \times 10^{-2}$  (C)  $2.7 \times 10^{-2}$  (D)  $0.27 \times 10^{-2}$

IE0166

### Paragraph for Question 21 & 22

A solution contains one mole each of HA & HB (both are weak acids) in one litre solution. Now one mole of NaOH is added to this solution so that both the acids are partially neutralised. Heat of neutralisation of HA & HB are  $-11.8$  and  $-12.4$  kcal per mole respectively and heat produced during partial neutralisation of HA & HB is  $-12.25$  kcal.

Q.21 Mole ratio of neutralisation of HA & HB is -

- (A) 1 : 4                      (B) 1 : 2                      (C) 1 : 3                      (D) 1 : 5

IE0226

Q.22 pH of solution containing mixture of 1 mol of HA & 1 mole of HB in 1 litre solution is

$$K_a(\text{HA}) = 1 \times 10^{-5}, K_a(\text{HB}) = 9 \times 10^{-5}$$

- (A) 2                      (B) 3                      (C) 4                      (D) 5

IE0227

### MATCH THE COLUMN :

Q.23 Match the effect of addition of 1 M NaOH to 100 mL 1 M  $\text{CH}_3\text{COOH}$  (in Column I) with pH (in Column II):

#### Column-I

- (A) 25 mL of NaOH  
(B) 50 mL of NaOH  
(C) 75 mL of NaOH  
(D) 100 mL of NaOH

#### Column-II

- (P)  $\text{p}K_a$   
(Q)  $\text{p}K_a + \log 3$   
(R)  $\text{p}K_a - \log 3$   
(S)  $\frac{1}{2} [\text{p}K_w + \text{p}K_a - \log 2]$

IE0167

Q.24

#### Column-I

#### pH

- (A) 3  
(B) 5  
(C) 8  
(D) 9

#### Column-II

#### Solution

- (P) When equal volumes of 0.2M  $\text{NH}_4\text{OH}$  ( $K_b = 10^{-5}$ ) & 0.2M HCl are mixed  
(Q) When equal volumes of 0.2M  $\text{CH}_3\text{COONa}$  & 0.2M HCl are mixed ( $K_{a(\text{CH}_3\text{COOH})} = 10^{-5}$ )  
(R) 0.1M  $\text{Na}_2\text{HPO}_4$   
(for  $\text{H}_3\text{PO}_4$ ;  $K_{a_1} = 10^{-4}$ ;  $K_{a_2} = 10^{-6}$ ;  $K_{a_3} = 10^{-10}$ )  
(S) At 1<sup>st</sup> half equivalence point of  $\text{H}_2\text{CO}_3$  when titrated against 0.1M NaOH,  $K_{a_1} = 10^{-5}$ ,  $K_{a_2} = 10^{-9}$   
(T)  $\text{Mg}(\text{OH})_2$ ;  $K_{sp} = 5 \times 10^{-16}$

IE0228

## TABLE TYPE QUESTION :

Column-I (solution)	Column-II (pH of solution)	Column-III (Introduction about solution)
(A) $\text{CH}_3\text{COOH}$ (0.2M, 1L) + NaOH (0.1M, 1 L )	(P) 1.3	(1) pH is determined by strong acid
(B) $\text{CH}_3\text{COOH}$ (0.1M ,1 L ) + HCl (0.1 M, 1 L )	(Q) 7	(2) Buffer solution at its maximum buffer capacity
(C) $\text{CH}_3\text{COOH}$ (0.1M, 1 L ) + $\text{NH}_4\text{OH}$ (0.1M, 1 L )	(R) 9	(3) pH is determined by salt hydrolysis.
(D) $\text{NH}_4\text{Cl}$ (200 ml, 0.1M) + NaOH (100 ml, 0.1M)	(S) 5	(4) pH is determined by buffer solution

(Given :  $(K_a)_{\text{CH}_3\text{COOH}} = 10^{-5}$  ,  $(K_b)_{\text{NH}_4\text{OH}} = 10^{-5}$ )

Q.25 Which of the following is incorrectly matched

- (A) A - S - 4                      (B) B - P - 1                      (C) D - R - 2                      (D) C - Q - 1

**IE0168**

Q.26 If 0.15 mole NaOH is added in solution (B) of column-I then which of the following is correct

- (A) S - 3                      (B) S - 4                      (C) R - 1                      (D) P - 3

**IE0169**

Q.27 If 0.1mole HCl is added in solution (A) of column-I then pH of the resulting solution will be

- (A) 7                      (B) 13                      (C) 3.0                      (D) 1

**IE0170**

## EXERCISE # J-MAIN

- Q.1 The solubility in water of a sparingly soluble salt  $AB_2$  is  $1.0 \times 10^{-5} \text{ mol L}^{-1}$ . Its solubility product will be [AIEEE-2003]  
 (1)  $1 \times 10^{-15}$  (2)  $1 \times 10^{-10}$  (3)  $4 \times 10^{-15}$  (4)  $4 \times 10^{-10}$  **IE0229**
- Q.2 The solubility of  $Mg(OH)_2$  is  $x$  mole/lit. then its solubility product is- [AIEEE-2002]  
 (1)  $x^3$  (2)  $5x^3$  (3)  $4x^3$  (4)  $2x^2$  **IE0230**
- Q.3 The molar solubility in  $\text{mol L}^{-1}$  of a sparingly soluble salt  $MX_4$  is 's'. The corresponding solubility product is  $K_{sp}$ . 's' is given in terms of  $K_{sp}$  by relation : [AIEEE-2004]  
 (1)  $s = (K_{sp} / 128)^{1/4}$  (2)  $s = (128K_{sp})^{1/4}$   
 (3)  $s = (256K_{sp})^{1/5}$  (4)  $s = (K_{sp}/256)^{1/5}$  **IE0231**
- Q.4 The solubility product of a salt having general formula  $MX_2$ , in water is :  $4 \times 10^{-12}$ . The concentration of  $M^{2+}$  ions in the aqueous solution of the salt is - [AIEEE-2005]  
 (1)  $1.0 \times 10^{-4} \text{ M}$  (2)  $2.0 \times 10^{-6} \text{ M}$  (3)  $4.0 \times 10^{-10} \text{ M}$  (4)  $1.6 \times 10^{-4} \text{ M}$  **IE0232**
- Q.5 Hydrogen ion concentration in mol/L in a solution of  $\text{pH} = 5.4$  will be - [AIEEE-2005]  
 (1)  $3.88 \times 10^6$  (2)  $3.98 \times 10^8$  (3)  $3.98 \times 10^{-6}$  (4)  $3.68 \times 10^{-6}$  **IE0233**
- Q.6 In a saturated solution of the sparingly soluble strong electrolyte  $AgIO_3$  (molecular mass = 283) the equilibrium which sets in is - [AIEEE-2007]  

$$AgIO_3 \rightleftharpoons Ag^+_{(aq)} + IO^-_{3(aq)}$$
 If the solubility product constant  $K_{sp}$  of  $AgIO_3$  at a given temperature is  $1.0 \times 10^{-8}$ , what is the mass of  $AgIO_3$  contained in 100 ml of its saturated solution ?  
 (1)  $28.3 \times 10^{-2} \text{ g}$  (2)  $2.83 \times 10^{-3} \text{ g}$  (3)  $1.0 \times 10^{-7} \text{ g}$  (4)  $1.0 \times 10^{-4} \text{ g}$  **IE0234**
- Q.7 The  $\text{pK}_a$  of a weak acid, HA, is 4.80. The  $\text{pK}_b$  of a weak base, BOH, is 4.78. The pH of an aqueous solution of the corresponding salt, BA, will be - [AIEEE-2008]  
 (1) 9.58 (2) 4.79 (3) 7.01 (4) 9.22 **IE0235**
- Q.8 Solid  $Ba(NO_3)_2$  is gradually dissolved in a  $1.0 \times 10^{-4} \text{ M}$   $Na_2CO_3$  solution. At what concentration of  $Ba^{2+}$  will a precipitate begin to form? ( $K_{sp}$  for  $BaCO_3 = 5.1 \times 10^{-9}$ ) [AIEEE-2009]  
 (1)  $8.1 \times 10^{-8} \text{ M}$  (2)  $8.1 \times 10^{-7} \text{ M}$  (3)  $4.1 \times 10^{-5} \text{ M}$  (4)  $5.1 \times 10^{-5} \text{ M}$  **IE0236**
- Q.9 Solubility product of silver bromide is  $5.0 \times 10^{-13}$ . The quantity of potassium bromide (molar mass taken as  $120 \text{ g mol}^{-1}$ ) to be added to 1 litre of 0.05 M solution of silver nitrate to start the precipitation of AgBr is :- [AIEEE-2010]  
 (1)  $5.0 \times 10^{-8} \text{ g}$  (2)  $1.2 \times 10^{-10} \text{ g}$  (3)  $1.2 \times 10^{-9} \text{ g}$  (4)  $6.2 \times 10^{-5} \text{ g}$  **IE0237**

Q.10 In aqueous solution the ionization constants for carbonic acid are

$$K_1 = 4.2 \times 10^{-7} \text{ and } K_2 = 4.8 \times 10^{-11}$$

[AIEEE-2010]

Select the correct statement for a saturated 0.034 M solution of the carbonic acid :-

- (1) The concentration of  $H^+$  is double that of  $CO_3^{2-}$
- (2) The concentration of  $CO_3^{2-}$  is 0.034 M
- (3) The concentration of  $CO_3^{2-}$  is greater than that of  $HCO_3^-$
- (4) The concentrations of  $H^+$  and  $HCO_3^-$  are approximately equal

IE0238

Q.11 At 25°C, the solubility product of  $Mg(OH)_2$  is  $1.0 \times 10^{-11}$ . At which pH, will  $Mg^{2+}$  ions start precipitating in the form of  $Mg(OH)_2$  from a solution of 0.001 M  $Mg^{2+}$  ions ?

[AIEEE-2010]

- (1) 8
- (2) 9
- (3) 10
- (4) 11

IE0239


Q.12 The  $K_{sp}$  for  $Cr(OH)_3$  is  $1.6 \times 10^{-30}$ . The molar solubility of this compound in water is :-

[AIEEE-2011]

- (1)  $\sqrt[3]{1.6 \times 10^{-30}}$
- (2)  $\sqrt[4]{1.6 \times 10^{-30}}$
- (3)  $\sqrt[4]{1.6 \times 10^{-30} / 27}$
- (4)  $1.6 \times 10^{-30} / 27$

IE0171

Q.13 An acid HA ionises as



The pH of 1.0 M solution is 5. Its dissociation constant would be :-

[AIEEE-2011]

- (1)  $1 \times 10^{-10}$
- (2) 5
- (3)  $5 \times 10^{-8}$
- (4)  $1 \times 10^{-5}$

IE0172

Q.14 If  $K_{sp}$  of  $CaF_2$  at 25°C is  $1.7 \times 10^{-10}$ , the combination amongst the following which gives a precipitate of  $CaF_2$  is :-

[JEE-MAIN(online)-2012]

- (1)  $1 \times 10^{-2}$  M  $Ca^{2+}$  and  $1 \times 10^{-5}$  M  $F^-$
- (2)  $1 \times 10^{-4}$  M  $Ca^{2+}$  and  $1 \times 10^{-4}$  M  $F^-$
- (3)  $1 \times 10^{-3}$  M  $Ca^{2+}$  and  $1 \times 10^{-5}$  M  $F^-$
- (4)  $1 \times 10^{-2}$  M  $Ca^{2+}$  and  $1 \times 10^{-3}$  M  $F^-$

IE0173

Q.15 The pH of a 0.1 molar solution of the acid HQ is 3. The value of the ionization constant,  $K_a$  of this acid is :-

[AIEEE-2012]

- (1)  $1 \times 10^{-7}$
- (2)  $3 \times 10^{-7}$
- (3)  $1 \times 10^{-3}$
- (4)  $1 \times 10^{-5}$

IE0174

Q.16 How many litres of water must be added to 1 litre of an aqueous solution of HCl with a pH of 1 to create an aqueous solution with pH of 2 ?

[AIEEE-2013]

- (1) 0.1 L
- (2) 0.9 L
- (3) 2.0 L
- (4) 9.0 L

IE0175

Q.17 Solid  $Ba(NO_3)_2$  is gradually dissolved in a  $1.0 \times 10^{-4}$  M  $Na_2CO_3$  solution. At which concentration of  $Ba^{2+}$ , precipitate of  $BaCO_3$  begins to form ? ( $K_{sp}$  for  $BaCO_3 = 5.1 \times 10^{-9}$ )

- (1)  $5.1 \times 10^{-5}$  M
- (2)  $8.1 \times 10^{-7}$  M
- (3)  $4.1 \times 10^{-5}$  M
- (4)  $7.1 \times 10^{-8}$  M

[JEE-MAIN(Online)-2013]

IE0176

Q.18 NaOH is a strong base. What will be pH of  $5.0 \times 10^{-2}$  M NaOH solution ? ( $\log 2 = 0.3$ )

[JEE-MAIN(Online)-2013]

- (1) 13.70                      (2) 13.00                      (3) 14.00                      (4) 12.70

IE0177

Q.19 Which one of the following arrangements represents the correct order of solubilities of sparingly soluble salts  $\text{Hg}_2\text{Cl}_2$ ,  $\text{Cr}_2(\text{SO}_4)_3$ ,  $\text{BaSO}_4$  and  $\text{CrCl}_3$  respectively ?

[JEE-MAIN(Online)-2013]

- (1)  $\left(\frac{K_{sp}}{4}\right)^{\frac{1}{3}}$ ,  $\left(\frac{K_{sp}}{108}\right)^{\frac{1}{5}}$ ,  $(K_{sp})^{\frac{1}{2}}$ ,  $\left(\frac{K_{sp}}{27}\right)^{\frac{1}{4}}$       (2)  $(K_{sp})^{\frac{1}{2}}$ ,  $\left(\frac{K_{sp}}{4}\right)^{\frac{1}{3}}$ ,  $\left(\frac{K_{sp}}{27}\right)^{\frac{1}{4}}$ ,  $\left(\frac{K_{sp}}{108}\right)^{\frac{1}{5}}$   
 (3)  $(K_{sp})^{\frac{1}{2}}$ ,  $\left(\frac{K_{sp}}{108}\right)^{\frac{1}{5}}$ ,  $\left(\frac{K_{sp}}{27}\right)^{\frac{1}{4}}$ ,  $\left(\frac{K_{sp}}{4}\right)^{\frac{1}{3}}$       (4)  $\left(\frac{K_{sp}}{108}\right)^{\frac{1}{5}}$ ,  $\left(\frac{K_{sp}}{27}\right)^{\frac{1}{4}}$ ,  $(K_{sp})^{\frac{1}{2}}$ ,  $\left(\frac{K_{sp}}{4}\right)^{\frac{1}{3}}$

IE0178

Q.20 What would be the pH of a solution obtained by mixing 5 g of acetic acid and 7.5 g of sodium acetate and making the volume equal to 500 mL?

[JEE-MAIN(Online)-2013]

( $K_a = 1.75 \times 10^{-5}$ ,  $pK_a = 4.76$ )

- (1)  $4.76 < \text{pH} < 5.0$     (2)  $\text{pH} < 4.70$   
 (3) pH of solution will be equal to pH of acetic acid      (4)  $\text{pH} = 4.70$

IE0179

Q.21 In some solutions, the concentration of  $\text{H}_3\text{O}^+$  remains constant even when small amounts of strong acid or strong base are added to them. These solutions are known as :-

[JEE-MAIN(Online)-2014]

- (1) Colloidal solutions    (2) True solutions  
 (3) Ideal solutions    (4) Buffer solutions

IE0180

Q.22 Zirconium phosphate  $[\text{Zr}_3(\text{PO}_4)_4]$  dissociates into three zirconium cations of charge +4 and four phosphate anions of charge -3. If molar solubility of zirconium phosphate is denoted by S and its solubility product by  $K_{sp}$  then which of the following relationship between S and  $K_{sp}$  is correct ?

- (1)  $S = \{K_{sp}/144\}^{1/7}$     (2)  $S = \{K_{sp}/(6912)^{1/7}\}$  [JEE-MAIN(Online)-2014]  
 (3)  $S = (K_{sp}/6912)^{1/7}$     (4)  $S = \{K_{sp}/6912\}^7$

IE0181

Q.23  $pK_a$  of a weak acid (HA) and  $pK_b$  of a weak base (BOH) are 3.2 and 3.4, respectively. The pH of their salt (AB) solution is

[JEE-MAIN(Offline)-2017]

- (1) 7.2                      (2) 6.9                      (3) 7.0                      (4) 1.0

IE0182

Q.24 Addition of sodium hydroxide solution to a weak acid (HA) results in a buffer of pH 6. If ionisation constant of HA is  $10^{-5}$ , the ratio of salt to acid concentration in the buffer solution will be :

[JEE-MAIN(Online)-2017]

- (1) 4 : 5                      (2) 1 : 10                      (3) 10 : 1                      (4) 5 : 4

IE0183



Q.25 50 mL of 0.2 M ammonia solution is treated with 25 mL of 0.2 M HCl. If  $pK_b$  of ammonia solution is 4.75, the pH of the mixture will be:- [JEE-MAIN(Online)-2017]

- (1) 8.25 (2) 4.75 (3) 9.25 (4) 3.75

IE0184

Q.26 Which of the following salts is the most basic in aqueous solution ? [JEE-MAIN(Offline)-2018]

- (1)  $\text{CH}_3\text{COOK}$  (2)  $\text{FeCl}_3$  (3)  $\text{Pb}(\text{CH}_3\text{COO})_2$  (4)  $\text{Al}(\text{CN})_3$

IE0185

Q.27 An alkali is titrated against an acid with methyl orange as indicator, which of the following is a correct combination ?

Base	Acid	End point	[JEE-MAIN(Offline)-2018]
(1) Strong	Strong	Pinkish red to yellow	
(2) Weak	Strong	Yellow to pinkish red	
(3) Strong	Strong	Pink to colourless	
(4) Weak	Strong	Colourless to pink	

IE0186

Q.28 An aqueous solution contains 0.10 M  $\text{H}_2\text{S}$  and 0.20 M HCl. If the equilibrium constants for the formation of  $\text{HS}^-$  from  $\text{H}_2\text{S}$  is  $1.0 \times 10^{-7}$  and that of  $\text{S}^{2-}$  from  $\text{HS}^-$  ions is  $1.2 \times 10^{-13}$  then the concentration of  $\text{S}^{2-}$  ions in aqueous solution is : [JEE-MAIN(Offline)-2018]

- (1)  $3 \times 10^{-20}$  (2)  $6 \times 10^{-21}$  (3)  $5 \times 10^{-19}$  (4)  $5 \times 10^{-8}$

IE0187

Q.29 A aqueous solution contains an unknown concentration of  $\text{Ba}^{2+}$ . When 50 mL of a 1 M solution of  $\text{Na}_2\text{SO}_4$  is added,  $\text{BaSO}_4$  just begins to precipitate. The final volume is 500 mL. The solubility product of  $\text{BaSO}_4$  is  $1 \times 10^{-10}$ . What is the original concentration of  $\text{Ba}^{2+}$  ? [JEE-MAIN(Offline)-2018]

- (1)  $2 \times 10^{-9} \text{ M}$  (2)  $1.1 \times 10^{-9} \text{ M}$  (3)  $1.0 \times 10^{-10} \text{ M}$  (4)  $5 \times 10^{-9} \text{ M}$

IE0188

Q.30 Following four solutions are prepared by mixing different volumes of NaOH and HCl of different concentrations, pH of which one of them will be equal to 1 ? [JEE-MAIN(Online)-2018]

- (1)  $75\text{mL} \frac{\text{M}}{5} \text{HCl} + 25\text{mL} \frac{\text{M}}{5} \text{NaOH}$  (2)  $100\text{mL} \frac{\text{M}}{10} \text{HCl} + 100\text{mL} \frac{\text{M}}{10} \text{NaOH}$   
 (3)  $55\text{mL} \frac{\text{M}}{10} \text{HCl} + 45\text{mL} \frac{\text{M}}{10} \text{NaOH}$  (4)  $60\text{mL} \frac{\text{M}}{10} \text{HCl} + 40\text{mL} \frac{\text{M}}{10} \text{NaOH}$

IE0189

Q.31 The minimum volume of water required to dissolve 0.1 g lead (II) chloride to get a saturated solution ( $K_{sp}$  of  $\text{PbCl}_2 = 3.2 \times 10^{-8}$ ; atomic mass of Pb = 207 u) is : [JEE-MAIN(Online)-2018]

- (1) 0.36 L (2) 0.18 L (3) 17.98 L (4) 1.798 L

IE0190

Q.32 If  $K_{sp}$  of  $Ag_2CO_3$  is  $8 \times 10^{-12}$ , the molar solubility of  $Ag_2CO_3$  in 0.1 M  $AgNO_3$  is :

- (1)  $8 \times 10^{-12}$  M (2)  $8 \times 10^{-10}$  M [JEE-MAIN(Online)-2019]  
 (3)  $8 \times 10^{-11}$  M (4)  $8 \times 10^{-13}$  M

IE0240

Q.33 25 ml of the given HCl solution requires 30 mL of 0.1 M sodium carbonate solution. What is the volume of this HCl solution required to titrate 30 mL of 0.2 M aqueous NaOH solution?

- (1) 25 mL (2) 50 mL [JEE-MAIN(Online)-2019]  
 (3) 12.5 mL (4) 75 mL

IE0241

Q.34 A mixture of 100 mmol of  $Ca(OH)_2$  and 2g of sodium sulphate was dissolved in water and the volume was made up to 100 mL. The mass of calcium sulphate formed and the concentration of  $OH^-$  in resulting solution, respectively, are : (Molar mass of  $Ca(OH)_2$ ,  $Na_2SO_4$  and  $CaSO_4$  are 74, 143 and  $136 \text{ g mol}^{-1}$ , respectively;  $K_{sp}$  of  $Ca(OH)_2$  is  $5.5 \times 10^{-6}$ )

- (1) 1.9 g,  $0.14 \text{ mol L}^{-1}$  (2) 13.6 g,  $0.14 \text{ mol L}^{-1}$  [JEE-MAIN(Online)-2019]  
 (3) 1.9 g,  $0.28 \text{ mol L}^{-1}$  (4) 13.6 g,  $0.28 \text{ mol L}^{-1}$

IE0242

Q.35 The pH of rain water, is approximately :

[JEE-MAIN(Online)-2019]

- (1) 6.5 (2) 7.5 (3) 5.6 (4) 7.0

IE0243

Q.36 20 mL of 0.1 M  $H_2SO_4$  solution is added to 30 mL of 0.2 M  $NH_4OH$  solution. The pH of the resultant mixture is : [ $pK_b$  of  $NH_4OH = 4.7$ ].

[JEE-MAIN(Online)-2019]

- (1) 9.4 (2) 5.0 (3) 9.0 (4) 5.2

IE0244

Q.37 If solubility product of  $Zr_3(PO_4)_4$  is denoted by  $K_{sp}$  and its molar solubility is denoted by  $S$ , then which of the following relation between  $S$  and  $K_{sp}$  is correct


[JEE-MAIN(Online)-2019]

- (1)  $S = \left( \frac{K_{sp}}{929} \right)^{1/9}$  (2)  $S = \left( \frac{K_{sp}}{216} \right)^{1/7}$  (3)  $S = \left( \frac{K_{sp}}{144} \right)^{1/6}$  (4)  $S = \left( \frac{K_{sp}}{6912} \right)^{1/7}$

IE0245

Q.38 In an acid-base titration, 0.1 M HCl solution was added to the NaOH solution of unknown strength. Which of the following correctly shows the change of pH of the titration mixture in this experiment?

[JEE-MAIN(Online)-2019]



- (1) (A) (2) (C) (3) (D) (4) (B)

IE0246

Q.39 The pH of a 0.02M  $\text{NH}_4\text{Cl}$  solution will be

[JEE-MAIN(Online)–2019]

[given  $K_b(\text{NH}_4\text{OH})=10^{-5}$  and  $\log 2=0.301$ ]

- (1) 4.65 (2) 5.35 (3) 4.35 (4) 2.65

IE0247

Q.40 The molar solubility of  $\text{Cd}(\text{OH})_2$  is  $1.84 \times 10^{-5}$  M in water. The expected solubility of  $\text{Cd}(\text{OH})_2$  in a buffer solution of pH = 12 is :

[JEE-MAIN(Online)–2019]

- (1)  $6.23 \times 10^{-11}$  M (2)  $1.84 \times 10^{-9}$  M (3)  $\frac{2.49}{1.84} \times 10^{-9}$  M (4)  $2.49 \times 10^{-10}$  M

IE0248

Q.41 Two solutions A and B, each of 100 L was made by dissolving 4g of NaOH and 9.8 g of  $\text{H}_2\text{SO}_4$  in water, respectively. The pH of the resultant solution obtained from mixing 40 L of solution A and 10 L of solution B is \_\_\_\_\_.

[JEE-MAIN(Online)–2020]

IE0249

Q.42 3g of acetic acid is added to 250 mL of 0.1 M HCl and the solution made up to 500 mL. To 20 mL of this solution  $\frac{1}{2}$  mL of 5 M NaOH is added. The pH of the solution is

[JEE-MAIN(Online)–2020]


[Given :  $\text{pK}_a$  of acetic acid = 4.75, molar mass of acetic acid = 60 g/mol,  $\log 3 = 0.4771$ ]

Neglect any changes in volume

IE0250

Q.43 The stoichiometry and solubility product of a salt with the solubility curve given below is, respectively :

[JEE-MAIN(Online)–2020]



- (1)  $\text{X}_2\text{Y}$ ,  $2 \times 10^{-9} \text{M}^3$  (2)  $\text{XY}_2$ ,  $1 \times 10^{-9} \text{M}^3$   
 (3)  $\text{XY}_2$ ,  $4 \times 10^{-9} \text{M}^3$  (4)  $\text{XY}$ ,  $2 \times 10^{-6} \text{M}^3$

IE0251

Q.44 For the following Assertion and Reason, the correct option is :

[JEE-MAIN(Online)–2020]

Assertion : The pH of water increases with increase in temperature.


Reason : The dissociation of water into  $\text{H}^+$  and  $\text{OH}^-$  is an exothermic reaction.

- (1) Both assertion and reason are true, but the reason is not the correct explanation for the assertion.  
 (2) Both assertion and reason are false.  
 (3) Assertion is not true, but reason is true.  
 (4) Both assertion and reason are true, and the reason is the correct explanation for the assertion.

IE0252

Q.45 The  $K_{sp}$  for the following dissociation is  $1.6 \times 10^{-5}$

[JEE-MAIN(Online)–2020]



Which of the following choices is correct for a mixture of 300 mL 0.134 M  $\text{Pb}(\text{NO}_3)_2$  and 100 mL 0.4 M NaCl ?

- (1)  $Q < K_{sp}$                       (2)  $Q > K_{sp}$                       (3)  $Q = K_{sp}$                       (4) Not enough data provided

IE0253

Q.46 The solubility product of  $\text{Cr}(\text{OH})_3$  at 298 K is  $6.0 \times 10^{-31}$ . The concentration of hydroxide ions in a saturated solution of  $\text{Cr}(\text{OH})_3$  will be :

- (1)  $(18 \times 10^{-31})^{1/4}$                       (2)  $(2.22 \times 10^{-31})^{1/4}$   
(3)  $(4.86 \times 10^{-29})^{1/4}$                       (4)  $(18 \times 10^{-31})^{1/2}$

[JEE-MAIN(Online)–2020]

IE0254

**EXERCISE # J-ADVANCE**

- Q.1 What will be the resultant pH when 200 ml of an aqueous solution of HCl (pH = 2.0) is mixed with 300 ml of an aqueous solution of NaOH (pH = 12.0) ? **[JEE '1998]**  
**IE0255**
- Q.2 The pH of 0.1 M solution of the following salts increases in the order **[JEE 1999]**  
(A)  $\text{NaCl} < \text{NH}_4\text{Cl} < \text{NaCN} < \text{HCl}$  (B)  $\text{HCl} < \text{NH}_4\text{Cl} < \text{NaCl} < \text{NaCN}$   
(C)  $\text{NaCN} < \text{NH}_4\text{Cl} < \text{NaCl} < \text{HCl}$  (D)  $\text{HCl} < \text{NaCl} < \text{NaCN} < \text{NH}_4\text{Cl}$   
**IE0256**
- Q.3 A buffer solution can be prepared from a mixture of **[JEE 1999]**  
(A) sodium acetate and acetic acid in water  
(B) sodium acetate and hydrochloric acid in water  
(C) ammonia and ammonium chloride in water  
(D) ammonia and sodium hydroxide in water.  
**IE0257**
- Q.4 The solubility of  $\text{Pb}(\text{OH})_2$  in water is  $6.7 \times 10^{-6}\text{M}$ . Calculate the solubility of  $\text{Pb}(\text{OH})_2$  in a buffer solution of pH = 8. **[JEE '1999]**  
**IE0258**
- Q.5 The average concentration of  $\text{SO}_2$  in the atmosphere over a city on a certain day is 10 ppm, when the average temperature is 298 K. Given that the solubility of  $\text{SO}_2$  in water at 298 K is 1.3653 moles  $\text{litre}^{-1}$  and the  $\text{pK}_a$  of  $\text{H}_2\text{SO}_3$  is 1.92, estimate the pH of rain on that day. **[JEE 2000]**  
[Given :  $10^{-1.92} = 1.2 \times 10^{-2}$ ,  $\sqrt{5.5678} = 2.5627$ ,  $\log(1.2213) = 0.0868$ ]  
**IE0259**
- Q.6 For sparingly soluble salt  $\text{ApBq}$ , the relationship of its solubility product ( $L_s$ ) with its solubility (S) is - **[JEE 2001]**  
(A)  $L_s = S^{p+q}$ .  $p^p$ .  $q^q$  (B)  $L_s = S^{p+q}$ .  $p^p$ .  $q^p$  (C)  $L_s = S^{pq}$ .  $p^p$ .  $q^q$  (D)  $L_s = S^{pq}$ .  $(p \cdot q)^{p+q}$   
**IE0260**
- Q.7 500 ml of 0.2 M aqueous solution of acetic acid is mixed with 500 mL of 0.2 M HCl at 25°C.  
(a) Calculate the degree of dissociation of acetic acid in the resulting solution and pH of the solution.  
(b) If 6 g of NaOH is added to the above solution, determine final pH. Assume there is no change in volume on mixing.  $K_a$  of acetic acid is  $1.75 \times 10^{-5}$  M. **[JEE 2002]**  
**IE0261**
- Q.8 A solution which is  $10^{-3}$  M each in  $\text{Mn}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Zn}^{2+}$  and  $\text{Hg}^{2+}$  is treated with  $10^{-16}\text{M}$  sulphide ion. If  $K_{sp}$ ,  $\text{MnS}$ ,  $\text{FeS}$ ,  $\text{ZnS}$  and  $\text{HgS}$  are  $10^{-15}$ ,  $10^{-23}$ ,  $10^{-20}$  and  $10^{-54}$  respectively, which one will precipitate first ? **[JEE 2003]**  
(A) FeS (B) MnS (C) HgS (D) ZnS  
**IE0262**
- Q.9 Will the pH of water be same at 4°C and 25°C ? Explain. **[JEE 2003]**  
**IE0263**

- Q.10 0.1 M of HA is titrated with 0.1 M NaOH, calculate the pH at end point. Given  $K_a(\text{HA}) = 5 \times 10^{-6}$  and  $\alpha \ll 1$ . [JEE 2004] IE0264
- Q.11 HX is a weak acid ( $K_a = 10^{-5}$ ). It forms a salt NaX (0.1 M) on reacting with caustic soda. The degree of hydrolysis of NaX is  
(A) 0.01% (B) 0.0001% (C) 0.1% (D) 0.5% [JEE 2004] IE0265
- Q.12  $\text{CH}_3\text{NH}_2$  (0.1 mole,  $K_b = 5 \times 10^{-4}$ ) is added to 0.08 moles of HCl and the solution is diluted to one litre, resulting hydrogen ion concentration is  
(A)  $1.6 \times 10^{-11}$  (B)  $8 \times 10^{-11}$  (C)  $5 \times 10^{-5}$  (D)  $2 \times 10^{-2}$  [JEE 2005] IE0266
- Q.13 If  $\text{Ag}^+ + \text{NH}_3 \rightleftharpoons [\text{Ag}(\text{NH}_3)]^+$ ;  $K_1 = 1.6 \times 10^3$  and  $[\text{Ag}(\text{NH}_3)]^+ + \text{NH}_3 \rightleftharpoons [\text{Ag}(\text{NH}_3)_2]^+$ ;  $K_2 = 6.8 \times 10^3$ . The formation constant of  $[\text{Ag}(\text{NH}_3)_2]^+$  is :  
(A)  $6.08 \times 10^{-6}$  (B)  $6.8 \times 10^{-6}$  (C)  $1.6 \times 10^3$  (D)  $1.088 \times 10^7$  [JEE 2006] IE0267
- Q.14 The species present in solution when  $\text{CO}_2$  is dissolved in water :  
(A)  $\text{CO}_2$ ,  $\text{H}_2\text{CO}_3$ ,  $\text{HCO}_3^-$ ,  $\text{CO}_3^{2-}$  (B)  $\text{H}_2\text{CO}_3$ ,  $\text{CO}_3^{2-}$  [JEE 2006] IE0268  
(C)  $\text{CO}_3^{2-}$ ,  $\text{HCO}_3^-$  (D)  $\text{CO}_2$ ,  $\text{H}_2\text{CO}_3$
- Q.15 2.5 mL of  $\frac{2}{5}$  M weak monoacidic base ( $K_b = 1 \times 10^{-12}$  at  $25^\circ\text{C}$ ) is titrated with  $\frac{2}{15}$  M HCl in water at  $25^\circ\text{C}$ . The concentration of  $\text{H}^+$  at equivalence point is [JEE 2008] IE0269  
( $K_w = 1 \times 10^{-14}$  at  $25^\circ\text{C}$ )  
(A)  $3.7 \times 10^{-13}$  M (B)  $3.2 \times 10^{-7}$  M (C)  $3.2 \times 10^{-2}$  M (D)  $2.7 \times 10^{-2}$  M
- Q.16 Solubility product constants ( $K_{sp}$ ) of salts of types MX,  $\text{MX}_2$  and  $\text{M}_3\text{X}$  at temperature 'T' are  $4.0 \times 10^{-8}$ ,  $3.2 \times 10^{-14}$  and  $2.7 \times 10^{-15}$ , respectively. Solubilities ( $\text{mol dm}^{-3}$ ) of the salts at temperature 'T' are in the order :  
(A)  $\text{MX} > \text{MX}_2 > \text{M}_3\text{X}$  (B)  $\text{M}_3\text{X} > \text{MX}_2 > \text{MX}$  [JEE 2008] IE0270  
(C)  $\text{MX}_2 > \text{M}_3\text{X} > \text{MX}$  (D)  $\text{MX} > \text{M}_3\text{X} > \text{MX}_2$
- Q.17 The dissociation constant of a substituted benzoic acid at  $25^\circ\text{C}$  is  $1.0 \times 10^{-4}$ . The pH of a 0.01 M solution of its sodium salt is [JEE 2009] IE0271
- Q.18 Aqueous solutions of  $\text{HNO}_3$ , KOH,  $\text{CH}_3\text{COOH}$  and  $\text{CH}_3\text{COONa}$  of identical concentrations are provided. The pair(s) of solutions which form a buffer upon mixing is(are) –  
(A)  $\text{HNO}_3$  and  $\text{CH}_3\text{COOH}$  (B) KOH and  $\text{CH}_3\text{COONa}$   
(C)  $\text{HNO}_3$  and  $\text{CH}_3\text{COONa}$  (D)  $\text{CH}_3\text{COOH}$  and  $\text{CH}_3\text{COONa}$  IE0272

- Q.19 In 1 L saturated solution of AgCl [ $K_{sp}(\text{AgCl}) = 1.6 \times 10^{-10}$ ], 0.1 mol of CuCl [ $K_{sp}(\text{CuCl}) = 1.0 \times 10^{-6}$ ] is added. The resultant concentration of  $\text{Ag}^+$  in the solution is  $1.6 \times 10^{-x}$ . The value of 'x' is. [JEE -2011] **IE0196**
- Q.20 The initial rate of hydrolysis of methyl acetate (1M) by a weak acid (HA, 1M) is  $1/100^{\text{th}}$  of that of a strong acid (HX, 1M), at  $25^\circ\text{C}$ . The  $K_a$  of HA is [JEE 2013] **IE0197**
- (A)  $1 \times 10^{-4}$  (B)  $1 \times 10^{-5}$  (C)  $1 \times 10^{-6}$  (D)  $1 \times 10^{-3}$
- Q.21 The  $K_{sp}$  of  $\text{Ag}_2\text{CrO}_4$  is  $1.1 \times 10^{-12}$  at 298 K. The solubility (in mol/L) of  $\text{Ag}_2\text{CrO}_4$  in a 0.1 M  $\text{AgNO}_3$  solution is [JEE 2013] **IE0198**
- (A)  $1.1 \times 10^{-11}$  (B)  $1.1 \times 10^{-10}$  (C)  $1.1 \times 10^{-12}$  (D)  $1.1 \times 10^{-9}$

### Paragraph For Questions 22 and 23

When 100 mL of 1.0 M HCl was mixed with 100 mL of 1.0 M NaOH in an insulated beaker at constant pressure, a temperature increase of  $5.7^\circ\text{C}$  was measured for the beaker and its contents. (**Expt-1**). Because the enthalpy of neutralisation of a strong acid with a strong base is a constant ( $-57.0 \text{ kJ mol}^{-1}$ ), this experiment could be used to measure the calorimeter constant. In a second experiment (**Expt-2**), 100 mL of 2.0 M acetic acid ( $K_a = 2.0 \times 10^{-5}$ ) was mixed with 100 mL of 1.0M NaOH (under identical conditions to (**Expt-1**)) where a temperature rise of  $5.6^\circ\text{C}$  was measured.

(Consider heat capacity of all solutions as  $4.2 \text{ J g}^{-1}\text{K}^{-1}$  and density of all solutions as  $1.0 \text{ g mL}^{-1}$ )

- Q.22 Enthalpy of dissociation (in  $\text{kJ mol}^{-1}$ ) of acetic acid obtained from the **Expt-2** is [JEE 2015] **IE0199**
- (A) 1.0 (B) 10.0 (C) 24.5 (D) 51.4
- Q.23 The pH of the solution after **Expt-2** **IE0200**
- (A) 2.8 (B) 4.7 (C) 5.0 (D) 7.0
- Q.24 The solubility of a salt of weak acid (AB) at pH 3 is  $Y \times 10^{-3} \text{ mol L}^{-1}$ . The value of Y is \_\_\_\_\_.  
(Given that the value of solubility product of AB ( $K_{sp}$ ) =  $2 \times 10^{-10}$  and the value of ionization constant of HB ( $K_a$ ) =  $1 \times 10^{-8}$ ) [JEE 2018] **IE0201**

**ANSWER KEY****EXERCISE # S-I**

- Q.1 Ans.  $6.022 \times 10^7$       Q.2 Ans. (i) 6.53 ; (ii) (a) Basic , (b) Acidic
- Q.3 Ans. 6.8
- Q.4 Ans. (a) 1, (b) 2.87, (c) 11.13 (d) 6.97, (e) 7, (f) 6 , (g) 6.97, (h) 11.30 (i) 9 , (j) 3
- Q.5 Ans. (a)  $K_a = 10^{-8}$ , (b)  $K_b = 10^{-6}$       Q.6 Ans. 10
- Q.7 Ans. 173.2 : 1      Q.8 Ans. 0.009
- Q.9 Ans.  $2.32 \times 10^{-8}$  M      Q.10 Ans.  $1.1 \times 10^{-3}$  M
- Q.11 Ans.  $1.11 \times 10^{-4}$       Q.12 Ans. 4.7
- Q.13 Ans. 3.3      Q.14 Ans. (a) 0.522 , (b) 2.522
- Q.15. Ans. (1)
- Q.16 Ans.  $[H^+] = 1.65 \times 10^{-2}$  M,  $[CHCl_2COO^-] = 6.5 \times 10^{-3}$  M
- Q.17 Ans.  $[H^+] = 3 \times 10^{-3}$  M,  $[CH_3COO^-] = 1.2 \times 10^{-3}$  M,  $[C_7H_5O_2^-] = 1.8 \times 10^{-3}$  M
- Q.18 Ans. 0.027 M, 0.073 M, 0.027 M,  $10^{-5}$  M
- Q.19 Ans.  $[H^+] = [H_2PO_4^-] = 2.7 \times 10^{-3}$  M,  $[HPO_4^{2-}] = 10^{-8}$  M,  $[PO_4^{3-}] = 3.7 \times 10^{-19}$  M
- Q.20 Ans. 11.3      Q.21 Ans.  $10^{-5}$  M
- Q.22 Ans. pH = 4.5      Q.23 Ans.  $K_b = 6.25 \times 10^{-10}$
- Q.24 Ans. 0.56%, pH = 7      Q.25 Ans. 1.667%
- Q.26 Ans. 0.25 %      Q.27 Ans.  $10^{-6}$  ;  $10^{-8}$
- Q.28 Ans. 8.3      Q.29 Ans. 4.19
- Q.30 Ans. (a) 6, (b)  $1 \times 10^{-5}$       Q.31 Ans. 9.0
- Q.32 Ans. 9.56      Q.33 Ans. 5.04
- Q.34 Ans. 0.05 mol      Q.35 Ans.  $[OH^-] = 9.0 \times 10^{-6}$  M
- Q.36 Ans. (10.1)      Q.37 Ans. 4.74
- Q.38 Ans. 9.56      Q.39 Ans. 8.7782
- Q.40 Ans. (3.33)      Q.41 Ans.  $10^{-5}$  M
- Q.42 Ans. 8.7 ,  $[H_3O^+] = 2 \times 10^{-9}$  M      Q.43 Ans. 5 ,  $10^{-5}$  M
- Q.44 Ans. (i) 2.85, (ii) 4.0969, (iii) 4.5229, (iv) 4.699, (v) 5.301, (vi) 8.699
- Q.45 Ans.  $[HI_n] = 80$  %      Q.46 Ans. 85.71%
- Q.47 Ans. (b), (c)      Q.48 Ans.  $\Delta pH = 0.954$
- Q.49 Ans.  $QX_2$  is more soluble      Q.50 Ans.  $4 \times 10^{-8}$
- Q.51 Ans.  $5 \times 10^{-11}$       Q.52 Ans.  $3.2 \times 10^{-11}$
- Q.53 Ans.  $2.56 \times 10^{-16}$       Q.54 Ans.  $1.0 \times 10^{-5}$  mol/lit
- Q.55 Ans.  $5 \times 10^{-10}$  M
- Q.56 Ans.  $5 \times 10^{-7}$
- Q.57 Ans.  $4 \times 10^{-7}$  mol/L AgBr,  $1.6 \times 10^{-6}$  mol/L AgSCN



Q.58 Ans.  $[F^-] = 3 \times 10^{-3}M$ Q.59 Ans.  $8 \times 10^{-3} M$ Q.60 Ans. (i)  $2 \times 10^{-9}$  , (ii)  $8 \times 10^{-3}$ Q.61 Ans.  $S = 2 \times 10^{-4} M$ , pH = 8.0Q.62 Ans.  $4 \times 10^{-2}M$ 

Q.63 Ans. (a) no precipitation will occur, (b) a precipitate will form

Q.64 Ans. No.

Q.65 Ans. 0.284 gm

**EXERCISE # S-II**Q.1 Ans. 0.209 M, 0.191 M,  $9.13 \times 10^{-3} M$ , 0Q.2 Ans. pH = 11.48 ,  $[enH_2^{2+}] = 7.1 \times 10^{-8} M$ 

Q.3 Ans. 10.6

Q.4 Ans.  $[S^{2-}] = 2.5 \times 10^{-15}M$ Q.5 Ans. ( $2 \times 10^{-4}M$ )

Q.6 Ans. pH = 10.52

Q.7 Ans.  $[OH^-] = 3.73 \times 10^{-2}M$ ,  $[H_3PO_4] = 6 \times 10^{-18}M$ 

Q.8 Ans. 8.35, 9.60, 4.66

Q.9 Ans.(9.6)

Q.10 Ans.  $K_b \approx 1.73 \times 10^{-5}$ , 5.27Q.11 Ans.  $K_a \approx 1.73 \times 10^{-5}$ , 8.73Q.12 Ans. ( $10^{-5}M$ )Q.13. Ans. ( $10^{-3}M$ )

Q.14 Ans. pH = 7.9 , pH = 7.3

Q.15 Ans.  $2.8 \times 10^{-3}$  moleQ.16 Ans.  $1.6 \times 10^{-3}$ Q.17 Ans.  $K_d = 1/K_f = 4.8 \times 10^{-4}$ **EXERCISE # O-I**

Q.1 Ans.(A)

Q.2 Ans.(B)

Q.3 Ans.(C)

Q.4 Ans.(C)

Q.5 Ans.(B)

Q.6 Ans.(C)

Q.7 Ans.(D)

Q.8 Ans.(D)

Q.9 Ans.(C)

Q.10 Ans.(C)

Q.11. Ans (C)

Q.12 Ans.(D)

Q.13 Ans.(B)

Q.14 Ans.(B)

Q.15 Ans.(C)

Q.16 Ans.(C)

Q.17 Ans.(B)

Q.18. Ans.(C)

Q.19 Ans.(C)

Q.20 Ans.(B)

Q.21 Ans.(B)

Q.22 Ans.(D)

Q.23 Ans.(D)

Q.24 Ans.(C)

Q.25 Ans.(C)

Q.26 Ans.(C)

Q.27 Ans.(B)

Q.28 Ans.(B)

Q.29 Ans.(A)

Q.30 Ans.(D)

Q.31 Ans.(B)

Q.32 Ans.(C)

Q.33 Ans.(C)

Q.34 Ans.(C)

Q.35 Ans.(B)

Q.36 Ans.(C)

Q.37 Ans.(D)

Q.38 Ans.(A)

Q.39 Ans.(C)

Q.40 Ans.(C)

Q.41 Ans.(C)

Q.42 Ans.(B)

Q.43 Ans.(C)

Q.44 Ans.(D)

Q.45 Ans.(A)

Q.46 Ans.(B)

Q.47 Ans.(D)

Q.48 Ans.(B)

Q.49 Ans.(C)

Q.50 Ans.(A)

Q.51 Ans.(C)

Q.52 Ans.(A)

Q.53 Ans.(A)

Q.54. Ans.(A)

Q.55. Ans.(B)

Q.56 Ans.(D)

Q.57 Ans.(A)

Q.58 Ans.(A)

Q.61 Ans.(D)

Q.64 Ans.(C)

Q.67 Ans.(D)

Q.70.Sol.(C)

Q.73 Ans.(D)

Q.59 Ans.(B)

Q.62 Ans.(A)

Q.65 Ans.(D)

Q.68 Ans.(C)

Q.71. Ans.(B)

Q.74 Ans.(B)

Q.60 Ans.(C)

Q.63 Ans.(D)

Q.66 Ans.(D)

Q.69 Ans.(A)

Q.72. Ans.(B)

Q.75 Ans.(B)

**EXERCISE # O-II**

Q.1 Ans.(A)

Q.4 Ans.(B)

Q.7 Ans.(A)

Q.10 Ans.(A)

Q.13 Ans.(B,C)

Q.16 Ans.(A)

Q.19 Ans (B)

Q.22 Ans. (A)

Q.23 Ans. A - (R), B - (P), C - (Q), D - (S)

Q.24 Ans (A) - Q; (B) - P, S; (C) - R; (D) -T

Q.25 Ans.(D)

Q.2 Ans.(C)

Q.5 Ans (C)

Q.8 Ans.(B)

Q.11 Ans. (A,B,C)

Q.14 Ans. (A, B, D)

Q.17 Ans. (A)

Q.20 Ans (B)

Q.3 Ans.(D)

Q.6 Ans.(A)

Q.9 Ans.(D)

Q.12 Ans.(A,B,C,D)

Q.15 Ans. (D)

Q.18 Ans (B)

Q.21 Ans. (C)

**EXERCISE # J-MAIN**

Q.1 Ans.(3)

Q.4 Ans.(1)

Q.7 Ans.(3)

Q.10 Ans.(4)

Q.13 Ans.(1)

Q.16 Ans.(4)

Q.19 Ans.(1)

Q.22 Ans.(3)

Q.25 Ans.(3)

Q.28 Ans.(1)

Q.31 Ans.(2)

Q.34 Ans.(3)

Q.37 Ans.(4)

Q.40 Ans.(4)

Q.43 Ans.(3)

Q.46 Ans.(1)

Q.2 Ans.(3)

Q.5 Ans.(3)

Q.8 Ans.(4)

Q.11 Ans.(3)

Q.14 Ans.(4)

Q.17 Ans.(1)

Q.20 Ans.(1)

Q.23 Ans. (2)

Q.26 Ans.(1)

Q.29 Ans.(2)

Q.32 Ans.(2)

Q.35 Ans.(3)

Q.38 Ans.(1)

Q.41 Ans.(10.60)

Q.44 Ans(2)

Q.3 Ans.(4)

Q.6 Ans.(2)

Q.9 Ans.(3)

Q.12 Ans.(3)

Q.15 Ans.(4)

Q.18 Ans.(4)

Q.21 Ans.(4)

Q.24 Ans.(3)

Q.27 Ans.(2)

Q.30 Ans.(1)

Q.33 Ans.(1)

Q.36 Ans.(3)

Q.39 Ans.(2)

Q.42 Ans. (5.22 to 5.24)

Q.45 Ans.(2)

---

**EXERCISE # J-ADVANCED**

---

**Q.1 Ans. pH = 11.3010****Q.2 Ans.(B)****Q.3 Ans.(A, B, C)****Q.4 Ans.  $s = 1.203 \times 10^{-3} \text{M}$** **Q.5 Ans. pH = 0.91325****Q.6 Ans.(A)****Q.7 Ans.(a) 0.0175% , (b) 4.757****Q.8 Ans.(C)****Q.9 Ans.No, it will be > 7 at 0°C.****Q.10 Ans.pH = 9****Q.11 Ans.(A)****Q.12 Ans.(B)****Q.13 Ans.(D)****Q.14 Ans.(A)****Q.15 Ans.(D)****Q.16 Ans.(D)****Q.17 Ans.(8)****Q.18 Ans.(C), (D)****Q.19 Ans.(7)****Q.20 Ans.(A)****Q.21 Ans.(B)****Q.22 Ans.(A)****Q.23 Ans.(B)****Q.24 Ans. (4.47)**