

CBSE Board Class 10 Chapter 4- Quadratic Equations

Objective Questions

Introduction to Quadratic Equations

1. What is the degree of a quadratic equation?

- (A) 0
- (B) 2
- (C) 3
- (D) 1

Answer: (B) 2

Solution: The standard form of quadratic equation is $ax^2 + bx + c = 0$, $a \neq 0$. So the degree of a quadratic equation is 2

2. Find the sum of the roots of the equation $x^2 - 8x + 2 = 0$

- (A) 8
- (B) -8
- (C) 2
- (D) -6

Answer: (A) 8

Solution: For general quadratic equation $ax^2 + bx + c = 0$.

Sum of the roots $= -b/a$

For $x^2 - 8x + 2 = 0$

Sum of the roots $= -(-8/1) = 8$

Sum of the roots of the equation is 8

3. Which of the following is not quadratic equation?

- (A) $x(2x + 3) = x^2 + 1$
- (B) $x(x + 1) + 8 = (x + 2)(x - 2)$
- (C) $(x + 2)^3 = x^3 - 4$
- (D) $(x - 2)^2 + 1 = 2x - 3$

Answer: (B) $x(x + 1) + 8 = (x + 2)(x - 2)$

Solution: (a) $(x-2)^2 + 1 = (2x - 3)$

$$x^2 - 4x + 4 + 1 = 2x - 3$$

$$x^2 - 4x + 4 + 1 - 2x + 3 = 0$$

$$x^2 - 6x + 8 = 0$$

This is a quadratic equation.

(b) $(x+2)^3 = x^3 - 4$

$$x^3 + 6x^2 + 12x + 8 = x^3 - 4$$

$$6x^2 + 12x + 12 = 0$$

This is a quadratic equation.

(c) $x(2x + 3) = x^2 + 1$

$$2x^2 + 3x = x^2 + 1$$

$$x^2 + 3x - 1 = 0$$

This is a quadratic equation.

(d) $x(x + 1) + 8 = (x + 2)(x - 2)$

$$x^2 + x + 8 = x^2 - 4$$

$$x + 12 = 0$$

This is not a Quadratic equation.

4. If the sum of the roots of a quadratic equation is 5 and the product of the roots is also 5, then the equation is

(A) $x^2 + 10x + 5 = 0$

(B) $x^2 - 5x + 5 = 0$

(C) $x^2 + 5x - 5 = 0$

(D) $x^2 - 5x + 10 = 0$

Answer: (B) $x^2 - 5x + 5 = 0$

Solution: For a quadratic equation $ax^2+bx+c=0$,

sum of roots = $-b/a$

product of roots = c/a .

sum of roots = 5 = $-b/a$

product of roots = 5 = c/a ,

Thus, quadratic equation is $x^2-5x+5=0$

5. A rectangular field has an area of 3 sq. units. The length is one more than twice the breadth 'x'. Frame an equation to represent this.

(A) $x^2 - 2x + 6 = 0$

(B) $x^2 - 2x + 3 = 0$

(C) $2x^2 + x - 3 = 0$

(D) $2x^2 + x - 6 = 0$

Answer: (C) $2x^2 + x - 3 = 0$

Solution: Area of rectangle = length \times breadth

Given, length = $(2 \times \text{breadth} + 1)$

Let the breadth of the field be x.

Length of the field = $2x + 1$

Area of the rectangular field = $x(2x + 1) = 3$

$$2x^2 + x = 3$$

$$2x^2 + x - 3 = 0$$

Solving QE by factorisation

6. The roots of the quadratic equation $x^2+5x-14=0$ is

- (A) 2, 7
- (B) -2, 7
- (C) -2, -7
- (D) 2, -7

Answer: (D) 2, -7

Solutions: $x^2+5x-14=0$

We need to split the coefficient of x such that the sum of the factors is 5 and their product is -14.

So we will find the coefficient as 7 and -2.

The sum of 7 and -2 is 5 and product is -14.

So now re-write the equation

$$x^2+7x-2x-14=0$$

Taking common terms out

$$x(x+7)-2(x+7)=0$$

Again taking out the common terms

$$(x-2)(x+7)=0$$

Now equate the factors to zero to find the roots.

So the roots of the equation are 2,-7

7. Factorize $x^2 +5x+6 =0$

- (A) $(x-1)(x-3)$
- (B) $(x+1)(x+3)$
- (C) $(x-2)(x-3)$
- (D) $(x+2)(x+3)$

Answer: (D) $(x+2)(x+3)$

Solution: Comparing $x^2 + 5x + 6 = 0$ to $ax^2 + bx + c = 0$, we have $a=1$, $b=5$ and $c=6$

Now, we need to find two numbers whose product is 6 and whose sum is 5

Pairs of numbers whose product is 6

1, 6

-1, -6

2, 3

-2, -3

Of these pairs, the pair that gives the sum 5 is the third pair

Identifying the pair, we rewrite the given quadratic equation as

$$x^2 + 5x + 6 = x^2 + 2x + 3x + 6 = x(x+2) + 3(x+2)$$


$$= (x+2)(x+3)$$

8. The altitude of a right triangle is 7 cm less than its base. If the hypotenuse is 13 cm, find the other two sides (in cm).

- (A) 12, 5
- (B) 7, 2
- (C) 5, 3
- (D) 2, 5

Answer: (A) 12, 5

Solutions:



Let the base = x cm

Given that the altitude of a right triangle is 7 cm less than its base

Altitude is = $x - 7$ cm

Given that hypotenuse = 13cm

Applying Pythagoras theorem,

$$\text{base}^2 + \text{altitude}^2 = \text{hypotenuse}^2$$

Substituting the values, we get

$$\Rightarrow x^2 + (x-7)^2 = 13^2$$

$$\Rightarrow x^2 + x^2 + 49 - 14x = 169$$

$$\Rightarrow 2x^2 - 14x + 49 - 169 = 0$$

$$\Rightarrow 2x^2 - 14x - 120 = 0$$

Dividing with 2 on both sides the above equation simplifies to

$$\Rightarrow x^2 - 7x - 60 = 0$$

$$\Rightarrow x^2 - 12x + 5x - 60 = 0$$

$$\Rightarrow x(x-12) + 5(x-12) = 0$$

$$\Rightarrow (x-12)(x+5) = 0$$

$$\Rightarrow x - 12 = 0 \text{ or } x + 5 = 0$$

$$\Rightarrow x = 12 \text{ or } x = -5$$

Length cannot be negative so x cannot be equal to -5

base $x = 12\text{cm}$; altitude $= 12 - 7 = 5\text{cm}$

9. If a train travelled 5 km/hr faster, it would take one hour less to travel 210 km.
The speed of the train is :

- (A) 60 km/hr
- (B) 70 km/hr
- (C) 35 km/hr
- (D) 30 km/hr

Answer: (D) 30 km/hr

Solution: Let the speed of the train be x km/hr.
Distance travelled = 210 km

Time taken to travel 210 km = $210/x$ hours

When the speed is increased by 5 km/h, the new speed is $(x+5)$

Time taken to travel 210 km with the new speed is $210 / (x+5)$ hours

According to the question,

$$210/x - 210/(x+5) = 1$$

$$\Rightarrow 210(x+5) - 210x = x(x+5)$$

$$\Rightarrow 210x + 1050 - 210x = x^2 + 5x$$

$$\Rightarrow x^2 + 5x - 1050 = 0$$

$$\Rightarrow (x+35)(x-30) = 0$$

$$\Rightarrow x = -35, 30$$

The speed cannot be negative. Thus, the speed of the train is 30 km/hr

10. If the solutions of the equation $x^2 + 3x - 18 = 0$ are -6, 3 then the roots of the equation $2(x^2 + 3x - 18) = 0$ are

- (A) 3, 3
- (B) -6, 3
- (C) -12, 6
- (D) -6, 6

Answer: (B) -6, 3

Solution: The roots of a quadratic equation do not change when it is multiplied by a constant non-zero real number. So when the equation $x^2+3x-18=0$ is multiplied by 2, the roots still remain the same i.e. -6, 3.

Solving QE by completing square

11. The square of $(5x + 1)$ is equal to 16. What is x ?

- (A) $x = 4, \frac{1}{4}$
- (B) $x = -1, \frac{3}{5}$
- (C) $x = 1, \frac{3}{2}$
- (D) $x = -1, \frac{4}{5}$

Answer: (B) $x = -1, \frac{3}{5}$

Solution: Converting statement into an equation-

$$\Rightarrow (5x+1)^2 = 16 \text{ (Applying } (a+b)^2 \text{ formula)}$$

$$\Rightarrow 5x + 1 = \pm 4 \text{ (Taking square root on both sides)}$$

$$\Rightarrow 5x = -5, 3$$

$$\Rightarrow x = -1, \frac{3}{5}$$

12. Using the method of completion of squares find one of the roots of the equation $2x^2-7x+3=0$. Also, find the equation obtained after completion of the square.

- (A) 6, $(x-\frac{7}{4})^2-\frac{25}{16}=0$
- (B) 3, $(x-\frac{7}{4})^2-\frac{25}{16}=0$
- (C) 3, $(x-\frac{7}{2})^2-\frac{25}{16}=0$
- (D) 13, $(x-\frac{7}{2})^2-\frac{25}{16}=0$

Answer: (B) 3, $(x-\frac{7}{4})^2-\frac{25}{16}=0$

Solution: $2x^2-7x+3=0$

Dividing by the coefficient of x^2 , we get

$$x^2-\frac{7}{2}x+\frac{3}{2}=0; a=1, b=\frac{7}{2}, c=\frac{3}{2}$$

x)

Adding and subtracting the square of $b/2=7/4$, (half of coefficient of

$$\text{We get, } [x^2 - 2(7/4)x + (7/4)^2] - (7/4)^2 + 3/2 = 0$$

The equation after completing the square is:

$$(x - 7/4)^2 - 25/16 = 0$$

Taking square root, $(x - 7/4) = (\pm 5/4)$

Taking positive sign $5/4$, $x = 3$

Taking negative sign $-5/4$, $x = 1/2$

13. Find the roots of the equation $5x^2 - 6x - 2 = 0$ by the method of completing the square.

(A) $x = 3$

(B) $x = (5 \pm \sqrt{19})/3$

(C) $x = (3 \pm \sqrt{19})/5$

(D) $x = 5$

Answer: (C) $x = (3 \pm \sqrt{19})/5$

Solution: Multiplying the equation throughout by 5, we get $25x^2 - 30x - 10 = 0$

This is the same as:

$$(5x)^2 - [2 \times (5x) \times 3] + 3^2 - 3^2 - 10 = 0$$

$$\Rightarrow (5x - 3)^2 - 9 - 10 = 0$$

$$\Rightarrow (5x - 3)^2 - 19 = 0$$

$$\Rightarrow (5x - 3)^2 = 19$$

$$\Rightarrow 5x - 3 = \pm \sqrt{19}$$

$$\Rightarrow x = (3 \pm \sqrt{19})/5$$

14. There is a natural number x . Write down the expression for the product of x and its next natural number.

(A) $2x^2 + 1$

(B) $x^2 - x$

(C) $x^2 + x$

(D) $(x + 1)(x + 2)$

Answer: (C) $x^2 + x$

Solution: If a natural number is x , the next natural number is greater than x by 1 and hence $x+1$. For eg. For 3, next natural number is 4. The product of the 2 numbers is $x(x+1) = x^2 + x$

15. What number should be added to x^2+6x to make it a perfect square?

- (A) 36
- (B) 18
- (C) 9
- (D) 72

Answer: (C) 9

Solution: The identity $(a+b)^2 = (a^2+2ab+b^2)$ represents a perfect square.

If we observe carefully we can see that x^2+6x can be written in the form of $(a^2+2ab+b^2)$ by adding a constant.

$x^2+2(x)(3) + \text{constant}.$

To make x^2+6x a perfect square, divide the coefficient of x by 2 and then add the square of the result to make this a perfect square.

Hence, $6/2=3$ and $3^2 = 9$

We should add 9 to make x^2+6x a perfect square.

Solving QE using quadratic formula

16. The equation $x^2+4x+c=0$ has real roots, then

- (A) $C \geq 6$
- (B) $C \leq 8$
- (C) $C \leq 4$
- (D) $C \geq 4$

Answer: (C) $C \leq 4$

Solution: Step 1:- For, $x^2+4x+c=0$, value of discriminant $D=4^2-4c=16-4c$

Step 2:- The roots of quadratic equation are real only when $D \geq 0$

$$16-4c \geq 0$$

Step 3:- $c \leq 4$

17. Find the discriminant of the quadratic equation $3x^2 - 5x + 2 = 0$ and hence, find the nature of the roots.

- (A) -1 , no real roots
- (B) 1 , two equal roots
- (C) -1 , two distinct real roots
- (D) 1 , two distinct real roots

Answer: (D) 1 , two distinct real roots

Solution: $D = b^2 - 4ac = (-5)^2 - 4 \times 3 \times 2 = 1 > 0$

$D = 1 > 0 \Rightarrow$ Two distinct real roots.

18. Taylor purchased a rectangular plot of area 634 m^2 . The length of the plot is 2 m more than thrice its breadth. Find the length and breadth (approximate values).

- (A) 34.6 m & 11.20 m
- (B) 44.6 m & 14.20 m
- (C) 32 m & 16 m
- (D) 88 m & 24 m

Answer: (B) 44.6 m & 14.20 m

Solution: Let x and y be the length and breadth of the rectangle respectively.

Given, $x = 2 + 3y$

Area of the rectangle = length \times breadth

$= xy$

$\Rightarrow 634 = (2 + 3y)y$

$\Rightarrow 634 = 2y + 3y^2$

So, $3y^2 + 2y - 634 = 0$

The roots of the above quadratic equation will be

$$y = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$y = \frac{-2 \pm \sqrt{2^2 - 4(3)(-634)}}{2(3)}$$

⇒

$$y = \frac{-2 \pm \sqrt{4 + 7608}}{6}$$

⇒

$$y = \frac{-2 \pm \sqrt{7612}}{6}$$

⇒

$$y = \frac{-2 \pm 87.246}{6}$$

⇒

$$y = \frac{-2 + 87.246}{6}$$

⇒

OR

$$y = \frac{-2 - 87.246}{6}$$

⇒

$$\Rightarrow y = 14.20 \text{ or}$$

$$y = -14.87$$

Considering positive value for breadth, we have $y = 14.20$.

Using $x = 2 + 3y$, we have

$$x = 2 + 3(14.20) = 44.6$$

Now, we have

$x = 44.6$ and $y = 14.20$ (approximately).

19. If the equation $x^2 + 2(k+2)x + 9k = 0$ has equal roots, then values of k are _____.

- (A) 1,4
- (B) -1,5
- (C) -1,-4
- (D) 1,-5

Answer: (A) 1, 4

Solution: Step 1:- For, $x^2 + 2(k+2)x + 9k = 0$, value of discriminant $D = [2(k+2)]^2 - 4(9k)$
 $= 4(k^2 + 4 - 5k)$

Step 2:- The roots of quadratic equation are real and equal only when $D=0$

$$k^2 + 4 - 5k = 0$$

$$\Rightarrow k^2 - 5k + 4 = 0$$

$$\Rightarrow k^2 - k - 4k + 4 = 0$$

$$\Rightarrow k(k-1) - 4(k-1) = 0$$

$$\Rightarrow (k-1)(k-4) = 0$$

Step 3:- $k=4$ or 1

20. Find the roots of the $3x^2 - 5x + 2 = 0$ quadratic equation, using the quadratic formula.

- (A) $(7 \pm 1)/6$
- (B) $(4 \pm 1)/6$
- (C) $(5 \pm 2)/6$
- (D) $(5 \pm 1)/6$

Answer: (D) $(5 \pm 1)/6$

Solution: Quadratic equation of the form $ax^2 + bx + c = 0$

The roots of the above quadratic equation will

$$\text{be } \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$a=3, b=-5 \text{ and } c=2$$

$$x = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \times 3 \times 2}}{2 \times 3}$$

$$x = \frac{(5) \pm \sqrt{25 - 24}}{6} = \frac{5 \pm 1}{6}$$

$$x = \frac{5+1}{6}, \frac{5-1}{6} = 1, \frac{2}{3}$$