

- Q.1** If  $A = \{1, 2, 3\}$ ,  $B = \{4, 5, 6\}$ , which of the following are relations from A to B?  
Give reasons in support of your answer.
- (i)  $\{(1, 6), (3, 4), (5, 2)\}$
  - (ii)  $\{(1, 5), (2, 6), (3, 4), (3, 6)\}$
  - (iii)  $\{(4, 2), (4, 3), (5, 1)\}$
  - (iv)  $A \times B$
- Q.2** A relation R is defined from a set  $A = \{2, 3, 4, 5\}$  to a set  $B = \{3, 6, 7, 10\}$  as follows:  $(x, y) \in R$  if x is relatively prime to y. Express R as a set of ordered pairs and determine its domain and range.
- Q.3** Let A be the set of first five natural and let R be a relation on A defined as follows:  $(x, y) \in R$  if  $x \leq y$ . Express R and  $R^{-1}$  as sets of ordered pairs. Determine also
- (i) the domain of  $R^{-1}$
  - (ii) The Range of R.
- Q.4** Find the inverse relation  $R^{-1}$  in each of the following cases:
- (i)  $R = \{(1, 2), (1, 3), (2, 3), (3, 2), (5, 6)\}$
  - (ii)  $R = \{(x, y) : x, y \in \mathbb{N}; x + 2y = 8\}$
  - (iii) R is a relation from  $\{11, 12, 13\}$  to  $\{8, 10, 12\}$  defined by  $y = x - 3$
- Q.5** Write the following relations as the sets of ordered pairs:
- (i) A relation R from the set  $\{2, 3, 4, 5, 6\}$  to the set  $\{1, 2, 3\}$  defined by  $x = 2y$ .
  - (ii) A relation R on the set  $\{1, 2, 3, 4, 5, 6, 7\}$  defined by  $(x, y) \in R \Leftrightarrow x$  is relatively prime to y.
  - (iii) A relation R on the set  $\{0, 1, 2, \dots, 10\}$  defined by  $2x + 3y = 12$ .
  - (iv) A relation R from a set  $A = \{5, 6, 7, 8\}$  to the set  $B = \{10, 12, 15, 16, 18\}$  defined by  $(x, y) \in R$  if x divides y.
- Q.6** Let R be a relation in  $\mathbb{N}$  defined by  $(x, y) \in R \Leftrightarrow x + 2y = 8$ . Express R and  $R^{-1}$  as sets of ordered pairs.
- Q.7** Let  $A = \{3, 5\}$  and  $B = \{7, 11\}$ . Let  $R = \{(a, b) : a \in A, b \in B, a-b \text{ is odd}\}$ . Show that R is an empty relation from A into B.
- Q.8** Let  $A = \{1, 2\}$  and  $B = \{3, 4\}$ . Find the total number of relations from A into B.
- Q.9** Determine the domain and range of the relation R defined by
- (i)  $R = \{(x, x+5) : x \in \{0, 1, 2, 3, 4, 5\}\}$
  - (ii)  $R = \{(x, x^3) : x \text{ is a prime number less than } 10\}$
- Q.10** Determine the domain and range of the following relations:
- (i)  $R = \{(a, b) : a \in \mathbb{N}, a < 5, b = 4\}$
  - (ii)  $S = \{(a, b) : b = |a-1|, a \in \mathbb{Z} \text{ and } |a| \leq 3\}$
- Q.11** Let  $A = \{a, b\}$ . List all relations on A and find their number.



# SOLUTION

(MATH)

## RELATION AND FUNCTION

DPP – 03

CLASS – 12<sup>th</sup>

TOPIC – RELATION AND FUNCTION

**Sol.1.**  $A = \{1, 2, 3\}$ ,  $B = \{4, 5, 6\}$

A relation from A to B can be defined as:

$$A \times B = \{1, 2, 3\} \times \{4, 5, 6\}$$

$$= \{(1, 4), (1, 5), (1, 6), (2, 4), (2, 5), (2, 6), (3, 4), (3, 5), (3, 6)\}$$

(i)  $\{(1, 6), (3, 4), (5, 2)\}$

No, it is not a relation from A to B. The given set is not a subset of  $A \times B$  as  $(5, 2)$  is not a part of the relation from A to B.

(ii)  $\{(1, 5), (2, 6), (3, 4), (3, 6)\}$

Yes, it is a relation from A to B. The given set is a subset of  $A \times B$ .

(iii)  $\{(4, 2), (4, 3), (5, 1)\}$

No, it is not a relation from A to B. The given set is not a subset of  $A \times B$ .

(iv)  $A \times B$

$A \times B$  is a relation from A to B and can be defined as:

$$\{(1, 4), (1, 5), (1, 6), (2, 4), (2, 5), (2, 6), (3, 4), (3, 5), (3, 6)\}$$

**Sol.2.** Relatively prime numbers are also known as co-prime numbers. If there is no integer greater than one that divides both (that is, their greatest common divisor is one).

Given:  $(x, y) \in R = x$  is relatively prime to  $y$

Here,

2 is co-prime to 3 and 7.

3 is co-prime to 7 and 10.

4 is co-prime to 3 and 7.

5 is co-prime to 3, 6 and 7.

$$\therefore R = \{(2, 3), (2, 7), (3, 7), (3, 10), (4, 3), (4, 7), (5, 3), (5, 6), (5, 7)\}$$

Domain of relation  $R = \{2, 3, 4, 5\}$

Range of relation  $R = \{3, 6, 7, 10\}$

**Sol.3.** A is a set of the first five natural numbers.

So,  $A = \{1, 2, 3, 4, 5\}$

Given:  $(x, y) R x \leq y$

1 is less than 2, 3, 4 and 5.

2 is less than 3, 4 and 5.

3 is less than 4 and 5.

4 is less than 5.

5 is not less than any number A

$$\therefore R = \{(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (2, 2), (2, 3), (2, 4), (2, 5), (3, 3), (3, 4), (3, 5), (4, 4), (4, 5), (5, 5)\}$$

“An inverse relation is the set of ordered pairs obtained by interchanging the first and second elements of each pair in the original relation. If the graph of a function contains a point  $(a, b)$ , then the graph of the inverse relation of this function contains the point  $(b, a)$ ”.

$$\therefore R^{-1} = \{(1, 1), (2, 1), (3, 1), (4, 1), (5, 1), (2, 2), (3, 2), (4, 2), (5, 2), (3, 3), (4, 3), (5, 3), (4, 4), (5, 4), (5, 5)\}$$

(i) Domain of  $R^{-1} = \{1, 2, 3, 4, 5\}$

(ii) Range of  $R = \{1, 2, 3, 4, 5\}$

**Sol.4.** (i) Given:

$$R = \{(1, 2), (1, 3), (2, 3), (3, 2), (5, 6)\}$$

$$\text{So, } R^{-1} = \{(2, 1), (3, 1), (3, 2), (2, 3), (6, 5)\}$$

(ii) Given,



$$R = \{(x, y) : x, y \in \mathbb{N}; x + 2y = 8\}$$

$$\text{Here, } x + 2y = 8$$

$$x = 8 - 2y$$

As  $y \in \mathbb{N}$ , Put the values of  $y = 1, 2, 3, \dots$  till  $x \in \mathbb{N}$

$$\text{When, } y = 1, x = 8 - 2(1) = 8 - 2 = 6$$

$$\text{When, } y = 2, x = 8 - 2(2) = 8 - 4 = 4$$

$$\text{When, } y = 3, x = 8 - 2(3) = 8 - 6 = 2$$

$$\text{When, } y = 4, x = 8 - 2(4) = 8 - 8 = 0$$

Now,  $y$  cannot hold the value 4 because  $x = 0$  for  $y = 4$ , which is not a natural number.

$$\therefore R = \{(2, 3), (4, 2), (6, 1)\}$$

$$R^{-1} = \{(3, 2), (2, 4), (1, 6)\}$$

(iii) Given,

$R$  is a relation from  $\{11, 12, 13\}$  to  $\{8, 10, 12\}$  defined by  $y = x - 3$

Here,

$$x = \{11, 12, 13\} \text{ and } y = \{8, 10, 12\}$$

$$y = x - 3$$

$$\text{When, } x = 11, y = 11 - 3 = 8 \in \{8, 10, 12\}$$

$$\text{When, } x = 12, y = 12 - 3 = 9 \notin \{8, 10, 12\}$$

$$\text{When, } x = 13, y = 13 - 3 = 10 \in \{8, 10, 12\}$$

$$\therefore R = \{(11, 8), (13, 10)\}$$

$$R^{-1} = \{(8, 11), (10, 13)\}$$

**Sol.5. (i)** A relation  $R$  from the set  $\{2, 3, 4, 5, 6\}$  to the set  $\{1, 2, 3\}$  defined by  $x = 2y$ .

$$\text{Let } A = \{2, 3, 4, 5, 6\} \text{ and } B = \{1, 2, 3\}$$

$$\text{Given, } x = 2y \text{ where } y = \{1, 2, 3\}$$

$$\text{When, } y = 1, x = 2(1) = 2$$

$$\text{When, } y = 2, x = 2(2) = 4$$

$$\text{When, } y = 3, x = 2(3) = 6$$

$$\therefore R = \{(2, 1), (4, 2), (6, 3)\}$$

(ii) A relation  $R$  on the set  $\{1, 2, 3, 4, 5, 6, 7\}$  defined by  $(x, y) \in R \Leftrightarrow x$  is relatively prime to  $y$ .

Given:

$(x, y) \in R$  if  $x$  is relatively prime to  $y$

Here,

2 is co-prime to 3, 5 and 7.

3 is co-prime to 2, 4, 5 and 7.

4 is co-prime to 3, 5 and 7.

5 is co-prime to 2, 3, 4, 6 and 7.

6 is co-prime to 5 and 7.

7 is co-prime to 2, 3, 4, 5 and 6.

$$\therefore R = \{(2, 3), (2, 5), (2, 7), (3, 2), (3, 4), (3, 5), (3, 7), (4, 3), (4, 5), (4, 7), (5, 2), (5, 3), (5, 4), (5, 6), (5, 7), (6, 5), (6, 7), (7, 2), (7, 3), (7, 4), (7, 5), (7, 6), (7, 7)\}$$

(iii) A relation  $R$  on the set  $\{0, 1, 2, \dots, 10\}$  defined by  $2x + 3y = 12$ .

Given,

$$(x, y) \in R \text{ if } 2x + 3y = 12$$

$$\text{Where } x \text{ and } y = \{0, 1, 2, \dots, 10\}$$

$$2x + 3y = 12$$

$$2x = 12 - 3y$$

$$x = (12 - 3y)/2$$

$$\text{When, } y = 0, x = (12 - 3(0))/2 = 12/2 = 6$$

$$\text{When, } y = 2, x = (12 - 3(2))/2 = (12 - 6)/2 = 6/2 = 3$$

$$\text{When, } y = 4, x = (12 - 3(4))/2 = (12 - 12)/2 = 0/2 = 0$$

$$\therefore R = \{(0, 4), (3, 2), (6, 0)\}$$



**(iv)** A relation  $R$  from a set  $A = \{5, 6, 7, 8\}$  to the set  $B = \{10, 12, 15, 16, 18\}$  defined by  $(x, y) \in R \Leftrightarrow x$  divides  $y$ .

Given,

$(x, y) \in R$   $x$  divides  $y$

Where,  $x = \{5, 6, 7, 8\}$  and  $y = \{10, 12, 15, 16, 18\}$

Here,

5 divides 10 and 15.

6 divides 12 and 18.

7 divides none of the values of set  $B$ .

8 divides 16.

$\therefore R = \{(5, 10), (5, 15), (6, 12), (6, 18), (8, 16)\}$

**Sol.6.** Given,

$(x, y) \in R$   $x + 2y = 8$  where  $x \in \mathbb{N}$  and  $y \in \mathbb{N}$

$x + 2y = 8$

$x = 8 - 2y$

Putting the values  $y = 1, 2, 3, \dots$  till  $x \in \mathbb{N}$

When,  $y = 1$ ,  $x = 8 - 2(1) = 8 - 2 = 6$

When,  $y = 2$ ,  $x = 8 - 2(2) = 8 - 4 = 4$

When,  $y = 3$ ,  $x = 8 - 2(3) = 8 - 6 = 2$

When,  $y = 4$ ,  $x = 8 - 2(4) = 8 - 8 = 0$

Now,  $y$  cannot hold the value 4 because  $x = 0$  for  $y = 4$ , which is not a natural number.

$\therefore R = \{(2, 3), (4, 2), (6, 1)\}$

$R^{-1} = \{(3, 2), (2, 4), (1, 6)\}$

**Sol.7.** Given

$A = \{3, 5\}$  and  $B = \{7, 11\}$

$R = \{(a, b) : a \in A, b \in B, a-b \text{ is odd}\}$

On putting  $a = 3$  and  $b = 7$ ,

$a - b = 3 - 7 = -4$  which is not odd

On putting  $a = 3$  and  $b = 11$ ,

$a - b = 3 - 11 = -8$  which is not odd

On putting  $a = 5$  and  $b = 7$ :

$a - b = 5 - 7 = -2$  which is not odd

On putting  $a = 5$  and  $b = 11$ :

$a - b = 5 - 11 = -6$  which is not odd

$\therefore R = \{\} = \Phi$

$R$  is an empty relation from  $A$  into  $B$ .

Hence proved.

**Sol.8.** Given,

$A = \{1, 2\}$ ,  $B = \{3, 4\}$

$n(A) = 2$  (Number of elements in set  $A$ ).

$n(B) = 2$  (Number of elements in set  $B$ ).

We know,

$n(A \times B) = n(A) \times n(B)$

$= 2 \times 2$

$= 4$  [since,  $n(x) = a$ ,  $n(y) = b$ . total number of relations  $= 2ab$ ]

$\therefore$  The number of relations from  $A$  to  $B$  is  $24 = 16$ .

**Sol.9. (i)**  $R = \{(x, x+5) : x \in \{0, 1, 2, 3, 4, 5\}\}$

Given,

$R = \{(x, x+5) : x \in \{0, 1, 2, 3, 4, 5\}\}$

$\therefore R = \{(0, 0+5), (1, 1+5), (2, 2+5), (3, 3+5), (4, 4+5), (5, 5+5)\}$

$R = \{(0, 5), (1, 6), (2, 7), (3, 8), (4, 9), (5, 10)\}$



So,

Domain of relation  $R = \{0, 1, 2, 3, 4, 5\}$

Range of relation  $R = \{5, 6, 7, 8, 9, 10\}$

(ii)  $R = \{(x, x^3) : x \text{ is a prime number less than } 10\}$

Given,

$R = \{(x, x^3) : x \text{ is a prime number less than } 10\}$

Prime numbers less than 10 are 2, 3, 5 and 7

$\therefore R = \{(2, 2^3), (3, 3^3), (5, 5^3), (7, 7^3)\}$

$R = \{(2, 8), (3, 27), (5, 125), (7, 343)\}$

So,

Domain of relation  $R = \{2, 3, 5, 7\}$

Range of relation  $R = \{8, 27, 125, 343\}$

**Sol.10. (i)**  $R = \{a, b\} : a \in \mathbb{N}, a < 5, b = 4\}$

Given,

$R = \{a, b\} : a \in \mathbb{N}, a < 5, b = 4\}$

Natural numbers less than 5 are 1, 2, 3 and 4

$a = \{1, 2, 3, 4\}$  and  $b = \{4\}$

$R = \{(1, 4), (2, 4), (3, 4), (4, 4)\}$

So,

Domain of relation  $R = \{1, 2, 3, 4\}$

Range of relation  $R = \{4\}$

(ii)  $S = \{a, b\} : b = |a-1|, a \in \mathbb{Z} \text{ and } |a| \leq 3\}$

Given,

$S = \{a, b\} : b = |a-1|, a \in \mathbb{Z} \text{ and } |a| \leq 3\}$

$\mathbb{Z}$  denotes an integer which can be positive as well as negative

Now,  $|a| \leq 3$  and  $b = |a-1|$

$\therefore a = \{-3, -2, -1, 0, 1, 2, 3\}$

For,  $a = -3, -2, -1, 0, 1, 2, 3$  we get,

$S = \{(-3, |-3-1|), (-2, |-2-1|), (-1, |-1-1|), (0, |0-1|), (1, |1-1|), (2, |2-1|), (3, |3-1|)\}$

$S = \{(-3, |-4|), (-2, |-3|), (-1, |-2|), (0, |-1|), (1, |0|), (2, |1|), (3, |2|)\}$

$S = \{(-3, 4), (-2, 3), (-1, 2), (0, 1), (1, 0), (2, 1), (3, 2)\}$

$b = 4, 3, 2, 1, 0, 1, 2$

So,

Domain of relation  $S = \{0, -1, -2, -3, 1, 2, 3\}$

Range of relation  $S = \{0, 1, 2, 3, 4\}$

**Sol.11.** The total number of relations that can be defined from a set  $A$  to a set  $B$  is the number of possible subsets of  $A \times B$ . If  $n(A) = p$  and  $n(B) = q$ , then  $n(A \times B) = pq$ .

So, the total number of relations is  $2pq$ .

Now,

$A \times A = \{(a, a), (a, b), (b, a), (b, b)\}$

The total number of relations are all possible subsets of  $A \times A$ :

$[\{(a, a), (a, b), (b, a), (b, b)\}, \{(a, a), (a, b)\}, \{(a, a), (b, a)\}, \{(a, a), (b, b)\}, \{(a, b), (b, a)\}, \{(a, b), (b, b)\}, \{(b, a), (b, b)\}, \{(a, a), (a, b), (b, a)\}, \{(a, b), (b, a), (b, b)\}, \{(a, a), (b, a), (b, b)\}, \{(a, a), (a, b), (b, b)\}, \{(a, a), (a, b), (b, a)\}, \{(a, a), (a, b), (b, a), (b, b)\}]$

$n(A) = 2 \Rightarrow n(A \times A) = 2 \times 2 = 4$

$\therefore$  Total number of relations  $= 2^4 = 16$