

Ø

NITROGEN CONTAINING COMPOUNDS (Amines)

Each of these questions has 4 choices (a), (b), (c) and (d) for its answer, out of which ONLY ONE is correct.







4. Which statement is true regarding the following structure?

$$CH_3 \stackrel{N}{\underset{C_2H_5}{}} C_3H_7$$

- (a) It is a chiral molecule
- (b) It exists in two resolvable optically active forms
- (c) Both a) and b) (d) Neither a) nor b)
- Introduction of a methyl group in ammonia markedly increases the basic strength of ammonia in aq. solution, introduction of the second methyl group increases only marginally the basic strength of methyl amine in water. This is due to
 - (a) different type of hybridisation in the two amines
 - (b) protonated dimethyl amine is more solvated than methyl amine
 - (c) protonated dimethyl amine is more solvated than the protonated methyl amine
 - (d) protonated dimethyl amine is less stable than the protonated methyl amine
- 6. The basic character of ethyl amine, diethyl amine and triethyl amine in chlorobenzene is
 - (a) $C_2H_5NH_2 < (C_2H_5)_2NH < (C_2H_5)_3N$
 - (b) $C_2H_5NH_2 < (C_2H_5)_3N < (C_2H_5)_2 NH$
 - (c) $(C_2H_5)_3N < (C_2H_5)_2NH < C_2H_5NH_2$
 - (d) $(C_2H_5)_3N < C_2H_5NH_2 < (C_2H_5)_2NH_3$

-					
Mark Your	1. abcd	2. abcd	3. abcd	4. abcd	5. abcd
Response	6. abcd				

5.

7. The correct order of decreasing basic character of the three aliphatic primary amines is

$$\label{eq:holdsystem} \underbrace{\begin{array}{c} & & \\ I & & \\ I & & \\ (a) & I > II > III \\ (c) & I > II \approx III \\ \end{array}}_{II} \underbrace{\begin{array}{c} & & \\ NH_2 \\ III \\ (b) & III > II > II \\ (c) & I = II \\ III \\ (d) & I = II \\ III \\ IIII \\ III \\ IIII$$

Which of the statement is true regarding the basicity of 8. the following two primary amines ?



- (a) Both are equally basic because both are 1° amines
- (b) I > II because it is an aromatic amine
- (c) II > I because it is an aliphatic amine
- (d) I < II because of difference in the nature of β -carbon 9.
 - The correct order of decreasing basic character is

$$\begin{array}{cc} C_6H_5NH_2\\ I \end{array}, \qquad \begin{array}{cc} C_6H_5CH_2NH_2\\ II \end{array},$$

$$(C_6H_5)_2 NH$$
, $C_6H_{11}NH_2$

- III (a) II > I > III > IV(b) IV > II > I > III(c) IV > III > II > I(d) IV > II > III > I
- 10. The correct order of increasing basicity is

The relative order of basic character of the following 11. compounds is



/ > III

(A)

- When aniline is treated with acetyl chloride in presence of anhydrous aluminium chloride, the main product is
 - (a) o-aminoacetophenone (b) p-aminoacetophenone
 - (c) both (a) and (b) (d) *m*-aminoacetophenone

13.
$$CH_3CH_2CH_2NH_2 \xrightarrow{NaNO_2, HCl} P$$
. P is

12.

- (a) $CH_3CH_2CH_2OH$ (b) $(CH_3)_2 CHCl$
- (c) Both (a) and (b) (d) Reaction not possible
- 14. Benzenediazonium chloride when treated with phenols gives azo dyes, to get best result the pH of the medium should be
 - (a) around 4 (b) around 8
 - (c) around 10 (d) 12
- 15. In the following reaction, the reagent X should be

$$RCOOH + [X] \xrightarrow{conc. H_2SO_4} RNH_2$$

- (a) NH₃ (b) HN_3
- (c) either of the two (d) None of the two
- 16. Which of the following can undergo electrophilic substitution when treated with nitrous acid at 0° C?
 - (a) $C_6H_5NH_2$ (b) C₆H₅NHCH₃ (c) $C_6H_5N(CH_3)_2$ (d) None
- Which of the following does not reduce $C_6H_5NO_2$ to 17. aniline?

- (c) Zn/HCl (d) LiAlH₄
- 18. Which of the following amines can be resolved into two enantiomers?



F -					
	7. abcd	8. abcd	9. abcd	10. abcd	11. abcd
MARK YOUR Response	12.abcd	13.abcd	14. abcd	15. abcd	16. abcd
	17.abcd	18. abcd			

- **19.** Which of the following statement is false?
 - (a) Dimethyl amine as well as trimethyl amine are soluble in water
 - (b) Trimethyl amine forms hydrogen bond neither with itself nor with water.
 - (c) Trimethyl amine can act as hydrogen bond acceptor only, while dimethyl amine can serve as both a hydrogen bond donor and acceptor
 - (d) All the three statements are false
- **20.** Ephedrine is a secondary amine. It is widely used in cold and allergy conditions in the form of its hydrochloride but not as such because
 - (a) the amine itself has unpleasant smell, while its salt is odourless
 - (b) the amine is insoluble in water, while the salt is soluble in water
 - (c) the amine is unstable and easily oxidised by air, while the salt is resistant to atmospheric oxidation
 - (d) of all the above facts
- 21. Benzamide and benzyl amine can be distinguished by
 - (a) cold dil. NaOH
 - (b) cold dil. HCl
 - (c) both a & b
 - (d) NaNO₂, HCl, 0°C, then β -naphthol
- 22. The correct order for the basic character of the compounds I to IV should be



- (a) IV < III < I < II (b) IV < I < III < II
- (c) IV < II < III < I (d) IV < III < II < I
- **23.** Which of the following is true regarding basic character of pyridine and pyrrole?
 - (a) Pyrrole is more basic because its nonbonding electrons occupy sp^3 orbital
 - (b) Pyridine is more basic because its nonbonding electrons is not a part of aromatic sextet.
 - (c) Both are equally basic

(d) Pyridine is less basic because it is 3° amine

24. Pyrrole and pyridine both are basic and form salts with acids?

Which of the following statement is true regarding the aromatic character of the four species?

- (a) All the four are aromatic
- (b) I, III and IV are aromatic
- (c) I, II and III are aromatic
- (d) I and III are aromatic
- 25. Ethylene can be prepared in good yield by
 - (a) $CH_3CH_2N^+(CH_3)_3I^- \xrightarrow{heat} \rightarrow$

$$CH_2 = CH_2 + (CH_3)_3 N + HI$$

(b) $CH_3CH_2N^+(CH_3)_3OH^- \xrightarrow{heat} \rightarrow$

$$CH_2 = CH_2 + (CH_3)_3 N + H_2O$$

- (c) Both a and b
- (d) $CH_3CH_2NH_2 \xrightarrow{heat} CH_2 = CH_2 + NH_3$

26. Which one of the following is not an oxidation product of a primary amine?

- (a) A hydroxylamine(b) A nitroso compound(c) A nitro compound(d) None of these
- 27. Which of the following method is used for eliminating nitrogen of an amine present outside the ring?
 - (a) Hofmann elimination (b) Cope elimination
 - (c) Both (d) Emde degradation
- 28. Which of the following does not react with nitrous acid?
 - (a) $C_6H_5NH_2$ (b) $C_6H_5NHCH_3$
 - (c) $C_6H_5N(CH_3)_2$ (d) None of these
- **29.** Which of the following leads to carbon-carbon double bond?
 - (a) 1° Amine + RCHO \rightarrow (b) 2° Amine + R₂CO \rightarrow
 - (c) 2° Amine + RCHO \rightarrow (d) both b & c
- **30.** Electrophilic aromatic substitution of pyridine resembles with
 - (a) benzene (b) aniline
 - (c) nitrobenzene (d) none of these

-					
	19.abcd	20. abcd	21. abcd	22. abcd	23. abcd
MARK YOUR Response	24. abcd	25. abcd	26. abcd	27. abcd	28. abcd
	29. abcd	30. abcd			

- **31.** Which of the following will react most readily with NaOH to form ethanol ?
 - (a) $(CH_3)_4 N^+ I^-$ (b) $(CH_3)_4 S^+ I^-$
 - (c) $(CH_3)_3CCl$ (d) CH_3OCH_3











Ø

- **33.** In Hofmann bromamide degradation, one of the important steps is the migration of
 - (a) an alkyl group without its electron pair to electron deficient N atom
 - (b) an alkyl group with its electron pair to electron deficient O atom
 - (c) an alkyl group with its electron pair to electron rich N atom
 - (d) an alkyl group with its electron pair to electron deficient N atom

- 34. $(CH_3)_3 C \overset{O}{C} NH_2 \xrightarrow{(i) OD^- / Br_2} Product P is$ (a) $(CH_3)_3 CNH_2$ (b) $(CH_3)_3 CNHD$ (c) $(CH_3)_3 CND_2$ (d) no reaction
- **35.** Reaction of ethyl amine with alkaline **chloroform** leads to the formation of carbylamine reaction. This reaction involves the attack of an electrophile on ethyl amine, the electrophile is

(a)
$$H_3O^+$$
 (b) H^+
(c) RNH^+_3 (d) : CCl_2

36. Predict the possible number of alkenes and the main alkene in the following reaction.



<i>v</i> = - -					
Mark Your	31.abcd	32. abcd	33. abcd	34. abcd	35. abcd
Response	36.abcd	37. abcd	38. abcd		

39.
$$\begin{array}{c} O \\ \parallel & \overrightarrow{} \\ R - C - C H \\ \overrightarrow{} \\ \overrightarrow{} \\ \end{array} \xrightarrow{+} N \xrightarrow{+}$$

What is the nature of its intermediate in this reaction ?

- (c) Carbene (d) Free radial
- **40.** The intermediate formed in the above reaction rearranges to

(a)
$$R - CH_2 - CH$$

(b) $R - CH = C = O$
(c) $R - C - CH_2$
(d) $R - C - CH_2$

41.
$$(H_3NO_2) \to [A] \longrightarrow [B]$$

 \cap



47.

Which reaction is/are involved in the above conversion?

- (a) Diazotisation and Beckmamn rearrangement
- (b) Diazotisation and Robinson annulation
- (c) Aldol condensation and diazotisation
- (d) Diazotisation and pinacol type of rearrangement
- **42.** Which of the reaction can not be used for introducing a C = C linkage?
 - (a) Hofmann elimination (b) Aldol condensation
 - (c) Wittig reaction (d) None of the three
- **43.** Following reaction is an example of

$$\begin{array}{c} O \\ \parallel \\ R - C - OH + HN_3 \xrightarrow{H_2SO_4} R - NH_2 \end{array}$$

- (a) Hofmann reaction(c) Schmidt reaction
 - (b) Curtius reaction(d) Lossen reaction

- 44. Which of the following is one of the intermediates in Hofmann, Curtius, Schmidt and Lossen reaction ?
 - (a) $R \stackrel{+}{C}O$ (b) RNCO
 - (c) RCNO (d) RCONH
- **45.** Which of the following involves migration to electron-deficient nitrogen?
 - (a) Benzilic acid rearrangement
 - (b) Wolf rearrangement
 - (c) Allylic rearrangement
 - (d) Beckmann rearrangement
- **46.** Which of the following is not formed as an intermediate in the Hofmann rearrangement?

$$R-CONH_{2} \xrightarrow{Br_{2}/NaOH} RNH_{2}$$

$$(a) R \xrightarrow{\parallel} H \\ (b) R \xrightarrow{\parallel} H \\ (c) R \xrightarrow{\parallel} H$$

(d) All the three are formed

Identify (C) and (D) in the following series of reactions

$$CH_{3}NH_{2} \xrightarrow{excess of} [A] \xrightarrow{AgOH} [B] \xrightarrow{heat} [C]+[D]$$

- (a) $(CH_3)_3COH, CH_3NH_2$
- (b) $(CH_3)_2C = CH_2, CH_3NH_2$
- (c) $(CH_3)_3N$, CH_3OH
- (d) $(CH_3)_2C=CH_2$, CH_3OH

48. Identify (D) and (E) in the given reaction

$$CH_{3}COOH \xrightarrow{(i) \text{ LiAlH}_{4}} (A) \xrightarrow{(i) \text{ KCN}} (B)$$

 $\xrightarrow{\text{CHCl}_3/\text{OH}^-} (\text{C}) \xrightarrow{\text{hydrolysis}} (\text{D}) + (\text{E})$

- (a) CH₃CH₂CH₂NHCH₃
- (b) (CH₃)₂CHNH₂, HCOOH
- (c) $CH_3CH_2CH_2NH_2$, HCOOH
- (d) CH₃CH₂CH₂COOH

Mark Your	39.@bcd	40. abcd	41. abcd	42. abcd	43. abcd
Response	44. abcd	45. abcd	46. abcd	47. abcd	48. abcd





51. (i)
$$CHN_2COOC_2H_5$$
 [Y] $\xrightarrow{\text{oxidation}}$ [Z]

Compound Z should be

1



52. Which of the following undergoes bromination most easily?



- 53. How many structural isomers of a Grignard reagent are possible for preparing *n*-butane by reaction with ethyl amine?
 - (a) 1 (b) 2
 - (c) 3 (d) 4
- 54. For the diazonium ions, the order of reactivity towards diazo-coupling with phenol in the presence of dil.NaOH is









(a) I < IV < II < III(c) I < II < IV < III

(b) IV < II < I < III

(d) III < IV < II < I

— <i>k</i> ı—					
Mark Your	49. abcd	50.abcd	51.abcd	52. abcd	53. abcd
Response	54.abcd				

 \equiv Comprehension Type \equiv

B

This section contains groups of questions. Each group is followed by some multiple choice questions based on a paragraph. Each question has 4 choices (a), (b), (c) and (d) for its answer, out of which ONLY ONE is correct.

PASSAGE-1

When a quaternary ammonium halide is treated with moist silver oxide (a slurry of Ag_2O in water), a quaternary ammonium hydroxide is formed as the main product. Thermal decomposition of the quaternary ammonium hydroxide gives an alkene. This reaction is known as Hofmann elimination and has most of the characteristics of an E2 reaction.

$$\begin{array}{cccc} I^{-} & N^{+}(CH_{3})_{3} & OH^{-} & N^{+}(CH_{3})_{3} \\ & & \downarrow \\ CH_{3}CH_{2} & CHCH_{3} & \xrightarrow{Ag_{2}O+H_{2}O} & CH_{3}CH_{2} & CHCH_{3} & \xrightarrow{heat} \end{array}$$

$$CH_{3}CH = CHCH_{3} + CH_{3}CH_{2}CH = CH_{2}$$

$$X$$

$$Y$$

- **1.** The Hofmann eliminations are
 - (i) concreted (ii) anti-stereoselective
 - (iii) occur in separate steps (iv) syn-stereoselective
 - (a) (i) and (iv) (b) (i) and (ii)
 - (c) Only (i) (d) Only (iii)
- 2. In the above reaction, which of the following is correct regarding yield of X and Y?
 - (a) X = Y(b) $X \approx Y$ (c) X > Y(d) X < Y

PASSAGE-2

The conversion of an amide to an amine with one carbon atom less by the action of alkaline hypohalite is known as Hofmann degradation.

$$\text{RCONH}_2 + \text{Br}_2 + 4\text{KOH} \longrightarrow \text{RNH}_2$$

The most important feature of the reaction is the rearrangement of N-bromamide anion to isocyanate :



Hofmann reaction is accelerated if the migrating group is more electron-releasing.

- **3.** Which step is the driving force in the above reaction to proceed in right direction?
 - (a) conversion of I to II
 - (b) conversion of II to III
 - (c) conversion of III to RNH₂
 - (d) All

CH₂

4.

Which of the following can undergo Hofmann reaction most easily?





6. Predict the product in the following reaction :



Understand carefully the following two reactions and answer the questions mentioned below.

Reaction (i)



Reaction (ii)



7. Which of the steps is(are) common in the two reactions?

- (a) 1st step(c) both
- (b) 2nd step
- (d) none

- 8. Which of the final products are medicinally important?
 - (a) Product from reaction (i)
 - (b) Product from reaction (ii)
 - (c) Products from both reactions
 - (d) None of the two products
- **9.** Which one of the following reagent can be used for the introduction of Cl and Na in reaction (ii)?
 - (a) Cl_2 / Na_2CO_3 (b) HCl / NaOH(c) NaCl / NaOH (d) NaOCl / NaOH
- 10. Saccharin, an important sweetening agent, of the following



products of the reaction (ii), that product should be



PASSAGE-4









11. What will be the product(s) when isopropylbenzene is oxidised with oxygen and product is acidified

CH2OH



(c)
$$OH + CH_3CHO$$

(d) $OH + (CH_2)_2C = O$

- 12. The final product in the above series of reactions is(a) an amine
 - (a) an annne

d'n

- (b) a mono substituted urea
- (c) a disubstituted urea (d) a urethane
- 13. Conversion of $C_6H_5CON_3$ to C_6H_5NCO is an example of rearrangement where an alkyl group migrates to
 - (a) electron deficient carbon atom
 - (b) electron deficient oxygen atom
 - (c) electron deficient nitrogen atom
 - (d) electron rich nitrogen atom

PASSAGE-5

Nitrous acid reacts with amines forming different products depending upon the type of amine. Aliphatic primary amines react with HONO forming alcohol as the major product, other products being alkene and alkyl halide. Certain cyclic primary amines can undergo ring expansion or ring contraction on treatment with HONO again forming alcohol as the major product.



Aromatic primary amines when treated with HONO undergo diazotization to form diazonium salt as the stable product.

- 14. Which of the following statement is not true ?
 - (a) All primary amines first form diazonium salt when treated with nitrous acid.
 - (b) All aliphatic primary amines when treated with nitrous acid form primary alcohols as the major product
 - (c) The three classes of amines give different products on treatment with nitrous acid.
 - (d) None of the three



- (a) enantiomer of the original compound
- (b) diastereomer of the original compound
- (c) racemic mixture of the original compound
- (d) (+) isomer

Bend I					
Mark Your	11. abcd	12. abcd	13. abcd	14. abcd	15. abcd
Response	16.abcd	17. abcd			

PASSAGE-6

Hofmann reaction is found to follow the following path.



- Number of moles of NaOH used in the complete reaction 18. is
 - (a) 1 (b) 2
 - (d) 4 (c) 3
- Hofmann reaction, when carried out in presence of 19. methanol containing sodium methoxide, instead of NaOH, product is

 $C_6H_5CONH_2 \xrightarrow{Br_2,CH_3ONa}$ Product is

(a)
$$C_6H_5NH_2$$
 (b) $C_6H_5NHCOOCH_3$
(c) $C_6H_5NHCH_3$ (d) C_6H_5NHBr

20.
$$CH_3 - \overset{O}{C} - ND_2 \xrightarrow{NaOH} Amine, Z is$$

- (a) $CH_3 ND_2$ (b) CH_3NH_2
- (c) mixture of (a) and (b) (d) Reaction not possible
- *21. Which of the following will not undergo Hofmann bromamide reaction ?



C₆H₅CONHCH₃

Ø





- Which of the following species in the above mechanism is 22. electron deficient?
 - (a) I (b) II (d) IV (c) III

* In presence of alkali, (b) undergoes hydrolysis to form

CH₂CONH₂ which when undergoes Hofmann reaction. CH₂COO⁻

PASSAGE-7

Beckmann rearrangement is used for determining the configuration of ketoxines since the two isomers form different N-substituted amides when heated with some acidic reagent. The different steps of the mechanism are as follows.



- Pick up the species which has an electron deficient centre? 23.
 - (a) I (b) II
 - (c) III (d) Two of the three
 - The mechanism involves migration of
 - R group with its bonding electron pair (a)
 - (b) R' group with its bonding electron pair
 - R' group without its bonding electron pair (c)
 - (d) OH group with its bonding electron pair
- 25. Which of the following reaction/step is involved in Beckmann rearrangement?
 - (a) Formation of conjugated acid
 - (b) Migration of an alkyl group to electron deficient N
 - (c) Enlolisation
 - (d) All the three

-					
Mark Your	18.@bCd	19. abcd	20. abcd	21. abcd	22. abcd
Response	23. abcd	24. abcd	25. abcd		

24.

Predict the product in the following case $\begin{array}{c} CH_{3} \\ C_{2}H_{5} \end{array} C = N \xrightarrow{OH} \begin{array}{c} (i) \ PCl_{5} \\ \hline (ii) \ D_{2}O \end{array} P$ (a) $CH_3CON(D)C_2H_5$ (b) $C_2H_5CON(D)CH_3$ (c) $CH_{3}CONHC_{2}H_{5}$ (d) $C_{2}H_{5}CONHCH_{3}$ $CH_{3} C = N \xrightarrow{(i) PCl_{5}} Q \text{ is}$ 27. (a) $C_2H_5NHCOCH_3$ (b) $C_2H_5NHCO^{18}CH_3$ (c) $\overline{CH_3}$ NHCO¹⁸C₂H₅ (d) $\overline{C_2}$ H₅CONHCH₃

26.

PASSAGE-8

Like ammonia, an amine (1°, 2° or 3°) can react with an alkyl halide to form next higher class of amine. Here, again it is the presence of electron pair on nitrogen which makes amines to behave as nucleophile and alkyl halides thus undergo nucleophilic substitutions.

$$\begin{array}{c} \overset{}{\operatorname{R}\overset{}{\operatorname{N}}\operatorname{H}_2}_{\operatorname{(Nucleophile)}} + \overset{R'\operatorname{CH}_2\operatorname{X}}{\overset{}{\operatorname{1^\circ}\operatorname{Alkyl}}\operatorname{halide}} & \longrightarrow \overset{}{\operatorname{RN}^+} - \operatorname{CH}_2\operatorname{R'} \\ \overset{}{\operatorname{H}}_{\operatorname{H}} \\ & \overset{}{\operatorname{H}} \\ \end{array} \\ & \longrightarrow \overset{}{\operatorname{R}\overset{}{\operatorname{N}}}_{\operatorname{H}} - \operatorname{CH}_2\operatorname{R'} + \operatorname{HX} \\ & \operatorname{H} \end{array}$$

A second alkylation may follow, converting the secondary amine to a tertiary amine which may be alkylated to give a quaternary ammonium salt.

28. Identify the product(s) obtained when $Br(CH_2)_ABr$ is heated with 1 equivalent of ethyl amine



2° Amine

29. Identify the product(s) obtained when $Br(CH_2)_4Br$ is heated with excess of methyl amine





(c) Both of these

- (d) None of the two
- Mark Your 26. abcd 27. (a)(b)(c)(d) 28. (a) (b) (c) (d) 29. abcd 30. abcd Response

PASSAGE-9

Summary of the four related name reactions for the preparation of primary amines are given below.

31.

32. In which of the reactions, leaving group is carboxylate anion?

(b) Cutius

(c) Lossen (d) Schmidt $\operatorname{RCONH}_2 \xrightarrow{\operatorname{Br}_2, \operatorname{OH}^-} \operatorname{RNH}_2$ 33. In which of the reactions, nitrogen constitutes the leaving Hofmann group? $\text{RCON}_3 \xrightarrow[\text{heat}]{\text{H}_2\text{O}} \text{RNH}_2$ Curtius (a) Curtius (b) Schmidt (c) Both (d) Hofmann $\text{RCONHOH} \xrightarrow{\text{OH}^-} \text{RNH}_2$ Lossen 34. Which of the reaction does not require alkaline medium? (a) Hofmann (b) Curtius $\text{RCOOH} \xrightarrow{\text{HN}_3} \text{RNH}_2$ Schmidt (c) Schmidt (d) Lossen Reactant of which reaction is not a derivative of carboxylic 35. Which of the reaction is of least importance ? acid? (a) Schmidt (b) Hofmann (b) Curtius (a) Hofmann (c) Curtius (d) Lossen (c) Lossen (d) None of the three

(a) Hofmann

#					
MARK YOUR	31 0000	32 @@@@@	33 @@@@@	34 ଉଦ୍ଭାରଣ	35 ଉତ୍ତର
Response					
					•

	REASC	DNING TYPE 🚃							
	C In the fo question responses (a) Bot (b) Bot (c) Stat (d) Stat	llowing questions has 4 choices (a), (t from the following h Statement-1 and Sta h Statement-1 and Sta tement-1 is true but St tement-1 is false but S	two Statement-1 (,)), (c) and (d) for it options: tement-2 are true and tement-2 are true and atement-2 is false. tatement-2 is true.	Asse s an l Sta l Sta	ertion) and Sta swer, out of w tement-2 is the co tement-2 is not th	ntem hich orrec ne co	ent-2 (Reason) ONLY ONE i et explanation of rrect explanation	are s con State r of S	provided. Each rrect. Mark your ement-1. Statement-1.
1.	Statement-1	: Sulphanilic acid ex whereas <i>p</i> -aminobe	tists as a dipolar ion enzoic acid does not.		Statement-	2 :	<i>m</i> -Nitroaniline and hence furthe	form er red	ed gets precipitated uction is prevented.
	Statement-2	: Carboxyl group, be -SO ₃ H group, can to the amino group	ing more acidic than easily transfer a H ⁺	4.	Statement-	1 :	Benzonitrile is of chloroben	prepa zene	ared by the reaction with potassium
2.	Statement-1	: Nitration of aniline done by protecting acetylation.	can be conveniently the amino group by	5.	Statement- Statement-	2 : 1 :	Cyanide (CN ⁻) 1° Amides read	is a s t wit	strong nucleophile. th Br ₂ + NaOH to
	Statement-2	: Acetylation incre density in the benz	eases the electron- ene ring.				give 1° amines we than the parent	vith o amid	ne carbon atom less le.
3.	Statement-1	: Reduction of <i>m</i> -dinitrobenzene with ammonium sulphide gives <i>m</i> - nitroaniline.			Statement-	2 :	The reactio intermediate fo	n (rmati	occurs through on of acylnitrene.
	- 🖾 ——								
	Mark Your Response	1. abcd	2. abcd	3.	abcd	4.	abcd	5.	abcd

6.	Statement-1	: In strongly acidic solutions, aniline becomes more reactive towards electrophilic reagents.	11.	Statement-1 Statement-2	 Pyridine is basic. Lone pair of electrons on its nitrogen is not involved in aromatic sextet.
	Statement-2	: The amino group being completely protonated in strongly acidic solution, the lone pair of electrons on the nitrogen is no longer available for resonance.	12.	Statement-1 Statement-2	 Pyridine is more basic than pyrrole. Lone pair of electrons on N in pyridine and pyrrole are different in nature, these form a part of aromatic sextet in pyrrole.
7.	Statement-1	: In order to convert R–Cl to pure R–NH ₂ , Gabriel-phthalimide synthesis can be used.	13.	Statement-1	while not in pyridine.All compounds containing an odd
	Statement-2	: With proper choice of alkyl halides, phthalimide synthesis can be used to prepare 1° 2° or 3° amines			number of nitrogen atoms have odd masses and those containing even number of N atoms have even masses.
8.	Statement-1	 Aniline does not undergo Friedel-Crafts reaction. 		Statement-2	: Nitrogen rule can be applied to both aliphatic and aromatic compounds.
	Statement-2	: -NH ₂ group of aniline reacts with AlCl ₃ (Lewis acid) to give acid-base reaction.	14.	Statement-1	: The main product of reaction of alcoholic silver nitrite and ethyl bromide
9.	Statement-1	: Acetamide reacts with Br ₂ in presence of methanolic CH ₃ ONa to form methyl N-methylcarbamate.		Statement-2	is nitroethane.Silver nitrite is predominantly covalent compound.
	Statement-2	: Methyl isocyanate is formed as an	15.	Statement-1	: HCN on hydrolysis gives formic acid
		intermediate which reacts with methanol		Statement-2	: HCN is a weak monoprotic acid
10	Statement-1	• 1° 2° and 3° amines are hydrogen-	16.	Statement-1	: Hofmann elimination gives alkene
10.	Statement 1	bonding donors, while 1° and 2° amines	17	Statement-2	: Alkenes are less acidic than NH ₃
		are hydrogen-bonding acceptors.	1/.	Statement-1	condensation
	Statement-2	: Low molecular weight amines are soluble in water.		Statement-2	: α-Hydrogen of nitromethane is acidic.
	¢				
	<i>y</i> — <i>y</i> =				

ManyVour	6. abcd	7. abcd	8. abcd	9. abcd	10. abcd
MARK YOUR Response	11.abcd	12. abcd	13. abcd	14. abcd	15. abcd
	16.abcd	17.abcd			

MULTIPLE CORRECT CHOICE TYPE Each of these questions has 4 choices (a), (b), (c) and (d) for its answer, out of which ONE OR MORE is/are correct.

 $N(CH_3)_2$

NH,

HNO₃

H₂SO₄

fuming

HNO₃

Nitro group can't be introduced easily by which 1. reaction?

HN

H_SO

HNO

CH_COOH

(b)

'(d)

NH₂

 $N(CH_3)_2$

(a)

(c)

E

2. Which of the following compound can be methylated by diazomethane?

- (a) C₂H₅COOH (b) $C_2H_5NH_2$ (c) C₆H₅OH
- (d) CH₃COCH₂COOC₂H₅

3. Which of the following amine can be prepared by Gabriel method ?

- (a) $CH_3CH_2NH_2$ (c) (CH₃)₃CNH₂
- (b) $(CH_3)_2CHNH_2$ (d) $C_6H_5NH_2$

-				
Mark Your Response	1. @bcd	2. abcd	3. abcd	

- 4. Which of the following can't be used as an alkylating agent for an amine ?
 - (a) CH_3CH_2Cl (b) $CH_2 = CHCl$
 - (c) C_6H_5Cl (d) $(CH_3)_3CCl$
- 5. $(CH_3)_3CNH_2$ can't be prepared by
 - (a) reductive amination (b) oxime reduction
 - (c) Hofmann degradation (d) Gabriel synthesis.
- **6.** Which of the following reaction can be used for preparing aniline ?
 - (a) $C_6H_5COOH \xrightarrow{N_3H, \text{ conc. } H_2SO_4}$
 - (b) $C_6H_5NC \xrightarrow{H_3O^+}$
 - (c) $C_6H_5NC \xrightarrow{\text{LiAlH}_4} \rightarrow$
 - (d) $C_6H_5CONH_2 \xrightarrow{Br_2/NaOH} \rightarrow$
- 7. Which of the following can exist as inner salt ?
 - (a) p-Aminobenzenesulphonic acid
 - (b) p-Aminobenzoic acid
 - (c) Aminoacetic acid
 - (d) Alanine

A

8. Which of the following pairs show coupling reaction ?





(d) Diazotised sulphanilic acid + Dimethylaniline

9. Which of the following statement is true regarding reaction of *p*-aminophenol with arenediazonium chloride?

HO
$$\longrightarrow 5 - 6$$
 NH₂ + ArN₂⁺Cl⁻ \longrightarrow

- (a) Reaction takes place at position 2 in presence of HCl.
- (b) Reaction takes place at position 3 in presence of NaOH.
- (c) Only two positions (2 and 6) can be coupled in presence of OH^-
- (d) Four azo groups can be introduced in the molecule.
- Which of the following reacts with nitrous acid?
 - (a) Acetamide

10.

- (b) 2-Nitrobutane
- (c) 2-Methyl-2-nitropropane
- (d) Diethylamine
- **11.** *p*-Nitroaniline can be obtained by









Been 1					
Mark Your	4. abcd	5. abcd	6. abcd	7. abcd	8. abcd
Response	9. abcd	10. abcd	11. abcd		

2.
$$\langle \bigcirc NH_2 + CH_3Cl \xrightarrow{AlO_3} ?$$

Which of the following statement is false regarding above reaction?



- (d) No substitution
- 13. The correct name for $CH_3 N \cong C$ is
 - (a) methyl carbylamine (b) methyl isocyanide
 - (c) methyl isonitrile (d) acetoisonitrile
 - Urea on heating with ethyl alcohol gives
 - (a) urethane (b) semicarbazide
 - (d) ethyl carbonate malonylurea (c)

15.

Ø

14.

(i) $O_2, V_2O_5, heat \rightarrow Product$ (ii) NH₃, heat (iii) NaOH / Br₂

Which of the following statement (s) is (are) true regarding above series of reactions ?



- Reaction involved is oxidation (c)
- (d) Reaction involved is Hofmann degradation

- 16. Oximes can be converted into amides by
 - (b) BF₃ (a) SO_3

(d) $C_6H_5SO_2Cl$ (c) PCl₅

- 17. Pick up the correct statement
 - (a) Toluene and ethane, both react at the same rate with chlorine in presence of light.



18. Which of the following is/are correct regarding nitration of aniline with conc HNO₃ and conc. H₂SO₄?





- (c) The substitution can be explained on the basis of inductive effect (- I)
- (d) The substitution can be influenced by +M and +E effects.

-					
Mark Your	12.@bcd	13. abcd	14. abcd	15. abcd	16. abcd
RESPONSE	17.abcd	18. abcd			

1

19. In which of the following reaction, Hofmann elimination is the major product ?

(a)
$$\xrightarrow{Me_3COK}$$
 Br





- 20. Which of the following intermediates are likely to be formed when 2-methylpropyl amine is treated with nitrous acid?
 - (a) $(CH_3)_2 CHCH_2$ (b) $(CH_3)_3 COH_2$
 - (c) $(CH_3)_2CHCH_2OH_2$ (d) $(CH_3)_2CHCH_3$

- 21. Carbenes are the reactive intermediates in
 - (a) Reimer Tiemann reaction
 - (b) Wittig reaction
 - (c) Hofmann bromamide reaction
 - (d) Carbylamine reaction
- 22. Which of the following are intermediates in Sandmeyer reaction?

(a)
$$C_6H_5N^+ \equiv NCl^-$$
 (b) $C_6H_5N^+ \equiv NCl^-$

(c)
$$\dot{C}_6 H_5$$
 (d) $C_6 H_5 Cl$

23.
$$C_4H_{11}N + HONO \longrightarrow C_4H_{10}O$$
(X) (3° alcohol)

X should

- (a) give carbylamine reaction
- (b) undergo diazotisation
- (c) react with water
- (d) give Hofmann bromamide reaction

— £ 1—					
Mark Your	19.@b©d	20. abcd	21. abcd	22. abcd	23. abcd
Response	19.0000	20. abcd	21. abcd	22. abcd	23. ab



Each question contains statements given in two columns, which have to be matched. The statements in Column-I are labeled A, B, C and D, while the statements in Column-II are labelled p, q, r, s and t. Any given statement in Column -I can have correct matching with ONE OR MORE statement(s) in Column-II. The appropriate bubbles corresponding to the answers to these questions have to be darkened as illustrated in the following example: If the correct matches are A-p, s and t; B-q and r; C-p and q; and D-s then the correct darkening of bubbles will look like the given.



1.	(A)	Column-I	n	Column-II Ethyl agotoggatata	4.		Column-I		Column-II
	(A) (B)	β-Amino acids	р. q.	Proteins		(A)	ArS _{N1}	p.	Carbocation
	(C)	H ₂ N-СООН	r.	Zwitterion			as intermediate		
	(D)	Aniline	s.	Diazotisation		(B)	ArS _{N2}	q.	Carbanion
2.		Column-I		Column-II			as intermediate		
		N				(C)	$OH^{-} > (CH_3)_3 CO^{-}$	r.	Basic character
	(A)		p.	One type of N		(D)	$OH^- < NH_2^-$	s.	Nucleophilic character
		Н			5.		Column-I		Column-II
	(B)	$H_{2}N + C - NH_{2}$	a	Two types of N		(A)	Allylic rearrangement	p.	Carbanions
	(D)	N N	ч. r	Aromatic		(B)	Hofmann rearrangement	q.	Electron deficient species
	(C)	N	1.	Aromatic		(C)	Wittig reaction	r.	Carbocations
3.	(D)	$C_6H_5N^+ \equiv NCI^-$ Column-I	S.	Non-aromatic Column-II		(D)	Carbylamine reaction	s.	Carbenes
	(A)	$\text{CHCl}_3 \xrightarrow{\text{OH}^-}$	p.	Hofmann product	6.		Column-I		Column-II
	(B)	$CHCl_2 - CF_3$ $OC_2H_5^{-}$	q.	E2 mechanism		(A)	Gabriel phthalimide reaction	p.	C ₆ H ₅ CH ₂ NH ₂
	(\mathbf{C})	Br	r	a Elimination		(B)	Reduction with LiAlH ₄	q.	C ₆ H ₅ NH ₂
	(C)	$(CH_3)_2 C - CH_2CH_3$ $\xrightarrow{Me_3CO^-}$		u-emination		(C)	Reaction with alc. KOH + CHCl ₃	r.	C ₆ H ₅ CN
	(D)	$CH_3 - CH - CH_2CH_3$	s.	E1cB		(D)	1° Amide with	s. C	H ₃ CH ₂ NH ₂
		$N^+(CH_3)_3$					Br ₂ + KOH		
		$\xrightarrow{\text{OH}^{-}}$							





Anemarkay

SINGLE CORRECT CHOICE TYPE

1.	(b)	7.	(a)	13.	(c)	19.	(b)	25.	(b)	31.	(a)	37.	(b)	43.	(c)	49.	(d)
2.	(c)	8.	(d)	14.	(b)	20.	(d)	26.	(d)	32.	(b)	38.	(d)	44.	(b)	50.	(c)
3.	(b)	9.	(b)	15.	(b)	21.	(b)	27.	(c)	33.	(d)	39.	(c)	45.	(d)	51.	(c)
4.	(a)	10.	(d)	16.	(c)	22.	(a)	28.	(d)	34.	(c)	40.	(b)	46.	(d)	52.	(a)
5.	(d)	11.	(c)	17.	(d)	23.	(b)	29.	(d)	35.	(d)	41.	(c)	47.	(c)	53.	(b)
6.	(a)	12.	(d)	18.	(d)	24.	(b)	30.	(c)	36.	(c)	42.	(d)	48.	(c)	54.	(b)

COMPREHENSION TYPE

A

B

1	(b)	7	(c)	13	(c)	19	(b)	25	(d)	31	(d)
2	(d)	8	(c)	14	(b)	20	(b)	26	(c)	32	(c)
3	(b)	9	(d)	15	(a,b)	21	(c)	27	(c)	33	(c)
4	(c)	10	(a)	16	(d)	22	(c)	28	(a, d)	34	(b)
5	(a)	11	(d)	17	(c)	23	(d)	29	(c,d)	35	(d)
6	(c)	12	(d)	18	(d)	24	(b)	30	(a)		

REASONING TYPE

1	(c)	4	(d)	7	(c)	10	(b)	13	(b)	16	(b)
2	(c)	5	(a)	8	(a)	11	(a)	14	(a)	17	(a)
3	(c)	6	(d)	9	(a)	12	(a)	15	(d)		

MULTIPLE CORRECT CHOICE TYPE

	ANSWER KEY													
1.	(a,b,d)	5.	(a,b,d)	9.	(a,b,d)	13.	(a,b,d)	17.	(c,d)	21.	(a,b,d)			
2.	(a,c,d)	6.	(a,b,d)	10.	(a,b,d)	14.	(a,d)	18.	(b,c)	22.	(b,c)			
3.	(a,b)	7.	(a,c,d)	11.	(c,d)	15.	(b,c,d)	19.	(a,c)	23.	(a,b,c)			
4.	(b,c,d)	8.	(b,c,d)	12.	(a,b,c)	16.	(a,b,c)	20.	(b,d)					

E

D

MATRIX-MATCH TYPE \equiv

- 1. A-p, q; B-r, s; C-r, s; D-s
- 3. A-r; B-s; C-p, q; D-p, q
- 5. A-q, r; B-q; C-p; D-q, s
- 7. A-s; B-q, r, s; C-p; D-r
- 9. A-r, s; B-r; C-p, q; D-p, q

- 2. A-q, r; B-p, s; C-p, r; D-p, r
- 4. A-p; B-q; C-s; D-r
- 6. A-p, s; B-p, s; C-p, q, s; D-p, q, s
- 8. A-q, s; B-q, r; C-q, s; D-p, r
- 10. A-p, q, r; B-p; C-q, s; D-q

A 📃 SINGLE CORRECT CHOICE TYPE 🗄

1. (b) LiAlH_4 does not reduce $-\text{NO}_2$ group to $-\text{NH}_2$ group.

2. (c)
$$\begin{pmatrix} & & \\ N & N \\ Cl^{-}H_{3}H_{3}Cl^{-} & H \end{pmatrix}$$
 +NH₄Cl

- (b) Amalgamated -Zn and HCl reduces carbonyl group to methylene group without affecting the double bond. However, Zn and HCl reduces -NO₂ to -NH₂ group.
- 4. (a) In amines, N is sp^3 hybridised and thus has pyramidal shape. In the given structure, since the three alkyl groups are different, and the fourth corner of the pyramid is occupied by lone pair of electrons, the molecule is chiral. However, the two enantiomers of the amine are not resolvable because of their rapid interconversion through a transition state having planar structure (sp^2 hybridised nitrogen)
- 5. (d) The basic character of an amine in water is determined by (i) electron availability on the N atom and (ii) the extent of stabilization of the cation (protonated amine) due to solvation by hydrogen bonding

- 6. (a) In presence of chlorobenzene, hydrogen bonding is not possible between the protonated amine and the solvent and thus the stabilization factor (solvation effect) is absent. Hence basicity is explained on the basis of the number of electron releasing groups in an amine.
- 7. (a) Note the point of difference in the given compounds which here lies at β -carbon. In I, II, III, the β -carbon atoms are sp^3 , sp^2 and sp hybridised respectively which in turn cause the difference in their *s* character. We know that more is the *s* character of an atom, greater will be its electron-withdrawing nature. Thus sp (50% *s* character) hybridised carbon is most electron-withdrawing, while sp^3 (25% *s* character) is least electron-withdrawing. Further, we know that presence of an electron-withdrawing group decreases basicity of an amine. Thus

$$\frac{\beta}{\beta} \frac{\beta}{NH_2} > \frac{\beta}{\beta} \frac{NH_2}{NH_2} > \frac{\beta}{\beta} \frac{NH_2}{NH_2}$$
I ($\beta - C sp^3$ hybridised) II ($\beta - C sp^2$) III ($\beta - C sp$)

8. (d) Here again the two amines differ in the nature of β -carbon atom



9. (b) Greater the delocalisation of electron pair on N, lesser is its availability for protonation leading to lesser basic character.



10. (d) II is most basic because delocalisation of electron pair leads to negative charge on N making it electron-rich and hence liable to be attacked by proton very easily. Moreover, the corresponding protonated species is very much stable because of equivalent contributing structures.



Species III is least basic because of delocalisation of electron pair on N, making it less available for protonation. Species I lies in mid-way, thus



- 12. (d) In presence of $AlCl_3$ (a Lewis acid), aniline is converted into anilinium cation, which being *m*-directing gives *m*-aminoacetophenone.
- (c) Although aliphatic 1° amines form diazonium salts, these are unstable and decompose to alkyl carbocation which can form variety of normal as well as rearranged products.

$$\operatorname{CH}_{3}\operatorname{CH}_{2}\operatorname{CH}_{2}\operatorname{NH}_{2} \xrightarrow{\operatorname{HONO}} \left[\operatorname{CH}_{3}\operatorname{CH}_{2}\operatorname{CH}_{2}\operatorname{N}_{2}^{+}\operatorname{Cl}^{-} \right] \rightarrow \operatorname{CH}_{3}\operatorname{CH}_{2}\overset{+}{\operatorname{C}}\operatorname{H}_{2} + \operatorname{N}_{2} + \operatorname{Cl}^{-}$$

$$\underbrace{\operatorname{Unstable}} \xrightarrow{\operatorname{HONO}} \left[\operatorname{CH}_{3}\operatorname{CH}_{2}\overset{+}{\operatorname{CH}} \operatorname{H}_{2} + \operatorname{N}_{2} + \operatorname{Cl}^{-} \right] \xrightarrow{\operatorname{HONO}} \left[\operatorname{CH}_{3}\operatorname{CH}_{2}\overset{+}{\operatorname{CH}} \operatorname{H}_{2} + \operatorname{N}_{2} + \operatorname{Cl}^{-} \right]$$

$$CH_{3}CH_{2}CH_{2}OH \xleftarrow{H_{2}O}{CH_{3}CH_{2}} \xrightarrow{+}{CH_{3}} CH_{2} \xrightarrow{+}{CH_{3}} CH_{3} \xrightarrow{+}{CH_{3}} CH_{3} \xrightarrow{-}{Cl^{-}} CH_{3}^{+}CH_{3}^{+}CH_{3}^{-}CH_{3}$$

14. (b) Couplings of arenediazonium cations and phenols take place most rapidly in slightly alkaline medium because

- (i) acidic medium (pH < 7) will suppress the ionisation of C_6H_5OH to the more reactive $C_6H_5O^-$, while the alkaline medium (pH > 7) will enhance ionisation of phenol to phenoxide ion.
- (ii) strongly alkaline medium (pH > 10) causes the arenediazonium salt to react with the OH⁻ ion to form a relatively unreactive diazohydroxide or diazoate ion.

$$Ar\overset{+}{N} \equiv \overset{OH}{\underset{\text{Diazohydroxide}}{}} Ar - \overset{OH}{\underset{\text{Diazote ion}}{}} Ar - \overset{OH}{\underset{\text{Diazote ion}}{} Ar - \overset{OH}{\underset{\text{Diazote ion}}{}} Ar - \overset{OH}{\underset{\text{Diazote ion}}{} Ar - \overset{OH}$$

(b) Reaction is an example of Schmidt reaction in which carboxylic acids are heated with hydrazoic acid in presence of a mineral acid to form primary amines.

$$RCOOH + HN_3 \xrightarrow{conc. H_2SO_4} RNH_2 + CO_2$$

16. (c) Nitrosonium ion, NO from HONO is a weak electrophile, hence it can attack only on highly activated benzene nucleus, provided proper position, *p*- or *o*- is free.



- 17. (d) LiAlH_4 does not reduce $-\text{NO}_2$ group.
- 18. (d) Most of the simple amines those having smaller alkyl (groups) can't be resolved, although they may contain three different alkyl groups.
- (b) Hydrogen bonding is a strong intermolecular attraction between an electrophilic O-H or N-H hydrogen atom and a pair of non-bonding electrons. Thus a hydrogen bond requires both a hydrogen bond donor (a molecule having O-H or N-H group) and a hydrogen bond acceptor (a molecule having lone pair of electrons). Thus dimethyl amine can serve as hydrogen bond donor due to N-H group as well as hydrogen bond acceptor due to lone pair of electrons on N. Such molecules can form hydrogen bond with themselves and also with water (or other hydrogen bond donor). On the other hand, trimethyl amine can serve only as hydrogen bond acceptor since it has lone pair of electrons but not hydrogen on N. Hence it can form hydrogen bond with water but not with themselves.



- 20. (d) Nearly all amines have unpleasant odour, while their salts are odourless, Amines having more than 6 carbon atoms are insoluble in water while their salts are soluble in water, hence the latter can be easily converted into syrup. Amines are susceptible to oxidation because of presence of lone pair of electrons on N.
- **21.** (b) Cold dil. NaOH does not attack to either of the compound, while cold dil. HCl reacts only with benzyl amine $C_6H_5CH_2NH_2$.
- 22. (a) The four compounds differ in two respects : Compounds II has sp^3 hybridised N, compounds I and III have sp^2 while compound IV has sp hybridised N. Now we know that greater the *s* character of an orbital, more tightly its electrons are held and hence lesser will be their availability for protonation causing weak basic character. Thus the basic character of the three N's is $sp^3N > sp^2 N > sp N$. However, in pyrrole electrons on N are part of aromatic sextet, i.e., these are delocalised and hence lesser available for protonation and thus pyrrole is a weaker base than pyridine because in pyridine nonbonding electrons present in sp^2 orbital do not form a part of aromatic sextet. Thus the basic character should be in the following order.



23. (b) Explained in the above question.



- (b) This is an example of Hofmann elimination which generally takes place by E2 mechanism and the latter requires a strong base (recall that OH⁻ is a strong base than I⁻). The NH₂⁻, being a strong base, can't be eliminiated easily.
- 26. (d) All of the three are oxidation produces of a 1° amine.



- 27. (c) Cope elimination is used for removing a nitrogen present outside the ring, while Hofmann elimination is used for removing nitrogen present inside as well as outside the ring.
- 28. (d) All the three react with HONO forming following respective products.

 $C_{6}H_{5} \stackrel{+}{N} \equiv NCl^{-}$ Diazonium chloride $C_{6}H_{5} \stackrel{\cdots}{N} - \stackrel{\cdots}{N} = O$ $C_{H_{3}}$ 2° N-Nirosoamine

p-Nitrosodimethylamine

NO

 $V(CH_3)_2$



- **30.** (c) Pyridine resembles a strongly deactivated benzene, here the deactivation is due to electron-withdrawing effect of the electrongative N atom from 2-and 4 positions making 3-positions to be more vulnerable for the attack of an electrophile. Remember that the nonbonding electrons on N are perpendicular to the π -system (they are present in sp^2 orbital), these are not involved in delocalisation, i.e., they can't stabilize the positively charged intermediate.
- 31. (a) Due to greater electronegativity of N over S, positive charge on N will make the methyl groups more electron- deficient than that on positively charged S. Therefore, $(CH_3)_4 N^+ I^-$ will undergo nucleophilic substitution more readily than the sulphur analogue.

$$HO^{-} + CH_{3} \rightarrow N^{+}(CH_{3})_{3} \longrightarrow CH_{3}OH + (CH_{3})_{3}N$$

32. (b) It is an example of Beckmann rearrangment ; where the group *anti* to –OH migrates

NOH NR NHR
$$O_2N$$

 $\| R - C.CH_3 \xrightarrow{Conc.} HOCCH_3 \xrightarrow{} O = CCH_3$; where $R =$

33. (d) For this one should remember that Hofmann's degradation involves migration to electron deficient nitrogen atom, hence the alkyl group will migrate with its bonding pair of electrons.

34. (c) Remember that in Hofmann rearrangement, the two original H atoms of the $-\text{CONH}_2$ group are removed by base (OH⁻) and new H's are derived from H₂O.

$$(CH_3)_3C - \overset{O}{\overset{\parallel}{C}} - NH_2 \xrightarrow{(i)OD^-/Br_2} (CH_3)_3C - ND_2$$

- **35.** (d) $\text{HCCl}_3 \xrightarrow{\text{OH}^-} : \overline{C} \text{Cl}_3 \xrightarrow{-\text{Cl}^-} : \frac{\text{CCl}_2}{\text{Dichlorocarbene}}$ (a neutral electron deficient species)
- 36. (c) The given amine has three β -hydrogens, hence it can form three alkenes of which least substituted (H₂C=CH₂) will be the major product (Hofmann product or Hofmann elimination).

$$\begin{array}{c} & \beta_2 & \beta_1 \\ & & \beta_3 & (i) CH_3I \\ & & (ii) Ag_2O \\ & & (iii) heat \end{array}$$

$$\begin{array}{c} & & (from \beta_1) \end{array} + \begin{pmatrix} from \beta_2 \end{pmatrix} + H_2C = CH_2 \\ & (from \beta_3) \end{array}$$

In Hofmann elimination H atom (in the form of proton) is eliminated from that β carbon atom which has maximum number of H atom(s). This is due to the fact that Hofmann elimination (an example of E2 reaction) requires anti-coplanar arrangement of the β H atom and the leaving group. Further, higher the chances for this arrangement to remain in staggered conformation more will be the ease of elimination. Thus greater the number of H atom at β position, higher will be the chances for Me₃N⁺– and H atom to remain in this particular conformation (anti-coplanar arragnement of Me₃N⁺– and H atom in staggered conformation).

37. **(b)**
$$\bigwedge_{H}$$
 $CH_3 \xrightarrow{(i) CH_3 I}_{H_3C} \alpha \bigwedge_{H}^{\beta} M \xrightarrow{(i) Ag_2 O}_{H_3C} M \xrightarrow{(i) Ag$

38. (d) The reaction involves Cope eliminiation (heating of a 3° amine oxide to form an alkene with the elimination of a 2° hydroxylamine).



42. (d) All the three reactions lead to the formation of C = C bonds under usual conditions.

43. (c) (a)
$$\mathbb{R} - (-N_{H_{2}} - \frac{H_{2} \cdot /N_{2}OH}{M_{2} - N_{H_{2}}} \mathbb{R} - NH_{2}$$
 (Hofmann reaction)
(b) $\mathbb{R} - (-N_{3} - \frac{h_{end}}{M_{2}} - \mathbb{R} - NH_{2}$ (Currius reaction)
Axis
(c) $\mathbb{R} - (-OH + HN_{3} - \frac{H_{2}SO_{4}}{M_{2}} - \mathbb{R} - NH_{2}$ (Losen reaction)
Hydroanic acid
44. (b) $\mathbb{R} - N - \mathbb{C} = O$ (Albyl isocymate) is the common intermediate in the four reactions.
45. (c) $\mathbb{R} - (-OH + HN_{3} - \frac{H_{2}SO_{4}}{M_{2}} - \mathbb{R} - NH_{2}$ (Losen reaction)
Hydroanic acid
44. (c) $\mathbb{R} - (-N_{3})H_{2} - (\frac{H_{3}}{H_{3}} - (\mathbb{C}H_{3})H_{2} - (\mathbb{C}H_{3})H_{2} - (\mathbb{C}H_{3})N + (\mathbb{H}_{3})H_{3} - (\mathbb{H}$

COMPREHENSION TYPE

- 1. (b) E2 eliminations are concreted and *anti*-stereoselective.
- 2. (d) Due to the bulk of the $-N^+Me_3$ group, the OH⁻ preferentially approaches and removes the least hindered β -hydrogen and thus form the least substituted alkene as the major product (difference from Saytzeff product).
- 3. (b) Species II (acylnitrene) is electron-deficient (N has only 6 electrons), hence it has a tendency to get its octet completed by the migration of alkyl group.
- 4. (c) -OCH₃ is more electron-releasing hence when the migrating aryl group has -OCH₃ in the para postion, its migration is accelerated.
- 5. (a) The migrating group (the group attached to C of the $-CONH_2$ part) never becomes free, so optical activity of the reactant is maintained.
- 6. (c) Since phenyl group (migrating group) does not become free, no cross product [i.e., (ii) and (iv)] is formed.



12. (d)
$$C_6H_5 - N = C = O \xrightarrow{C_2H_5OH} C_6H_5 - NHCOOC_2H_5$$

Urethane

13. (c)
$$C_6H_5 - C - N \xrightarrow{+} N \equiv N : \xrightarrow{-N_2} C_6H_5 - \underbrace{C - N :}_{Nitrene} \xrightarrow{O} C_6H_5 = N - C_6H_5$$

C

Reasoning Type

- 1. (c) The correct reason is : $-SO_3H$ group, being more acidic than $-CO_2H$ group, can easily transfer a proton to the amino group.
- 2. (c) The correct reason is : Acetylation decreases the electron-density in the benzene ring thereby preventing oxidation.

- 3. (c) The correct reason is : The overall electron deficiency in *m*-nitroaniline is much less (due to -R-effect of NO₂ group and +R-effect of NH₂ group) than in *m*-dinitrobenzene (-R-effect of the two NO₂ groups) and hence does not accept additional electrons from a weak reducing agent such as (NH₄)₂S and thus further reduction is prevented.
- 4. (d) Aryl halides (chlorobenzene) do not undergo nucleophilic substitution with KCN because of the low reactivity of the Cl atom, which is because of resonance in chlorobenzene. So assertion is wrong. Reason is correct.
- 5. (a) **R** is the correct explanation of **A**.
- 6. (d) In strongly acidic conditions, aniline becomes protonated with the result lone pair of electrons is not available to produce

+E and +M effects. On the other hand, the $-\overset{\oplus}{\mathrm{NH}_3}$ group exerts strong –I effect causing deactivation of the ring.

- 7. (c) Correct R : Only primary aliphatic amines can be prepared by Gabriel phthalimide reaction.
- 8. (a) **R** is the correct explanation of **A**.
- 9. (a) **R** is the correct explanation of **A**.

D MULTIPLE CORRECT CHOICE TYPE **=**

- (a,b,d) Tertiary amino group is a strong activator, nitration can be carried out under milder conditions (HNO₃ + CH₃COOH). In Tertiary amines, electron pair on N is shielded by two bulky methyl groups, so these are not easily oxidised by mild oxidising agents.
- (a,c,d) Diazomethane is used for methylating acidic groups ; compound IV has enolic —OH group, hence it can also be methylated by CH₂N₂.
- 3. (a,b) For the preparation of Me_3CNH_2 , the required alkyl halide is Me_3CX which will react with potassium phthalimide, a strong base, to form alkene rather than substituted product. For preparing $C_6H_5NH_2$, C_6H_5Cl will be the starting halide in which Cl is non-reactive.
- 4. (b,c,d) Vinyl and aryl halides do not undergo S_{N^2} reactions, unless activated by electron withdrawing group (in the *o* and *p*-position in case of aryl halides). 3° Halides form alkenes on elimination.
- 5. (a,b,d) In reductive amination method, products can have only CH_3 , a 1°C, or 2°C bonded to N as in CH_3NH_2 , RCH_2NH_2 or R_2CHNH_2 respectively. In the reduction of oxime, amine formed can have only 1° or 2°C bonded to N. In Gabriel phthalimide method, the required Me₃CCl would undergo elimination (E2) rather than substitution (S_{N^2}).
- 6. (a,b,d) Isonitriles (C_6H_5NC) on reduction give 2° amines ($C_6H_5NHCH_3$). All other three methods give aniline.
- 7. (a,c,d) In p-NH₂C₆H₄COOH, —COOH group is very weak so it can't transfer H⁺ to the weakly basic amino group. All other three form zwitterions.
- 8. (b,c,d) In $C_6H_5OCH_3$, $-OCH_3$ does not sufficiently increase electron density on the ring. Recall that C_6H_5OH undergoes coupling in weakly alkaline medium which converts C_6H_5OH to the more reactive $C_6H_5O^-$. In options (*b*) and (*c*), presence of electron-withdrawing $-NO_2$ groups increases electrophilic character to such an extent that these diazonium cations can couple even with the compounds having weak electron-releasing groups. Option (d) undergoes coupling reaction easily because $-NMe_2$ is sufficiently electron-releasing.
- **9.** (a,b,d) In *p*-aminophenol all the four positions (2, 3, 5 and 6) can be coupled (positions 2 and 6 in presence of H⁺ and positions 3 and 5 in presence of OH⁻).
- 10. (a,b,d) *tert*-Nitro compounds (Me₃CNO₂) do not react with HONO because they do not have any α -H. The three others react with HONO as usual.

$$\begin{array}{ccc} & & & & & & NO_2 \\ & & & & \\ CH_3 - CHCH_2CH_3 & \longrightarrow & CH_3 - CH_2CH_2CH_3 \\ & & & & \\ 2-Nitrobutane & & & & \\ NO \end{array}$$

Pseudonitrol (blue)

11. (c,d) $-SO_3H$ group present in *o*- and *p*-positions are easily replaced.

12. (a,b,c)
$$C_6H_5 \overset{\cdots}{N}H_2 + AlCl_3 \longrightarrow C_6H_5 \overset{+}{N}H_2 \xrightarrow{CH_3Cl} No reaction$$

substituent $(-\overset{+}{N}H_2 \overrightarrow{AlCl_3})$
a meta director

However, Friedel-Crafts reactions are the most sluggish electrophilic aromatic substitutions (halogenation occurs most easily), hence if a meta director is present on the ring, the compound does not undergo Friedel-Crafts alkylation/acylation.

MATRIX-MATCH ТҮРЕ

- 1. A-p, q; B-r, s; C-r, s; D-s
 - (A) Hydrogen bodning is possible in *ethyl acetoacetate* (due to the presence of enolic group) and in *proteins* (due to the presence of -CO-NH- grouping).
 - (B) β -Amino acids, like α -amino acids, form *zwitterions* and also undergo *diazotisation* although the diazo salts, so formed, are unstable.
 - (C) p-Amino benzoic acid, like β- and α-amino acids forms zwitterions and being 1° aromatic amine also forms, well-known, diazonium salt.
 - (D) Aniline undergoes diazotisation easily.
- 2. A-q, r; B-p, s; C-p, r; D-p, r



- (A) A is aromatic due to 10π electrons, N₁, N₃ and N₇ are sp^2 hybridised their ℓp present in sp^2 orbital are localised hence undergo protonation. The N₉ is also sp^2 hybridised but its ℓp is in *p*-orbital hence involved in delocalisation, and thus not basic.
- (B) Due to resonance, all the three nitrogens of guanidine are identical. It is not cyclic, hence non-aromatic.
- (C-D) As mentioned above in (A), here the two nitrogens are identical, and the compound is aromatic due to the presence of aromatic sextet.

(A)
$$CHCl_3 \xrightarrow{OH} : CCl_2 + HCl$$
 (α -elimination)

(B)
$$\operatorname{CHCl}_2 - \operatorname{CF}_3 \xleftarrow{\operatorname{OC}_2\operatorname{H}_5^- \text{ fast}} \operatorname{Cl}_2 \operatorname{C}^- \operatorname{CF}_3 \xrightarrow{\operatorname{Slow}} \operatorname{Slow}$$

 $Cl_2C = CF_2$

5.

This reaction, although involves two steps; the rate determining step involves only one molecule hence the reaction is labelled as E1. Further, it is the carbanion (conjugate base of $CHCl_2CF_3$) that determines the rate of reaction, hence it is labelled as E1CB.

(C) When the proton to be removed is on the sterically hindered carbon, and the base also has sterically hindered carbon, less substituted alkene is the main product (*Hofmann elimination*)

$$\begin{array}{c} CH_{3} & CH_{3} \\ | \\ CH_{3} - \overset{}{C} - CH_{2}CH_{3} \xrightarrow{Me_{3}CO^{-}} CH_{2} = \overset{}{C} - CH_{2}CH_{3} \\ | \\ Br \end{array}$$

(D) Elimination from quaternary ammonium ions usually gives the least substituted alkene (*Hofmann elimination*).

 (A) ArS_N¹ reactions are rare, and involve carbocation as intermediate. The inportant example is substitution in aromatic diazonium salts.

$$ArN^+ \equiv N \xrightarrow{slow} Ar^+ + N_2$$

Highly unstable Highly stable

$$Ar^+ + : Nu \xrightarrow{fast} Ar - Nu$$





(C) Higher the bulk of the group present on the nucleophilic centre, lesser is its nucleophilic character.

$$O-H > O-C (CH_3)_3$$

bulkyl group

(D) Stronger the conjugated acid, weaker will be the base.

$$H_2O > NH_3$$

stronger acid weaker acid

A-q, r; B-q; C-p; D-q, s

- (A) Allylic rearrangements involve the migration of the double bond (and the functional group, if present) from one position to other. These reactions proceed through SN¹, SN², SE¹ and SE² mechanism; SN¹ involves carbocations (electron-deficient species), while SE¹ involves carbanion.
- (B) Hofmann rearrangement involves nitrene O

(R - C - N) intermediate in which nitrogen is electron deficient.

- (C) Aldol condensation involves the attack of base on aldehyde or ketone, the reaction will involve the formation of carbanion.
- (D) Carbylamine reaction involves the formation of dichlorocarbene (:CCl₂) as intermediate which is electron deficient.