

Chapter 1

Some Basic Concepts of Chemistry

Solutions

SECTION – A

Objective Type Questions

(Mole Concept)

1. The total number of electrons in 4.2 g of N^{3-} ion is (N_A is the Avogadro's number)
 (1) $2.1 N_A$ (2) $4.2 N_A$ (3) $3 N_A$ (4) $3.2 N_A$

Sol. Answer (3)

$$\text{Total number of moles} = \frac{4.2}{14} = 0.3 \text{ mol}$$

$$1 \text{ mol of } \text{N}^{3-} \text{ have electrons} = 10 \times N_0.$$

$$\therefore \text{Number of } e^- \text{ in } 0.3 \text{ mol} = 0.3 \times 10 \times N_0 = 3 \times N_0$$

2. Suppose that A and B form the compounds B_2A_3 and B_2A . If 0.05 mole of B_2A_3 weighs 9 g and 0.1 mole of B_2A weighs 10 g, the atomic weight of A and B respectively are
 (1) 30 and 40 (2) 40 and 30 (3) 20 and 5 (4) 15 and 20

Sol. Answer (2)

Let the atomic mass of B = y g ; A = x g

In B_2A_3

$$2y + 3x = \text{mol. mass of } \text{B}_2\text{A}_3 = \frac{\text{given weight}}{\text{mole}}$$

$$2y + 3x = \frac{9}{0.05} \text{ g}$$

In B_2A

$$\therefore 2y + x = \frac{10}{0.1} \text{ g}$$

Solving x and y

$$\begin{cases} x = 40 \\ y = 30 \end{cases}$$

3. Number of Fe atoms in 100 g Haemoglobin if it contains 0.33% Fe. (Atomic mass of Fe = 56)
 (1) 0.035×10^{23} (2) 35 (3) 3.5×10^{23} (4) 7×10^8

Sol. Answer (1)

$$\text{Mass of Fe} = 100 \times \frac{0.33}{100} = 0.33 \text{ g}$$

$$\therefore \text{Moles of Fe} = \frac{0.33}{56} = 5.89 \times 10^{-3} \text{ mole}$$

$$\therefore \text{Number of atom of Fe} = 5.89 \times 10^{-3} \times 6.022 \times 10^{23} = 0.035 \times 10^{23} \text{ atom}$$

4. The number of electrons in 1.6 g of CH_4 is approximately

- (1) 25×10^{24} (2) 1.5×10^{24} (3) 6×10^{23} (4) 3.0×10^{24}

Sol. Answer (3)

$$\text{Moles of CH}_4 = \frac{1.6}{16} = 0.1 \text{ mol}$$

$$\begin{aligned} \text{Number of } e^- \text{ of CH}_4 &= 0.1 \times 10 \times N_0 \\ &= 6 \times 10^{23} \end{aligned}$$

5. Specific volume of cylindrical virus particle is $6.02 \times 10^{-2} \text{ cc/gm}$ whose radius and length are $7 \text{ \AA}$ and $10 \text{ \AA}$ respectively. If $N_A = 6.02 \times 10^{23}$, find molecular weight of virus.

- (1) 15.4 kg/mol (2) $1.54 \times 10^4 \text{ kg/mol}$
(3) $3.08 \times 10^4 \text{ kg/mol}$ (4) $3.08 \times 10^3 \text{ kg/mol}$

Sol. Answer (1)

One molecule (gm)

$$d = \frac{\text{M.wt.}}{V}$$

$$\frac{1}{6.02 \times 10^{-2}} = \frac{\text{M.wt.}}{\pi r^2 \times h}$$

$$\text{M.Wt. (One molecule in gm)} = \frac{\pi \times (7 \times 10^{-8})^2 \times 10 \times 10^{-8}}{6.02 \times 10^{-2}}$$

$$\text{M.Wt. (One mole in kg)} = \frac{22}{7} \times \frac{7 \times 7 \times 6.02 \times 10^{-3}}{6.02 \times 10^{-2}} = 2.2 \times 7 = 15.4 \text{ kg mol}^{-1}$$

6. The number of mole of nitrogen in one litre of air containing 10% nitrogen by volume, under standard conditions, is

- (1) 0.03 mole (2) 2.10 moles (3) 0.186 mole (4) 4.46×10^{-3} mole

Sol. Answer (4)

$$\text{Volume of N}_2 \text{ in 1 L i.e., 1000 ml of N}_2 = \frac{10}{1000} \times 1000 = 100 \text{ ml}$$

$$22400 \text{ ml at STP} = 1 \text{ mol.}$$

$$\therefore 100 \text{ ml at STP} = \frac{1}{22400} \times 100 = \frac{1}{224} = 4.46 \times 10^{-3} \text{ mol}$$

7. Number of significant figures in 6.62×10^{-34} .

- (1) Two (2) Three
(3) Four (4) One

Sol. Answer (2)


Na_2CO_3 is soda ash which does not decompose upon heating even to redness.

$\therefore \text{CO}_2$ will not be evolved.

12. One litre of CO_2 is passed through red hot coke. The volume becomes 1.4 litres at same temperature and pressure. The composition of products is

- (1) 0.8 litre of CO_2 and 0.6 litre of CO
 (2) 0.7 litre of CO_2 and 0.7 litre of CO
 (3) 0.6 litre of CO_2 and 0.8 litre of CO
 (4) 0.4 litre of CO_2 and 1.0 litre of CO

Sol. Answer (3)



Initial	1L	0	0
Final volume	(1 - x)		2x

$$\text{Final volume} = 1 - x + x + 2x = 1.4 \text{ L}$$

$$\therefore \boxed{x = 0.4 \text{ L}}$$

$$\therefore \text{Volume of CO} = 2x = 2 \times 0.4 = 0.8 \text{ L}$$

$$\text{Volume of CO}_2 = (1 - x) = 1 - 0.4 = 0.6 \text{ L}$$

13. An organic compound containing C and H gave the following analysis C = 40%, H = 6.7%. Its empirical formula would be

- (1) CH_4 (2) CH_2O
 (3) $\text{C}_2\text{H}_4\text{O}_2$ (4) C_2H_4

Sol. Answer (2)


	% age	Atomic mass	Moles	Simple ratio
C	40%	12	$\frac{40}{12} = 3.33$	1
H	6.7%	1	$\frac{6.7}{1} = 6.7$	2
O	53.3%	16	$\frac{53.3}{16} = 3.33$	1

$$\therefore \boxed{\text{EF} = \text{CH}_2\text{O}}$$

14. How many litre of oxygen at STP is required to burn 60 g C_2H_6 ?

- (1) 22.4 L (2) 11.2 L (3) $22.4 \times 7 \text{ L}$ (4) 8.5 L

Sol. Answer (3)




For 2 moles of C_2H_6 = 7 moles of O_2 required.

$$\boxed{\text{i.e., for 60 g of } \text{C}_2\text{H}_6 = 7 \times 22.4 \text{ L of } \text{O}_2 \text{ at STP required}}$$

15. For the formation of 3.65 g of HCl gas, what volume of hydrogen gas and chlorine gas are required at NTP conditions?

- (1) 1 L, 1 L (2) 1.12 L, 2.24 L (3) 3.65 L, 1.83 L (4) 1.12 L, 1.12 L

Sol. Answer (4)

$$1 \text{ mol} \quad 1 \text{ mol} \quad 36.5 \text{ g} \times 2$$

$$22.4 \text{ L} \quad 22.4 \text{ L} \quad 36.5 \text{ g} \times 2$$

$$1.12 \text{ L} \quad 1.12 \text{ L} \quad 3.65 \text{ g}$$

For (36.5×2) g of HCl volume of H_2 and Cl_2 required will be 22.4 L H_2 and 22.4 L of Cl_2

$$\therefore \text{ For } 3.65 \text{ g and } 1.12 \text{ L of } \text{H}_2 \left\{ \begin{array}{l} 1.12 \text{ L of } \text{Cl}_2 \end{array} \right\}$$

16. The crystalline salt $\text{Na}_2\text{SO}_4 \cdot x\text{H}_2\text{O}$ on heating loses 55.9% of its mass and becomes anhydrous. The formula of crystalline salt is

**Sol.** Answer (4)

17. A certain amount of a metal whose equivalent mass is 28 displaces 0.7 L of H_2 at S.T.P. from an acid hence mass of the element is

(1) 1.75 g

(2) 0.875 g

(3) 3.50 g

(4) 7.00 g

Sol. Answer (1)

Weight of metal which can displace 11.2 L of H_2 gas is equivalent mass.

$$\therefore 11.2 \text{ L of } \text{H}_2 (\text{g}) \text{ have mass} = 28 \text{ g}$$

$$0.7 \text{ L of } \text{H}_2 \text{ have mass} = \frac{28}{11.2} \times 0.7 = 1.75 \text{ g}$$

(Reactions in Solutions)

18. When 100 ml of $\frac{M}{10} \text{H}_2\text{SO}_4$ is mixed with 500 ml of $\frac{M}{10} \text{NaOH}$ then nature of resulting solution and normality of excess of reactant left is

(1) Acidic, $\frac{N}{5}$

(2) Basic, $\frac{N}{5}$

(3) Basic, $\frac{N}{20}$

(4) Acidic, $\frac{N}{10}$

Sol. Answer (3)

100 ml of $\frac{M}{10} \text{H}_2\text{SO}_4$ ↓ $\frac{N}{5} \text{H}_2\text{SO}_4$ [because $N = M \times n$, n factor of $\text{H}_2\text{SO}_4 = 2$]	500 ml of $\frac{M}{10} \text{NaOH}$ ↓ $\frac{N}{10}$ [because N factor is 1]
$\therefore \text{ meq of } \text{H}_2\text{SO}_4 = \left(100 \times \frac{1}{5} \right) = 20 \text{ meq}$	$\therefore \text{ meq of } \text{NaOH} = \left(500 \times \frac{1}{10} \right) = 50 \text{ meq}$

$$\begin{aligned}\text{For Neutralisation Reaction} &= \frac{\text{larger meq} - \text{smaller meq}}{\text{Total volume}} \\ &= \frac{50 - 20}{600} = \frac{1}{20} \text{ N NaOH} \quad [\text{because larger meq of NaOH will remain}]\end{aligned}$$

∴ Solution will be basic

19. Mole fraction of solvent in aqueous solution of NaOH having molality of 3 is

- (1) 0.3 (2) 0.05 (3) 0.7 (4) 0.95

Sol. Answer (4)

$$m = \frac{1000 \cdot x_B}{x_A \cdot M_A} \quad \left\{ \begin{array}{l} m = \text{molality} \\ x_B = \text{molality fraction of solute} \\ x_A = \text{molality fraction of solvent} \end{array} \right.$$

$$\boxed{x_A + x_B = 1}$$

$$\therefore x_A = (1 - x_B)$$

$$m = \frac{1000 \cdot x_B}{(1 - x_B) M_A}$$

Putting $m = 3$

$M_A = 18$ because aqueous solution is present

$$\begin{aligned}3 &= \frac{1000 \cdot x_B}{(1 - x_B) 18} \Rightarrow 54 (1 - x_B) = 1000 x_B \\ &= 54 - 54 x_B = 1000 x_B\end{aligned}$$

$$x_B = \frac{54}{1054} \Rightarrow x_B = 0.05$$

$$\therefore \boxed{x_A = (1 - x_B) = (1 - 0.05) = 0.95}$$


20. Concentrated aqueous sulphuric acid is 98% H_2SO_4 by mass and has a density of 1.80 g mL^{-1} . Volume of acid required to make one litre of $0.1 \text{ M H}_2\text{SO}_4$ solution is

- (1) 16.65 mL (2) 22.20 mL (3) 5.55 mL (4) 11.10 mL

Sol. Answer (3)

Molarity of 98% H_2SO_4 by mass having density 1.80 g/mL will be

$$M = \frac{\% \text{ w/w} \times d \times 10}{\text{M. mass}} = \frac{98 \times 1.80 \times 10}{98} = 18 \text{ M}$$



Applying $M_1 V_1 = M_2 V_2$

$$18 \times V_1 = 1000 \times 0.1$$

$$V_1 = \frac{100}{18} = 5.55 \text{ ml}$$

21. Ammonia gas is passed into water, yielding a solution of density 0.93 g/cm^3 and containing 18.6% NH_3 by weight. The mass of NH_3 per cc of the solution is

- (1) 0.17 g/cm^3
- (2) 0.34 g/cm^3
- (3) 0.51 g/cm^3
- (4) 0.68 g/cm^3

Sol. Answer (1)

$$\text{Molarity of } \text{NH}_3 \text{ solution} = \frac{18.6 \times 0.93 \times 10}{17} = 10.17 \text{ M}$$

$$\begin{aligned}\text{Strength (g/L)} &= \text{molarity} \times \text{mol. mass} \\ &= 10.17 \times 17 = 172.9 \text{ g/L}\end{aligned}$$

i.e., 1000 mL of solution contain $\text{NH}_3 = 172.9 \text{ g}$

$$1 \text{ mL or } 1 \text{ cm}^3 \text{ of solution contain } \text{NH}_3 = \frac{172.9}{1000} = 0.172 \text{ g} \approx 0.17 \text{ g}$$

22. 6.025×10^{20} molecules of acetic acid are present in 500 mL of its solution. The concentration of solution is

- (1) 0.002 M
- (2) 10.2 M
- (3) 0.012 M
- (4) 0.001 M

Sol. Answer (1)

$$\text{Moles of oxalic acid} = \frac{6.022 \times 10^{20}}{6.022 \times 10^{23}} = 10^{-3} \text{ moles}$$

$$\text{Molarity} = \frac{10^{-3}}{500} \times 1000 = 2 \times 10^{-3} \text{ M} = 0.002 \text{ M}$$

(Equivalent Mass)

23. 74.5 g of metal chloride contains 35.5 g of chlorine. The equivalent weight of metal is

- (1) 74.5
- (2) 39
- (3) 35.5
- (4) 7.45

Sol. Answer (2)

g eq. of metal = g eq. of chlorine

$$\frac{74.5 - 35.5}{E_M} = \frac{35.5}{35.5}$$

$$\therefore \text{eq. wt. of metal} = 39$$

24. Equivalent weight of crystalline oxalic acid is

- (1) 90
- (2) 63
- (3) 53
- (4) 45

Sol. Answer (2)

Crystalline oxalic acid = $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$

$$\text{Eq. wt.} = \frac{126}{2} = 63$$

25. Equivalent mass of H_3PO_4 (Molar mass = M) for the given reaction, $\text{H}_3\text{PO}_4 + \text{Ca}(\text{OH})_2 \rightarrow \text{CaHPO}_4 + 2\text{H}_2\text{O}$ is

- (1) $\frac{M}{1}$ (2) $\frac{M}{2}$ (3) $2M$ (4) $\frac{M}{4}$

Sol. Answer (2)

$$\text{Eq. wt.} = \frac{\text{Molar mass}}{n\text{-factor}} = \frac{M}{2}$$

26. Equivalent mass of a metal is 12. Hence, equivalent mass of its oxide is

- (1) 24 (2) 28 (3) 20 (4) 34

Sol. Answer (3)

$$\text{Eq. wt. of metal oxide} = E_{\text{metal}} + E_{\text{oxygen}} = 12 + 8 = 20$$

27. 4 g of a metal oxide contains 1.6 g oxygen, then equivalent mass of the metal is

- (1) 3.2 (2) 24 (3) 12 (4) 20

Sol. Answer (3)

g eq. of metal = g eq. of oxygen.

$$\frac{4 - 1.6}{E_M} = \frac{1.6}{8}$$

eq. wt. of metal = 12


SECTION – B

Previous Years Questions

1. A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc. H_2SO_4 . The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be **[NEET-2018]**

- (1) 1.4 (2) 3.0 (3) 4.4 (4) 2.8

Sol. Answer (4)



Gaseous mixture formed is CO and CO_2 when it is passed through KOH, only CO_2 is absorbed. So the remaining gas is CO.

So, weight of remaining gaseous product CO is

$$\frac{2}{20} \times 28 = 2.8\text{g}$$

So, the correct option is (4)

2. In which case is number of molecules of water maximum?

[NEET-2018]

- (1) 18 mL of water (2) 0.18 g of water
(3) 10^{-3} mol of water (4) 0.00224 L of water vapours at 1 atm and 273 K

Sol. Answer (1)

$$\text{Mass of water} = 18 \times 1 = 18 \text{ g}$$

$$\begin{aligned}\text{Molecules of water} &= \text{mole} \times N_A = \frac{18}{18} N_A \\ &= N_A\end{aligned}$$

$$\begin{aligned}(2) \text{ Molecules of water} &= \text{mole} \times N_A = \frac{0.18}{18} N_A \\ &= 10^{-2} N_A\end{aligned}$$

$$(3) \text{ Molecules of water} = \text{mole} \times N_A = 10^{-3} N_A$$

$$(4) \text{ Moles of water} = \frac{0.00224}{22.4} = 10^{-4}$$

$$\text{Molecules of water} = \text{mole} \times N_A = 10^{-4} N_A$$

3. Suppose the elements X and Y combine to form two compounds XY_2 and X_3Y_2 . When 0.1 mole of XY_2 weighs 10 g and 0.05 mole of X_3Y_2 weighs 9 g, the atomic weights of X and Y are **[NEET-Phase-2-2016]**

- (1) 40, 30 (2) 60, 40 (3) 20, 30 (4) 30, 20

Sol. Answer (1)For XY_2 ,

$$\therefore 0.1 \text{ mole } XY_2 \equiv 10 \text{ g}$$

$$\therefore 1 \text{ mole } XY_2 \equiv 100 \text{ g}$$

$$\text{and } X + 2Y = 100 \quad \dots(i)$$

For X_3Y_2 ,

$$\therefore 0.05 \text{ mole } X_3Y_2 \equiv 9 \text{ g}$$

$$\therefore 1 \text{ mole } X_3Y_2 \equiv 180 \text{ g}$$

$$\text{and } 3X + 2Y = 180 \quad \dots(ii)$$

On solving,

$$X = 40$$

$$\text{and } Y = 30$$

4. What is the mass of the precipitate formed when 50 mL of 16.9% solution of $AgNO_3$ is mixed with 50 mL of 5.8% $NaCl$ solution? ($Ag = 107.8$, $N = 14$, $O = 16$, $Na = 23$, $Cl = 35.5$) **[Re-AIPMT-2015]**

- (1) 7 g (2) 14 g
(3) 28 g (4) 3.5 g

Sol. Answer (1)

$$n_{AgNO_3} = \frac{50 \times 16.9}{100 \times 169.8} \approx 0.05 \text{ Mole}$$

$$n_{NaCl} = \frac{50 \times 5.8}{100 \times 58.5} \approx 0.05 \text{ Mole}$$



$$t = 0; \quad 0.05 \text{ mole} \quad 0.05 \text{ mole} \quad 0$$

$$t = t; \quad 0 \quad 0 \quad 0.05 \text{ mole}$$

$$\therefore \text{Mass of } AgCl = 0.05 \times 143.3 = 7.16$$

$$\approx 7 \text{ g}$$

5. If Avogadro number N_A , is changed from $6.022 \times 10^{23} \text{ mol}^{-1}$ to $6.022 \times 10^{20} \text{ mol}^{-1}$, this would change

- (1) The ratio of chemical species to each other in a balanced equation
- (2) The ratio of elements to each other in a compound
- (3) The definition of mass in units of grams
- (4) The mass of one mole of carbon

[Re-AIPMT-2015]

Sol. Answer (4)


Fact.

6. 20.0 g of a magnesium carbonate sample decomposes on heating to give carbon dioxide and 8.0 g magnesium oxide. What will be the percentage purity of magnesium carbonate in the sample? (At. wt. : Mg = 24)

[Re-AIPMT-2015]

- (1) 60 (2) 84 (3) 75 (4) 96

Sol. Answer (2)



$$\therefore x = \frac{84 \times 8}{40} = 16.8 \text{ g}$$

$$\therefore \% \text{ purity of MgCO}_3 = \frac{16.8}{20} \times 100 = 84\%$$

7. A mixture of gases contains H_2 and O_2 gases in the ratio of 1 : 4 (w/w). What is the molar ratio of the two gases in the mixture?

[AIPMT-2015]

- (1) 2 : 1 (2) 1 : 4 (3) 4 : 1 (4) 16 : 1

Sol. Answer (3)


$$\frac{n_{\text{H}_2}}{n_{\text{O}_2}} = \frac{1/2}{4/32} = \frac{32}{2 \times 4} = \frac{4}{1}$$

8. 1.0 g of magnesium is burnt with 0.56 g O_2 in a closed vessel. Which reactant is left in excess and how much? (At. wt. Mg = 24; O = 16)

[AIPMT-2014]

- (1) Mg, 0.16 g (2) O_2 , 0.16 g (3) Mg, 0.44 g (4) O_2 , 0.28 g

Sol. Answer (1)



$\therefore \text{O}_2$ is limiting reagent

$$\therefore 32 \text{ g of } \text{O}_2 \text{ will react with, } \frac{2 \times 24}{32} \text{ g}$$


$$\therefore 0.56 \text{ g of } \text{O}_2 \text{ will react with, } \frac{2 \times 24 \times 0.56}{32} = 0.84 \text{ g}$$

$$\therefore \text{Excess of Mg} = 1 - 0.84 = 0.16 \text{ g}$$

9. When 22.4 litres of $\text{H}_2(\text{g})$ is mixed with 11.2 litres of $\text{Cl}_2(\text{g})$, each at STP, the moles of $\text{HCl}(\text{g})$ formed is equal to [AIPMT-2014]

- (1) 1 mol of $\text{HCl}(\text{g})$ (2) 2 mol of $\text{HCl}(\text{g})$
 (3) 0.5 mol of $\text{HCl}(\text{g})$ (4) 1.5 mol of $\text{HCl}(\text{g})$

Sol. Answer (1)



$$\frac{n_{\text{Cl}_2}}{1} = \frac{n_{\text{HCl}}}{2} \quad \left| \quad \frac{11.2}{22.4} = \frac{n_{\text{HCl}}}{2} \right.$$

$$\boxed{n_{\text{HCl}} = 1}$$

10. 6.02×10^{20} molecules of urea are present in 100 mL of its solution. The concentration of solution is [NEET-2013]

- (1) 0.01 M (2) 0.001 M (3) 0.1 M (4) 0.02 M

Sol. Answer (1)

$$\text{Moles of urea} = \frac{6.02 \times 10^{20}}{6.02 \times 10^{23}} = 10^{-3} \text{ moles}$$

$$\therefore \text{Molarity} = \frac{\text{moles}}{\text{volume (mL)}} \times 1000 = \frac{10^{-3}}{100} \times 1000 = 10^{-2} \text{ M} = 0.01 \text{ M}$$

11. How many grams of concentrated nitric acid solution should be used to prepare 250 mL of 2 M HNO_3 ? The concentrated acid is 70% HNO_3 ? [NEET-2013]

- (1) 90.0 g conc. HNO_3 (2) 70.0 g conc. HNO_3
 (3) 54.0 g conc. HNO_3 (4) 45.0 g conc. HNO_3

Sol. Answer (4)

$$\therefore M = \frac{n_{\text{HNO}_3}}{V_{\text{mL}}} \times 1000,$$

$$\therefore n_{\text{HNO}_3} = \frac{2 \times 250}{1000} = 0.5 \text{ mole}$$

$$\therefore \text{Mass of concentrated acid required} = \frac{0.5 \times 63 \times 100}{70} = 45 \text{ g}$$

12. Mole fraction of the solute in a 1.00 molal aqueous solution is [AIPMT (Prelims)-2011]

- (1) 1.7700 (2) 0.1770 (3) 0.0177 (4) 0.0344

Sol. Answer (3)

$$\chi_{\text{solute}} = \frac{1}{1 + 55.55} = 0.0177$$

13. Which has the maximum number of molecules among the following? [AIPMT (Mains)-2011]

- (1) 8 g H_2 (2) 64 g SO_2 (3) 44 g CO_2 (4) 48 g O_3

Sol. Answer (1)

Number of molecules in

$$\text{H}_2 = \frac{8}{2} N_A = N_A, \quad \text{SO}_2 = \frac{64}{64} N_A = N_A$$

$$\text{CO}_2 = \frac{44}{44} N_A = N_A, \quad \text{O}_3 = \frac{48}{48} N_A = N_A$$

∴ Maximum number of molecules is present in 8 g H₂

14. The number of atoms in 0.1 mol of a triatomic gas is ($N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$)

[AIPMT (Prelims)-2010]

- (1) 6.026×10^{22} (2) 1.806×10^{23}
 (3) 3.600×10^{23} (4) 1.800×10^{22}

Sol. Answer (2)

$$\text{Number of atoms} = 3 \times 0.1 \times 6.022 \times 10^{23} = 1.806 \times 10^{23}$$

15. 25.3 g of sodium carbonate, Na₂CO₃ is dissolved in enough water to make 250 mL of solution. If sodium carbonate dissociates completely, molar concentration of sodium ion, Na⁺ and carbonate ions, CO₃²⁻ are respectively (Molar mass of Na₂CO₃ = 106 g mol⁻¹)

[AIPMT (Prelims)-2010]


- (1) 0.955 M and 1.910 M (2) 1.910 M and 0.955 M
 (3) 1.90 M and 1.910 M (4) 0.477 M and 0.477 M

Sol. Answer (2)

$$\text{Molarity of Na}_2\text{CO}_3 = \frac{\text{moles of Na}_2\text{CO}_3}{\text{volume (mL)}} \times 1000$$

$$\text{Moles of Na}_2\text{CO}_3 = \frac{25.3}{106}$$

$$\therefore \text{Molarity of Na}_2\text{CO}_3 = \frac{25.3}{106} \times 1000 = \frac{25.3}{106} \times 4 = 0.955 \text{ M}$$



As one mole Na₂CO₃ gives 2 mol Na[⊕] and 1 mol CO₃²⁻


$$\therefore \text{Molarity of Na}^{\oplus} = 2 \times \text{molarity of Na}_2\text{CO}_3 = \boxed{2 \times 0.955 = 1.910 \text{ M}}$$

$$\text{Molarity of CO}_3^{2-} = 1 \times \text{molarity of Na}_2\text{CO}_3 = \boxed{0.955 \text{ M}}$$

16. 10 g of hydrogen and 64 g of oxygen were filled in a steel vessel and exploded. Amount of water produced in this reaction will be

[AIPMT (Prelims)-2009]

- (1) 3 mol (2) 4 mol
 (3) 1 mol (4) 2 mol


Sol. Answer (2)

Amount of water produced = 4 mol

17. How many moles of lead (II) chloride will be formed from a reaction between 6.5g of PbO and 3.2 g of HCl?

[AIPMT (Prelims)-2008]

- (1) 0.029 (2) 0.044
(3) 0.333 (4) 0.011

Sol. Answer (1)

As 73 g HCl reacts with 224 g of PbO

$$\therefore \text{On reaction for 3.2 g of HCl with PbO the required amount of PbO} = \frac{224}{73} \times 3.2 = 9.81 \text{ g}$$

But only 6.5 g PbO is present

 $\therefore$ PbO will be the LR and calculation will be according to PbO.224 g of PbO gives = 1 mol PbCl₂

$$6.5 \text{ g of PbO gives} = \frac{1}{224} \times 6.5 = \boxed{0.029 \text{ mole}}$$
18. Volume occupied by one molecule of water (density = 1 g cm⁻³) is

[AIPMT (Prelims)-2008]


- (1) $5.5 \times 10^{-23} \text{ cm}^3$ (2) $9.0 \times 10^{-23} \text{ cm}^3$
(3) $6.023 \times 10^{-23} \text{ cm}^3$ (4) $3.0 \times 10^{-23} \text{ cm}^3$

Sol. Answer (4) $\therefore 18 \text{ g H}_2\text{O} \equiv 18 \text{ mL}$

$$\therefore 1 \text{ molecule} \equiv \frac{18}{6.022 \times 10^{23}} \approx 3.0 \times 10^{-23} \text{ cm}^3$$
19. What volume of oxygen gas (O₂) measured at 0°C and 1 atm, is needed to burn completely 1 L of propane gas (C₃H₈) measured under the same conditions?

[AIPMT ((Prelims)-2008]

- (1) 10 L (2) 7 L (3) 6 L (4) 5 L

Sol. Answer (4)

20. An organic compound contains carbon, hydrogen and oxygen. Its elemental analysis gave C, 38.71% and H, 9.67%. The empirical formula of the compound would be

[AIPMT (Prelims)-2008]

- (1) CH₄O (2) CH₃O (3) CH₂O (4) CHO

Sol. Answer (2)

Element	%/At. wt.	Simplest Ratio	No. of Atoms
C	$(38.71)/_{12}$	$(3.22)/_{3.22} = 1$	1
H	$(9.67)/_1$	$(9.67)/_{3.22} = 3$	3
O	$(51.62)/_{16}$	$(3.22)/_{3.22} = 1$	1
$\therefore$ Empirical formula = CH_3O			

21. An element, X has the following isotopic composition ; ^{200}X : 90% ; ^{199}X : 8.0% ; ^{202}X : 2.0%

The weighted average atomic mass of the naturally occurring element X is closest to :

[AIPMT (Prelims)-2007]

(1) 199 amu

(2) 200 amu

(3) 201 amu

(4) 202 amu

Sol. Answer (2)

$$\text{Average atomic mass} = \frac{200 \times 90 + 199 \times 8 + 202 \times 2}{100} = 199.96 \approx 200 \text{ amu}$$

22. Concentrated aqueous sulphuric acid is 98% H_2SO_4 by mass and has a density of 1.80 g mL^{-1} . Volume of acid required to make one litre of $0.1 \text{ M H}_2\text{SO}_4$ is

[AIPMT (Prelims)-2007]

(1) 5.55 mL

(2) 11.10 mL

(3) 16.65 mL

(4) 22.20 mL

Sol. Answer (1)

$$M = \frac{10\chi d}{M_b} = \frac{10 \times 98 \times 1.8}{98} = 18\text{M}$$

$$\therefore M_1V_1 = M_2V_2 \Rightarrow 18 \times V_1 = 0.1 \times 1000$$

$$\therefore V_1 = \frac{100}{18} = 5.55 \text{ mL}$$

Questions asked prior to Medical Ent. Exams. 2005

23. How many grams of CH_3OH should be added to water to prepare 150 ml solution of $2 \text{ M CH}_3\text{OH}$?(1) 9.6×10^3 (2) 2.4×10^3

(3) 9.6

(4) 2.4

Sol. Answer (3)

$$\text{Moles of CH}_3\text{OH} = \frac{M \times V \text{ mL}}{1000} = \frac{2 \times 150}{1000} = 0.3 \text{ mole}$$

$$\therefore \text{weight of CH}_3\text{OH} = \text{moles} \times \text{mol. mass}$$

$$= 0.3 \times 32 = 9.6 \text{ g}$$

24. The total number of valence electrons in 4.2 g of N_3^- ion is (N_A is the Avogadro's number)

- (1) $2.1 N_A$ (2) $4.2 N_A$
(3) $1.6 N_A$ (4) $3.2 N_A$

Sol. Answer (3)

$$\text{Moles of } \text{N}_3^- = \frac{4.2}{42} = 0.1 \text{ mol}$$

$$\therefore \text{Total number of valence electrons in } \text{N}_3^- = 0.1 N_A \times 16 = 1.6 N_A$$

25. The number of mole of oxygen in one litre of air containing 21% oxygen by volume, under standard conditions, is

- (1) 0.0093 mole (2) 2.10 moles
(3) 0.186 mole (4) 0.21 mole

Sol. Answer (1)


$$\text{Volume of } \text{O}_2 = \frac{21}{100} \times 1000 = 210 \text{ mL}$$

$$\text{Moles of } \text{O}_2 \text{ at STP} = \frac{210}{22400} = 0.0093 \text{ mole}$$

26. The amount of zinc required to produce 224 ml of H_2 at STP on treatment with dilute H_2SO_4 will be ($\text{Zn} = 65$)

- (1) 65 g (2) 0.065 g
(3) 0.65 g (4) 6.5 g

Sol. Answer (3)



22400 mL of H_2 is produced by 1 mol Zn i.e., = 65 g

$$224 \text{ mL of } \text{H}_2 \text{ is produced by 1 mol Zn i.e.,} = \frac{65}{22400} \times 224 = 0.65 \text{ g}$$

27. Given the numbers : 161 cm, 0.161 cm, 0.0161 cm. The number of significant figures for the three numbers is

- (1) 3, 3 and 4 respectively (2) 3, 4 and 4 respectively
(3) 3, 4 and 5 respectively (4) 3, 3 and 3 respectively


Sol. Answer (4)

All have same significant figures.

28. Change in volume when 100 mL PH_3 decomposed to solid phosphorus and H_2 gas.

- (1) Increase in 50 mL (2) Decrease in 50 mL
(3) Increase in 150 mL (4) Decrease in 200 mL

Sol. Answer (1)



As 4 moles of $\text{PH}_3 (\text{g})$ converts into 6 moles $\text{H}_2 (\text{g})$

$\therefore 4 \times 22.4 \text{ L of PH}_3 (\text{g})$ will produce $= 6 \times 22.4 \text{ L of H}_2$


89.6 L of $\text{PH}_3 (\text{g})$ will produce $= 134.4 \text{ L}$

1 L of $\text{PH}_3 (\text{g})$ will produce $= \frac{134.4}{89.6} = 1.5 \text{ L}$

100 mL of $\text{PH}_3 (\text{g})$ will produce $= \frac{134.4}{89.6} \times 100 = 150 \text{ mL}$

$\therefore$ Increase in volume will be 50 mL


29. In the reaction,



When 1 mole of ammonia and 1 mole of O_2 are made to react to completion

- | | |
|--|--------------------------------------|
| (1) All the oxygen will be consumed | (2) 1.0 mole of NO will be produced |
| (3) 1.0 mole of H_2O is produced | (4) All the ammonia will be consumed |

Sol. Answer (1)



4 mol NH_3 reacts with 5 mol O_2

1 mol NH_3 reacts with $\frac{5}{4} = 1.25$ mol of O_2

as 1 mol of O_2 is taken therefore all the O_2 will be consumed.

30. An organic compound containing C, H and N gave the following analysis C = 40%, H = 13.33%, N = 46.67%. Its empirical formula would be

- | | |
|--------------------------------------|------------------------------------|
| (1) CH_4N | (2) CH_5N |
| (3) $\text{C}_2\text{H}_7\text{N}_2$ | (4) $\text{C}_2\text{H}_7\text{N}$ |

Sol. Answer (1)

	Percentage	At mass	Moles	Simple ratio
C	40%	12	$\frac{40}{12} = 3.33$	$\frac{3.33}{3.33} = 1$
H	6.7%	1	$\frac{6.7}{1} = 6.7$	$\frac{13.33}{3.33} = 4$
O	53.3%	16	$\frac{53.3}{16} = 3.33$	$\frac{3.33}{3.33} = 1$

$\therefore$ Empirical formula $= \text{CH}_4\text{N}$

31. How many g of dibasic acid (mol. weight 200) should be present in 100 ml. of the aqueous solution to give strength of 0.1 N?

- (1) 10 g (2) 2 g
(3) 1 g (4) 20 g

Sol. Answer (3)

$$\text{Amount of acid (in gram)} = \frac{N \times E \times V_{\text{mL}}}{1000}$$

$$\text{Here, } E = \frac{200}{2} = 100$$

$$\therefore \text{Amount of acid} = \frac{0.1 \times 100 \times 100}{1000} = 1 \text{ g}$$

32. The number of atoms in 4.25 g of NH_3 is approximately

- (1) 4×10^{23} (2) 2×10^{23}
(3) 1×10^{23} (4) 6×10^{23}

Sol. Answer (4)

$$\text{moles of } \text{NH}_3 = \frac{4.25}{17} = 0.25 \text{ mol}$$


As 1 NH_3 have 4 atoms

$$\therefore \text{Total number of atoms} = 0.25 \times 4 \times 6.022 \times 10^{23} \text{ atoms} \\ = 6.022 \times 10^{23} \text{ atoms}$$

33. Volume of CO_2 obtained at STP by the complete decomposition of 9.85 gm BaCO_3 is (Mol. wt. of $\text{BaCO}_3 = 197$)

- (1) 2.24 litre (2) 1.12 litre (3) 0.85 litre (4) 0.56 litre

Sol. Answer (3)



$$\text{mol mass of } \text{BaCO}_3 \quad 197 + 12 + 48 = 257 \text{ g}$$

$$\text{moles of } \text{BaCO}_3 = \frac{9.85}{257} \text{ g} = 0.038 \text{ mol}$$

$$1 \text{ mol of } \text{BaCO}_3 \text{ gives } \text{CO}_2 = 22.4 \text{ L}$$

$$0.038 \text{ mol of } \text{BaCO}_3 \text{ gives } \text{CO}_2 = 22.4 \times 0.038 = 0.85 \text{ L}$$

34. Percentage of Se in peroxidase anhydrous enzyme is 0.5% by weight (at. wt. = 78.4) then minimum molecular weight of peroxidase anhydrous enzyme is

- (1) 1.568×10^4 (2) 1.568×10^3
(3) 15.68 (4) 2.136×10^4

Sol. Answer (1)

0.5% of Se by weight is present

∴ 0.5% of enzyme have weight = 78.4 g

$$\begin{aligned} 100\% \text{ of enzyme have wt} &= \frac{78.4}{0.5} \times 100 \\ &= 15680 \text{ g} = 1.568 \times 10^4 \text{ g} \end{aligned}$$

35. 2.5 litre of 1 M NaOH solution mixed with another 3 litre of 0.5 M NaOH solution. Then find out molarity of resultant solution.


- (1) 0.80 M (2) 1.0 M
(3) 0.73 M (4) 0.50 M

Sol. Answer (3)

for same solution

$$M_1 V_1 + M_2 V_2 = M_3 (V_1 + V_2)$$

$$(2.5 \times 1) + (3 \times 0.5) = M_3 (2.5 + 3)$$



$$2.5 + 1.5 = M_3 \times 5.5$$

$$M_3 = \frac{4}{5.5} = 0.727 \approx 0.73 \text{ M}$$

36. Which has maximum molecules?

- (1) 7 gm N_2 (2) 2 gm H_2
(3) 16 gm NO_2 (4) 16 gm O_2

Sol. Answer (2)

Maximum number of molecules of gas are present which are having maximum number of moles

$$\text{for } \text{N}_2 = \frac{7}{28} = 0.25 \text{ mol, for } \text{NO}_2 = \frac{16}{46} = 0.34 \text{ mol}$$

$$\text{for } \text{H}_2 = \frac{2}{2} = 1 \text{ mol, for } \text{O}_2 = \frac{16}{32} = 0.5 \text{ mol}$$


As H_2 have maximum number of moles

∴ H_2 will have maximum number of molecules.

37. In Haber process 30 litres of dihydrogen and 30 litres of dinitrogen were taken for reaction which yielded only 50% of the expected product. What will be the composition of gaseous mixture under the aforesaid condition in the end?

- (1) 20 litres ammonia, 20 litres nitrogen, 20 litres hydrogen
- (2) 10 litres ammonia, 25 litres nitrogen, 15 litres hydrogen
- (3) 20 litres ammonia, 10 litres nitrogen, 30 litres hydrogen
- (4) 20 litres ammonia, 25 litres nitrogen, 15 litres hydrogen


Sol. Answer (2)



At STP 10 L of N_2 reacts with 30 L of H_2 gives 20 L NH_3

H_2 will be the LR because 30 L N_2 and 30 L H_2 are taken

Now the expected product will be = 50% i.e. 10 L NH_3



38. The maximum number of molecules is present in

- (1) 15 L of water at STP
- (2) 15 L of H_2O gas at STP
- (3) 15 g of ice
- (4) Same in all

Sol. Answer (1)

As of H_2O is 1 g/mL or 1 kg/L

$\therefore$ 15 L of $\text{H}_2\text{O} \equiv 15 \text{ kg of } \text{H}_2\text{O}$

$$\therefore n_{\text{H}_2\text{O}} = \frac{15000}{18} \text{ moles.}$$

$$\text{Moles of 15 L of } \text{H}_2\text{O} \text{ gas at STP} = \frac{15}{22.4} \text{ mole}$$

$$\text{Moles of 15 g of ice} = \frac{15}{18} \text{ mole}$$

$\therefore$ Maximum number of moles are present in 15 L of H_2O

39. Concentrated aqueous sulphuric acid is 98% H_2SO_4 (w/v) and has a density of 1.80 g mL^{-1} . Molarity of solution
 (1) 1 M (2) 1.8 M (3) 10 M (4) 1.5 M

Sol. Answer (3)

$$M = \frac{\% \frac{w}{v} \times 10}{\text{mol. mass}} = \frac{98 \times 10}{98} = 10 \text{ M}$$

40. An element, X has the following isotopic composition $^{56}\text{X} : 90\%$ $^{57}\text{X} : 8\%$ $^{59}\text{X} : 2.0\%$. The weighted average atomic mass of the naturally occurring element X is closest to
 (1) 56.14 amu (2) 56.8 amu (3) 60 amu (4) 55 amu


Sol. Answer (1)

$$\frac{\sum \text{percentage} \times \text{atomic mass}}{100} = \frac{\sum \text{percentage abundance of each} \times \text{isotopic} \times \text{atomic mass}}{100}$$

$$= \frac{(56 \times 90) + (57 \times 8) + 59 \times 2}{100} = 56.14 \text{ amu}$$

41. 10 g of hydrogen and 64 g of oxygen were filled in a steel vessel and exploded. Volume of gaseous product after reaction
 (1) $1 \times 22.4 \text{ L}$ (2) $2 \times 22.4 \text{ L}$ (3) $3 \times 22.4 \text{ L}$ (4) $4 \times 22.4 \text{ L}$

Sol. Answer (1)



10 g 64 g

From the reaction 2 g of H_2 (g) combine with 16 g of O_2

$$\therefore 64 \text{ g of } \text{O}_2 \text{ will combine with } \text{H}_2 = \frac{2}{16} \times 64 = 8 \text{ g of } \text{H}_2 (\text{g})$$

After the reaction 2 g of H_2 (g) i.e., 1 mol of H_2 (g) will remain unreacted


$\therefore$ volume will be $1 \times 22.4 \text{ L}$ gaseous product.

Note: Volume of H_2O will not be considered as only volume of gas is asked in the question.

42. What is the $[\text{OH}^-]$ in the final solution prepared by mixing 20.0 mL of 0.050 M HCl with 30.0 mL of 0.10 M $\text{Ba}(\text{OH})_2$?
 (1) 0.12 M (2) 0.10 M (3) 0.40 M (4) 0.0050 M

Sol. Answer (2)

It is neutralisation reaction



$$\therefore [\text{OH}^-] = \frac{6-1}{50} = \frac{5}{50} = 0.1 \text{ M}$$

43. The number of atoms in 0.1 mol of a triatomic gas is ($N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$)

(1) 1.800×10^{22}

(2) 6.026×10^{22}

(3) 1.806×10^{23}

(4) 3.600×10^{23}

Sol. Answer (3)

As triatomic gas means 3 atoms are present in a molecule

$$\text{Number of atoms} = 0.1 \times 3 \times 6.022 \times 10^{23}$$

$$= 1.806 \times 10^{23} \text{ atoms}$$

44. The total number of electrons in 2.0 g of D_2O to that in 1.8 g of H_2O

(1) Double

(2) Same

(3) Triple

(4) One fourth

Sol. Answer (2)

Both have same number of e^- [Both 2.0 g D_2O and 1.8 g H_2O have same number of atom]

$$\text{Moles of } \text{D}_2\text{O} = \frac{2.0}{20} = 0.1$$

$$\text{Number of } e^- = 0.1 \times 10 \times N_0 = 1 \times N_0$$

$$\text{Moles of } \text{H}_2\text{O} = \frac{1.8}{18} = 0.1$$

$$\text{Number of } e^- = 0.1 \times 10 \times N_0 = 1 \times N_0$$

45. From 200 mg of CO_2 when x molecules are removed, 2.89×10^{-3} moles of CO_2 are left. x will be


(1) 10^{20} molecules

(2) 10^{10} molecules

(3) 21 molecules

(4) 10^{21} molecules

Sol. Answer (4)



From Equation

$$\begin{aligned} 200 \text{ mg of } \text{CO}_2 \text{ have molecule} &= \frac{200}{44} \times 10^{-3} \times 6.022 \times 10^{23} \\ &= 2.7 \times 10^{21} \end{aligned}$$

$$\begin{aligned} \therefore 2.89 \times 10^{-3} \text{ moles of } \text{CO}_2 \text{ have molecule} &= 2.89 \times 10^{-3} \times 6.022 \times 10^{23} \\ &= 1.7 \times 10^{21} \text{ molecule} \end{aligned}$$

$$\therefore 200 \text{ mg of } \text{CO}_2 - x \text{ molecule} = 2.89 \times 10^{-3} \text{ moles of } \text{CO}_2$$

$$2.7 \times 10^{21} - x \text{ molecule} = 1.7 \times 10^{21}$$

$$x = (2.7 - 1.7) \times 10^{21} \text{ molecule}$$

$$= 10^{21} \text{ molecule}$$

$\therefore$ The value of x will be 10^{21}

46. If the weight of metal oxide is x g containing y g of oxygen, the equivalent weight of metal will be

(1) $E = \frac{8x}{y}$


(2) $E = \frac{8(y-x)}{x}$

(3) $E = \frac{y}{8}$

(4) $E = \frac{8(x-y)}{y}$

Sol. Answer (4)

$$\text{Equivalent mass of metal in oxide} = \frac{\text{weight of metal}}{\text{weight of oxygen}} \times 8$$



$$\text{Equivalent weight of metal} = \frac{x-y}{y} \times 8$$

$$E = \frac{8(x-y)}{y}$$

47. The number of significant figures in 2.653×10^4 is

(1) 8

(2) 4

(3) 7

(4) 1

Sol. Answer (2)

2.653×10^4 have only 4 significant figure

48. Mole fraction of solute in aqueous solution of 30% NaOH.

(1) 0.16

(2) 0.05

(3) 0.25

(4) 0.95

Sol. Answer (1)

30% NaOH means 30% by mass

$$\text{i.e., } \frac{30 \text{ g of NaOH}}{100 \text{ g of solution}}$$

Weight of NaOH = 30 g

Mol. mass NaOH = 40

Weight of H_2O = $100 - 30 = 70$

Mol. mass = 18

$$\text{Mol. fraction of NaOH} = \frac{\text{moles of NaOH}}{\text{moles of H}_2\text{O} + \text{moles of NaOH}} = \frac{\frac{30}{40}}{\frac{70}{18} + \frac{30}{40}} = 0.16$$

SECTION – C

Assertion–Reason Type Questions

1. A : 1 a.m.u. = 1.66×10^{-24} gram.

R : Actual mass of one atom of C-12 is equal to 1.99×10^{-23} g.

Sol. Answer (2)

Both are correct but R is not explanation of A because $1 \text{ amu} = 1.66 \times 10^{-24} \text{ g}$

and mass of 1 atom of C = $1.99 \times 10^{-23} = \frac{12}{6.022 \times 10^{23}} = 1.99 \times 10^{-23} = \text{g}$

Both A and R are correct.

2. A : Unit of specific gravity is gram-cc^{-1} .

R : Specific gravity is same as density of a liquid in normal conditions.

Sol. Answer (4)

Specific gravity have number units because it specific gravity = $\frac{\text{density of substance}}{\text{density of H}_2\text{O at } 4^\circ\text{C}}$

Both A and R are incorrect.

3. A : Number of atoms in 2 mole of NH_3 is equal to number of atoms in 4 mole of CH_4 .

R : Both are chemically similar species.

Sol. Answer (4)


Number of atoms in $\text{NH}_3 = 2 \times 4 \times N_0 = 8 N_0$

Number of atoms in $\text{CH}_4 = 4 \times 5 \times N_0 = 20 N_0$

Both are chemically different

Both A and R are incorrect.


4. A : In the reaction



equivalent weight of H_3PO_4 is $\frac{M}{2}$, where M is its molecular weight.

R : Equivalent weight = $\frac{\text{Molecular weight}}{n - \text{factor}}$

Sol. Answer (1)



For above reaction n-factor = 2

$$\therefore \text{equivalent mass} = \frac{M}{2}$$

As tow 'H' are replaced.

Equivalent mass = $\frac{\text{mol. mass}}{n\text{-factor}}$

A and R are correct R is correct explanation of A.

5. A : Mass of 1 gram molecule of H_2SO_4 is 98 gram.

R : One gram atom contains N_A atoms.

Sol. Answer (2)

mass of 1 g molecule means 1 mol of $\text{H}_2\text{SO}_4 = 98 \text{ g}$


1 g atom = 1 mol atom = N_0

Both are correct but R is not explanation of A.

6. A : One mole of sucrose reacts completely with oxygen produces 268.8 litre of carbon dioxide at STP.

R : Amount of oxygen required for reaction is 268.8 litre.


Sol. Answer (2)



$$\left[\begin{array}{l} 1 \text{ mol sucrose produce } \text{CO}_2 = 12 \times 22.4 = 268.8 \text{ L} \\ 1 \text{ mol sucrose require } \text{O}_2 = 12 \times 22.4 = 268.8 \text{ L} \end{array} \right]$$

Both are correct but reason is not correct explanation of Assertion.

7. A : In the reaction




equivalents of NaOH, Na_2SO_4 and H_2SO_4 are equal.

R : Number of equivalents = number of moles $\times$ n-factor.

Sol. Answer (1)

All are having same equivalents




Both A and R are correct R is correct explanation of A.

8. A : When 4 moles of H_2 reacts with 2 moles of O_2 , then 4 moles of water is formed.

R : O_2 will act as limiting reagent.

Sol. Answer (3)



2 mol 1 mol 2 mol

4 mol 2 mol 4 mol

[4 mol of H_2 when reacts with 2 moles O_2 produces moles of H_2O]

When moles are taken in above reaction H_2 will act on LR.

A is correct but R is wrong.

9. A : 50 ml, decinormal HCl when mixed with 50 ml, decinormal H_2SO_4 , then normality of H^+ ion in resultant solution is 0.1 N.

R : Here, $MV = M_1V_1 + M_2V_2$.

Sol. Answer (3)

meq of HCl = $50 \times 0.1 = 5$

meq. of $\text{H}_2\text{SO}_4 = 50 \times 0.1 = 5$

$$\text{Normality} = \frac{\text{Total meq}}{\text{Total volume}} = \frac{10}{100} = 0.1 \text{ N}$$

when difference solutions of different n-factors are taken than $N_1V_1 + N_2V_2 = N_3(V_1 + V_2)$

A is correct but R is wrong.

10. A : 50 ml, decimolar H_2SO_4 when mixed with 50 ml, decimolar NaOH, then normality of resultant solution is 0.05 N.

R : Here, $NV = |N_1V_1 - N_2V_2|$.

Sol. Answer (1)

$$\text{Meq of } \text{H}_2\text{SO}_4 = N \times V = 50 \times 0.2 = 10$$

$$\text{Meq of NaOH} = N \times V = 50 \times 0.1 = 5$$

$$N = m \times n = 0.1 \times 2 = 0.2 \text{ N } \text{H}_2\text{SO}_4$$

$$N = m \times n = 0.1 \times 1 = 0.1 \text{ N NaOH}$$

$$N_{\text{solution}} = \frac{10 - 5}{100} = \frac{5}{100} = 0.05 \text{ N}$$

$$\text{for neutralisation} = \frac{\text{larger NV} - \text{smaller NV}}{\text{Total volume}} = \frac{N_1V_1 - N_2V_2}{\text{Total volume}}$$

R is correct exp. of A.

11. A : Ratio of empirical formula mass and molecular formula mass may be a whole number.

R : Molecular formula mass = $n \times$ empirical formula mass, where n is the simplest whole number.

Sol. Answer (2)

$$= \text{mol. formula} = n \times \text{Empirical formula.}$$

$$= \text{mol. mass} = n \times \text{formula mass (for ionic solids).}$$

Both A and R are correct but R is not correct explanation of A

12. A : For a given solution (density = 1 gm/ml), molality is greater than molarity.

R : Molarity involves volume of solution while molality involves mass of solvent.

Sol. Answer (1)

If density is 1 g/mL

$\therefore$ mass of solution > mass of solvent

$$\text{Molality} = \frac{\text{moles}}{\text{volume of solution}}$$

$$\text{Molality} = \frac{\text{moles}}{\text{mass of solvent}}$$

As weight of solvent is less therefore molality will be more.

Both A and R are correct R is the correct explanation of A.

13. A : 1 gram of salt in 1 m^3 of solution has concentration of 1 ppm.

R : ppm is defined as number of parts by mass of solute per million parts of solution.

Sol. Answer (1)

$$\text{ppm} = \frac{\text{weight of solvent}}{\text{weight of solution}} \times 10^6 = \frac{1}{1} \times 10^6 = \text{ppm} \approx 1 \text{ ppm}$$

Both are A and R correct R is correct explanation of A.

14. A : Total charge on N_A ions of CO_3^{2-} is 1.93×10^5 coulomb.

R : Charge on one electron is 96500 coulomb.

Sol. Answer (3)

CO_3^{2-} as one CO_3^{2-} ion have two unit charge

$\therefore$ 1 mole i.e., N_A CO_3^{2-} have charge = $2 \times 96500 \text{ C} = 1.93 \times 10^5 \text{ C}$

Charge on one mole electron = $1.602 \times 10^{-19} \times 6.022 \times 10^{23} \approx 96478 = 96500 \text{ C}$

A is true but R is false.

15. A : Number of ions in 9 gram of NH_4^+ is equal to Avogadro's number (N_A).

R : Number of ions is equal to number of atoms.

Sol. Answer (4)

$$\text{Formula units} = \frac{9}{18} = 0.5 \times N_A$$

No formula units = moles = Number of molecules.

Both A and R are incorrect.

