

CBSE Board
Class XI Mathematics
Sample Paper – 9

Time: 3 hrs

Total Marks: 100

General Instructions:

1. All questions are compulsory.
 2. The question paper consist of 29 questions.
 3. Questions 1 – 4 in Section A are very short answer type questions carrying 1 mark each.
 4. Questions 5 – 12 in Section B are short-answer type questions carrying 2 mark each.
 5. Questions 13 – 23 in Section C are long-answer I type questions carrying 4 mark each.
 6. Questions 24 – 29 in Section D are long-answer type II questions carrying 6 mark each.
-

SECTION – A

1. Find $\lim_{x \rightarrow 0} \frac{\sin ax}{bx}$

2. Is the given sentence statement? Justify. “There are 35 days in a month.”

3. Write in the form of $a + bi$: $\frac{1}{i-1}$

OR

Find modulus of $2i$.

4. If variance of 20 observations is 5. If each observation is multiplied by 2, then find variance of the new observations.

SECTION – B

5. Let $A = \{1, 2\}$, $B = \{1, 2, 3, 4\}$, $C = \{5, 6\}$ and $D = \{5, 6, 7, 8\}$ verify that $A \times C$ is a subset of $B \times D$.

6. Let f be defined by $f(x) = x - 4$ and g be defined by

$$g(x) = \frac{x^2 - 16}{x + 4} \quad x \neq -4$$
$$= \lambda \quad x = -4$$

Find λ such that $f(x) = g(x)$ for all x .

OR

Find domain and range of the function $f(x) = \frac{x^2 - 9}{x - 3}$

7. Assuming that a person of normal sight can read print at such a distance that the letters subtend an angle of $5'$ at this eye, find the height of the letters that he can read at a distance of 12 m.

OR

If the arcs of the same length in two circles subtend angles of 60° and 75° at their centres. Find the ratio of their radii.

8. If $n(U) = 600$, $n(A) = 460$, $n(B) = 390$ and $n(A \cap B) = 325$ then find $n(A \cup B)$ and $n(A \cup B)'$

9. Prove that $\sin(\theta + 30^\circ) = \cos \theta + \sin(\theta - 30^\circ)$

OR

Prove that $\frac{\sin 7A - \sin 5A}{\cos 5A + \cos 7A} = \tan A$

10. Find compound statements of the "It is raining and it is cold."

11. If $f(x) = x^2$ find $\frac{f(1.1) - f(1)}{1.1 - 1}$

12. Find the equation of line joining the points $(-1, 3)$ and $(4, -2)$.

SECTION - C

13. Prove that $\frac{\cos^2 33^\circ - \cos^2 57^\circ}{\sin^2 \frac{21^\circ}{2} - \sin^2 \frac{69^\circ}{2}} = -\sqrt{2}$

14. If f is a real function defined by $f(x) = \frac{x-1}{x+1}$ then prove that $f(2x) = \frac{3f(x)+1}{f(x)+3}$

15. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be given by $f(x) = x^2 + 3$. Find

- $\{x : f(x) = 28\}$
- The pre-image of 39 and 2 under f .

16. A man accepts a position with an initial salary of Rs. 5200 per week. It is understood that he will receive an automatic increase of Rs. 320 in the very next and each month.

- find his salary for the tenth month
- his total earning during the first year.

17. If $(x + yi)^3 = u + vi$ prove that $\frac{u}{x} + \frac{v}{y} = 4(x^2 + y^2)$

18. Two cards are drawn from a pack of cards. What is the probability that either both are red or both are kings?

19. Determine the number n in a geometric progression $\{a_n\}$, if $a_1 = 3$, $a_n = 96$ and $S_n = 189$.

20. Find n , if ${}^{2n}C_1$, ${}^{2n}C_2$ and ${}^{2n}C_3$ are in A. P.

OR

Prove that the product of $2n$ consecutive negative integers is divisible by $(2n)!$.

21. Find the equation of the straight line through the origin making angle of 60° with the straight line $x + \sqrt{3}y + 3\sqrt{3} = 0$

OR

Find the equations of the lines, which cut off intercepts on the axes whose sum and product are 1 and -6 respectively.

22. Differentiate $x^{-3/2}$ with respect to x using first principle.

OR

Differentiate $\frac{x+2}{x^2+3}$ and find the value of derivative at $x = 0$.

23. Find the equation of hyperbola whose foci are $(8, 3)$ and $(0, 3)$ and $e = 4/3$

SECTION - D

24. Prove that $\cot \theta \cot 2\theta + \cot 2\theta \cot 3\theta + 2 = \cot \theta (\cot \theta - \cot 3\theta)$

OR

Prove that $5\cos \theta + 3\cos \left(\theta + \frac{\pi}{3} \right) + 3$ lies between -4 and 10.

25. Find the mean and variance of the following data

Classes	0 - 30	30 - 60	60 - 90	90 - 120	120 - 150	150 - 180	180 - 210
Frequency	2	3	5	10	3	5	2

26. If Find $\sin \frac{x}{2}$, $\cos \frac{x}{2}$ and $\tan \frac{x}{2}$ where $\tan x = -\frac{4}{3}$, x is in quadrant II

27. Plot the given linear inequations and shade the region which is common to the solution of all inequations $x \geq 0$, $y \geq 0$, $5x + 3y \leq 500$; $x \leq 70$ and $y \leq 125$.

OR

How many litres of water will have to be added to 1125 litres of a 45% solution of acid so that the resulting mixture will contain more than 25% but less than 30% acid content?

28. Using principle of mathematical induction prove that $5^n - 5$ is divisible by 4 for all $n \in \mathbb{N}$.
Hence, prove that $2 \times 7^n + 3 \times 5^n - 5$ is divisible by 24 for all $n \in \mathbb{N}$.

29. If a , b , and c are in A.P.; b , c , and d are in G.P. and $\frac{1}{c}$, $\frac{1}{d}$, and $\frac{1}{e}$ are in A.P., prove that a , c , and e are in G.P.

OR

Show that:

$$\frac{1 \times 2^2 + 2 \times 3^2 + \dots + n \times (n+1)^2}{1^2 \times 2 + 2^2 \times 3 + \dots + n^2 \times (n+1)} = \frac{3n+5}{3n+1}$$

CBSE Board
Class XI Mathematics
Sample Paper – 9 Solution

SECTION – A

1.

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\sin ax}{bx} \\&= \lim_{x \rightarrow 0} \frac{\sin ax}{ax} \times \frac{a}{b} \\&= \frac{a}{b}\end{aligned}$$

2. The sentence is false because a month can't have more than 31 days. Hence it is a statement.

3.

$$\begin{aligned}\frac{1}{i-1} \\&= \frac{1}{-1+i} \\&= \frac{1}{-1+i} \times \frac{-1-i}{-1-i} \\&= \frac{-1-i}{1-i^2} \\&= \frac{-1-i}{1-(-1)} \\&= \frac{-1-i}{2} \\&= \frac{-1}{2} - \frac{i}{2}\end{aligned}$$

$$a = -1/2 \text{ and } b = -1/2$$

OR

$$z = 2i$$

Comparing with $a + bi$ we get $a = 0$ and $b = 2$

$$|z| = \sqrt{0+2^2} = 2$$

4. According to the question,
 Variance of 20 observations is 5.
 New variance = $2^2 \times 5 = 20$

SECTION - B

5. $A \times C = \{(1, 5), (1, 6), (2, 5), (2, 6)\} \dots (i)$
 $B \times D = \{(1, 5), (1, 6), (1, 7), (1, 8), (2, 5), (2, 6), (2, 7), (2, 8), (3, 5), (3, 6), (3, 7), (3, 8), (4, 5), (4, 6), (4, 7), (4, 8)\} \dots (ii)$
 $A \times C$ is a subset of $B \times D$ all the elements of $A \times C$ are contained in $B \times D$.

6. According to the question,

$$f(x) = g(x)$$

$$f(-4) = g(-4)$$

$$-4 - 4 = \lambda$$

$$\lambda = -8$$

OR

$$f(x) = \frac{x^2 - 9}{x - 3}$$

$f(x)$ is not defined for $x - 3 = 0$ i. e. $x = 3$. Therefore, domain $(f) = R - \{3\}$

Let $f(x) = y$ then

$$\frac{x^2 - 9}{x - 3} = y$$

$$x + 3 = y$$

It follows from the above relation that y takes all real values except 6 when x takes values in the set $R - \{3\}$. Therefore, Range $(f) = R - \{6\}$

7. Let h be the required height in metres. Here h can be considered as the arc of a circle of radius 12 m, which subtends an angle of $5'$ at its centre.

$$\theta = 5' = \left(\frac{5}{60}\right)^\circ = \left(\frac{1}{12} \times \frac{\pi}{180}\right)^c \text{ and } r = 12 \text{ m}$$

$$\theta = \frac{\text{arc}}{r} = \frac{\pi}{12 \times 180} = \frac{h}{12}$$

$$h = \frac{\pi}{180} = 1.7 \text{ m}$$

OR

Let r_1 and r_2 be the radii of the given circles and let their arcs of same length s subtend angles of 60° and 75° at their centres.

$$60^\circ = 60^\circ \times \frac{\pi}{180} = \frac{\pi}{3}$$

$$75^\circ = 75^\circ \times \frac{\pi}{180} = \frac{5\pi}{12}$$

$$s = r_1\theta_1 = r_2\theta_2$$

$$\frac{\pi}{3}r_1 = \frac{5\pi}{12}r_2$$

$$\frac{r_1}{r_2} = \frac{5}{12} \times 3$$

$$\frac{r_1}{r_2} = \frac{5}{4}$$

8. $n(U) = 600$, $n(A) = 460$, $n(B) = 390$ and $n(A \cap B) = 325$

$$n(A \cup B) = n(A) + n(B) - n(A \cap B)$$

$$= 460 + 390 - 325$$

$$= 525$$

$$n(A \cup B)' = n(U) - n(A \cup B)$$

$$= 600 - 525$$

$$= 75$$

9. $\sin(\theta + 30^\circ) = \sin \theta \cos 30^\circ + \cos 30^\circ \sin \theta$
 $= (\sin \theta \cos 30^\circ - \cos \theta \sin 30^\circ) + 2\cos \theta \sin 30^\circ$
 $= \sin(\theta - 30^\circ) + 2\cos \theta \times \frac{1}{2}$
 $= \cos \theta + \sin(\theta - 30^\circ)$
 $\therefore \sin(\theta + 30^\circ) = \cos \theta + \sin(\theta - 30^\circ)$

OR

$$\begin{aligned} & \frac{\sin 7A - \sin 5A}{\cos 5A + \cos 7A} \\ &= \frac{2\cos \frac{7A+5A}{2} \sin \frac{7A-5A}{2}}{2\cos \frac{7A+5A}{2} \cos \frac{7A-5A}{2}} \\ &= \frac{\cos 6A \sin A}{\cos 6A \cos A} \\ &= \frac{\sin A}{\cos A} \\ &= \tan A \end{aligned}$$

10. The compound statements are

p : It is raining

q : It is cold.

11. $f(x) = x^2$

$$\frac{f(1.1) - f(1)}{1.1 - 1}$$

$$= \frac{1.1^2 - 1^2}{1.1 - 1} \quad s$$

$$= \frac{1.21 - 1}{0.1}$$

$$= 2.1$$

12. $(x_1, y_1) = (-1, 3)$ and $(x_2, y_2) = (4, -2)$

The equation of line is $y - 3 = \frac{3 - (-2)}{-1 - 4} (x + 1)$

$$y - 3 = -x - 1$$

$$x + y - 2 = 0$$

SECTION - C

13.

$$\begin{aligned} & \frac{\cos^2 33^\circ - \cos^2 57^\circ}{\sin^2 \frac{21^\circ}{2} - \sin^2 \frac{69^\circ}{2}} \\ &= \frac{1 - \sin^2 33^\circ - (1 - \sin^2 57^\circ)}{\sin^2 \frac{21^\circ}{2} - \sin^2 \frac{69^\circ}{2}} \\ &= \frac{\sin^2 57^\circ - \sin^2 33^\circ}{\sin^2 \frac{21^\circ}{2} - \sin^2 \frac{69^\circ}{2}} \\ &= \frac{\sin(57^\circ + 33^\circ) \sin(57^\circ - 33^\circ)}{\sin\left(\frac{21^\circ}{2} + \frac{69^\circ}{2}\right) \sin\left(\frac{21^\circ}{2} - \frac{69^\circ}{2}\right)} \\ &= \frac{\sin 90^\circ \sin 24^\circ}{\sin 45^\circ \sin(-24^\circ)} \\ &= \frac{\sin 24^\circ}{\frac{-1}{\sqrt{2}} \sin 24^\circ} = -\sqrt{2} \end{aligned}$$

$$14. f(x) = \frac{x-1}{x+1}$$

$$\frac{f(x)}{1} = \frac{x-1}{x+1}$$

$$\frac{f(x)+1}{f(x)-1} = \frac{x-1+x+1}{x-1-x-1}$$

$$\frac{f(x)+1}{f(x)-1} = -x$$

$$\frac{f(x)+1}{1-f(x)} = x$$

$$f(2x) = \frac{2x-1}{2x+1}$$

$$f(2x) = \frac{2 \times \frac{f(x)+1}{1-f(x)} - 1}{2 \times \frac{f(x)+1}{1-f(x)} + 1}$$

$$f(2x) = \frac{2 \times f(x) + 2 - 1 + f(x)}{2 \times f(x) + 2 + 1 - f(x)}$$

$$f(2x) = \frac{3f(x)+1}{f(x)+3}$$

$$15. i. f(x) = x^2 + 3$$

$$f(x) = 28$$

$$x^2 + 3 = 28$$

$$x^2 = 25$$

$$x = \pm 5$$

$$\{x : f(x) = 28\} = \{5, -5\}$$

ii. Let x be the pre-image of 39. Then,

$$f(x) = 39$$

$$x^2 + 3 = 39$$

$$x^2 = 36$$

$$x = \pm 6$$

So, pre-image of 39 are -6 and 6.

Let x be the pre-image of 2. Then,

$$f(x) = 2$$

$$x^2 + 3 = 2$$

$$x^2 = -1$$

We find that no real value of x satisfies the equation. Therefore, 2 does not have any pre-image under f.

16. Salary in the first week = $5200 \times 4 = \text{Rs. } 20800$

Increase = Rs. 320 per week

$a = 20800$ and $d = 320$

$$\begin{aligned}\text{i. Salary in the } 10^{\text{th}} \text{ month} &= t_{10} \\ &= a + 9d \\ &= 20800 + 9 \times 320 \\ &= \text{Rs. } 23680\end{aligned}$$

ii. Total earning during the first year.

$$\begin{aligned}&= \frac{12}{2} [2 \times 20800 + (12-1) \times 320] \\ &= 6 (41600 + 3520) \\ &= \text{Rs. } 270720\end{aligned}$$

17. $(x + yi)^3 = u + vi$

$$x^3 + 3x^2yi + 3xy^2i^2 + y^3i^3 = u + vi$$

$$x^3 + 3x^2yi - 3xy^2 - y^3i = u + vi$$

$$x^3 - 3xy^2 + (3x^2y - y^3)i = u + vi$$

$$x^3 - 3xy^2 = u \text{ and } 3x^2y - y^3 = v$$

$$x(x^2 - 3y^2) = u \text{ and } y(3x^2 - y^2) = v$$

$$\frac{u}{x} + \frac{v}{y} = x^2 - 3y^2 + 3x^2 - y^2$$

$$\frac{u}{x} + \frac{v}{y} = 4x^2 - 4y^2$$

$$\frac{u}{x} + \frac{v}{y} = 4(x^2 - y^2)$$

18. Two cards out of 52 cards. Can be drawn in ${}^{52}C_2$ ways.

$$n(S) = {}^{52}C_2$$

Let A be the event that two cards are drawn are red cards and B be the event that two cards are drawn are kings.

$$n(A) = {}^{26}C_2 \text{ and } n(B) = {}^4C_2$$

$$\text{Required probability} = P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A) = \frac{n(A)}{n(S)} = \frac{{}^{26}C_2}{{}^{52}C_2} \text{ and } P(B) = \frac{n(B)}{n(S)} = \frac{{}^4C_2}{{}^{52}C_2}$$

Also there are two cards which are both red and kings.

$$P(A \cap B) = \text{Probability of getting two red kings} = \frac{{}^2C_2}{{}^{52}C_2}$$

$$\text{Required probability} = \frac{{}^{26}C_2}{{}^{52}C_2} + \frac{{}^4C_2}{{}^{52}C_2} - \frac{{}^2C_2}{{}^{52}C_2} = \frac{325}{1326} + \frac{1}{221} - \frac{1}{1326} = \frac{55}{221}$$

19. Let r be the common ratio.

$$a_n = a_1 r^{n-1} = 3 r^{n-1}$$

$$3 r^{n-1} = 96$$

$$r^{n-1} = 32 \dots (i)$$

$$S_n = \frac{a_1(r^n - 1)}{r - 1} = \frac{3(r^n - 1)}{r - 1} = 189$$

$$\frac{r^n - 1}{r - 1} = 63$$

$$r^n - 1 = 63r - 63$$

$$r^n = 63r - 62$$

$$r^{n-1} \cdot r = 63r - 62$$

$$32r = 63r - 62 \quad \text{from (i)}$$

$$31r = 62$$

$$r = 2$$

$$2^{n-1} = 32 = 2^5$$

$$n - 1 = 5$$

$$n = 6$$

20. Since, ${}^{2n}C_1$, ${}^{2n}C_2$ and ${}^{2n}C_3$ are in A. P.

$$2^{2n}C_2 = {}^{2n}C_1 + {}^{2n}C_3$$

$$2 \frac{2n(2n-1)}{2 \times 1} = 2n + \frac{2n(2n-1)(2n-2)}{3 \times 2 \times 1}$$

$$2n(2n-1) = 2n + \frac{1}{3}n(2n-1)(2n-2)$$

$$6n(2n-1) = 6n + n(2n-1)(2n-2)$$

$$6(2n-1) = 6 + (2n-1)(2n-2) \quad \because n \neq 0$$

$$12n - 6 = 6 + 4n^2 - 2n - 4n + 2$$

$$4n^2 - 18n + 14 = 0$$

$$2n^2 - 9n + 7 = 0$$

$$n = \frac{9 \pm \sqrt{81 - 56}}{4} = \frac{9 \pm 5}{4} = \frac{7}{2}, 1$$

But $n \neq 1$ as 2C_3 not possible hence, $n = 7/2$

OR

Let $-r, -r - 1, -r - 2, \dots, (-r - 2n + 1)$ be $2n$ consecutive negative integers.

Then their product $= (-r)(-r - 1)(-r - 2) \dots (-r - 2n + 1)$

$$= (-1)^{2n} r(r + 1)(r + 2) \dots (r + 2n - 1)$$

$$= \frac{(r - 1)! r(r + 1)(r + 2) \dots (r + 2n - 1)}{(r - 1)!}$$

$$= \frac{(r + 2n - 1)!}{(r - 1)!}$$

$$= \frac{(r + 2n - 1)!(2n)!}{(r - 1)!(2n)!}$$

$$= {}^{r+2n-1}C_{2n} (2n)!$$

Hence, the product is divisible by $(2n)!$.

21. The equation of any straight line through $(0, 0)$ is

$$y = 0 = m(x - 0)$$

$$y = mx \dots \dots (i)$$

The given straight line is $x + \sqrt{3}y + 3\sqrt{3} = 0 \dots \dots (ii)$

$$\text{Thus } m_1 = m \text{ and } m_2 = \frac{-\text{coefficient of } x}{\text{coefficient of } y} = \frac{-1}{\sqrt{3}}$$

By the question, angle between (i) and (ii) is 60° .

$$\left| \frac{m - \left(\frac{-1}{\sqrt{3}} \right)}{1 + m \left(\frac{-1}{\sqrt{3}} \right)} \right| = \tan 60^\circ$$

$$\pm \frac{\sqrt{3}m + 1}{\sqrt{3} - m} = \sqrt{3}$$

$$\frac{\sqrt{3}m + 1}{\sqrt{3} - m} = \sqrt{3}$$

$$\sqrt{3}m + 1 = \sqrt{3}(\sqrt{3} - m)$$

$$\sqrt{3}m + 1 = 3 - \sqrt{3}m$$

$$2\sqrt{3}m = 2$$

$$m = \frac{1}{\sqrt{3}}$$

$$-\frac{\sqrt{3}m + 1}{\sqrt{3} - m} = \sqrt{3}$$

$$-\sqrt{3}m - 1 = \sqrt{3}(\sqrt{3} - m)$$

$$-\sqrt{3}m - 1 = 3 - \sqrt{3}m$$

Here, m is undefined.

Hence, the equation of the straight line is $y = \frac{1}{\sqrt{3}}x$

OR

Let the intercepts be a and b so that $a + b = 1$(i)

$ab = -6$(ii)

$b = 1 - a$(iii)

$a(1 - a) = -6$

$a^2 - a + 6 = 0$ wrong step

$(a - 3)(a + 2) = 0$

$a = 3$ or $a = -2$

When $a = 3$ then $b = 1 - 3 = -2$

When $a = -2$ then $b = 1 - (-2) = 3$

The equations of the straight lines are

$$\frac{x}{3} + \frac{y}{-2} = 1 \text{ or } \frac{x}{-2} + \frac{y}{3} = 1$$

$$2x - 3y - 6 = 0 \text{ or } 3x - 2y + 6 = 0$$

22. $f(x) = x^{-3/2} \therefore f(x + h) = (x + h)^{-3/2}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^{-\frac{3}{2}} - x^{-\frac{3}{2}}}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^{-\frac{3}{2}} - x^{-\frac{3}{2}}}{x+h-x}$$

Put $z = x + h$ and $z \rightarrow x$ as $h \rightarrow 0$

$$f'(x) = \lim_{z \rightarrow x} \frac{z^{-\frac{3}{2}} - x^{-\frac{3}{2}}}{z - x}$$

$$f'(x) = -\frac{3}{2}x^{-\frac{3}{2}-1}$$

$$f'(x) = -\frac{3}{2}x^{-\frac{5}{2}}$$

OR

$$y = \frac{x+2}{x^2+3}$$

$$\frac{dy}{dx} = \frac{(x^2+3) \frac{d}{dx}(x+2) - (x+2) \frac{d}{dx}(x^2+3)}{(x^2+3)^2}$$

$$\frac{dy}{dx} = \frac{x^2+3 - (x+2) \times 2x}{(x^2+3)^2}$$

$$\frac{dy}{dx} = \frac{x^2+3 - 2x^2 - 4x}{(x^2+3)^2}$$

$$\frac{dy}{dx} = \frac{-x^2 - 4x + 3}{(x^2+3)^2}$$

$$\left. \frac{dy}{dx} \right|_{x=0} = \frac{-0+0+3}{(0+3)^2} = \frac{3}{9} = \frac{1}{3}$$

23. The centre of the hyperbola is the mid-point of the line joining the two foci.

$$\text{Centre} = \left(\frac{8+0}{2}, \frac{3+3}{2} \right) \text{ i. e. } (4, 3)$$

Since the foci lie on the line $y = 3$ the equation of the hyperbola is of the form

$$\frac{(x-4)^2}{a^2} - \frac{(y-3)^2}{b^2} = 1 \text{ where } 2a \text{ and } 2b \text{ are the transverse and conjugate axis}$$

respectively.

$$\text{Distance between foci} = \sqrt{(8-0)^2 + (3-3)^2} = 8 = 2ae$$

$$2a \times \frac{4}{3} = 8$$

$$a = 3$$

$$b^2 = a^2(e^2 - 1) = 9 \left(\frac{16}{9} - 1 \right) = 7$$

The equation of hyperbola is

$$\frac{(x-4)^2}{9} - \frac{(y-3)^2}{7} = 1$$

$$7x^2 - 9y^2 - 56x + 54y - 32 = 0$$

SECTION - D

$$\begin{aligned}
 24. & \cot \theta \cot 2\theta + \cot 2\theta \cot 3\theta + 2 \\
 &= \cot \theta \cot 2\theta + 1 + \cot 2\theta \cot 3\theta + 1 \\
 &= \frac{\cos \theta \cos 2\theta}{\sin \theta \sin 2\theta} + 1 + \frac{\cos 2\theta \cos 3\theta}{\sin 2\theta \sin 3\theta} + 1 \\
 &= \frac{\cos \theta \cos 2\theta + \sin \theta \sin 2\theta}{\sin \theta \sin 2\theta} + \frac{\cos 2\theta \cos 3\theta + \sin 2\theta \sin 3\theta}{\sin 2\theta \sin 3\theta} \\
 &= \frac{\cos(2\theta - \theta)}{\sin \theta \sin 2\theta} + \frac{\cos(3\theta - 2\theta)}{\sin 2\theta \sin 3\theta} \\
 &= \frac{\cos \theta}{\sin \theta \sin 2\theta} + \frac{\cos \theta}{\sin 2\theta \sin 3\theta} \\
 &= \cos \theta \left(\frac{1}{\sin \theta \sin 2\theta} + \frac{1}{\sin 2\theta \sin 3\theta} \right) \\
 &= \frac{\cos \theta}{\sin \theta} \left(\frac{\sin \theta}{\sin \theta \sin 2\theta} + \frac{\sin \theta}{\sin 2\theta \sin 3\theta} \right) \\
 &= \cot \theta \left(\frac{\sin(2\theta - \theta)}{\sin \theta \sin 2\theta} + \frac{\sin(3\theta - 2\theta)}{\sin 2\theta \sin 3\theta} \right) \\
 &= \cot \theta \left(\frac{\sin 2\theta \cos \theta - \cos 2\theta \sin \theta}{\sin \theta \sin 2\theta} + \frac{\sin 3\theta \cos 2\theta - \cos 3\theta \sin 2\theta}{\sin 2\theta \sin 3\theta} \right) \\
 &= \cot \theta (\cot \theta - \cot 2\theta + \cot 2\theta - \cot 3\theta) \\
 &= \cot \theta (\cot \theta - \cot 3\theta) \\
 \cot \theta \cot 2\theta + \cot 2\theta \cot 3\theta + 2 &= \cot \theta (\cot \theta - \cot 3\theta)
 \end{aligned}$$

OR

$$\begin{aligned}
 \text{Let } f(\theta) &= 5\cos \theta + 3\cos \left(\theta + \frac{\pi}{3} \right) + 3 \\
 f(\theta) &= 5\cos \theta + 3\cos \left(\cos \theta \cos \frac{\pi}{3} - \sin \theta \sin \frac{\pi}{3} \right) + 3 \\
 &= 5\cos \theta + 3\cos \left(\frac{3}{2}\cos \theta - \frac{3\sqrt{3}}{2}\sin \theta \right) + 3 \\
 &= \frac{13}{2}\cos \theta - \frac{3\sqrt{3}}{2}\sin \theta + 3 \\
 -\sqrt{\left(\frac{13}{2} \right)^2 + \left(\frac{3\sqrt{3}}{2} \right)^2} &\leq \frac{13}{2}\cos \theta - \frac{3\sqrt{3}}{2}\sin \theta \leq \sqrt{\left(\frac{13}{2} \right)^2 + \left(\frac{3\sqrt{3}}{2} \right)^2} \\
 -7 &\leq \frac{13}{2}\cos \theta - \frac{3\sqrt{3}}{2}\sin \theta \leq 7
 \end{aligned}$$

$$-7+3 \leq \frac{13}{2} \cos \theta - \frac{3\sqrt{3}}{2} \sin \theta + 3 \leq 7+3$$

$$-4 \leq \frac{13}{2} \cos \theta - \frac{3\sqrt{3}}{2} \sin \theta + 3 \leq 10$$

25. Let assumed mean be $A = 105$

Classes	f_i	x_i	$u_i = \frac{x_i - 105}{30}$	$f_i u_i$	$f_i u_i^2$
0 - 30	2	15	-3	-6	18
30 - 60	3	45	-2	-6	12
60 - 90	5	75	-1	-5	5
90 - 120	10	105	0	0	0
120 - 150	3	135	1	3	3
150 - 180	5	165	2	10	20
180 - 210	2	195	3	6	18
	$\sum f_i = 30$			$\sum f_i u_i = 2$	$\sum f_i u_i^2 = 76$

$$\begin{aligned} \text{Mean } \bar{X} &= A + \frac{\sum f_i u_i}{\sum f_i} \times h \\ &= 105 + \frac{2}{30} \times 30 \\ &= 107 \end{aligned}$$

$$\begin{aligned} \text{Variance } \sigma^2 &= h^2 \times \left[\frac{1}{N} \sum f_i u_i^2 - \left(\frac{1}{N} \sum f_i u_i \right)^2 \right] \\ &= (30)^2 \left[\frac{1}{30} \times 76 - \left(\frac{2}{30} \right)^2 \right] \\ &= 900 \left[\frac{76}{30} - \frac{4}{900} \right] = 2276 \end{aligned}$$

26.

$$\tan x = -\frac{4}{3}; \frac{\pi}{2} \leq x \leq \pi$$

$$\tan x = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}} \quad \left(\because \tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta} \right)$$

$$\Rightarrow -\frac{4}{3} = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}} \Rightarrow 4 \left(1 - \tan^2 \frac{x}{2} \right) = -6 \tan \frac{x}{2}$$

$$\Rightarrow 4 \tan^2 \frac{x}{2} - 6 \tan \frac{x}{2} - 4 = 0$$

$$\Rightarrow 2 \tan^2 \frac{x}{2} - 3 \tan \frac{x}{2} - 2 = 0$$

The equation is quadratic in $\tan \frac{x}{2}$

$$\Rightarrow \tan \frac{x}{2} = \frac{-(-3) \pm \sqrt{9+16}}{2.2} = \frac{3 \pm 5}{4} = 2, -\frac{1}{2}$$

Given $\frac{\pi}{2} \leq x \leq \pi \Rightarrow \frac{\pi}{4} \leq \frac{x}{2} \leq \frac{\pi}{2} \Rightarrow \frac{x}{2} \in \text{II}^{\text{st}} \text{ quadrant}$

In IIst quadrant, $\tan \frac{x}{2} \geq 0 \Rightarrow \tan \frac{x}{2} = 2$

We know, $1 + \tan^2 \theta = \sec^2 \theta$

$$\Rightarrow 1 + \tan^2 \frac{x}{2} = \sec^2 \frac{x}{2} \Rightarrow 1 + (2)^2 = \sec^2 \frac{x}{2}$$

$$\Rightarrow \sec^2 \frac{x}{2} = 5 \Rightarrow \sec \frac{x}{2} = \pm \sqrt{5} \Rightarrow \cos \frac{x}{2} = \pm \frac{1}{\sqrt{5}}$$

In IIst quadrant, $\cos \frac{x}{2} \leq 0 \Rightarrow \cos \frac{x}{2} = -\frac{1}{\sqrt{5}}$

We know $\sin \theta = \pm \sqrt{1 - \cos^2 \theta}$

$$\sin \frac{x}{2} = \pm \sqrt{1 - \cos^2 \frac{x}{2}} = \pm \sqrt{1 - \frac{1}{5}} = \pm \sqrt{\frac{4}{5}} = \pm \frac{2}{\sqrt{5}}$$

In IIst quadrant, $\sin \frac{x}{2} \geq 0 \Rightarrow \sin \frac{x}{2} = \frac{2}{\sqrt{5}}$

$$\therefore \text{(i) } \sin \frac{x}{2} = \frac{2}{\sqrt{5}} \quad \text{(ii) } \cos \frac{x}{2} = -\frac{1}{\sqrt{5}} \quad \text{(iii) } \tan \frac{x}{2} = 2$$

27. System of inequations

$$x \geq 0, y \geq 0, 5x + 3y \leq 500; x \leq 70 \text{ and } y \leq 125.$$


Converting inequations to equations

$$5x + 3y = 500 \Rightarrow y = \frac{500 - 5x}{3}$$

x	100	40	-80
y	0	100	300

$x \leq 70$ is $x = 70$ and $y \leq 125$ is $y = 125$.

Plotting these lines and determining the area of each line we get



OR

The amount of acid in 1125 lt of the 45% solution = $45\% \text{ of } 1125 = \frac{45 \times 1125}{100}$

Let x lt of the water be added to it to obtain a solution between 25% and 30% solution

$$\begin{aligned}
&\Rightarrow 25\% < \frac{1125 \times \frac{45}{100}}{1125 + x} < 30\% \\
&\Rightarrow \frac{25}{100} < \frac{1125 \times \frac{45}{100}}{1125 + x} < \frac{30}{100} \\
&\Rightarrow \frac{25}{100} < \frac{1125 \times 45}{(1125 + x) \times 100} < \frac{30}{100} \\
&\Rightarrow 25 < \frac{1125 \times 45}{(1125 + x)} < 30 \\
&\Rightarrow \frac{1}{25} > \frac{(1125 + x)}{1125 \times 45} > \frac{1}{30} \\
&\Rightarrow \frac{1125 \times 45}{25} > (1125 + x) > \frac{1125 \times 45}{30} \\
&\Rightarrow \frac{50625}{25} > (1125 + x) > \frac{50625}{30} \\
&\Rightarrow 2025 > (1125 + x) > 1687.5 \\
&\Rightarrow 2025 > (1125 + x) > 1687.5 \\
&\Rightarrow 2025 - 1125 > x > 1687.5 - 1125 \\
&\Rightarrow 900 > x > 562.5 \\
&\Rightarrow 562.5 < x < 900
\end{aligned}$$

So the amount of water to be added must be between 562.5 to 900 lt

28. Let $P(n)$: $5^n - 5$ is divisible by 4 be the given statement.

Let $n = 1$, $5 - 5 = 0$ is divisible by 4

$\Rightarrow P(1)$ is true

Let $P(k)$ be true i.e. $5^k - 5$ is divisible by 4

Let $5^k - 5 = 4m$

To prove the result for $n = k + 1$ we need to show that $5^{k+1} - 5$ is divisible by 4

$$\begin{aligned}
5^{k+1} - 5 &= 5^k \times 5 - 5 \\
&= (4m + 5) \times 5 - 5 \\
&= 20m + 25 - 5 \\
&= 4(5m + 5) \text{ is divisible by } 4
\end{aligned}$$

$\therefore$ Result holds for $n = k + 1$

$\therefore 5^n - 5$ is divisible by 4 for all n .

Let us take another statement

$P'(n)$: $2 \times 7^n + 3 \times 5^n - 5$ is divisible by 24.

For $n = 1$, $2 \times 7 + 3 \times 5 - 5 = 24$ is divisible by 24

$\therefore P'(1)$ is true

Let $P'(k)$ be true i.e., $2 \times 7^k + 3 \times 5^k - 5 = 24q$

To prove: $P'(k + 1)$ is also true

$$\begin{aligned}
\text{Consider } 2 \times 7^{k+1} + 3 \times 5^{k+1} - 5 &= 2 \times 7^k \times 7 + 3 \times 5^k \times 5 - 5 \\
&= (24q - 3 \times 5^k + 5)7 + 3 \times 5^k \times 5 - 5 \\
&= 24 \times 7q - 21 \times 5^k + 35 + 15 \times 5^k - 5 \\
&= 24 \times 7q - 6(5^k - 5) \quad (\text{Since } 5^k - 5 = 4p) \\
&= 24 \times 7q - 6 \times 4p \\
&= 24 (7q - p) \text{ which is a multiple of 24} \\
\therefore P'(k+1) \text{ is true.}
\end{aligned}$$

Hence by the principle of Mathematical Induction, the result holds true for all $n \in \mathbb{N}$.

29. a, b, and c are in A.P. $\Rightarrow 2b = a + c$

b, c, and d are in G.P. $\Rightarrow c^2 = bd$

$\frac{1}{c}, \frac{1}{d}$, and $\frac{1}{e}$ are in A.P. $\Rightarrow \frac{2}{d} = \frac{1}{c} + \frac{1}{e}$

$$d = \frac{2ce}{c+e}$$

$$c^2 = bd \Rightarrow c^2 = \frac{1}{2} (2b) d$$

$$= \frac{1}{2} (a+c) \left(\frac{2ce}{c+e} \right)$$

$$\Rightarrow (c+e) c^2 = ce (a+c)$$

$$\Rightarrow (c+e) c = e (a+c)$$

$$\Rightarrow c^2 + ec = ea + ec$$

$$\Rightarrow c^2 = ea$$

$\Rightarrow a, c$, and e are in G.P.

OR

$$\begin{aligned}
&\frac{1 \times 2^2 + 2 \times 3^2 + \dots + n \times (n+1)^2}{1^2 \times 2 + 2^2 \times 3 + \dots + n^2 \times (n+1)} \\
&= \frac{\sum_{k=1}^n k(k+1)^2}{\sum_{k=1}^n k^2(k+1)} \\
&= \frac{\sum k^3 + 2 \sum k^2 + \sum k}{\sum k^3 + \sum k^2} \\
&= \frac{\left[\frac{n(n+1)}{2} \right]^2 + 2 \left(\frac{n(n+1)(2n+1)}{6} \right) + \frac{n(n+1)}{2}}{\left(\frac{n(n+1)}{2} \right)^2 + \frac{n(n+1)(2n+1)}{6}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\frac{n(n+1)}{2} \left[\frac{n(n+1)}{2} + \frac{2(2n+1)}{3} + 1 \right]}{\frac{(n(n+1))}{2} \left[\frac{n(n+1)}{2} + \frac{2n+1}{3} \right]} \\
&= \frac{3n^2 + 3n + 8n + 4 + 6}{3n^2 + 3n + 4n + 2} \\
&= \frac{3n^2 + 11n + 10}{3n^2 + 7n + 2} \\
&= \frac{(n+2)(3n+5)}{(n+2)(3n+1)} = \frac{3n+5}{3n+1}
\end{aligned}$$