

# Arithmetic Progressions


---

- **The Concept of Arithmetic Progression**


- An arithmetic progression is a list of numbers in which the difference between any two consecutive terms is equal.
- In an AP, each term, except the first term, is obtained by adding a fixed number called common difference to the preceding term.
- The common difference of an AP can be positive, negative or zero.

**Example 1:**


1.

1.  is an AP whose first term and common difference are 3 and 3 respectively.

2.

2.  is an AP whose first term and common difference are 7 and -2 respectively.

4.

3.  is an AP whose first term and common difference are -7 and 0 respectively.

•

- The general form of an AP can be written as  $a, a + d, a + 2d, a + 3d \dots$ , where  $a$  is the first term and  $d$  is the common difference.
- A given list of numbers i.e.,  $a_1, a_2, a_3 \dots$  forms an AP if  $a_{k+1} - a_k$  is the same for all values of  $k$ .

**Example 2:**

Which of the following lists of numbers forms an AP? If it forms an AP, then write its next three terms.

(a) -4, 0, 4, 8, ...

(b) 2, 4, 8, 16, ...

**Solution:**

(a) -4, 0, 4, 8, ...

$$a_2 - a_1 = 0 - (-4) = 4$$

$$a_3 - a_2 = 4 - 0 = 4$$

$$a_4 - a_3 = 8 - 4 = 4$$

$$a_{n+1} - a_n = 4; \text{ for all values of } n$$

Therefore, the given list of numbers forms an AP with 4 being its common difference.

The next three terms of the AP are  $8 + 4 = 12$ ,  $12 + 4 = 16$ ,  $16 + 4 = 20$

Hence, AP:  $-4, 0, 4, 8, 12, 16, 20 \dots$

(b)  $2, 4, 8, 16, \dots$

$$a_2 - a_1 = 4 - 2 = 2$$

$$a_3 - a_2 = 8 - 4 = 4$$

$$a_3 - a_2 \neq a_2 - a_1$$

Therefore, the given list of numbers does not form an AP.

- **The terminology related to arithmetic progression**

- An arithmetic progression is a list of numbers in which each term is obtained by adding a fixed number to the preceding term except the first term.
- The fixed number is called the common difference ( $d$ ) of the A.P. The common difference can be either positive or negative or zero.

- **The general form of an A.P.**

- $a, (a + d), (a + 2d), (a + 3d), \dots, [a + (n - 1)d], \dots$  where  $a$  is the first term and  $d$  is common difference

- **Type of AP**

- Finite AP: The APs have finite number of terms.
- Infinite AP: The APs have not finite number of terms.

- In an A.P., except the first term, all the terms can be obtained by adding the common difference to the previous term.
- In an A.P., except the last term, all the terms can be obtained by subtracting the common difference from its subsequent term.

**Example:**

Find the first four terms of an A.P. whose first term is 9 and the common difference is 6.

**Solution:**

$$a = 9, d = 6$$

$$a_2 = a + d = 9 + 6 = 15$$

$$a_3 = a + 2d = 9 + 2 \times 6 = 9 + 12 = 21$$

$$a_4 = a + 3d = 9 + 3 \times 6 = 9 + 18 = 27$$

The first four terms are 9, 15, 21, 27.

- **$n^{\text{th}}$  term of an AP**

The  $n^{\text{th}}$  term ( $a_n$ ) of an AP with first term  $a$  and common difference  $d$  is given by  $a_n = a + (n - 1)d$ .

Here,  $a_n$  is called the general term of the AP.

- **$n^{\text{th}}$  term from the end of an AP**

The  $n^{\text{th}}$  term from the end of an AP with last term  $l$  and common difference  $d$  is given by  $l - (n - 1)d$ .

**Example:**

Find the 12<sup>th</sup> term of the AP 5, 9, 13 ...

**Solution:**

Here,  $a = 5$ ,  $d = 9 - 5 = 4$ ,  $n = 12$

$$\begin{aligned}
 a_{12} &= a + (n - 1) d \\
 &= 5 + (12 - 1) 4 \\
 &= 5 + 11 \times 4 \\
 &= 5 + 44 \\
 &= 49
 \end{aligned}$$

- **Sum of  $n$  terms of an AP**

- The sum of the first  $n$  terms of an AP is given by  $S_n = \frac{n}{2}(2a + n - 1)d$ , where  $a$  is the first term and  $d$  is the common difference.

- 

- If there are only  $n$  terms in an AP, then  $S_n = \frac{n}{2}(a + l)$ , where  $l = a_n$  is the last term.

**Example :**

Find the value of  $2 + 10 + 18 + \dots + 802$ .

**Solution:**

2, 10, 18... 802 is an AP where  $a = 2$ ,  $d = 8$ , and  $l = 802$ .

Let there be  $n$  terms in the series. Then,

$$\begin{aligned}
 a_n &= 802 \\
 \Rightarrow a + (n - 1) d &= 802 \\
 \Rightarrow 2 + (n - 1) 8 &= 802 \\
 \Rightarrow 8(n - 1) &= 800 \\
 \Rightarrow n - 1 &= 100 \\
 \Rightarrow n &= 101
 \end{aligned}$$

Thus, required sum  $= \frac{n}{2}(a + l) = \frac{101}{2}(2 + 802) = 40602$

- **Properties of an Arithmetic progression**

- If a constant is added or subtracted or multiplied to each term of an A.P. then the resulting sequence is also an A.P.
- If each term of an A.P. is divided by a non-zero constant then the resulting sequence is also an A.P.

- **Arithmetic mean**

- For any two numbers  $a$  and  $b$ , we can insert a number  $A$  between them such that  $a$ ,  $A$ ,  $b$  is an A.P. Such a number i.e.,  $A$  is called the arithmetic mean (A.M) of numbers  $a$  and  $b$  and it is given by  $A = \frac{a+b}{2}$ .

- For any two given numbers  $a$  and  $b$ , we can insert as many numbers between them as we want such that the resulting sequence becomes an A.P.

Let  $A_1, A_2, \dots, A_n$  be  $n$  numbers between  $a$  and  $b$  such that  $a, A_1, A_2, \dots, A_n, b$  is an A.P.

Here, common difference ( $d$ ) is given by  $\frac{b-a}{n+1}$ .

**Example:**

Insert three numbers between  $-2$  and  $18$  such that the resulting sequence is an A.P.

**Solution:**

Let  $A_1, A_2$ , and  $A_3$  be three numbers between  $-2$  and  $18$  such that  $-2, A_1, A_2, A_3, 18$  are in an A.P.

Here,  $a = -2, b = 18, n = 5$

$$\therefore 18 = -2 + (5 - 1) d$$

$$\Rightarrow 20 = 4 d$$

$$\Rightarrow d = 5$$

$$\text{Thus, } A_1 = a + d = -2 + 5 = 3$$

$$A_2 = a + 2d = -2 + 10 = 8$$

$$A_3 = a + 3d = -2 + 15 = 13$$

Hence, the required three numbers between  $-2$  and  $18$  are  $3, 8$ , and  $13$ .