



OXIDATION

(I) OXIDATION OF ALKENES



- * Cold dil. alkaline KMnO_4 is called as Bayer's reagent, use as a test of unsaturation.
- * Overall syn addition
- * Given by alkenes & alkynes
- * Benzene & Cyclopropane can not give this reaction.


If we use acidic KMnO_4 or warm KMnO_4 or concentrated KMnO_4 the oxidative cleavage of Glycol occurs resulting in mixture of Carboxylic acids & Ketones.






Hot acidic KMnO_4 , Hot acidic $\text{K}_2\text{Cr}_2\text{O}_7$ & hot acidic NaIO_4 gives same result with alkene. The effect is similar to that of oxidative ozonolysis on alkenes.






Preilschaive reaction :

Epoxidation of alkenes is reaction of alkenes with peroxyacids.




- Rate $\propto$ nucleophilicity of alkene.
- With the increase in e^- withdrawing substituents in leaving group, rate increases.
- Hydrolysis of epoxides form anti diols.




OXO PROCESS :**Hydroformylation/rolyen synthesis :****Carbonylation / Koch's reaction :****(II) OXIDATION OF ALCOHOLS****Oxidising agents**

- (1) $\text{H}^\oplus / \text{K}_2\text{Cr}_2\text{O}_7, \Delta$ (Strong oxidising agent)
- (2) $\text{H}^\oplus / \text{KMnO}_4, \Delta$ (Strong oxidising agent)
- (3) **Jones reagent** : $\text{CrO}_3 / \text{H}_2\text{SO}_4$ treated with alcohol usually taken in acetone (strong oxidising agent)
- (4) $\text{Cu} / 300^\circ\text{C}$ (or Red hot Cu tube)




- (5) PCC (Pyridinium chloro chromate)  $\text{CrO}_3^\ominus \text{Cl}^\ominus$ or  $\text{CrO}_3 + \text{HCl}$

- (6) Collin's reagent ( (2 mol) + $\text{CrO}_3 + \text{CH}_2\text{Cl}_2$)

- (7) Sarett reagent (i.e. PCC in CH_2Cl_2)  + $\text{CrO}_3 + \text{HCl} + \text{CH}_2\text{Cl}_2$

- (8) PDC
(Pyridinium dichromate)  $2^\ominus \text{Cr}_2\text{O}_7$


- (9) $\text{TsCl} + \text{DMSO} + \text{NaHCO}_3$



(10) MnO_2 -Oxidises only allylic or benzylic- OH. 1° Allylic or benzylic OH $\xrightarrow{\text{MnO}_2}$ Aldehyde


2° Allylic or benzylic OH $\xrightarrow{\text{MnO}_2}$ Ketone

No effect on 3° ROH and on Carbon-Carbon multiple bond.

(11) NBS (N-Bromosuccinamide) 



(12) **Periodic cleavage**

A similar oxidation is obtained in case of HIO_4 known as periodic cleavage.






Reaction is observed for Vic-diols.


(13) **Openaur oxidation**



Oxidation of alcohol with aluminium tertiary butoxide is Openaur oxidation.

Ex. Different oxidising agents are used to oxidise alcohols in corresponding carbonyl compounds and carboxylic acids.






(V) Double bond or Tripple bond is not affected by 4,5,6,7,8,9,10

(VI) No effect on 3° alcohol by 1,2,3,5,6,7,8,9,10,12,13

(III) OXIDATION OF CARBONYL COMPOUNDS

1. Tollen's Test (Silver mirror test)



Aldehyde acts as reducing agent, they can reduce mild oxidizing agents like Tollen's Reagent.

Tollen's test gentle Heating for 20 to 25 mins.


2. Fehling's Solutions

Fehling's A

aq. CuSO_4


Fehling's B

Alkaline solution of Roschelye
salt (sodium potassium tartrate)



It act's a carrier for Cu^{2+} as it make reversible complex with Cu^{2+}


This test is also used is Blood and Urine test.




Note : Aromatic aldehyde shows negative test with fehling reagent.

3. Benedict's solution

Sodium Citrate + NaOH + NaHCO_3 + CuSO_4




It is similar to Fehling test



5. Schiff's Reagent

Schiff's Reagent is aq. solution of following base decolourised by passing SO_2 .

Aldehyde restore pink colour of Schiff's reagent.




Ketons are not easy to oxidize so they do not give these 5 tests. These five tests can be used to distinguish aldehyde and ketones. Both gives 2,4 DNP test

Oxidation of ketones : Ketones undergo oxidation only in drastic conditions.


During the oxidation of ketones there is breaking of carbon-carbon bond between α -carbon and carbonyl carbon. In this process both carbons convert into carboxylic groups. This leads to the formation of two moles of monocarboxylic acids.

Case - I : Oxidation of Symmetrical ketones :




Thus number of carbons in any product is less than the number of carbons in ketone.


Case - II : Oxidation of unsymmetrical ketones : In case of unsymmetrical ketones, those ---C(=O)---R bond break in which alkyl group has more number of carbons. This rule is known as **Poff's rule**.



Case - III : Oxidation of cyclic ketones : Formation of dibasic acid takes place from cyclic ketones. In this case number of carbons in ketone and dibasic carboxylic acid is always same.




Note : If both α -carbons are not identical then ;




Allylic oxidation

SeO_2 is a selective oxidizing agent which converts $-\text{CH}_2-$ group adjacent to carbonyl group into carbonyl group. The reagent, in general, oxidises active methylene and methyl groups to ketonic and aldehydic groups respectively.






Double bonds, triple bonds and aromatic rings may also activate the methylene group. The methylene or methyl group α to the most highly substituted end of the double bond is hydroxylated according to the order of preference of oxidation $\text{CH}_2 > \text{CH}_3 > \text{CH}$ groups.




$2^\circ \text{C} - \text{H} > 1^\circ \text{C} - \text{H} > 3^\circ \text{C} - \text{H}$

Rate of reactivity order











Baeyer villiger oxidation



Migratory order for above reaction







$-\text{H} > 3^\circ > 2^\circ > -\text{Ph} > 1^\circ > -\text{Me}$

Oxidation of Aromatic Compounds.




(10) $\text{Ph}-(\text{CH}_2)_n\text{CH}_3$, PhCHMe_2 , $\text{Ph}-\text{CH}_2\text{OH}$, $\text{Ph}-\text{CH}_2\text{Br}$, PhCHCl_2 , PhCHO , $\text{Ph}-\triangle$ & all the compounds having at least one α H give PhCOOH .






Etard oxidation




Elb's persulphate oxidation:



EXERCISE#O-I

1. The given reaction




is known as :

- (A) Kolbe reaction (B) Tischenko reaction (C) MPV reaction (D) Oppenauer oxidation

OX0001

2. In the given reaction :



(a) and (b) respectively be :

- (A) CH₃CHO and CH₃CHO (B) CH₃COCH₃ and CH₃CHO
(C) CH₃COCH₃ and CH₃COCH₃ (D) CH₃COOH and CH₃COCH₃


OX0002

3. CH₃-C(=O)-H and CH₃-C(=O)-CH₃ is differentiated by

- (A) Tollen's reagent (B) Lucas test (C) Iodoform test (D) NaHSO₃ test

OX0003

4. In the given reaction



[X] will be:

- (A) (B) (C) (D)

OX0004

5. Which will give silver mirror test with Tollens reagent :

- (A) C₆H₅CHO (B) CH₃-CHO (C) HCOOH (D) All of these

OX0005

6. Which one of the following compounds will give dimethyl glyoxal with SeO_2 :

- (A) Acetone (B) Acetophenone
(C) Ethyl methyl ketone (D) Propanaldehyde

OX0006


7. Which one of the following alcohols can be oxidised by K_2CrO_4 ?

- (A) Ethanol (B) Tert butyl alcohol
(C) Isopropyl alcohol (D) Allyl alcohol

OX0007

8. Fehling solution gives red precipitate with:


- (A) Aromatic aldehyde (B) Aliphatic aldehyde (C) Ketones




OX0008

9. Which of the following compound will give positive Tollens test

- (A) CH_3CHO



- (C) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\underset{\text{OH}}{\text{CH}} - \text{OCH}_3$



OX0009

10. Schiff's reagent gives pink colour with :

- (A) Acetaldehyde (B) Formic acid (C) Acetic acid (D) Methyl acetate

OX0010

11. Which one of the following alcohols are oxidised by MnO_2 ?

- (A) $\text{C}_6\text{H}_5 - \text{CH}_2 - \text{CH}_2 - \text{OH}$






- (C) $\text{CH}_3 - \text{CH} = \underset{\text{OH}}{\text{CH}} - \text{CH}_3$

- (D) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{OH}$


OX0011

12. $\xrightarrow{\text{MCPBA}}$ Major product is :





OX0012

13. Which of the following can be oxidised by MnO_2 :




OX0013



OX0014





15. $\text{A} \xrightarrow{\text{H}^+ / \text{KMnO}_4} \text{B} \xrightarrow{\Delta} \text{Phthalic Anhydride}$; A is :



OX0015


EXERCISE#O-II

1. Which will give the Tollen test.




OX0016


2. Which of the following is an example of oxidation reaction :



OX0017




OX0018




OX0019




OX0020




OX0020




OX0021




OX0021




OX0022




OX0022



OX0023



OX0023




OX0023


4. A to F alkenes with minimum possible carbon.



OX0024




OX0024




OX0025




OX0025




OX0026




OX0026




OX0027




OX0027




OX0028




OX0028




OX0029




OX0029




OX0030




OX0030




OX0030




OX0031




OX0031




OX0031




OX0032




OX0032




OX0032




OX0032




OX0033




OX0033




OX0033




OX0034




OX0034



OX0034



OX0035



OX0036


10. (a) How will you differentiate HCHO and PhCHO ?
 (b) How will you differentiate HCHO and MeCHO ?

OX0037

EXERCISE# (MAIN & ADVANCE)

1. In the reaction, P is

[IIT 1995]



- (A)
- CH_3COCHO
- (B)
- $\text{CH}_3\text{COOCH}_3$
- (C)
- $\text{CH}_3\text{COCH}_2\text{OH}$
- (D) None

OX0038

2. The best reagent to convert pent-3-en-2-ol into pent-3-en-2-one is -


[AIEEE-2005]

- (A) Acidic dichromate (B) Acidic permanganate
(C) Pyridinium chloro-chromate (D) Chromic anhydride in glacial acetic acid


OX0039

3. Which of the following will change the colour of acidic dichromate solution.


[JEE-MAINS 2013]




OX0040

- 4.
- 
- $\xrightarrow{\text{MCPBA}}$
- Product


[JEE-MAINS 2013]



OX0041

- 5.
- 
- $\xrightarrow[\Delta]{\text{KMnO}_4}$
- Major product of reaction is?

[JEE-MAINS 2013]



OX0042

6. The most suitable reagent for the conversion of
- $\text{R-CH}_2\text{-OH} \rightarrow \text{R-CHO}$
- is :-


[JEE-MAINS 2014]

- (A) CrO_3 (B) PCC (Pyridinium chlorochromate)
(C) KMnO_4 (D) $\text{K}_2\text{Cr}_2\text{O}_7$

OX0043

7. Positive Tollen's test is observed for


[JEE-Advance 2016]



OX0044

8. The major product formed in the following reaction is :-


[Jee Main - Onl_line Evening 2018]



OX0045

9. The major product of the following reaction is:






[JEE-MAIN-On-line-(Jan)-2019]



OX0046

10. The major product of the following reaction


[JEE-MAIN-On-line-(April)-2019]



OX0047

11. Total number of hydroxyl groups present in a molecule of the major product P is ____


[JEE-Advance 2019]



OX0048





12. An unsaturated hydrocarbon X absorbs two hydrogen molecules on catalytic hydrogenation, and also gives following reaction :

[JEE-MAIN-On-line-(Jan)-2020]



B(3-oxo-hexanedicarboxylic acid)






X will be :-



OX0049

13. Identify (A) in the following reaction sequence :

[JEE-MAIN-On-line-(Jan)-2020]



OX0050

ANSWER-KEY









EXERCISE#O-I

- | | | | |
|-----------------|----------------|-----------------|---------------|
| 1. Ans. (D) | 2. Ans. (B) | 3. Ans. (A) | 4. Ans. (B) |
| 5. Ans. (D) | 6. Ans. (C) | 7. Ans. (A,C,D) | 8. Ans. (B,D) |
| 9. Ans. (A,B,C) | 10. Ans. (A,B) | 11. Ans. (C) | 12. Ans. (B) |
| 13. Ans. (C) | 14. Ans. (D) | 15. Ans. (C) | |


EXERCISE#O-II

1. Ans. (A,C,D) 2. Ans. (C)





3. Ans.




4. Ans.









5. Ans.






6. Ans.








7.



8. Ans.



9. Ans.



10. (a) Ans. F and B test


(b) Ans. Iodoform test

EXERCISE# (MAIN & ADVANCE)

1. Ans. (A)


2. Ans. (D)

3. Ans. (A)



3° alcohols do not oxidize
they undergo dehydration

4. Ans. (C)






5. Ans. (A)

6. Ans. (B)

7. Ans. (A,B,C)

Sol. Tollens's test is given by compounds having aldehyde group. Also α -hydroxy carbonyl gives positive tollens's test.



8. Ans. (D)


9. Ans. (2)

10. Ans. (4)

11. Ans. (6.00)


12. Ans. (1)

Sol.



13. Ans. (4)

Sol.



REDUCING AGENTS AND THEIR ROLE

Group	Product	LAH in Ether	NaBH ₄ in H ₂ O	B ₂ H ₆ in THF	H ₂ / Catalyst Δ
–CHO	–CH ₂ OH	+	+	+	+
>C = O	>CH – OH	+	+	+	+
–CO ₂ H	–CH ₂ OH	+	–	+	–
–CO ₂ R	–CH ₂ OH	+	–	+	+
–COCl	–CH ₂ OH	+	+	–	+
–CONH ₂	–CH ₂ NH ₂	+	–	+	+
(RCO) ₂ O	RCH ₂ OH	+	–	+	+
–CN	–CH ₂ NH ₂	+	–	+	+
>C = NOH	–CH ₂ NH ₂	+	–	–	+
>C = C<	>CH – CH<	–	–	+	+
–C ≡ C–	–CH = CH–	–	–	+	+
1° RX	RH	+	–	–	+

Note :



1. B₂H₆ in THF not able to reduce cyclic ester.
2. NaBH₄ can also reduce imine group (>C = NH).
3. Reactivity order towards Ni / H₂ , Δ ⇒ $-\text{C} \equiv \text{N} > -\overset{\text{O}}{\underset{\text{O}}{\text{C}}} - \text{O} - \text{R}$

Name	Reagent	Function
Wolf Kishner Reduction	(i) N_2H_4 / (ii) KOH, Δ	$\text{>C=O} \longrightarrow \text{>CH-CH}_3$
Clemenson Reduction	$\text{Zn-Hg} / \text{HCl}$	$\text{>C=O} \longrightarrow \text{>CH-CH}_3$
Mozingo Reduction	$\begin{array}{ c} \text{SH} \\ \text{---} \\ \text{SH} \end{array}$ Dry HCl, followed by Raney Ni	$\text{>C=O} \longrightarrow \text{>CH-CH}_3$
Stephen's Reduction	$\text{SnCl}_2 / \text{HCl}$ followed by H_3O^+	$\text{R-C}\equiv\text{N} \longrightarrow \text{R-CH=O}$
MPV Reduction	$\text{Al}\left(\text{O-CH}\begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}\right)_3 / \text{HO-CH}\begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}$	$\text{>C=O} \longrightarrow \text{>CH-OH}$
Hydroboration Reduction	$\text{B}_2\text{H}_6 / \text{AcOH}, \text{H}_2\text{O}$	$\text{>C=C<} \longrightarrow \text{>CH-CH<}$ $\text{>C=O} \longrightarrow \text{>CH-OH}$
Bouvault Blank Reduction	Na / EtOH	$\text{R-COO-R} \longrightarrow \text{RCH}_2\text{OH} + \text{ROH}$
Transfer Hydrogenation	$\text{N}_2\text{H}_4 / \text{H}_2\text{O}_2$	$\text{>C=C<} \longrightarrow \text{>CH-CH<}$
Rosenmund Reduction	$\text{H}_2, \text{Pd-BaSO}_4$	$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{Cl} \longrightarrow \text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H}$ $*-\text{C}\equiv\text{C}- \longrightarrow \text{>CH-CH<}$
Birch Reduction	$\text{Na} / \text{Liq. NH}_3$	$-\text{C}\equiv\text{C}- \longrightarrow \text{>CH=CH<}$
DIBAL-H (-78°C)	$\text{H-Al}\left(\text{CH}_2\text{-CH}\begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}\right)_2$ followed by H_3O^+	$\left. \begin{array}{l} -\text{COOR} \\ -\text{C}\equiv\text{N} \\ -\text{COCl} \\ -\text{C}(=\text{O})-\text{O}-\text{C}(=\text{O})- \end{array} \right\} \longrightarrow -\text{CHO}$ $\left. \begin{array}{l} -\text{CHO} \end{array} \right\} \longrightarrow -\text{CH}_2\text{OH}$
Red phosphorus in presence of HI	Red P + HI	$\text{R-CO}_2\text{H} \longrightarrow \text{RCH}_3$ $\text{R-CH=O} \longrightarrow \text{RCH}_3$ $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{R} \longrightarrow \text{RCH}_2\text{R}$ $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH} \longrightarrow \text{R-H}$


EXERCISE # O-I



The end products of the reaction are :



RD0001



Products of the above reaction will be :





(A) Racemic mixture

(B) Diastereomers

(C) Meso

(D) Structural isomer

RD0002



(D) Both (B) and (C)

RD0003

4. On catalytic reduction with H_2/Pt how many alkenes will give n-butane ?


(A) 1

(B) 2

(C) 3

(D) 4

RD0004



Product of above reaction will be :


(A) Racemic mixture

(B) Diastereomers

(C) Meso

(D) Constitutional isomers

RD0005



Product of above reaction will be :


(A) Racemic mixture

(B) Diastereomers

(C) Meso

(D) Constitutional isomers


RD0006







Which reagent will be used for the above conversion ?


- (A) Na/Liq. NH_3 (B) H_2 , Pd- CaCO_3 (C) Li, Ph- NH_2 (D) H_2 , Pt




RD0007




- (A)  (B)  (C)  (D) 

RD0008










- (A)  (B)  (C)  (D) None of these

RD0009




A and B respectively are :

- (A) Both  (B)  , 
 (C)  ,  (D)  , 





RD0010

11. If the following compound is treated with Pd in excess of H_2 gas, how many stereoisomers of the product will be obtained ?



- (A) 1 (B) 2 (C) 3 (D) 4




RD0011



(D) All of these


RD0012

13. Which alkyne gives 3-ethylhexane on catalytic hydrogenation ?







(D) All of these


RD0013






The product is :



RD0014




The final product (B) is :




RD0015

EXERCISE # O-II

1. Column-I (Conversion)	Column-II (Required Reagent)
(A) $\text{H}_3\text{C}-\text{C}\equiv\text{C}-\text{CH}_3 \longrightarrow \begin{array}{c} \text{H}_3\text{C} \quad \text{CH}_3 \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$	(P) H_2 , Pd-BaSO ₄
(B) $\text{H}_3\text{C}-\text{C}\equiv\text{C}-\text{CH}_3 \longrightarrow \begin{array}{c} \text{H}_3\text{C} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{CH}_3 \end{array}$	(Q) Li, Liq.NH ₃
(C) $\text{H}_3\text{C}-\text{CH}=\text{CH}-\text{CH}_3 \longrightarrow \text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}_3$	(R) N ₂ H ₄ / H ₂ O ₂ (1 eq.)
(D) 	(S) B ₂ H ₆ , CH ₃ COOH

RD0016



Identify numbers of reagent that can be used for above conversion.






- (a) Zn-Hg / HCl (b) LiAlH₄ (c) N₂H₄ / $\text{OH}^\ominus$ (d) $\begin{array}{c} \text{CH}_2-\text{SH} \\ | \\ \text{CH}_2-\text{SH} \end{array}$, H₂ / Ni

RD0017

3. Column-I (Conversion)	Column-II (Required Reagent)
(A) $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH} \longrightarrow \text{R}-\text{CH}_2\text{OH}$	(P) LiAlH ₄
(B) $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{Cl} \longrightarrow \text{R}-\text{CH}_2\text{OH}$	(Q) NaBH ₄
(C) $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OR}' \longrightarrow \text{R}-\text{CH}_2\text{OH}$	(R) B ₂ H ₆ / THF
(D) $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{R} \longrightarrow \text{R}-\text{CH}_2\text{OH}$	(S) H ₂ / Catalyst, Δ


RD0018

4. How many alkene on catalytic reduction gives following product.




RD0019


5. Give the expected major product for each reaction, including stereochemistry where applicable.




RD0020




RD0021




RD0022




RD0023



RD0024




RD0025




RD0026


7. (a) Identify the product ?




RD0027



RD0028



RD0029




RD0030

(b) Identify the product ?

RD0031




RD0031




RD0031



RD0032




RD0032




RD0032

8. Give product in following reactions.


RD0033




RD0034




RD0033




RD0034




RD0033



RD0034




RD0033




RD0034

9. Give product in following reactions.




RD0035


10. Give product in following reactions.




RD0036



RD0037




RD0038




RD0039


11. Give product in following reactions.




RD0040




RD0041




RD0042




RD0043




RD0043






RD0043




RD0044



RD0045



RD0046



above conversion can be achieved by

(A) LiAlH_4

(B) $\text{H}_2 / \text{Ni}, \Delta$

(C) NaBH_4

(D) All


RD0047

EXERCISE # (MAIN & ADVANCE)

1. Hydrogenation of benzoyl chloride in the presence of Pd and BaSO₄ gives : [IIT 1992]
 (A) Benzyl alcohol (B) Benzaldehyde (C) Benzoic acid (D) Phenol

RD0048

2. Draw the stereochemical structure of the product in the following reactions. [IIT 1994]







RD0049

3. Under Wolff Kishner reduction conditions, the conversions which may be brought about is?
 (A) Benzaldehyde into Benzyl alcohol (B) Cyclohexanol into Cyclohexane
 (C) Cyclohexanone into Cyclohexanol (D) Benzophenone into Diphenylmethane [IIT 1995]


RD0050

4. Which one of the following alkenes will react fastest with H₂ under catalytic hydrogenation condition [IIT 2000]




RD0051

5. The appropriate reagent for the following transformation: [IIT-2000]



- (A) Zn(Hg), HCl (B) NH₂NH₂, OH⁻ (C) H₂/Ni (D) NaBH₄

RD0052



Hydrogenation of the above compound in the presence of poisoned palladium catalyst gives-

- (A) An optically active compound (B) An optically inactive compound [IIT-2001]
 (C) A racemic mixture (D) A diastereomeric mixture

RD0053

7. When CH₂=CH-COOH is reduced with LiAlH₄, the compound obtained will be -[AIEEE-2003]
 (A) CH₃-CH₂-CH₂OH (B) CH₃-CH₂-CHO
 (C) CH₃-CH₂-COOH (D) CH₂=CH-CH₂OH


RD0054

8. 2-hexyne can be converted into trans-2-hexene by the action of : [IIT-2004]
 (A) H₂-Pd-BaSO₄ (B) Li in liq. NH₃ (C) H₂-PtO₂ (D) NaBH₄

RD0055





Paragraph for Question No. 09 to 11

In the following reaction sequence, products **I**, **J** and **L** are formed. **K** represents a reagent.



9. The structure of the product **I** is





[JEE-2008]



RD0056

10. The structures of compounds **J** and **K**, respectively, are





[JEE-2008]



RD0056

11. The structure of product **L** is :

[JEE-2008]



RD0056


12. Consider all possible isomeric ketones including stereoisomers of MW = 100, All these isomers are independently reacted with NaBH_4 (**NOTE** : stereoisomers are also reacted separately). The total number of ketones that give a racemic product(s) is/are.

[JEE-2014]

RD0057

13. In the reaction,

[JEE-2014]




- (A) Ethylene (B) Acetyl chloride (C) Acetaldehyde (D) Acetylene

RD0058

14. The reagent needed for converting is :

[JEE-2014]



- (A) H_2 / Lindlar Cat.

- (B) Cat. hydrogenation


- (C) LiAlH_4

- (D) Li / NH_3

RD0059

15. The major product obtained in the following reaction is :

[JEE MAIN-2017]



RD0060

16. Glucose on prolonged heating with HI gives :

[Jee Main - 2018]

- (1) 1-Hexene (2) Hexanoic acid (3) 6-iodohexanal (4) n-Hexane

RD0061

17. The *trans*-alkenes are formed by the reduction of alkynes with :


[Jee Main - 2018]

- (1) NaBH_4 (2) Na/liq. NH_3 (3) Sn-HCl (4) $\text{H}_2\text{-Pd/C, BaSO}_4$

RD0062

18. The main reduction product of the following compound with NaBH_4 in methanol is :-


[Jee Main - Onl_line Morning 2018]



RD0063

19. The major product of following reaction is :

[JEE-MAIN-On-line-(Jan)-2019]



- (1) RCHO (2) RCOOH (3) RCH_2NH_2 (4) RCONH_2

RD0064

20. Wilkinson catalyst is :


[JEE-MAIN-On-line-(Jan)-2019]

- (1) $[(\text{Ph}_3\text{P})_3\text{RhCl}]$ ($\text{Et} = \text{C}_2\text{H}_5$) (2) $[\text{Et}_3\text{P}]_3\text{IrCl}]$
(3) $[\text{Et}_3\text{P}]_3\text{RhCl}]$ (4) $[\text{Ph}_3\text{P}]_3\text{IrCl}]$

RD0065

21. The major product 'X' formed in the following reaction is :

[JEE-MAIN-On-line-(Jan)-2019]




- (1) (2) (3) (4)

RD0066

22. The major product of the following reaction is:


[JEE-MAIN-On-line-(Jan)-2019]



RD0067

23. The major product of the following reaction is:


[JEE-MAIN-On-line-(Jan)-2019]



RD0068

24. The major product obtained in the following reaction is :-






[JEE-MAIN-On-line-(Jan)-2019]



RD0069

25. The major product of the following reaction is:





[JEE-MAIN-On-line-(Jan)-2019]



RD0070

26. The increasing order of the reactivity of the following with LiAlH_4 is :

[JEE-MAIN-On-line-(Jan)-2019]



(1) (A) < (B) < (D) < (C)

(2) (A) < (B) < (C) < (D)






(3) (B) < (A) < (D) < (C)

(4) (B) < (A) < (C) < (D)

RD0071

27. The major product of the following reaction is:






[JEE-MAIN-On-line-(Jan)-2019]



RD0072

28. The major product of the following reaction is:





[JEE-MAIN-On-line-(April)-2019]



RD0073

29. The major product of the following reaction is :

[JEE-MAIN-On-line-(April)-2019]







RD0074

30. The major products A and B for the following reactions are, respectively:







[JEE-MAIN-On-line-(April)-2019]



RD0075

31. What is the product of following reaction ?



[JEE-MAIN-On-line-(Jan)-2020]



RD0076

32. The most suitable reagent for the given conversion is :

[JEE-MAIN-On-line-(Jan)-2020]



(1) LiAlH_4

(2) NaBH_4

(3) H_2/Pd

(4) B_2H_6

RD0077

ANSWER KEY


EXERCISE # O-I

1. Ans. (D)	2. Ans. (B)	3. Ans. (C)	4. Ans. (C)	5. Ans. (A)
6. Ans. (C)	7. Ans. (B)	8. Ans. (B)	9. Ans. (C)	10. Ans. (B)
11. Ans. (C)	12. Ans. (C)	13. Ans. (D)	14. Ans. (D)	15. Ans. (B)

EXERCISE # O-II

1. Ans. (A)→P, R, S; (B)→Q; (C)→R, S; (D)→Q 2. Ans. (3)
3. Ans. (A)→P, R, S; (B)→P, Q, S; (C)→P, R, S; (D)→P, R, S
4. Ans.
- (i) Cis & trans 2-butene & 1-butene ; (ii) $\begin{array}{c} \text{C}-\text{C}-\text{C}=\text{C} \\ | \\ \text{C} \end{array}$ $\begin{array}{c} \text{C}-\text{C}=\text{C}-\text{C} \\ | \\ \text{C} \end{array}$ $\begin{array}{c} \text{C}-\text{C}-\text{C}-\text{C} \\ || \\ \text{C} \end{array}$
- (iii) Zero (Neo-pentane can not be prepared by catalytic hydrogenation of alkene) ; (iv) One
- (v) Including optical = 4  , Excluding optical = 3.





5. Ans.






6. Ans. (i) $\text{P}_1 = \text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$

(ii) $\text{P}_2 = \text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}_2$






8. Ans.



10. Ans.



12. Ans.






15. Ans.(C)

EXERCISE # (MAIN & ADVANCE)

1. Ans. (B) 2. Ans. $\begin{array}{c} \text{R} \\ \diagdown \\ \text{C} = \text{C} \\ \diagup \\ \text{H} \end{array}$ 3. Ans. (D) 4. Ans. (A)
5. Ans. (B) 6. Ans. (B) 7. Ans. (D) 8. Ans. (B)
9. Ans. (D) 10. Ans. (A) 11. Ans. (C)
12. Ans. (5)


M. wt 100 of ketone

So m. formula = $\text{C}_6\text{H}_{12}\text{O}$



(1 ; 2 ; 3 ; 6 ; 7)

13. Ans. (A)



Reduction of CH_3COOH will produce $\text{CH}_3\text{CH}_2\text{OH}$ (A)

Now reaction of $\text{CH}_3\text{CH}_2\text{OH}$ (A) with PCl_5 will produce $\text{CH}_3\text{CH}_2\text{Cl}$ (B).

Now alcoholic KOH with Et-Cl will produce $\text{CH}_2=\text{CH}_2$ (C) Ethylene by E_2 elimination.


Thus Ans. is (1) Ethylene.

14. Ans. (D)


15. Ans. (1)

16. Ans. (4)

Sol.



17. Ans. (2)



Birch reduction is anti addition. So trans alkene will be produced.

18. Ans. (4) 19. Ans. (1) 20. Ans. (1) 21. Ans. (4)


22. Ans. (3) 23. Ans. (2) 24. Ans. (2) 25. Ans. (3)

26. Ans. (1) 27. Ans. (4) 28. Ans. (4) 29. Ans. (2)

30. Ans. (2)


31. Ans. (3)

Sol.



32. Ans. (4)

Sol.



Most suitable reagent for given conversion is B_2H_6 (electrophilic reducing agent)