

DPP - Daily Practice Problems

Name :

Date :

Start Time :

End Time :

CHEMISTRY

15

SYLLABUS : Redox Reactions

Max. Marks : 120

Time : 60 min.

GENERAL INSTRUCTIONS

- The Daily Practice Problem Sheet contains 30 MCQ's. For each question only one option is correct. Darken the correct circle/bubble in the Response Grid provided on each page.
- You have to evaluate your Response Grids yourself with the help of solution booklet.
- Each correct answer will get you 4 marks and 1 mark shall be deducted for each incorrect answer. No mark will be given/ deducted if no bubble is filled. Keep a timer in front of you and stop immediately at the end of 60 min.
- The sheet follows a particular syllabus. Do not attempt the sheet before you have completed your preparation for that syllabus. Refer syllabus sheet in the starting of the book for the syllabus of all the DPP sheets.
- After completing the sheet check your answers with the solution booklet and complete the Result Grid. Finally spend time to analyse your performance and revise the areas which emerge out as weak in your evaluation.

DIRECTIONS (Q.1-Q.21) : There are 21 multiple choice questions. Each question has 4 choices (a), (b), (c) and (d), out of which ONLY ONE choice is correct.


Q.1 When $K_2Cr_2O_7$ is converted into K_2CrO_4 the change in oxidation number of Cr is—

- (a) 0 (b) 3 (c) 4 (d) 6

Q.2 The oxidation number of S in $(CH_3)_2SO$ is—


- (a) 1 (b) 2 (c) 0 (d) 3

Q.3 What will be the value of a, b, c, d, e and f in the following equation ?




- (a) 2, 2, 10, 8, 5, 16 (b) 2, 5, 16, 2, 10, 8
(c) 2, 5, 10, 2, 8, 16 (d) 2, 8, 16, 2, 5, 10

Q.4 What will be the equivalent weight of permanganate ion in the following redox reaction ?



- (a) M/5 (b) M/6
(c) M/8 (d) 5M

Q.5 What will be the equivalent weight of the reducing agent which donates one electron in the following chemical reaction?




- (a) 2M (b) 3M
(c) M/2 (d) M

RESPONSE GRID

1. (a)(b)(c)(d) 2. (a)(b)(c)(d) 3. (a)(b)(c)(d) 4. (a)(b)(c)(d) 5. (a)(b)(c)(d)

Q.6 In the following reaction, what is the equivalent weight of oxidant and reductant respectively ?



[V = 50.94, Zn = 65.38 and O = 16]

- (a) 18.2, 32.69 (b) 30, 20
(c) 34.10, 20.2 (d) 40, 10


Q.7 Which of the following acts as both oxidant and reductant?

- (a) HNO_3
(b) HNO_2
(c) Both HNO_3 & HNO_2
(d) Neither HNO_3 nor HNO_2

Q.8 State which of the following reactions is neither oxidation nor reduction ?

- (a) $\text{Na} \rightarrow \text{NaOH}$
(b) $\text{Cl}_2 \rightarrow \text{Cl}^- + \text{ClO}_3^-$
(c) $\text{P}_2\text{O}_5 \rightarrow \text{H}_4\text{P}_2\text{O}_7$
(d) $\text{Zn} + \text{H}_2\text{SO}_4 \rightarrow \text{ZnSO}_4 + \text{H}_2$

Q.9 In the reaction



the reductant is –

- (a) $\text{C}_2\text{O}_4^{2-}$ (b) H^+
(c) MnO_4^- (d) None of the above

Q.10 What is the oxidation state of nitrogen in NaN_3 ?

- (a) $-3/1$ (b) 3
(c) -3 (d) $-1/3$

Q.11 What is the oxidation number of oxygen in OF_2 ?

- (a) +2 (b) +4
(c) +3 (d) None

Q.12 Oxidation number of cobalt in $[\text{Co}(\text{NH}_3)_6]\text{Cl}_2\text{Br}$ is –

- (a) +6 (b) zero
(c) +3 (d) +2

Q.13 The order of increasing O.N. of S in S_8 , $\text{S}_2\text{O}_8^{2-}$, $\text{S}_2\text{O}_3^{2-}$, $\text{S}_4\text{O}_6^{2-}$ is :

- (a) $\text{S}_8 < \text{S}_2\text{O}_8^{2-} < \text{S}_2\text{O}_3^{2-} < \text{S}_4\text{O}_6^{2-}$
(b) $\text{S}_2\text{O}_8^{2-} < \text{S}_2\text{O}_3^{2-} < \text{S}_4\text{O}_6^{2-} < \text{S}_8$
(c) $\text{S}_2\text{O}_8^{2-} < \text{S}_8 < \text{S}_4\text{O}_6^{2-} < \text{S}_2\text{O}_3^{2-}$
(d) $\text{S}_8 < \text{S}_2\text{O}_3^{2-} < \text{S}_4\text{O}_6^{2-} < \text{S}_2\text{O}_8^{2-}$

Q.14 The composition of a sample of wustite is $\text{Fe}_{0.93}\text{O}_{1.00}$. What percentage of iron is present in the form of Fe (III) ?

- (a) 13.05 (b) 14.05
(c) 15.05 (d) 16.05

Q.15 The O.N. of Cl in NOClO_4 is –

- (a) +11 (b) +9
(c) +7 (d) +5

Q.16 The two possible oxidation numbers of N atoms in NH_4NO_3 are respectively –

- (a) +3, +5 (b) +3, -5
(c) -3, +5 (d) -3, -5

Q.17 The oxidation number of S in $\text{H}_2\text{S}_2\text{O}_8$ is –

- (a) +8 (b) -8
(c) +6 (d) +4

Q.18 What will be the oxidation number of I in the KI_3 ?

- (a) $-\frac{1}{3}$ (b) $-\frac{1}{4}$ (c) +4 (d) +3

Q.19 Oxidation number of Fe in $[\text{Fe}(\text{CN})_6]^{-3}$, $[\text{Fe}(\text{CN})_6]^{-4}$, $[\text{Fe}(\text{SCN})]^{+2}$ and $[\text{Fe}(\text{H}_2\text{O})_6]^{+3}$ respectively would be –

- (a) +3, +2, +3 and +3 (b) +3, +3, +3 and +3
(c) +3, +2, +2 and +2 (d) +2, +2, +2 and +2

**RESPONSE
GRID**

6. (a)(b)(c)(d) 7. (a)(b)(c)(d) 8. (a)(b)(c)(d) 9. (a)(b)(c)(d) 10. (a)(b)(c)(d)
11. (a)(b)(c)(d) 12. (a)(b)(c)(d) 13. (a)(b)(c)(d) 14. (a)(b)(c)(d) 15. (a)(b)(c)(d)
16. (a)(b)(c)(d) 17. (a)(b)(c)(d) 18. (a)(b)(c)(d) 19. (a)(b)(c)(d)

Q.20 In the redox reaction –



The values of x and y are respectively –

- (a) 6,3 (b) 3,6 (c) 3,3 (d) 6,6

Q.21 A solution containing 2.68×10^{-3} mol of A^{+n} ions requires

1.61×10^{-3} mole of MnO_4^- for the oxidation of A^{+n} to AO_3^- in acidic medium. What is the value of n ?


- (a) 1 (b) 2 (c) 3 (d) 4

DIRECTIONS (Q.22-Q.24) : In the following questions, more than one of the answers given are correct. Select the correct answers and mark it according to the following codes:

Codes :


- (a) 1, 2 and 3 are correct (b) 1 and 2 are correct
(c) 2 and 4 are correct (d) 1 and 3 are correct

Q.22 Choose the correct value of x , y and z in the following equation –



- (1) $x=1$ (2) $y=2$ (3) $z=2$ (4) $z=4$

Q.23 Consider the redox reaction :




- (1) $\text{S}_2\text{O}_3^{2-}$ gets oxidised to $\text{S}_4\text{O}_6^{2-}$
(2) $\text{S}_2\text{O}_3^{2-}$ gets reduced to $\text{S}_4\text{O}_6^{2-}$
(3) I_2 gets reduced to I^-
(4) I_2 gets oxidised to I^-

Q.24 Which of the following are redox reactions?

- (1) $\frac{1}{2}\text{H}_2 + \frac{1}{2}\text{I}_2 \rightarrow \text{HI}$
(2) $\text{PCl}_5 \rightarrow \text{PCl}_3 + \text{Cl}_2$
(3) $2\text{CuSO}_4 + 4\text{KI} \rightarrow \text{Cu}_2\text{I}_2 + 2\text{K}_2\text{SO}_4 + \text{I}_2$
(4) $\text{CaOCl}_2 \rightarrow \text{Ca}^{+2} + \text{OCl}^- + \text{Cl}^-$

DIRECTIONS (Q.25-Q.27) : Read the passage given below and answer the questions that follows :

Redox reactions play a pivotal role in chemistry and biology. The values of standard redox potential (E°) of two half-cell reactions decide which way the reaction is expected to proceed. A simple example is a Daniel cell in which zinc goes into solution and copper gets deposited. Given below are a set of half-cell reactions (acidic medium) along with their E° (V with respect to normal hydrogen electrode) values. Using this data obtain the correct explanations to questions given.



Q.25 Among the following, identify the correct statement :

- (a) Chloride ion is oxidised by O_2
(b) Fe^{2+} is oxidised by iodine
(c) Iodide ion is oxidised by chlorine
(d) Mn^{2+} is oxidised by chlorine

Q.26 While Fe^{3+} is stable, Mn^{3+} is not stable in acid solution because

- (a) O_2 oxidises Mn^{2+} to Mn^{3+}
(b) O_2 oxidises both Mn^{2+} to Mn^{3+} and Fe^{2+} to Fe^{3+}
(c) Fe^{3+} oxidises H_2O to O_2
(d) Mn^{3+} oxidises H_2O to O_2

Q.27 Sodium fusion extract, obtained from aniline, on treatment with iron (II) sulphate and H_2SO_4 in presence of air gives a prussian blue precipitate. The blue colour is due to the formation of

- (a) $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$
(b) $\text{Fe}_3[\text{Fe}(\text{CN})_6]_2$
(c) $\text{Fe}_4[\text{Fe}(\text{CN})_6]_2$
(d) $\text{Fe}_3[\text{Fe}(\text{CN})_6]_3$

**RESPONSE
GRID**

20. (a) (b) (c) (d)

21. (a) (b) (c) (d)

22. (a) (b) (c) (d)

23. (a) (b) (c) (d)

24. (a) (b) (c) (d)

25. (a) (b) (c) (d)

26. (a) (b) (c) (d)

27. (a) (b) (c) (d)

DIRECTIONS (Q. 28-Q.30) : Each of these questions contains two statements: Statement-1 (Assertion) and Statement-2 (Reason). Each of these questions has four alternative choices, only one of which is the correct answer. You have to select the correct choice.

- (a) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (b) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1..
 (c) Statement -1 is False, Statement-2 is True.
 (d) Statement -1 is True, Statement-2 is False.

Q.28 Statement 1 : Oxidation number of carbon in CH_2O is zero.

Statement 2 : CH_2O formaldehyde, is a covalent compound.

Q.29 Statement 1 : H_2SO_4 cannot act as reducing agent.

Statement 2 : Sulphur cannot increase its oxidation number beyond +6.

Q.30 Statement 1 : HClO_4 is a stronger acid than HClO_3 .

Statement 2 : Oxidation state of Cl in HClO_4 is + VII and in HClO_3 + V.

RESPONSE GRID

28. (a) (b) (c) (d) 29. (a) (b) (c) (d) 30. (a) (b) (c) (d)

DAILY PRACTICE PROBLEM SHEET 15 - CHEMISTRY

Total Questions	30	Total Marks	120
Attempted		Correct	
Incorrect		Net Score	
Cut-off Score	44	Qualifying Score	64
Success Gap = Net Score – Qualifying Score			
Net Score = (Correct × 4) – (Incorrect × 1)			

- (1) (a) When $\text{Cr}_2\text{O}_7^{2-}$ is converted into CrO_4^{2-} the change in oxidation number of Cr is zero
 $\text{Cr}_2\text{O}_7^{2-} \rightarrow \text{CrO}_4^{2-}$
 $+6 \quad +6$

There is no change in oxidation state of Cr, hence it is neither oxidised nor reduced and remains in the same oxidation state.

- (2) (c) Let the oxidation no. of S is 'a'

$$\text{O.N. of CH}_3 = +1$$

$$\text{O.N. of O} = -2$$


$$2(+1) + a + (-2) = 0$$

$$a = 0$$

Hence the oxidation no. of S in dimethyl sulphoxide is zero.

- (3) (b)


- (i) The half reaction for reduction is,




Balancing with respect to oxygen by adding $4\text{H}_2\text{O}$ on R.H.S.,




Balancing with respect to hydrogen by adding 8H^+ on L.H.S.,



Balancing charge by adding electrons,






- (ii) The half-reaction for oxidation is,



Balancing with respect to electrical charge by adding electrons on R.H.S.



Now, to equalise the number of electrons, the reduction half reaction is multiplied by 2 and oxidation half reaction by 5, so on adding, we get




This is the balanced equation.

- (4) (a) Equivalent weight of MnO_4^-

$$= \frac{\text{Molecular weight of MnO}_4^-}{5} = \frac{M}{5}$$

($\because$ Change in oxi. state = 5)

- (5) (d) Equivalent weight of $\text{S}_2\text{O}_3^{2-} = \frac{2M}{2} = M$.



Zn undergoes oxidation and is $\therefore$ acting as reductant.

Change in O.S. of Zn : $\text{Zn} \rightarrow \text{ZnO}$




i.e. O.S. of Zn changes by 2

$$\therefore \text{Eq. wt. of reductant i.e., Zn} = \frac{\text{mol/wt.}}{n}$$

$$= \frac{65.38}{2} = 32.69\text{g}$$

V_2O_5 undergoes reduction and is $\therefore$ acting as oxidant

Change in O.S. of V : $\text{V}_2\text{O}_5 \rightarrow 2\text{V}$



i.e. O.S. of 2V changes by 10

$$\therefore \text{Eq. wt. of oxidant i.e., V}_2\text{O}_5 = \frac{\text{mol/wt.}}{n}$$

$$= \frac{50.94 \times 2 + 80}{10} = 18.2\text{g}$$

- (7) (b) O.N. of N in HNO_2 is +3

Max. O.N. of N is +5




Min. O.N. of N is -3

Thus O.N. of N in HNO_2 can show an increase or decrease as the case may be. That is why HNO_2 acts as oxidant and reductant both.

O.N. of N in HNO_3 is +5,


Hence it can act only as an oxidant.

- (8) (c) $\text{Na}^0 \rightarrow \text{NaOH}^{+1}$



Thus, O.S. of P remains the same in P_2O_5 and $\text{H}_4\text{P}_2\text{O}_7$. In rest of the reactions, there occurs a change of O.S. Hence, all other reactions except (c) are redox reactions.

- (9) (a) In the above reaction $C_2O_4^{2-}$ acts as a reductant because it is oxidised to CO_2 as :



$C_2O_4^{2-}$ reduces MnO_4^- to Mn^{2+} ion in solution.

- (10) (d) $NaN_3 \Rightarrow +1 + 3x = 0$
 $\Rightarrow 3x = -1 \Rightarrow x = -\frac{1}{3}$
 So, oxidation number of nitrogen in NaN_3 is $-\frac{1}{3}$.

- (11) (a) In OF_2 , O.N.(O) + 2 O.N.(F) = 0

$$\Rightarrow x + 2(-1) = 0, \quad x = +2$$

Hence, oxidation number of oxygen in $OF_2 = +2$.

- (12) (c) Let the O.N. of Co be x

O.N. of NH_3 is zero

O.N. of Cl is -1

O.N. of Br is -1


$$\therefore \text{O.N. (Co)} + 6 \text{O.N. (NH}_3\text{)} + 2 \text{O.N. (Cl)} + \text{O.N. (Br)} = 0$$

$$\Rightarrow x + 6(0) - 1 \times 2 - 1 = 0$$


$$\therefore x = +3$$

So, the oxidation number of cobalt in the given complex compound is +3.

- (13) (d) The oxidation number of S are shown below along with the compounds



Hence the order of increasing O.N. of S is



- (14) (c) O.N. of Fe in wustite is $= \frac{200}{93} = 2.15$

It is an intermediate value between Fe(II) & Fe(III)

Let % of Fe(III) be a, then

$$2 \times (100 - a) + 3 \times a = 2.15 \times 100$$

$$a = 15.05$$

$$\therefore \% \text{ of Fe(III)} = 15.05\%$$

- (15) (c) The compound may be written as $NO^+ ClO_4^-$

For ClO_4^- , Let O.N. of Cl = a

$$\therefore \text{in } ClO_4^-, \text{O.N. (Cl)} + 4 \text{O.N. (O)} = -1$$

$$a + 4 \times (-2) = -1$$

$$a = +7$$

Hence, the oxidation no. of Cl in $NOClO_4$ is +7.

- (16) (c) There are two N atoms in NH_4NO_3 , but one N atom has negative oxidation number (attached to H) and the other has positive O.N. (attached to O). Therefore, evaluation should be made separately as –

O.N. of N is NH_4^+

O.N. of N in NO_3^-

$$a + 4 \times (+1) = +1$$


$$a + 3(-2) = -1$$

$$\therefore a = -3$$

$$\therefore a = +5$$

Here the two O.N. are -3 and +5 respectively.

- (17) (c) In $H_2S_2O_8$, two O atoms form peroxide linkage i.e.



$$\Rightarrow 2 \text{ON (H)} + 2 \text{ON (S)} + 6 \text{ON (O}^{2-}\text{)} + 2 \text{ON (O}_2^{2-}\text{)} = 0$$

$$\Rightarrow 2 \times 1 + 2a + 6(-2) + 2(-1) = 0$$

$$\therefore a = +6$$

Thus the O.N. of S in $H_2S_2O_8$ is +6

- (18) (a) In KI_3 , $1 + 3 \times (a) = 0$

$$a = -\frac{1}{3}$$

or KI_3 is $KI + I_2$

$\therefore I$ has two oxidation no. -1 and 0 respectively.

However factually speaking oxidation number of I in KI_3 is an average of two values -1 and 0.

$$\text{Average O.N.} = \frac{-1 + 2 \times 0}{3} = -\frac{1}{3}$$

- (19) (a) O.N. of Fe in : $[Fe(CN)_6]^{3-} \Rightarrow x - 6 = -3 \Rightarrow x = +3$

$$[Fe(CN)_6]^{4-} \Rightarrow x - 6 = -4 \Rightarrow x = +2$$

$$[Fe(SCN)]^{2+} \Rightarrow x - 1 = +2 \Rightarrow x = +3$$

$$[Fe(II_2O)_6]^{3+} \Rightarrow x + 0 = +3 \Rightarrow x = +3$$



Thus, option (a) is correct.

- (20) (d) The balanced redox reaction given above can be written as:



So the value of x = 6 and y = 6

- (21) (b) The reaction is




Amount of electrons involved in the given amount of $MnO_4^- = 5 \times 1.61 \times 10^{-3} \text{ mol}$.

Equating these two we get


$$5 \times 1.61 \times 10^{-3} = (5-n) 2.68 \times 10^{-3}$$

$$\therefore n = 2 \text{ (approx.)}$$


- (22) (a) (i) The half reaction for oxidation is,



Balancing carbon atoms on both sides,




Balancing hydrogen atoms on both sides,



Balancing the charge on both sides,



(ii) The half-reaction for reduction is—




Balancing oxygen atoms on both sides,




Balancing hydrogen atoms,





Balancing the charge,






Now, adding both equations,




This is balanced equation.



Hence, $\text{S}_2\text{O}_3^{2-}$ is getting oxidised to $\text{S}_4\text{O}_6^{2-}$, while I_2 is getting reduced to 2I^- . So, (d) is the correct answer.




Except (4), there occurs a change in O.S. of the reactants and products. Hence, except (4), all other



$$E^\circ = E^\circ_{\text{I}^-/\text{I}_2} + E^\circ_{\text{Cl}_2/\text{Cl}^-} = -0.54 + 1.36$$


$$E^\circ = 0.82\text{V}$$

E° is positive hence, iodide ion is oxidized by chlorine.




$$E^\circ_{\text{Mn}^{3+}/\text{Mn}^{2+}} + E^\circ_{\text{H}_2\text{O}/\text{O}_2} = 1.50 + (-1.23) = 0.27\text{V}$$


Reaction is feasible. [$\because E^\circ$ is positive]



In presence of air, Fe^{2+} gets oxidised to Fe^{3+} , i.e.,



Fe^{3+} then combines with $[\text{Fe}(\text{CN})_6]^{4-}$ to form ferric ferrocyanide which is Prussian blue in colour, i.e.



(28) (b) Oxidation number can be calculated using some rules. H is assigned +1 oxidation state and O has oxidation number -2

$\therefore$ O. No. of C in CH_2O :

$$\text{O. no. of C} + 2(+1) + (-2) = 0$$

$\therefore$ O. No. of C = 0

(29) (a) In H_2SO_4 , the O.N. of S is +6, which is maximum. Therefore, H_2SO_4 can only decrease its O.N. and can act only as an oxidising agent.

(30) (a) Acid strength of oxoacids of the same halogen increases with increase in O.N. of the halogen, i.e.

