


Chapter - 9

Force And Laws Of Motion

CONCEPT MAPPING



Forces and Laws of Motion :

Force : It is the force that enables us to do any work. To do anything, either we pull or push the object. Therefore, pull or push is called force.

Example, to open a door, either we push or pull it. A drawer is pulled to open and pushed to close.

Effect of Force

- (i) Force can move a stationary body or object. For example, a football can be set to move by kicking it, *i.e.*, by applying a force.
- (ii) Force can stop a moving body. For example, by applying brakes, a running cycle or a running vehicle can be stopped.
- (iii) Force can change the direction of a moving object. For example, by applying force, *i.e.*, by moving handle, the direction of a running bicycle can be changed. Similarly by moving steering, the direction of a running vehicle is changed.
- (iv) Force can change the speed of a moving body. By accelerating, the speed of a running vehicle can be increased or by applying brakes the speed of a running vehicle can be decreased.
- (v) Force can change the shape and size of an object. For example, by hammering, a block of metal can be turned into a thin sheet. By hammering, a stone can be broken into pieces.

Forces are mainly of two types :

- (A) Balanced forces
- (B) Unbalanced forces

(A) Balanced Forces

- If the resultant of applied forces is equal to zero, it is called balanced forces. an

Example, In the tug of war if both the team apply similar magnitude of forces in opposite directions, rope does not move in either side. This happens because of balanced forces in which resultant of applied forces become zero.

- Balanced forces do not cause any change of state of an object. Balanced forces are equal in magnitude and opposite in direction.
- Balanced forces can change the shape and size of an object. For example, when forces are applied from both sides over a balloon, the size and shape of balloon is changed.

(B) Unbalanced Forces

- If the resultant of applied forces are greater than zero, the forces are called unbalanced forces. An object in rest can be moved because of applying Unbalanced forces.
- Unbalanced forces can do the following :
 - * Move a stationary object
 - * Increase the speed of a moving object
 - * Decrease the speed of a moving object
 - * Stop a moving object
 - * Change the shape and size of an object

Laws of Motion :

Galileo Galilei : Galileo first of all said that object move with a constant speed when no forces act on them. This means if an object is moving on a frictionless path and no other force is acting upon it, the object would be moving forever. That is, there is no unbalanced force working on the object.

- But practically it is not possible for any object. Because to attain the condition of zero, unbalanced force is impossible. Force of friction, force of air and many other forces are always acting upon an object.

Newton's Laws of Motion :

Newton studied the ideas of Galileo and gave the three laws of motion. These laws are known as Newton's laws of motion.

Newton's First Law of Motion (Law of Inertia) :

Any object remains in the state of rest or in uniform motion along a straight line, until it is compelled to change the state by applying external force.

Explanation : If any object is in the state of rest, then it will remain in rest until an external force is applied to change its state. Similarly, an object will remain in motion until any external force is applied over it to change its state. This means all objects resist to in changing their state. The state of any object can be changed by applying external forces only.

Newton's First Law of Motion in Everyday Life :

- (a) A person standing in a bus falls backward when bus starts moving suddenly. This happens because the person and bus both are in rest while bus is not moving, but as the bus starts moving, the legs of the person start moving along with bus but rest portion of his body has the tendency to remain in rest. Because of this, the person falls backward; if he is not alert.
- (b) A person standing in a moving bus falls forward if driver applies brakes suddenly. This happens because when bus is moving, the person standing in it is also in motion along with bus. But when driver applies brakes the speed of bus decreases suddenly or bus comes in the state of rest suddenly in this condition the legs of the person which are in contact with the bus come in rest while the rest part of his/her body have the tendency to remain in motion. Because of this person falls forward if he/ She is not alert.
- (c) Before hanging the wet clothes over laundry line, usually many jerks are given to the clothes to get them dried quickly. Because of jerks, droplets of water from the pores of the cloth falls on the ground and reduced amount of water in clothes dries them quickly. This happens because when suddenly clothes are made in motion by giving jerks, the water droplets in it have the tendency to remain in rest and they are separated from clothes and fall on the ground.
- (d) When the pile of coin on the carom-board is hit by a striker, only the coin at the bottom moves away leaving rest of the pile of coin at same place. This happens because when the pile is struck with a striker, the coin at the bottom comes in motion while rest of the coin in the pile has the tendency to remain in the rest and they vertically falls on the carom-board and remain at same place.

Mass and Inertia

- The property of an object because of which it resists to get disturb ^{the} its state is called inertia. Inertia of an object is measured by its mass. Inertia is directly proportional to the mass. This means inertia increases with increase in mass and decreases with decrease in mass. A heavy object will have more inertia than the lighter one.
- In other words, the natural tendency of an object that resists the change in state of motion or rest of the object is called inertia.

- Since a heavy object has more inertia, thus it is difficult to push or pull a heavy box over the ground than the lighter one.

Momentum

- Momentum is the power of motion of an object.
- The product of velocity and mass is called the momentum. Momentum is denoted by 'p'.

Therefore, Momentum of the object = Mass \times Velocity

Or,
$$p = m \times v$$

Where, p = momentum, m = mass of the object and v = velocity of the object.

Consider the following explanations to understand the momentum :

- A person get injured in the case of hitting by a moving object, such as stone, pebbles or anything because of momentum of the object.
- Even a small bullet is able to kill a person when it is fired from a gun because of its momentum due to great velocity.
- A person get injured severely when hit by a moving vehicle because of momentum of vehicle due to mass and velocity.

Momentum and Mass and Velocity

- Since momentum is the product of mass and velocity ($p = m \times v$) of an object. This means momentum is directly proportional to mass and velocity. Momentum increases with increase of either mass or velocity of an object.
- This means if a lighter and a heavier object is moving with same velocity, then heavier object will have more momentum than the lighter one.
- If a small object is moving with great velocity, it has tremendous momentum. And because of momentum, it can harm an object more severely. For example, a small bullet having a little mass even kills a person when it is fired from a gun.
- Usually, road accidents prove more fatal because of high speed than in slower speed. This happens because vehicles running with high speed have greater momentum compared to a vehicle running with slower speed.

Momentum of an object which is in the state of rest :

Let an object with mass ' m ' is in the rest.

Since, object is in rest, therefore, its velocity, $v = 0$

Now, we know that

$$\text{Momentum} = \text{mass} \times \text{velocity}$$

Or
$$p = m \times 0 = 0$$

Thus, the momentum of an object in the rest *i.e.*, non-moving, is equal to zero.

Unit of momentum :

$$\text{SI unit of mass} = \text{kg}$$

$$\text{SI unit of velocity} = \text{meter per second } i.e., \text{ m/s}$$

We know that
$$\text{Momentum } (p) = m \times v$$

Therefore,
$$p = \text{kg} \times \text{m/s}$$

Or
$$p = \text{kg m/s}$$

Therefore, SI unit of momentum
$$= \text{kg m/s}$$

Numerical Problems Based on Momentum

Type I. Calculation of Momentum

Example 1. *What will be the momentum of a stone having mass of 10 kg when it is thrown with a velocity of 2 m/s ?*

Solution :
$$\text{Mass } (m) = 10 \text{ kg}$$

$$\text{Velocity } (v) = 2 \text{ m/s}$$

$$\text{Momentum } (p) = ?$$

We know that,
$$\text{Momentum } (p) = \text{Mass } (m) \times \text{Velocity } (v)$$

Therefore,
$$p = 10 \text{ kg} \times 2 \text{ m/s} = 20 \text{ kg m/s}$$

Thus, the momentum of the stone
$$= 20 \text{ kg m/s.} \quad \text{Ans.}$$

Example 2. *The mass of a goods lorry is 4000 kg and the mass of goods loaded on it is 20000 kg. If the lorry is moving with a velocity of 2 m/s, what will be its momentum ?*

Solution : Given, Velocity (v) = 2 m/s

Mass of lorry = 4000 kg, Mass of goods on the lorry = 20000 kg

Therefore, Total mass (m) on the lorry = 4000 kg + 20000 kg = 24000 kg

Momentum (p) = ?

We know that, Momentum (p) = Mass (m) \times Velocity (v)

Therefore, $p = 24000 \text{ kg} \times 2 \text{ m/s}$

Or $p = 48000 \text{ kg m/s}$

Thus, the momentum of the lorry = 48000 kg m/s. **Ans.**

Example 3. *A car having mass of 1000 kg is moving with a velocity of 0.5 m/s. What will be its momentum ?*

Solution : Given, Velocity of the car (v) = 0.5 m/s

Mass of the car (m) = 1000 kg

Momentum (p) = ?

We know that, Momentum (p) = Mass (m) \times Velocity (v)

Therefore, $p = 1000 \text{ kg} \times 0.5 \text{ m/s} = 500 \text{ kg m/s}$

Thus, momentum of the car = 500 kg m/s. **Ans.**

Statement of Second Law

Rate of change of momentum of an object is proportional to applied unbalanced force in the direction of force.

Mathematical expression

Suppose, Mass of an object = m kg

Initial velocity of an object = u m/s

Final velocity of an object = v m/s

So, Initial momentum, $p_1 = mu$, Final momentum, $p_2 = mv$

\therefore Change in momentum = Final momentum – Initial momentum

$$= mv - mu$$

$$= m(v - u)$$

\therefore Rate of change of momentum = $\frac{\text{Change in momentum}}{\text{Time taken}}$

$$= \frac{m(v - u)}{t}$$

- According to IInd law, this rate of change of momentum is directly proportional to force.

$$\therefore F \propto \frac{m(v - u)}{t}$$

We know that, $\frac{v - u}{t} = a$ (From 1st equation of motion)

So, $F = kma$

Where k is a constant. Its value = 1.

$$\therefore F = 1 \times m \times a = ma$$

SI unit = kg m/s² or Newton

Q. Define 1 Newton.

Ans. When an acceleration of 1 m/s² is seen in a body of mass 1 kg, then the force applied on the body is said to be 1 Newton.

Proof of Newton's First Law of Motion from Second Law

First law states that if external force $F = 0$, then a moving body keeps moving with the same velocity, or a body at rest continues to be at rest.

So, $F = 0$

We know $F = \frac{m(v-u)}{t}$

(a) A body is moving with initial velocity u , then

$$0 = \frac{m(v-u)}{t} \Rightarrow v - u = 0$$

So, $v = u$

Thus, final velocity is also same.

(b) A body is at rest *i.e.*, $u = 0$.

Therefore, from above $u = v = 0$

So, the body will continue to be at rest.

Third Law of Motion

To every action there is an equal and opposite reaction.

Applications :


- (i) Walking is enabled by IIIrd law.
- (ii) A boat moves back when we deboard it.
- (iii) A gun recoils.
- (iv) Rowing of a boat.

Law of Conservation of Momentum

When two (or more) bodies act upon one another, their total momentum remains constant (or conserved) provided no external forces are acting.

$$\text{Initial momentum} = \text{Final momentum}$$

Suppose, two objects A and B each of mass m_1 and mass m_2 are moving initially with velocities u_1 and u_2 , strike each other after time t and start moving with velocities v_1 and v_2 respectively.



Now,

Initial momentum of object A = $m_1 u_1$

Initial momentum of object B = $m_2 u_2$

Final momentum of object A = $m_1 v_1$

Final momentum of object B = $m_2 v_2$

So,

$$\begin{aligned} \text{Rate of change of momentum in A, } F_1 &= \frac{m_1 v_1 - m_1 u_1}{t} \\ &= \frac{m_1 (v_1 - u_1)}{t} \quad \dots(i) \end{aligned}$$

And

$$\begin{aligned} \text{Rate of change of momentum in B, } F_2 &= \frac{m_2 v_2 - m_2 u_2}{t} \\ &= \frac{m_2 (v_2 - u_2)}{t} \quad \dots(ii) \end{aligned}$$

We know from IIIrd law of motion,

$$F_1 = -F_2$$

$$\text{So, } \frac{m_1 (v_1 - u_1)}{t} = -\frac{m_2 (v_2 - u_2)}{t} \quad [\text{From equations (i) \& (ii)}]$$

Or

$$m_1 v_1 - m_1 u_1 = -m_2 v_2 + m_2 u_2$$

So

$$m_1 u_1 + m_2 u_2 = m_1 v_1 + m_2 v_2$$

Thus,

Initial momentum = Final momentum

Example 1. A bullet of mass 20 g is fired horizontally with a velocity of 150 m/s from a pistol of mass 2 kg. Find the recoil velocity of the pistol.

Solution : Given, Mass (m_1) of bullet = 20 g = 0.02 kg

Mass (m_2) of pistol = 2 kg

Initially bullet is inside the gun and it is not moving.

So, Mass = $m_1 + m_2 = (0.02 + 2) \text{ kg} = 2.02 \text{ kg}$

And $u_1 = 0$

So, Initial momentum = $2.02 \times 0 = 0$...**(i)**

Finally let the velocity of pistol be v_2 and v_1 for bullet = 150

So, Final momentum = $m_1v_1 + m_2v_2$
 $= 0.02 \times 150 + 2v_2$...**(ii)**

We know that Initial momentum = Final momentum

So, $0 = \frac{0.02 \times 150}{100} + 2v_2$ [From equations (i) and (ii)]

$\Rightarrow 3 + 2v_2 = 0$

Or $2v_2 = -3$

Or $v_2 = -1.5 \text{ m/s}$ **Ans.**

(-)**ve** sign indicates that gun recoils in direction opposite to that of the bullet.

Example 2. Two hockey players viz A of mass 50 kg is moving with a velocity of 4 m/s and another one B belonging to opposite team with mass 60 kg is moving with 3 m/s, get entangled while chasing and fall down. Find the velocity with which they fall down and in which direction ?

Solution : Given, $m_A = 50 \text{ kg}$, $u_A = 4 \text{ m/s}$

$m_B = 60 \text{ kg}$, $u_B = 3 \text{ m/s}$

$$\begin{aligned}\text{Initial momentum}_A &= m_A u_A \\ &= 50 \times 4 = 200 \text{ kg m/s}\end{aligned}$$

$$\begin{aligned}\text{Initial momentum}_B &= m_B u_B \\ &= 60 \times 3 = 180 \text{ kg m/s}\end{aligned}$$

So, Total initial momentum = $200 + 180 = 380 \text{ kg m/s}$...**(i)**

$$\begin{aligned}\text{Final momentum} &= (m_A + m_B)v = (50 + 60)v \\ &= 110v \quad \dots\text{(ii)}\end{aligned}$$

According to the law of conservation of momentum,

$$380 = 110v$$

Or $v = \frac{380}{110} = 3.454 \text{ m/s}$ **Ans.**

QUESTIONS

VERY SHORT ANSWER TYPE QUESTIONS

1. Can force be (-)ve ? When ?
2. What is the tendency of a body to resist its change of state called ?
3. Inertia is also measured by.....of an object.
4. Higher the mass of an object, higher is its..... .
5. $\frac{\text{Force}}{\text{Acceleration}}$ is determined by.....which is also mass of the object.
6. Why does the load from the cage above the seats in a bus falls down when suddenly brakes are applied ?
7. When a tree is shaken, its fruits and leaves fall down. Why?
8. Define Momentum of a body.
9. On what factors does the momentum of a body depend?
10. Why it is difficult to walk on a slippery road?

SHORT ANSWER TYPE QUESTIONS

1. Quantity of motion contained in a body is..... .
2. Unit of momentum is..... .
3. Define 1 Newton.
4. Although we know that a moving body keeps moving indefinitely until an external force is applied on it, then why does a ball stops when we slide it on ground (without stopping it) ?
5. Why is it difficult to stop a truck suddenly than a motorbike ?
6. When a metro suddenly stops all the passengers fell forward on its floor. Why do this happen ?

7. We have a huge atmosphere above us that exerts a huge pressure on our shoulders, head and whole body. Why don't we get crushed under it ?
8. A coin of mass 1 kg and a stone of mass 5 kg are thrown down the Eiffel Tower with an acceleration of 10 m/s^2 . Which one would reach the ground early and why ?
9. Give applications of 1st law of motion i.e., inertia.
10. (a) Friction is measured in..... .
(b) Distinguish between balanced and unbalanced forces.

LONG ANSWER TYPE QUESTIONS

1. (a) Derive first law of Newton from second law.
(b) Find the force required to stop a car of mass 100 kg with two passengers each of 50 kg sitting inside, if it is moving at 60 km/hr speed and takes 5 s to stop.
2. Two balls A and B of masses 40 g and 50 g are moving at speeds of 40 m/s and 30 m/s respectively. If after colliding, B starts moving with a velocity of 25 m/s, what is the velocity of A ?
3. A girl of mass 30 kg jumps on a cart of mass 5 kg with a velocity of 10 m/s. Find the velocity with which she and cart start moving after she jumps on it.
4. (a) Why does a gunman get a jerk on firing a bullet?
(b) Calculate the momentum of a toy car of mass 200 gm moving with a speed of 5 m/s. [Hint - convert mass into kg].
(c) State the law of conservation of momentum.
5. For how long should a force of 100 N acts on a body of 20 kg so that it acquires a velocity of 100 ms? [Hint - using formula $f = ma$. $V = u + at$]
6. (a) Find the acceleration produced by a force of 5 N acting on a mass of 10 kg.

- (b) Which would require a greater force : (a) accelerating a 10 gm mass of 5 m/s^2 or (b) a 20 gm mass at 2 m/s^2 ?
[convert mass into kg].
7. The velocity of a body of mass 10 kg increases from 4 m/s to 8 m/s when a force acts on it for 2s.
- (a) What is the momentum before the force acts?
 (b) What is the momentum after the force acts?
 (c) What is the gain in momentum per second?
 (d) What is the value of force?

$$\left[\text{Hint} - a = \frac{v-u}{t} \text{ and } f = ma \right]$$

Answers to Long Answer Type Questions

1. (b) - 2000/3 N
2. 46.25 m/s
3. 8.57 m/s
4. (b) 1 kg/ms
5. 20 sec.
6. (a) 0.5 m/s^2
 (b) A greater force of 0.05 N is required for accelerating a 10 gm mass.
7. (a) 40 kg.m/s
 (b) 80 kg.m/s
 (c) 20 kg.m/s^2
 (d) 20 N.

OBJECTIVE TYPE QUESTIONS :

MCQ.

- 1. A truck and a car are moving with equal velocity, on applying brakes, both will stop after certain distance and then :**
 - (a) Truck will cover less distance before stopping.
 - (b) Car will cover less distance before stopping.
 - (c) Both will cover equal distance.
 - (d) None of the above.
- 2. In which of the following cases, the net force is not zero ?**
 - (a) An object floating in air
 - (b) A ball freely falling from a certain height.
 - (c) A cork floating on the surface of water
 - (d) All the cases.
- 3. A force acts on a body of mass 3kg such that its velocity changes from 4ms^{-1} to 10ms^{-1} . The change in momentum of the body is :**
 - (a) 42Kgms^{-1}
 - (b) 2Kgms^{-1}
 - (c) 18Kgms^{-1}
 - (d) 14Kgms^{-1}
- 4. While opening a top with two fingers, the force applied are:**
 - (a) equal in magnitude
 - (b) Parallel to each other
 - (c) opposite in direction
 - (d) All of the above

5. The engine of a car produces an acceleration of 4ms^{-2} in a car, if this Car pulls another car of same mass, what is the acceleration produces ?

- (a) 8ms^{-2} (b) 2ms^{-2} (c) 4ms^{-2} (d) 0.5ms^{-2}

6. A force 100N acts in a body mass 2kg for 10 sec. The change in the velocity of the body is.

- (a) 100ms^{-1} (b) 250ms^{-1} (c) 500ms^{-1} (d) 1000ms^{-1}

Assertion and Reason type questions :

Choose the appropriate answer from the following choices :

• **Assertion Reasoning Questions :-**

Read the Assertion (A) and Reason (R) statements carefully and mark the correct option out of the following :

- a) Both Assertion (A) and Reason (R) are true, and (R) is correct explanation of the assertion (A)
- b) Both (A) and (R) are true, but (R) is not current explanation of the (A).
- c) (A) is True but (R) is false.
- d) (A) is False but (R) is true.

1. **Assestion (A) :** If the external force on the body is Zero, then its accelerations is Zero.

Reason (B) : Acceleration does not depend on force.

2. **Assertion (A) :** If two objects of different masses have same momentum, the lighter

body possess greater velocity.

Reason (B) : For all bodies momentum always remains same.

3. **Assertion (A) :** Newton's third law of motion is applicable only when bodies are in motion.

Reason (B) : Newton's third law applies to all types of forces eg., gravitaional, electric or magnetic force etc.