

S.No.	CONTENTS	Page
1.	Introduction	115
2.	Conductors and Conductance terms	115
3.	Factors affecting electrolytic conductance	118
4.	Kohlrausch's law and its applications	119
5.	Electrode potential	122
6.	Reference electrode	122
	(a) Primary reference electrode	124
	(b) Secondary reference electrode	124
7.	Electrochemical series and its applications	125
8.	Galvanic cell	125
	(a) Working	125
	(b) Construction of cell	125
	(c) Salt bridge and its functions	125
	(d) Cell representation	126
9.	Nernst equation and its applications	126
10.	Some special cells	130
	(a) Concentration cell	130
	(b) Reversible cell	131
11.	Electrolysis	132
	(a) Qualitative aspect	132
	(b) Quantitative aspect (Faraday's law)	134
12.	Some commercial voltaic cells	137
13.	Corrosion	141
14.	Exercise-I (Conceptual Questions)	143
15.	Exercise-II (Previous Years Questions)	150
16.	Exercise-III (Analytical Questions)	154
17.	Exercise-IV (Assertion & Reason)	157

NEET SYLLABUS

Electrochemistry : Conductance in electrolytic solutions, specific and molar conductivity variation of conductivity with concentration, Kohlrausch's Law, Galvanic cells, Redox reactions, EMF of a cell, standard electrode potential, Relation between Gibbs energy change and EMF of a cell, electrolysis and electrolytic cells, Laws of electrolysis (elementary idea), dry cell, lead accumulator, fuel cells, corrosion.

OBJECTIVES

After studying this unit, we will be able to :

- describe an electrochemical cell and differentiate between galvanic and electrolytic cells;
- apply Nernst equation for calculating the emf of galvanic cell and define standard potential of the cell;
- derive relation between standard potential of the cell, Gibbs energy of cell reaction and its equilibrium constant;
- define resistivity (ρ), conductivity (κ) and molar conductivity (Λ_m) of ionic solutions;
- differentiate between ionic (electrolytic) and electronic conductivity;
- describe the methods for measurement of conductivity of electrolytic solutions and calculation of their molar conductivity;
- justify the variation of conductivity and molar conductivity of solutions with change in their concentration and define Λ_m^∞ (molar conductivity at zero concentration or infinite dilution);
- enunciate Kohlrausch law and learn its applications;
- understand quantitative aspects of electrolysis;
- describe the construction of some primary and secondary batteries and fuel cells;
- explain corrosion as an electrochemical process.

In arranging the bodies in order of their electrical nature, there is formed an electro-chemical system which, in my opinion is more fit than any other to give an idea of Chemistry.


Jons Jacob Berzelius

ELECTROCHEMISTRY

4.0 INTRODUCTION

Electrochemistry is the branch of physical chemistry which deals with the study of interconversion of chemical energy and electrical energy

ELECTROCHEMICAL CELL



4.1 CONDUCTORS

Substances which allow electric current to flow through them are called conductors.

Examples - Metals, Aqueous solution of acids, bases and salts, fused salts and impure water etc.

Conductors are of two types - (i) Metallic conductors or electronic conductors
(ii) Electrolytic conductors or ionic conductors

(i) Metallic conductors : The conductors which conduct electric current by the movement of electrons without undergoing any chemical change are known as metallic conductors.

Metals - Cu, Ag, Fe, Al etc, non metals - graphite

(ii) Electrolytic conductors : Those substances which conduct the electric current through ions in aqueous solution or in molten state are called electrolytic conductors.

- **Strong electrolyte :** Electrolytes which are completely ionized in aqueous solution are called strong electrolytes.

Ex. : Salts, strong acids and strong bases.

- **Weak electrolyte :-** Electrolytes which are not completely ionized in aqueous solution are called weak electrolytes.

Ex. : CH_3COOH , HCN , NH_3 , amine, etc.

Difference between metallic and electrolytic conductor

Metallic conductor	Electrolytic conductor
(i) Charge carriers are free electrons.	Charges carriers are free ions.
(ii) Flow of electricity takes place without any chemical change.	Flow of electricity takes place by chemical changes at electrodes.
(iii) No transfer of matter takes place.	Transfer of matter takes place in the form of ions.
(iv) Resistance is due to vibration of Kernels.	Resistance is due to inter ionic attraction and viscosity of medium.
(v) The resistance increases with the increase in temperature.	The resistance decreases with the increase in temperature.
(vi) Faraday's laws of electrolysis are not followed.	Faraday's laws of electrolysis are followed.

4.2 ELECTROLYTIC CONDUCTANCE

- (a) **Resistance (R)** :- Metallic and electrolytic conductors obey ohm's law according to which the resistance of a conductor is the ratio of the applied potential difference (V) to the current (I) flowing .

$$R = \frac{V}{I}$$


R is expressed in ohms.

- (b) **Conductance (G)** :- It is the property by virtue of which it favours flow of current. The conductance of a conductor is equal to reciprocal of resistance $G = \frac{1}{R}$

unit of G is **mho** or **ohm⁻¹** or **Siemens (S)**.

- (c) **Specific resistance/Resistivity (ρ)** :- The resistance (R) of a conductor is directly proportional to its length (ℓ) and inversely proportional to its area of cross section (A).

$$R \propto \frac{\ell}{A} \quad ; \quad R = \rho \frac{\ell}{A}$$



$$R = \rho \cdot \frac{\ell}{A} \quad ; \quad \rho = R \cdot \frac{A}{\ell}$$

If $\ell = 1 \text{ cm}$, $A = 1 \text{ cm}^2$, therefore $\rho = R$

$$\therefore V = A \times \ell = 1 \text{ cm}^3$$

Therefore resistance offered by 1 cm^3 electrolytic solution is known as resistivity.

Unit of $\rho \rightarrow \text{ohm cm}$

- (d) **Specific conductance/ Conductivity (κ)** :-

It is defined as the reciprocal of specific resistance, $\kappa = \frac{1}{\rho}$

$$R = \rho \frac{\ell}{A}$$

$$\frac{1}{\rho} = \frac{1}{R} \cdot \frac{\ell}{A}$$

$$\kappa = G \times G^*$$

Specific conductance = Conductance $\times$ Cell constant

Hence specific conductivity of a solution is defined as the conductance offered by 1 cm^3 of electrolytic solution. Unit of κ is $\text{ohm}^{-1} \text{ cm}^{-1}$

Cell constant :

$$G^* = \frac{\ell}{A} \quad ; \text{ Its unit is } \text{cm}^{-1}$$

(e) **Molar conductivity or Molar conductance** :- (Λ_m , λ_m or μ) : It is defined as the conductance of all the ions produced by one mole of electrolyte present in the given volume of solution.

$$\Lambda_m = \kappa \times V$$

V = Volume of solution containing 1 mol of electrolyte.

If concentration of solution is M - mol per litre then

$$\Lambda_m = \frac{\kappa \times 1000}{M} \quad \text{Unit} \rightarrow \text{ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

(f) **Equivalent conductivity or Equivalent conductance** (Λ_{eq} or λ_{eq}) :- It is defined as the conductance of all the ions produced by one gram equivalent of the electrolyte in the given volume of solution.

$$\therefore \Lambda_{eq} = \kappa \times V$$

V = Volume of solution containing 1 g-eq of electrolyte.

If concentration of solution is N - gram equivalent per litre then

$$\Lambda_{eq} = \frac{\kappa \times 1000}{N} \quad \text{Unit} \rightarrow \text{ohm}^{-1} \text{ cm}^2 \text{ eq}^{-1}$$

● **Relation between Λ_{eq} and Λ_m**

$$\Lambda_m = \frac{\kappa \times 1000}{M} \quad \text{and} \quad \Lambda_{eq} = \frac{\kappa \times 1000}{N}$$

We know that

Normality = Valency Factor $\times$ Molarity . So

$$\Lambda_{eq} = \frac{\Lambda_m}{V.F.}$$

Illustrations

Illustration 1. The resistance of a 1N solution of salt is 50 Ω . Calculate the equivalent conductance of the solution, if the two platinum electrodes in solution are 2.1 cm apart and each having an area of 4.2 cm².

Solution.

$$\kappa = \frac{1}{\rho} = \frac{1}{R} \left(\frac{\ell}{A} \right) = \frac{1}{50} \times \frac{2.1}{4.2} = \frac{1}{100} \quad \text{and} \quad \Lambda_{eq} = \frac{\kappa \times 1000}{N} = \frac{1}{100} \times \frac{1000}{1} = 10 \text{ ohm}^{-1} \text{ cm}^2 \text{ eq}^{-1}$$

Illustration 2. Which of the following have maximum molar conductivity.

- (i) 0.08 M solution and its specific conductivity is $2 \times 10^{-2} \Omega^{-1} \text{ cm}^{-1}$.
(ii) 0.1 M solution and its resistivity is 50 $\Omega \text{ cm}$.

Solution.


$$(i) \quad \Lambda_m = \frac{\kappa \times 1000}{M} = 2 \times 10^{-2} \times \frac{1000}{0.08} = 250 \Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

$$(ii) \quad \Lambda_m = \frac{\kappa \times 1000}{M}, \quad \therefore \kappa = \frac{1}{\rho}, \quad \therefore \Lambda_m = \frac{1}{50} \times \frac{1000}{0.1} = 200 \Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

So, the molar conductivity of 0.08 M solution will be greater than 0.1 M solution

Illustration 3. The conductance of a salt solution (AB) measured by two parallel electrodes of area 100 cm² separated by 10 cm was found to be 0.0001 Ω^{-1} . If volume enclosed between two electrodes contain 0.5 mol of salt. What is the molar conductivity ($\text{Scm}^2 \text{ mol}^{-1}$) of salt at same concentration.

- (A) 0.01 (B) 0.02 (C) 2×10^{-5} (D) none of these



Solution

$$G^* = \frac{\ell}{a} = \frac{10}{100} = 0.1; \quad G = 0.0001 \text{ S}; \quad V = 100 \times 10 = 1000 \text{ cm}^3 = 1 \text{ litre}$$

$$\kappa = G G^* = 0.1 \times 0.0001 = 10^{-5}$$

$$\Lambda_m = \frac{\kappa \times 1000}{M} = \frac{(0.1 \times 0.0001) \times 1000}{0.5} = 0.02 \text{ S cm}^2 \text{ mol}^{-1}$$

4.3 FACTORS AFFECTING ELECTROLYTIC CONDUCTANCE

- (a) **Inter ionic attraction :-** If inter ionic attraction between ions of solute is more, then the conductance will be less.
- (b) **Polarity of solvent :-** If solvent has high-dielectric constant then the ionization and conductance will be higher.
- (c) **Viscosity of medium :-** On increasing the viscosity of medium, the conductance decreases.
- (d) **Temperature :-** As the temperature of electrolytic solution is increased, the conductance increases because K.E. of the ions increases and all types of attraction forces decrease and the viscosity of medium decreases.
- (e) **Hydrated size :** Due to hydration of ions conductance decreases.
- (f) **Dilution :-**
- On increasing the dilution conductance (G) increases.
For strong electrolyte on dilution interionic force of attraction decreases therefore conductance increases.
For weak electrolyte with dilution degree of dissociation (α) increases therefore conductance increases.
 - On dilution specific conductance decreases because on dilution number of ions in 1 ml solution decreases.
 - On dilution equivalent and molar conductance increases because with dilution normality or molarity decreases

DETERMINATION OF MOLAR CONDUCTANCE OF ELECTROLYTES AT INFINITE DILUTION


Debye Huckel Onsager equation –

$$\Lambda_m = \Lambda_m^\infty - b\sqrt{C} \quad [\text{only for strong electrolyte}]$$

where Λ_m = molar conductance at concentration C ,

Λ_m^∞ = molar conductance at ∞ dilution,

b = constant and its value is same for a particular type of electrolyte.



If a graph is plotted between Λ_m vs $\sqrt{C}$ a straight line is obtained for strong electrolyte. On extrapolation this line touches Λ_m axis. Therefore Λ_m^∞ for strong electrolyte can be found out from intercept. For weak electrolyte the slope of curve is not constant so it can not be extrapolated to obtain Λ_m^∞ .

4.4 KOHLRAUSCH'S LAW OF INDEPENDENT MIGRATION OF IONS

At infinite dilution when dissociation of electrolyte is complete; each ion makes a definite contribution towards equivalent conductivity of any electrolyte irrespective of the nature of other ion associated with it.

This law states that the equivalent conductivity of any electrolyte at infinite dilution (Λ_{eq}^{∞}) is the sum of equivalent ionic conductivities of the cation and anion given by the electrolytes at infinite dilution.

$$\Lambda_{eq}^{\infty} = \lambda_c^{\infty} + \lambda_a^{\infty}$$

Where λ_c^{∞} = equivalent conductance of cation at infinite dilution.

λ_a^{∞} = equivalent conductance of anion at infinite dilution.

For $A_x B_y \rightarrow xA^{y+} + yB^{x-}$

In terms of molar conductances of cation and anion

$$\Lambda_{eq}^{\infty} = \frac{1}{y} \lambda_{m(c)}^{\infty} + \frac{1}{x} \lambda_{m(a)}^{\infty}$$

Where $\lambda_{m(c)}^{\infty}$ = Limiting molar conductance of cation .

$\lambda_{m(a)}^{\infty}$ = Limiting molar conductance of anion .

y = charge on cation

x = charge on anion

$$\Lambda_m^{\infty} = x\lambda_{m(c)}^{\infty} + y\lambda_{m(a)}^{\infty}$$

Where x = Stoichiometric coefficient of cation

y = Stoichiometric coefficient of anion

Illustrations

Illustration 4. Calculate Λ_m^{∞} of oxalic acid, given that

$$\Lambda_{eq}^{\infty} \text{Na}_2\text{C}_2\text{O}_4 = 400 \Omega^{-1} \text{cm}^2 \text{eq}^{-1}$$

$$\Lambda_m^{\infty} \text{H}_2\text{SO}_4 = 700 \Omega^{-1} \text{cm}^2 \text{mol}^{-1}$$

$$\Lambda_{eq}^{\infty} \text{Na}_2\text{SO}_4 = 450 \Omega^{-1} \text{cm}^2 \text{eq}^{-1}$$

Solution

$$\Lambda_m^{\infty} \text{H}_2\text{C}_2\text{O}_4 = 700 + 800 - 900 = 600 \Omega^{-1} \text{cm}^2 \text{mol}^{-1}$$

4.5 APPLICATIONS OF KOHLRAUSCH'S LAW :

(a) To calculate conductance of weak electrolyte.

(b) To calculate degree of dissociation :

$$\text{Degree of dissociation } \alpha = \frac{\Lambda_{eq}}{\Lambda_{eq}^{\infty}} = \frac{\text{equivalent conductance at a given concentration}}{\text{equivalent conductance at infinite dilution}}$$

$$\text{or } \alpha = \frac{\Lambda_m}{\Lambda_m^{\infty}} = \frac{\text{molar conductance at a given concentration}}{\text{molar conductance at infinite dilution}}$$

(c) To calculate dissociation constant of weak electrolyte

$$K_a = \frac{C\alpha^2}{(1-\alpha)}$$


If $\alpha \leq 5\%$ then $K_a = C\alpha^2$

Illustration 7. The conductivity of a saturated solution of Ag_3PO_4 is $9 \times 10^{-6} \text{ S m}^{-1}$ and its equivalent conductivity is $1.50 \times 10^{-4} \text{ S m}^2 \text{ eq}^{-1}$. The K_{sp} of Ag_3PO_4 is

- (1) 4.32×10^{-18} (2) 1.8×10^{-9} (3) 8.64×10^{-13} (4) None of these

Solution $\Lambda_{\text{eq}} = \kappa \times \frac{1000}{N}$; $1.50 \times 10^{-4} \times 10^4 = 9 \times 10^{-6} \times 10^{-2} \times \frac{1000}{N}$; $N = 6 \times 10^{-5}$

$$S = M = \frac{N}{n_f} = \frac{6 \times 10^{-5}}{3} = 2 \times 10^{-5} \text{ mol L}^{-1}$$



$$K_{\text{sp}} = (3S)^3 \cdot S = 27 \cdot S^4$$

$$= 27 \times (2 \times 10^{-5})^4 = 4.32 \times 10^{-18}$$

BEGINNER'S BOX-1

1. The resistance of 1 N solution of CH_3COOH is 250Ω , when measured in a cell of cell constant 1.15 cm^{-1} . The equivalent conductance will be

- (1) $4.6 \Omega^{-1} \text{ cm}^2 \text{ eq}^{-1}$ (2) $9.2 \Omega^{-1} \text{ cm}^2 \text{ eq}^{-1}$
(3) $18.4 \Omega^{-1} \text{ cm}^2 \text{ eq}^{-1}$ (4) $0.023 \Omega^{-1} \text{ cm}^2 \text{ eq}^{-1}$

2. The specific conductivity of N/10 KCl solution at 20°C is $0.012 \Omega^{-1} \text{ cm}^{-1}$ and the resistance of the cell containing this solution at 20°C is 56Ω . The cell constant is

- (1) 4.616 cm^{-1} (2) 0.672 cm^{-1} (3) 2.173 cm^{-1} (4) 3.324 cm^{-1}

3. Molar conductances of BaCl_2 , H_2SO_4 and HCl at infinite dilutions are x_1 , x_2 and x_3 , respectively. Equivalent conductance of BaSO_4 at infinite dilution will be :

- (1) $\frac{[x_1 + x_2 - x_3]}{2}$ (2) $\frac{[x_1 - x_2 - x_3]}{2}$
(3) $2(x_1 + x_2 - 2x_3)$ (4) $\frac{[x_1 + x_2 - 2x_3]}{2}$

4. The ionization constant of a weak electrolyte is 25×10^{-6} while the equivalent conductance of its 0.01 M solution is $19.6 \text{ S cm}^2 \text{ eq}^{-1}$. The equivalent conductance of the electrolyte at infinite dilution (in $\text{S cm}^2 \text{ eq}^{-1}$) will be

- (1) 39.2 (2) 78.4 (3) 392 (4) 196

5. Which has maximum conductivity :

- (1) $[\text{Cr}(\text{NH}_3)_3 \text{Cl}_3]$ (2) $[\text{Cr}(\text{NH}_3)_4 \text{Cl}_2]\text{Cl}$ (3) $[\text{Cr}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ (4) $[\text{Cr}(\text{NH}_3)_6]\text{Cl}_3$

4.6 ELECTRODE POTENTIAL :


- The potential developed between metal electrode and its ions in solution is known as electrode potential.
- The potential developed between metal electrodes and the solution of its ions at 1 M concentration at 1 bar pressure and 298 K is known as **standard electrode potential**.
- There are two types of electrode potential :-

(a) Oxidation Potential (O.P.)

- The electrode potential for oxidation half reaction



- Tendency to get oxidised.
- **Greater the O.P. greater will be the tendency to get oxidised.**




(b) Reduction Potential (R.P.)

- The electrode potential for reduction half reaction



- Tendency to get reduced.
- **Greater the R.P. greater will be the tendency to get reduced.**



- Electrode potential depends upon :
 - Concentration of the solution.
 - Nature of the metal.
 - Pressure temperature conditions.


4.7 REFERENCE ELECTRODE :

- The potential of a single electrode cannot be determined but the potential difference between two electrodes can be accurately measured using a reference electrode.
- An electrode is chosen as a reference with respect to which all other electrodes are valued.
- There are two types of reference electrodes

(a) **Primary reference electrode : Standard hydrogen electrode (SHE)**


Representation : $\text{Pt}, \text{H}_2(\text{g}) \mid \text{H}^+ (1\text{M})$

When acts as anode




$$E^0_{\text{H}_2(\text{g})/\text{H}^+(\text{aq})} = \text{SOP}$$

When acts as cathode



$$E^0_{\text{H}^+/\text{H}_2(\text{g})} = \text{SRP}$$



- For SHE electrode potential is arbitrarily assumed to be zero at all temperature.



$$\text{SOP} = -\text{SRP} = 0 \text{ for SHE.}$$

- To calculate standard potential of any other electrode a cell is coupled with standard hydrogen electrode (SHE) and its potential is measured that gives the value of electrode potential of that electrode.

(b) **Secondary reference electrode :**


(i) **Calomel electrode (Hg_2Cl_2) -**

Representation : $\text{Pt} \mid \text{Hg}-\text{Hg}_2\text{Cl}_{2(\text{s})} \mid \text{KCl}_{(\text{aq})}$



(ii) **Silver-silver chloride electrode -**

Representation : $\text{Pt} \mid \text{Ag}-\text{AgCl}_{(\text{s})} \mid \text{KCl}_{(\text{aq})}$



4.8 CELL POTENTIAL (E^0_{cell}) :

$$E^0_{\text{cell}} = \text{SRP of cathode} - \text{SRP of anode}$$

$$E^0_{\text{cell}} = \text{SRP of cathode} + \text{SOP of anode}$$

$$E^0_{\text{cell}} = \text{SOP of anode} - \text{SOP of cathode}$$

(ii) **Reactivity of metals –**

$$\text{Reactivity of metal} \propto \text{SOP} \propto \frac{1}{\text{SRP}}$$

- In ECS reactivity of metal decreases from top to bottom.
- Li is most reactive metal.

(iii) **Reactivity of non-metals –**

$$\text{Reactivity of non-metal} \propto \text{SRP} \propto \frac{1}{\text{SOP}}$$

- In ECS reactivity of non-metal increases from top to bottom.
- F_2 is most reactive non-metal.

(iv) **Displacement reactions in solution –**

More reactive metal / non-metal displaces less reactive metal / non-metal in their solution.


- $\text{Cu} + 2\text{AgNO}_3 \rightarrow \text{Cu}(\text{NO}_3)_2 + 2\text{Ag}$
 $\text{Ag} + \text{Cu}(\text{NO}_3)_2 \rightarrow \text{No reaction}$
- $2\text{KI} + \text{Cl}_2 \rightarrow 2\text{KCl} + \text{I}_2$
 $\text{I}_2 + 2\text{KCl} \rightarrow \text{No reaction}$

(v) **Metal above hydrogen displaces H_2 from dilute acid solution.**

- $\text{Zn} + \text{H}_2\text{SO}_4(\text{aq}) \rightarrow \text{ZnSO}_4 + \text{H}_2(\text{g})$
 $\text{Cu} + \text{H}_2\text{SO}_4(\text{dilute}) \rightarrow \text{No reaction}$

4.11 GALVANIC CELL :

- It has two **half-cells**, each having a beaker containing a metal strip that dips in its aqueous salt solution.
- The metal strips are called **electrodes** and are connected by a conducting wire.
- Two solutions are connected by a **salt bridge**.
- The oxidation and reduction half reactions occur at separate electrodes and electric current flows through the wire.



Salt Bridge and its functions :

Salt bridge is inverted U-tube containing solution of inert electrolyte with agar-agar or gelatin to convert into semi solid form.

Ions of inert electrolyte do not involve in any chemical change.

The electrolyte in salt bridge should be such that speed of it's cation is nearly equal to speed of it's anion.

Ex. KCl , KNO_3 , NH_4NO_3

If Ag^+ , Hg_2^{+2} , Pb^{+2} , Tl^+ ions are present in a cell then KCl is not used because there can be formation of precipitate of AgCl , Hg_2Cl_2 , PbCl_2 or TlCl .


Functions

- It connects the solution of two half cell to complete the circuit.
- It maintains the electrical neutrality of the solution in order to give continuous flow or generation of current.
- If the salt bridge is removed then voltage drops to zero.
- It prevents the liquid-liquid junction potential. The potential difference which arises between two solutions when they brought in contact with each other.

Cell Representation :

We require two half cells to produce an electrochemical cell, which can be represented by following few rules;

- The anode half-cell is always written on the left followed on the right by cathode half cell.
- The separation of two phases (state of matter) is shown by a vertical line.
- The various materials present in the same phase are shown together using commas.
- The salt bridge is represented by a double slash ($\parallel$).
- The significant features of the substance like pressure of a gas, concentration of ions etc. are indicated in brackets immediately after writing the substance.



4.12 NERNST EQUATION :

It gives relation between electrode potential/ EMF of cell and concentration of electrolytic solution.

Walther Nernst derived a relation between free energy change (ΔG) and Reaction quotient (Q).

$$\Delta G = \Delta G^\circ + RT \ln Q \quad \dots(1)$$

where ΔG and ΔG° are free energy change and standard free energy change; 'Q' is reaction quotient.

$$\therefore -\Delta G = nFE \quad \text{and} \quad -\Delta G^\circ = nFE^\circ$$

Thus from Eq. (1), $-nFE = -nFE^\circ + RT \ln Q$

$$E = E^\circ - \frac{2.303 RT}{nF} \log Q$$

Where - E° = standard electrode potential,

R = gas constant,

T = temperature (in K)

F = Faraday (96500 coulomb mol^{-1}),


n = number of moles of e^- gained lost or transferred in balanced equation.

At 25°C, above equation may be written as

$$E = E^\circ - \frac{0.0591}{n} \log Q$$

$$E = E^\circ - \frac{0.0591}{n} \log \frac{[P]}{[R]}$$


In general, for a redox cell reaction involving the transference of n electrons



$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.0591}{n} \log \frac{[C]^c [D]^d}{[A]^a [B]^b} \quad \text{or} \quad E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.0591}{n} \log \frac{[\text{Product}]}{[\text{Reactant}]}$$

4.13 APPLICATIONS OF NERNST EQUATION

(i) Calculation of electrode potential (E_{RP} or E_{OP}) -




$$E_{RP} = E_{RP}^0 - \frac{0.0591}{n} \log \frac{[M]}{[M^{+n}]}$$

$$E_{RP} = E_{RP}^0 - \frac{0.0591}{n} \log \frac{1}{[M^{+n}]}$$

If $[M^{+n}]$ increases, then E_{RP} increases

(ii) Calculation of electrode potential and pH of hydrogen electrode -



$$E_{RP} = E_{RP}^0 - \frac{0.0591}{2} \log \frac{P_{H_2}}{[H^{+}]^2}$$


$$\therefore E_{RP}^0 = 0; \quad E_{RP} = E_{RP}^0 - \frac{0.0591}{2} \log \frac{1}{[H^{+}]^2} \quad [P_{H_2} \text{ is taken 1 atm}]$$

$$E_{RP} = 0.0591 \log [H^{+}]$$

$$E_{RP} = -0.0591 \text{ pH}$$

$$E_{OP} = +0.0591 \text{ pH}$$

(iii) Calculation of EMF of cell (E_{cell}) -



$$E_{cell} = E_{cell}^0 - \frac{0.0591}{2} \log \frac{[Zn^{+2}]}{[Cu^{+2}]}$$

If $[Cu^{+2}]$ increases, then E_{cell} increases

If $[Zn^{+2}]$ increases, then E_{cell} decreases

(iv) Prediction and feasibility of a cell reaction -

For a feasible cell reaction

$$\Delta G = -ve \quad (\Delta G = -nFE_{cell})$$

$$E_{cell} = +ve$$

(v) Calculation of equilibrium constant (K_{eq}) and ΔG° -

From Nernst equation -

$$E_{cell} = E_{cell}^0 - \frac{0.0591}{n} \log \frac{[P]}{[R]}$$

At equilibrium, $E_{cell} = 0$ and $\frac{[P]}{[R]} = K_{eq}$

$$E_{\text{cell}}^0 = \frac{0.0591}{n} \log K_{\text{eq}}$$


$$E_{\text{cell}}^0 = \frac{2.303RT}{nF} \log K_{\text{eq}}$$

$$n F E_{\text{cell}}^0 = 2.303 RT \log K_{\text{eq}}$$

$$\Delta G^\circ = -2.303RT \log K_{\text{eq}}$$

Illustrations

Illustration 8. Write cell representation for the following redox reaction,



Solution The cell consists of a platinum wire anode dipped in Sn^{2+} solution and a silver cathode dipped in Ag^+ solution therefore $\text{Pt}(\text{s}) | \text{Sn}^{2+}(\text{aq}), \text{Sn}^{4+}(\text{aq}) || \text{Ag}^+(\text{aq}) | \text{Ag}(\text{s})$.

Illustration 9. Calculate the EMF of a Daniel cell when the concentration of ZnSO_4 and CuSO_4 are 0.001 M and 0.1 M respectively. The standard EMF of the cell is 1.1 V.

Solution $\text{Zn}(\text{s}) + \text{CuSO}_4 \rightarrow \text{ZnSO}_4 + \text{Cu}$

$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.0591}{2} \log \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]} = 1.1 - \frac{0.0591}{2} \log \frac{10^{-3}}{10^{-1}} = 1.159 \text{ V}$$

Illustration 10. Calculate E^0 and E for the cell $\text{Sn} | \text{Sn}^{2+} (1\text{M}) || \text{Pb}^{2+} (10^{-3}\text{M}) | \text{Pb}$, $E^0 (\text{Sn}^{2+} | \text{Sn}) = -0.14\text{V}$, $E^0 (\text{Pb}^{2+} | \text{Pb}) = -0.13 \text{ V}$. Is cell representation correct?

Solution $E_{\text{cell}}^0 = E_{\text{Pb}^{2+}/\text{Pb}}^0 - E_{\text{Sn}^{2+}/\text{Sn}}^0 = -0.13 - (-0.14) = +0.01 \text{ V}$

$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.0591}{2} \log \frac{[\text{Sn}^{2+}]}{[\text{Pb}^{2+}]} = +0.01 - \frac{0.0591}{2} \log \left[\frac{1}{10^{-3}} \right] = -0.078 \text{ V cell with not work}$$

Illustration 11. Calculate the equilibrium constant for the reaction $\text{Fe}^{2+} + \text{Ce}^{4+} \rightleftharpoons \text{Fe}^{3+} + \text{Ce}^{3+}$,

[Given : $E_{\text{Ce}^{4+}/\text{Ce}^{3+}}^0 = 1.44 \text{ V}$; $E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^0 = 0.68 \text{ V}$]

$$\text{Take } \frac{2.303 RT}{F} = 0.06 \text{ at } 25^\circ\text{C}, \log 4.68 = 0.67$$

Solution $\text{Fe}^{2+} + \text{Ce}^{4+} \rightleftharpoons \text{Fe}^{3+} + \text{Ce}^{3+}$

$$E^0 = 1.44 - 0.68 = 0.76\text{V} = \frac{0.06}{1} \log K_{\text{c}}; K_{\text{c}} = 4.64 \times 10^{12}$$

Illustration 12. The 0.1 M copper sulphate solution in which copper electrode is dipped at 25°C . Calculate the electrode potential of copper electrode [Given $E_{\text{Cu}^{2+}/\text{Cu}}^0 = 0.34\text{V}$]

Solution. $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$

$$E_{\text{red}} = E_{\text{red}}^0 - \frac{0.059}{n} \log \frac{[\text{Product}]}{[\text{Reactant}]}$$

$$\text{Here } n = 2 \text{ so } E = 0.34 - \frac{0.059}{2} \log 10$$

$$E = 0.34 - 0.03 = 0.31 \text{ V}$$

Illustration 13. Calculate the EMF of the cell $\text{Cr} \mid \text{Cr}^{+3} (0.1\text{M}) \parallel \text{Fe}^{+2} (0.01\text{M}) \mid \text{Fe}$

(Given $E^\circ_{\text{Cr}^{+3}/\text{Cr}} = -0.75\text{V}$, $E^\circ_{\text{Fe}^{+2}/\text{Fe}} = -0.45\text{V}$)

Solution.

Half cell reactions are –

At Anode $[\text{Cr} \rightarrow \text{Cr}^{+3} + 3e^-] \times 2$

At Cathode $[\text{Fe}^{+2} + 2e^- \rightarrow \text{Fe}] \times 3$

Cell reaction $2\text{Cr} + 3\text{Fe}^{+2} \rightarrow 2\text{Cr}^{+3} + 3\text{Fe}$

$E^\circ_{\text{cell}} = \text{Oxidation potential} + \text{Reduction potential} = 0.75 + (-0.45) = 0.30\text{V}$

$$E_{\text{cell}} = E^\circ - \frac{0.059}{n} \log \frac{[\text{Product}]}{[\text{Reactant}]} = 0.30 - \frac{0.059}{6} \log \frac{[\text{Cr}^{+3}]^2}{[\text{Fe}^{+2}]^3} = 0.30 - \frac{0.059}{6} \log \frac{[0.1]^2}{[0.01]^3}$$

$$= 0.30 + \frac{0.24}{6} = 0.34\text{V}$$

Illustration 14. For the cell $\text{Pt(s)} \mid \text{H}_2(0.4\text{ atm}) \mid \text{H}^+(\text{pH}=1) \parallel \text{H}^+(\text{pH}=2) \mid \text{H}_2(0.1\text{ atm}) \mid \text{Pt}$
The measured potential at 25°C is

(1) -0.1V (2) -0.5V (3) -0.041V (4) -0.030V

Solution

Anode reaction : $\text{H}_2(0.4\text{ atm}) \rightarrow 2\text{H}^+(10^{-1}\text{M}) + 2e^-$

Cathode reaction : $2\text{H}^+(10^{-2}\text{M}) + 2e^- \rightarrow \text{H}_2(0.1\text{ atm})$

Cell reaction $\text{H}_2(0.4\text{ atm}) + 2\text{H}^+(10^{-2}\text{M}) \rightarrow 2\text{H}^+(10^{-1}\text{M}) + \text{H}_2(0.1\text{ atm})$

$$E = 0 - \frac{0.059}{2} \log \frac{(10^{-1})^2(0.1)}{(0.4)(10^{-4})} = -0.041\text{V}$$

Illustration 15. If $E^\circ_{\text{Au}^+/\text{Au}}$ is 1.69V & $E^\circ_{\text{Au}^{3+}/\text{Au}}$ is 1.40V , then $E^\circ_{\text{Au}^{3+}/\text{Au}^+}$ will be

(1) 0.19V (2) 2.945V (3) 1.255V (4) none

Solution

$\text{Au}^+ + e^- \longrightarrow \text{Au(s)}$ $E^\circ = 1.69\text{V}$... (1); ΔG_1°

$\text{Au}^{3+} + 3e^- \longrightarrow \text{Au(s)}$ $E^\circ = 1.40\text{V}$... (2); ΔG_2°

From (2) – (1)

$\text{Au}^{3+} + 2e^- \longrightarrow \text{Au}^+$... (3); ΔG_3°

$$\Delta G_3^\circ = \Delta G_2^\circ - \Delta G_1^\circ$$

$$-2 \times F \times E^\circ = -3 \times F \times 1.40 + 1 \times 1.69 \times F$$

$$E^\circ = 1.255\text{V}$$

GOLDEN KEY POINTS

- The value of electrode potential does not depend on stoichiometry of half reactions as it is an intensive property.
- Gibb's free energy is extensive property.
- In electrolytic cell, cathode is negative terminal whereas in Galvanic cell cathode is positive terminal.
- Cell reaction is spontaneous, if $\Delta G = -ve$ or $E_{\text{cell}} = +ve$
- Cell reaction will be in equilibrium, if e.g. $\Delta G = 0$ or $-nFE_{\text{cell}} = 0$ or $E_{\text{cell}} = 0$

● **Work done by a cell :**

- (i) Let 'n' faraday charge be taken out of a cell of EMF 'E'; then work done by the cell will be calculated as:
 Work = Charge $\times$ Potential = $nF \times E$
- (ii) Work done by cell = Decrease in free energy
 so $-\Delta G = nFE$ or $W_{\max} = +nFE^\circ$, where E° is standard EMF of the cell.

BEGINNER'S BOX-2

1. Calculate E°_{cell} for the following cell




- (1) 0.04 V (2) - 1.56 V (3) 1.56 V (4) 0.84 V

2. Standard EMF of a galvanic cell involving $2e^-$ transfer is found to be 0.591 V at 25°C . Calculate equilibrium constant for the cell reaction

- (1) 20^{10} (2) 10^{20} (3) 10^{200} (4) 200^{10}


3. Calculate cell potential at 298 K for the following galvanic cell



$$E^\circ_{\text{Cd}^{+2}/\text{Cd}} = -0.40\text{V}$$

- (1) 0.38 (2) - 0.38 (3) 0.36 (4) - 0.36

4. Calculate E°_{cell} of the following galvanic cell at 298 K ;



$$E^\circ_{\text{Ca}^{+2}/\text{Ca}} = -2.87\text{V}; E^\circ_{\text{Fe}/\text{Fe}^{+2}} = 0.41\text{V}$$

- (1) 2.46 V (2) - 2.46 V (3) 3.28 V (4) - 3.28 V

5. Given $E^\circ_{\text{Ag}^{+}/\text{Ag}} = 0.80\text{V}$, $E^\circ_{\text{Mg}^{2+}/\text{Mg}} = -2.37\text{V}$, $E^\circ_{\text{Cu}^{2+}/\text{Cu}} = 0.34\text{V}$, $E^\circ_{\text{Hg}^{2+}/\text{Hg}} = 0.79\text{V}$.

Which of the following statement is correct

- (1) AgNO_3 can be stored in copper vessel
 (2) $\text{Mg}(\text{NO}_3)_2$ can not be stored in copper vessel
 (3) CuCl_2 can be stored in silver vessel
 (4) HgCl_2 can be stored in copper vessel

6. $1/2 \text{H}_2(\text{s}) + \text{AgCl(s)} \longrightarrow \text{H}^{+}(\text{aq}) + \text{Cl}^{-}(\text{aq}) + \text{Ag(s)}$ occurs in the galvanic cell :

- (1) $\text{Ag(g)} | \text{AgCl(s)} | \text{KCl(sol)} || \text{AgNO}_3(\text{sol}) | \text{Ag(s)}$
 (2) $\text{Pt(s)} | \text{H}_2(\text{g}) | \text{HCl(sol)} || \text{AgNO}_3(\text{sol}) | \text{Ag(s)}$
 (3) $\text{Pt(s)} | \text{H}_2(\text{g}) | \text{HCl(sol)} || \text{Cl}^{-}(\text{aq}) | \text{AgCl(sol)} | \text{Ag(s)}$
 (4) $\text{Pt(s)} | \text{H}_2(\text{g}) | \text{KCl(sol)} || \text{AgCl(sol)} | \text{Ag(s)}$

4.14 SOME SPECIAL CELLS

(A) CONCENTRATION CELLS

A concentration cell is a form of galvanic cell that has two equivalent half cells of the same material having difference only in concentrations. For such cells anode and cathode are same so $E^\circ_{\text{cell}} = 0$.

eg. $\text{Pt}, \text{H}_2(\text{g}) (P_1 \text{ atm}) | \text{H}^{+} (C_1) || \text{H}^{+} (C_2) | \text{H}_2(\text{g}) (P_2 \text{ atm}), \text{Pt}$

Anode Reaction : $\text{H}_2(\text{P}_1) \longrightarrow 2\text{H}^+(\text{C}_1) + 2\text{e}^-$

Cathode Reaction : $2\text{H}^+(\text{C}_2) + 2\text{e}^- \longrightarrow \text{H}_2(\text{P}_2)$

Cell Reaction : $\text{H}_2(\text{P}_1) + 2\text{H}^+(\text{C}_2) \longrightarrow 2\text{H}^+(\text{C}_1) + \text{H}_2(\text{P}_2)$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{0.0591}{n} \log \frac{[\text{P}]}{[\text{R}]}$$

$$\text{So, } E_{\text{cell}} = 0 - \frac{0.0591}{2} \log \left[\frac{C_1}{C_2} \right]^2 \frac{P_2}{P_1}$$

Case 1 : If $P_1 = P_2 = 1 \text{ atm}$; $E_{\text{cell}} = -\frac{0.059}{2} \log \left[\frac{C_1}{C_2} \right]^2$

$$E_{\text{cell}} = -0.0591 \log \left[\frac{C_1}{C_2} \right]$$

If $C_2 > C_1$ then $E_{\text{cell}} > 0$ and cell will be spontaneous.

Case 2 : If $C_1 = C_2$ $E_{\text{cell}} = -\frac{0.0591}{2} \log \left[\frac{P_2}{P_1} \right]$


$P_1 \neq P_2$

If $P_1 > P_2$ then $E_{\text{cell}} > 0$ and cell will be spontaneous.

(B) REVERSIBLE CELLS :


Those cell's in which chemical reactions are reversed by applying external emf in opposite directions known as reversible cells. A daniel cell is said to be reversible if it follows two conditions :

- (i) When $E_{\text{ext}} > 1.1 \text{ V}$ (emf of daniel cell) : cell reactions reversed
 - (ii) When $E_{\text{ext}} = 1.1 \text{ V}$ (emf of daniel cell) ; equilibrium will be established.
- In general secondary cells are reversible in nature.



4.15 ELECTROLYSIS

The process of chemical decomposition of an electrolyte by the passage of electricity is called electrolysis or electrolytic dissociation. It is carried out in **electrolytic cell** where electrical energy is converted into chemical energy. For electrolysis two suitable electrodes are immersed in the solution of an electrolyte. When an electric potential is applied between the electrodes, the positive ions move towards the cathode and negative ions move towards the anode. The gain of electrons (decrease in oxidation number) means reduction takes place at the cathode and loss of electrons (increase in oxidation number) means oxidation takes place at anode.




There are two aspects of electrolysis :

(A) Qualitative aspects of electrolysis :-

In qualitative aspect products formed at electrode are identified. At electrode product formation depends up on the following :


- (i) Nature of electrolyte
- (ii) Nature of electrodes : The metal strip at which current enters is called **anode**; anode is positively charged in electrolytic cell. On the other hand, the electrode at which current leaves is called **cathode**. Cathode is negatively charged. There are two types of electrodes
 - (a) Attackable : The attackable electrodes participate in the electrode reaction. They are made up of reactive metals like Zn, Cu, Ag etc in their respective solutions. In such electrodes, atom of the metal gets oxidised into the corresponding cation, which is passed into the solution. Thus, such anode gets dissolved and their mass decreases.
 - If electrode is active at cathode, metal goes on depositing and at anode metal is dissolved. The process is called electroplating.
 - (b) Non - attackable : Non-attackable electrodes do not participate in the electrode reaction as they are made up of unreactive elements like Pt, graphite etc. Such electrodes do not dissolve and their mass remain same.
 - Among the two cations, that cation is discharged at the cathode which has higher reduction potential. Among the two anions, that anion will be discharged at the anode which has lower reduction potential or higher oxidation potential.
 - The discharge of NO_3^- , PO_4^{3-} and SO_4^{2-} ions at anode does not commonly take place from aqueous solution.
- (iii) Concentration of electrolytic solution.

Examples of electrolysis :**(a) Electrolysis of Molten Sodium Chloride**

(b) **Electrolysis of Sodium Chloride Solution**


On electrolysis of aqueous solution of NaCl, the possible species involved in half-reactions are Na^+ , Cl^- and H_2O .

The possible cathode half-reactions are



Under the standard conditions, H_2O is to be reduced in preference to Na^+ and Hydrogen is evolved at the cathode.



The possible anode half-reactions are



Under the standard-state conditions, H_2O is to be oxidized in preference to Cl^- . But since electrode potentials, depend on concentrations it turns out that when the solution is concentrated enough in Cl^- , Cl_2 is the product; but in dilute solution, O_2 is the product.




Starting with very dilute NaCl solutions, the oxidation potential of Cl^- is very negative, so H_2O is oxidised in preference to Cl^- . But as the NaCl concentration increase, the oxidation potential of Cl^- increases until eventually Cl^- is oxidized in preference to H_2O . The product changes from O_2 to Cl_2 .

The half-reactions and cell reaction for the electrolysis of aqueous sodium chloride to chlorine and hydroxide ion are as follows:



Because the electrolysis started with sodium chloride, the cation in the electrolyte solution is Na^+ . On evaporation of the electrolyte solution, sodium hydroxide NaOH is obtained.



(c) **Electrolysis of Molten lead bromide using inert (pt/graphite) electrodes.**



$$E_{\text{cell}} = -0.13 - 1.09 = -1.22\text{ V}$$



$$E_{\text{ext}} > 1.22\text{ V}$$

(d) **Electrolysis of aq CuSO_4 using inert (pt/graphite) electrode**



At cathode reduction of $\text{Cu}^{2+}(\text{aq})$ will be preferred over reduction of water as its standard reduction potential is more, than that of water.


Thus reaction at cathode : $\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu}(\text{s})$



At anode oxidation of water is preferred over oxidation of sulphate ion as its standard oxidation potential is more, than that of sulphate ion.



Thus reaction at anode : $2\text{H}_2\text{O}(\text{l}) \rightarrow \text{O}_2(\text{g}) + 4\text{H}^+ + 4e^-$

After electrolysis solution will become acidic ($\text{pH} < 7$)

(e) Electrolysis of aq. CuSO₄ using Cu electrode.

At cathode reduction of Cu²⁺ (aq) will be preferred over reduction of water as its standard reduction potential is more, than that of water.

Thus reaction at cathode :




At anode oxidation of copper solid is preferred over oxidation of water and sulphate ion, as its standard oxidation potential is more than that of water and sulphate ion.


Thus reaction at anode :




It is an example of electrolytic refining and there will be no change in the pH and concentration of electrolyte.

(f) Electrolysis of dilute H₂SO₄ :


due to less availability of H⁺ ions, it will not participate in formation of H₂ gas therefore reduction of water will take place.



oxidation of water is preferred over oxidation of sulphate ion as its standard oxidation potential is more, than that of sulphate ion.

(g) Electrolysis of concentrated H₂SO₄




due to reduction of H⁺ ions, formation of H₂ gas takes place.



due to oxidation of SO₄²⁻ ions, formation of persulphate ion (S₂O₈²⁻) takes place.

(B) QUANTITATIVE ASPECTS OF ELECTROLYSIS

1 mole electron = N_A electron = 96500 coulomb = 1 Faraday



nF charge discharges = 1 mol of M⁺ⁿ ion

∴ 1 F charge discharges = $\frac{1}{n}$ mol of M⁺ⁿ ion

$$= \frac{1}{n} \times n = 1 \text{ g equivalent of M}^{+n} \text{ ion}$$

Hence 1 mole e⁻ = 1 F = 1 g equivalent

FARADAY'S LAWS OF ELECTROLYSIS

(a) First law of electrolysis :

Amount of substance deposited or liberated at an electrode is directly proportional to amount of charge passed through the solution.

$$W \propto Q$$

W = amount of substance deposited, Q = charge in coulomb

$$W = ZQ$$

Z = electrochemical equivalent

when Q = 1 coulomb, then W = Z

Thus, amount of substance deposited or liberated by 1 coulomb charge is called **electrochemical equivalent**.

Let I ampere current is passed till 't' seconds .

$$\text{Then, } Q = It \quad \therefore W = ZIt$$

1 Faraday = 96500 coulomb = Charge on one mole electrons

One faraday is the charge required to liberate or deposit one gram equivalent of a substance at corresponding electrode.

Let 'E' is equivalent weight then 'E' gram will be liberated by 96500 coulomb.

$$\therefore 1 \text{ Coulomb will liberate } \frac{E}{96500} \text{ gram ;}$$

$$\text{By definition, } Z = \frac{E}{96500} \quad \therefore W = \frac{EIt}{96500} \quad \boxed{\frac{W}{E} = \frac{It}{96500} = \text{number of g eq} = \text{number of faraday}}$$

(b) Second law of electrolysis :

When same amount of charge is passed through different electrolyte solutions connected in series then weight of substances deposited or liberated at electrodes are in ratio of their respective equivalent weights.

$$\text{i.e. } \frac{W_1}{W_2} = \frac{E_1}{E_2}$$

GOLDEN KEY POINTS

PRODUCTS OF ELECTROLYSIS OF SOME ELECTROLYTES


S. No.	Electrolyte	Electrode	Product obtained at anode	Product obtained at cathode
(i)	Aqueous NaCl	Pt or Graphite	Cl ₂	H ₂
(ii)	Fused NaCl	Pt or Graphite	Cl ₂	Na
(iii)	Aqueous NaOH	Pt or Graphite	O ₂	H ₂
(iv)	Fused NaOH	Pt or Graphite	O ₂	Na
(v)	Aqueous CuSO ₄	Pt or Graphite	O ₂	Cu
(vi)	Aqueous CuSO ₄	Copper	Cu oxidises to Cu ⁺² Ions	Cu
(vii)	Aqueous CuCl ₂	Copper	Cu oxidises to Cu ⁺² ions	Cu
(viii)	Aqueous HCl	Pt or Graphite	Cl ₂	H ₂
(ix)	Dilute H ₂ SO ₄	Pt or Graphite	O ₂	H ₂
(x)	Conc. H ₂ SO ₄	Pt or Graphite	S ₂ O ₈ ⁻²	H ₂
(xi)	Aqueous AgNO ₃	Pt of Graphite	O ₂	Ag
(xii)	Dilute NaCl	Pt or Graphite	O ₂	H ₂

- For electrolysis ΔG = +ve, So it is non-spontaneous process.
- In electrolytic cell D.C. current is used.

Illustrations

Illustration 16 How much electric charge is required to oxidise :(a) 1 mol of H_2O to O_2 (b) 1 mol of FeO to Fe_2O_3 ?**Solution**

(a) The oxidation reaction is :



1 mol 2 mol

$$Q = 2F = 2 \times 96500 = 1.93 \times 10^5 \text{ C}$$

(b) The oxidation reaction is $\text{FeO} + \frac{1}{2}\text{H}_2\text{O} \longrightarrow \frac{1}{2}\text{Fe}_2\text{O}_3 + \text{H}^+ + \text{e}^-$

$$Q = F = 96500 \text{ C}$$

Illustration 17 One mole of electron passes through each of the solution of AgNO_3 , CuSO_4 and AlCl_3 when Ag, Cu and Al are deposited at cathode. The molar ratio of Ag, Cu and Al deposited are

(1) 1 : 1 : 1

(2) 6 : 3 : 2

(3) 6 : 3 : 1

(4) 1 : 3 : 6

Solution

Deposited equivalent will be same

$$1 \text{ eq. Ag} = 1 \text{ mol Ag}$$


$$1 \text{ eq. Cu} = \frac{1}{2} \text{ mol Cu}$$

$$1 \text{ eq. Al} = \frac{1}{3} \text{ mol Al}$$

$$\Rightarrow \text{ratio} \quad 1 : \frac{1}{2} : \frac{1}{3} \Rightarrow 6 : 3 : 2$$


Illustration 18 Exactly 0.4 F electric charge is passed through three electrolytic cells connected in a series. First cell containing AgNO_3 , second cell containing CuSO_4 and third cell containing FeCl_3 solution. How many grams of each metal will be deposited assuming only cathodic reaction in each cell.**Solution**

The cathodic reaction in the cell are respectively,




1 mol 1 mol

108 g 1 F



1 mol 1 mol

63.5 g 2 F



1 mol 1 mol

56 g 3 F


Hence, Ag deposited = $108 \times 0.4 = 43.2 \text{ g}$

$$\text{Cu deposited} = \frac{63.5}{2} \times 0.4 = 12.7 \text{ g}$$

$$\text{and} \quad \text{Fe deposited} = \frac{56}{3} \times 0.4 = 7.47 \text{ g}$$

Illustration 19 An electric current of 100 A is passed through a molten liquid of sodium chloride for 5 hours. Calculate the volume of chlorine gas liberated at the electrode at NTP.**Solution**

The reaction taking place at anode is

71.0 g 71.0 g $2 \times 96500 \text{ C}$

= 1 mol

$$Q = i \times t = 100 \times 5 \times 60 \times 60 \text{ C}$$

The amount of chlorine liberated by passing $100 \times 5 \times 60 \times 60 \text{ C}$ of electric charge.

$$= \frac{1}{2 \times 96500} \times 100 \times 5 \times 60 \times 60 = 9.3264 \text{ mole}$$

$$\text{Volume of } \text{Cl}_2 \text{ liberated at NTP} = 9.3264 \times 22.4 = 208.91 \text{ L}$$

Illustration 20 How long a current of 2 A has to be passed through a solution of AgNO_3 to coat a metal surface of 80 cm^2 with $5 \mu\text{m}$ thick layer? Density of silver = 10.8 g/cm^3 .

Solution

$$d = \frac{M}{V} \Rightarrow 10.8 = \frac{M}{80 \times 5 \times 10^{-4}} \Rightarrow M = 10.8 \times 400 \times 10^{-4}$$

$$W = \frac{E \times I \times t}{96500} \Rightarrow 10.8 \times 400 \times 10^{-4} = \frac{108 \times 2 \times t}{96500} \Rightarrow t = 193 \text{ s}$$

BEGINNER'S BOX-3



- 1 L of 1 M CuSO_4 solution is electrolysed. After passing 2 F charge, the molarity of CuSO_4 will be
(1) M/2 (2) M/4 (3) M (4) zero
- Same quantity of current is passed through molten NaCl and molten Al_2O_3 . If 4.6 g of Na was deposited in one cell, the mass of Al deposited in other cell is :-
(1) 0.9 g (2) 1.8 g (3) 2.7 g (4) 3.6 g
- A certain current liberates 0.5 g of hydrogen in 2 hr. How many grams of copper can be deposited by the same current flowing for the same time in a copper sulphate solution?
(1) 12.7 g (2) 15.9 g (3) 31.8 g (4) 63.5 g
- A quantity of electric charge that brings about the deposition of 4.5 g Al from Al^{+3} at the cathode will also produce the following volume (STP) of $\text{H}_2(\text{g})$ from H^+ at the cathode
(1) 44.8 L (2) 22.4 L (3) 11.2 L (4) 5.6 L
- A current of 9.65 A is passed through the aqueous solution of NaCl using suitable electrodes for 1000 s. The amount of NaOH formed during electrolysis is
(1) 2.0 g (2) 4.0 g (3) 6.0 g (4) 8.0 g
- If 0.224 L of H_2 gas is formed at the cathode, the volume of O_2 gas formed at the anode under identical conditions, is
(1) 0.224 L (2) 0.448 L (3) 0.112 L (4) 1.12 L

4.16 COMMERCIAL VOLTAIC CELLS


Batteries can be classified as primary and secondary. Primary batteries can not be returned to their original state by recharging, so when the reactants are consumed, the battery is "dead" and must be discarded. Secondary batteries are often called storage batteries or rechargeable batteries. The reactions in these batteries can be reversed; thus, the batteries can be recharged.

(A) Primary cells : Dry cells and mercury cells


The most familiar type of battery is the dry cell which is a compact of Leclanche cell known after its discoverer Leclanche (fig.). In this cell, the anode consists of a zinc container and the cathode is a graphite rod surrounded by powdered MnO_2 and carbon. The space between the electrodes is filled with a moist paste of NH_4Cl and ZnCl_2 . The electrode reactions are complex, but they can be written approximately as follows.




In the cathode reaction, manganese is reduced from the 4+ oxidation state to the 3+ state. Ammonia is not liberated as a gas but combines with Zn^{2+} to form $\text{Zn}(\text{NH}_3)_4^{2+}$ ion. Dry cells do not have an indefinite life as acidic NH_4Cl corrodes the zinc container even when not in use. **Dry cells have a potential of approximately 1.25 to 1.5V.**




Mercury cell, suitable for low current devices like hearing aids, watches, etc. consists of zinc – mercury amalgam as anode and a paste of HgO and carbon as the cathode. The electrolyte is a paste of KOH and ZnO. The electrode reactions for the cell are given below :



The overall reaction is represented by



The cell potential is **approximately 1.35 V** and remains constant during its life as the overall reaction does not involve any ion in solution whose concentration can change during its life time.



(B) SECONDARY OR RECHARGEABLE BATTERIES

The most important secondary cell is the lead storage battery. It consists of a **lead anode** and a grid of lead packed with **lead dioxide as cathode**. A solution of sulphuric acid (38 percent by mass or having density 1.30 gm cm^{-3}) is used as an electrolyte.





Fig : The Lead storage battery

When the cell discharges, it operates as a voltaic cell. The lead is oxidised to Pb^{+2} ion and lead plates acquire a negative charge.






Pb^{+2} ions combine with sulphate ions of solution to form insoluble PbSO_4






The electrons given by Pb are used at cathode (PbO_2 electrode)




At Cathode :



Therefore the overall cell reaction during discharging




When a potential slightly greater than the potential of battery is applied, the battery can be recharged. During recharging the cell is operated like an **electrolytic cell** i.e. now electrical energy is supplied to it from an external source. The electrode reactions are reverse of those that occur during discharge :



It is clear that H_2SO_4 is used up during the discharge & produced during recharging reaction of cell.




The discharging process of the storage cell is based on the principles of electrochemical cell, whereas the charging process is based upon the principles of electrolytic cells.

- n factor for H_2SO_4 during charging and discharging is equal to one.



(C) Nickel - Cadmium ("Ni - Cd") batteries :

They are used in variety of cordless appliances such as telephones, video camcorders, and cordless power tools, are lightweight and rechargeable. The chemistry of the cell utilizes the oxidation of cadmium and the reduction of nickel (III) oxide under basic conditions.




As the active masses of all reaction components are independent of their amount. The cell delivers constant EMF throughout.




(D) Fuel Cells

It is possible to make batteries in which the reactants are fed continuously to the electrodes. Electrical cells that are designated to convert the free energy from the combustion of fuels such as hydrogen, carbon monoxide or methane directly into electrical energy are called **fuel cells**.

One of the most successful fuel cells uses the reaction of hydrogen with oxygen to form water (fig.). This cell has been used for electric power in the Appollo space programme. The water vapours produced were condensed and added to the drinking water supply for the astronauts.



In the cell shown in figure hydrogen and oxygen are bubbled through a porous carbon electrode into concentrated aqueous sodium hydroxide. Catalysts are incorporated in the electrode. The electrode reactions are :



This cell runs continuously as long as the reactants are supplied. Since fuel cells convert the energy of a fuel directly to electricity, they are potentially more efficient than the conventional method of generating electricity on a large scale by burning hydrogen, carbon fuels or by using nuclear reactor. Although **theoretically we expect 100 percent efficiency in fuel cells, so far an efficiency of only 60-70 per cent has been attained**. Since fuel cells are efficient and free from pollution, efforts are being made to produce better commercially practical fuel cells.

Efficiency of a Fuel Cell : $\eta = \frac{(\Delta G)}{(\Delta H)} \times 100$


Main features of different cells.

	Name of the cell/Battery	Anode	Cathode	Electrolyte
(i)	Dry cell	Zinc	Graphite surrounded by MnO_2 & C.	$\text{NH}_4\text{Cl} + \text{ZnCl}_2$ (touching anode)
(ii)	Mercury cell	Zinc/Mercury Amalgum	Paste of HgO and carbon	Paste of KOH and ZnO
(iii)	Lead storage battery	Lead	Lead dioxide	H_2SO_4 (38%)
(iv)	Ni-Cd cell	Cadmium	Nickel dioxide	KOH sol.
(v)	H_2 - O_2 fuel cell	Porous carbon containing catalysts (H_2 passed)	Porous carbon containing catalysts (O_2 passed)	Conc. aq. NaOH sol.


4.17 CORROSION

Corrosion is basically an electrochemical phenomenon. The rusting of iron, the tarnishing of silver, development of a green coating on copper and bronze are some of the examples of corrosion. Corrosion causes enormous damage to building, bridges, ships and to many other articles made of iron.


In corrosion, a metal is oxidised by loss of electrons to oxygen and forms metal oxide. Corrosion of iron (which is commonly known as rusting) occurs in presence of water and oxygen (air). Although the chemistry of corrosion is complex but it is understood that at one spot of an iron object oxidation occurs and that spot behaves as an anode.




Electrons released at anode move through the metal and go to another spot on the metal and reduce the oxygen in presence of H^+ ions (which is believed to be available from H_2CO_3 formed due to dissolution of CO_2 in water). This spot works as a cathode :



Here Fe^{2+} ions move through water on the surface of the iron object. (If water present is saline, it will help more in carrying the current in the miniature cell thus formed and will enhance corrosion). The overall reaction of the miniature cell is the sum of the cathode and anode reactions as follows :



The Fe^{2+} ions are further oxidised by atmospheric oxygen to Fe^{3+} (as Fe_2O_3) and comes out as rust in the form of a hydrated iron (III) oxide expressed as $\text{Fe}_2\text{O}_3 \cdot x\text{H}_2\text{O}$.



H^+ ions produced in the above reaction help further in rusting.

Impurities present in iron also enhance rusting by setting a number of miniature cells. Very pure iron does not rust quickly.



4.18 PREVENTION FROM CORROSION

Corrosion of a metal is prevented by applying protective coating (such as grease, paint, bisphenol or metal coatings) on metal surface. In case of iron this coating is done in two ways : (i) electrolysis (Cr, Ni and Cd coating) (ii) dipping iron objects in a molten metal (Zn and Sn coating). Use of zinc coating to protect iron is called Galvanisation. In galvanising iron, zinc being more reactive than iron serves as an anode and is oxidised




Here it is important to note that iron does not rust even after zinc coating is broken which is not true in case of tin coating over iron

($E^0_{\text{Sn}^{2+}/\text{Sn}} = -0.14\text{V}$). Now if the coating is broken, iron is exposed and iron being more active than both copper and tin, is corroded. Here iron corrodes more rapidly than it does in the absence of tin. But tin protects copper in the same way as zinc protects iron ($E^0_{\text{Cu}^{2+}/\text{Cu}} = 0.34\text{V}$).



Instead of coating more reactive metal on iron, the use of such metals is made as sacrificial anode. This method of preventing iron from corrosion is called CATHODIC PROTECTION.

In this method, a plate of reactive metal (Zn or Mg) is buried beside the iron pipe or tank and connected to it by wires. Here iron becomes **cathode** and more reactive metal becomes **anode**. The reactive metal anode is sacrificed to protect the iron. Since these reactive metal plates are oxidised quickly, they are replaced from time to time which is easy to do.



ANSWER KEY

BEGINNER'S BOX-1	Que.	1	2	3	4	5					
	Ans.	1	2	4	3	4					
BEGINNER'S BOX-2	Que.	1	2	3	4	5	6				
	Ans.	3	2	1	1	3	3				
BEGINNER'S BOX-3	Que.	1	2	3	4	5	6				
	Ans.	4	2	2	4	2	3				

EXERCISE-I (Conceptual Questions)

Build Up Your Understanding

ELECTROLYTIC CONDUCTANCE

- Strong electrolytes are those which :
(1) dissolve readily in water
(2) conduct electricity
(3) dissociate into ions even at high concentration
(4) dissociate into ions at high dilution.
- Molten sodium chloride conducts electricity due to the presence of :
(1) free electrons
(2) free ions
(3) free molecules
(4) free atoms of Na and Cl
- Electrolytic conduction is due to the movement of :
(1) molecules (2) atoms
(3) ions (4) electrons
- Which of the following solutions of KCl has the lowest value of equivalent conductance ?
(1) 1 M (2) 0.1 M
(3) .01 M (4) .001 M
- In the equation $\Lambda = \text{sp. cond.} \times V$, If V is the volume in cc containing 1 equivalent of the electrolyte then V for a $\frac{N}{10}$ solution will be :
(1) 10 c.c. (2) 100 c.c.
(3) 1000 c.c. (4) 10,000 c.c.
- If the specific resistance of a solution of concentration C g eq L^{-1} is R , then its equivalent conductance is :
(1) $\frac{100R}{C}$ (2) $\frac{RC}{1000}$
(3) $\frac{1000}{RC}$ (4) $\frac{C}{1000R}$
- The specific conductances in $\text{ohm}^{-1} \text{cm}^{-1}$ of four electrolytes P, Q, R and S are given in brackets :
P (5.0×10^{-5}) Q (7.0×10^{-8})
R (1.0×10^{-10}) S (9.2×10^{-3})
The one that offers highest resistance to the passage of electric current is
(1) P (2) S
(3) R (4) Q
- The specific conductance of a salt of 0.01 M concentration is $1.061 \times 10^{-4} \text{ S cm}^{-1}$. Molar conductance of the same solution will be :
(1) 1.061×10^{-4} (2) 1.061
(3) 10.61 (4) 106.1

- Which of the following solutions of NaCl will have the highest specific conductance ?
(1) 0.001 N (2) 0.1 N
(3) 0.01 N (4) 1.0 N
- The value of molar conductivity of HCl is greater than that of NaCl at a particular temperature because :
(1) Molecular mass of HCl is less than that of NaCl.
(2) Velocity of H^+ ions is more than that of Na^+ ions
(3) HCl is strongly acidic
(4) Ionisation of HCl is larger than that of NaCl
- Which statement is not correct :—
(1) Conductance of an electrolytic solution increases with dilution
(2) Conductance of an electrolytic solution decreases with dilution
(3) Specific conductance of an electrolytic solution decreases with dilution
(4) Equivalent conductance of an electrolytic solution increases with dilution.
- The resistance of 0.01 N solution of an electrolyte was found to be 210 ohm at 298 K using a conductivity cell of cell constant 0.66 cm^{-1} . The equivalent conductance of solution is :—
(1) $314.28 \text{ mho cm}^2 \text{ eq}^{-1}$
(2) $3.14 \text{ mho cm}^2 \text{ eq}^{-1}$
(3) $314.28 \text{ mho}^{-1} \text{ cm}^2 \text{ eq}^{-1}$
(4) $3.14 \text{ mho}^{-1} \text{ cm}^2 \text{ eq}^{-1}$
- Electrolytic conduction differs from metallic conduction from the fact that in the former
(1) The resistance increases with increasing temperature
(2) The resistance decreases with increasing temperature
(3) The resistance remains constant with increasing temperature
(4) The resistance is independent of the length of the conductor
- The specific conductance of a 0.01 M solution of KCl is $0.0014 \text{ ohm}^{-1} \text{cm}^{-1}$ at 25°C . Its equivalent conductance ($\text{cm}^2 \text{ ohm}^{-1} \text{eq}^{-1}$) is :—
(1) 140 (2) 14
(3) 1.4 (4) 0.14

15. Which one of the following is wrong :-
 (1) Specific conductance increases on dilution.
 (2) Specific conductance decreases on dilution.
 (3) Equivalent conductance increases on dilution.
 (4) Molar conductance increases on dilution.

KOHLRAUSCH LAW

16. At infinite dilution, the equivalent conductances of CH_3COONa , HCl and CH_3COOH are 91, 426 and 391 $\text{mho cm}^2 \text{eq}^{-1}$ respectively at 25 °C. The eq. conductance of NaCl at infinite dilution will be :
 (1) 126 (2) 209
 (3) 391 (4) 908
17. The equivalent conductivity of 0.1 N CH_3COOH at 25 °C is 80 and at infinite dilution it is 400. The degree of dissociation of CH_3COOH is :
 (1) 1 (2) 0.2
 (3) 0.1 (4) 0.5
18. The limiting molar conductivities Λ^0 for NaCl , KBr and KCl are 126, 152 and 150 $\text{S cm}^2 \text{mol}^{-1}$ respectively. The Λ^0 for NaBr is :
 (1) 278 $\text{S cm}^2 \text{mol}^{-1}$ (2) 176 $\text{S cm}^2 \text{mol}^{-1}$
 (3) 128 $\text{S cm}^2 \text{mol}^{-1}$ (4) 302 $\text{S cm}^2 \text{mol}^{-1}$
19. For HCl solution at 25 °C, equivalent conductance at infinite dilution, is 425 $\text{ohm}^{-1} \text{cm}^2 \text{eq}^{-1}$. The specific conductance of a solution of HCl is 3.825 $\text{ohm}^{-1} \text{cm}^{-1}$. If the degree of dissociation is 90%. The normality of the solution is :-
 (1) 0.90 N (2) 1.0 N
 (3) 10 N (4) 1.2 N
20. The molar conductivities Λ_{NaOAc}^0 and Λ_{HCl}^0 at infinite dilution in water at 25°C are 91.0 and 426.2 $\text{S cm}^2 \text{mol}^{-1}$ respectively. To calculate Λ_{HOAc}^0 the additional value required is :
 (1) Λ_{NaCl}^0 (2) $\Lambda_{\text{H}_2\text{O}}^0$
 (3) Λ_{KCl}^0 (4) Λ_{NaOH}^0
21. The molar conductance at infinite dilution of AgNO_3 , AgCl and NaCl are 116.5, 121.6 and 110.3 $\text{S cm}^2 \text{mol}^{-1}$ respectively. The molar conductance of NaNO_3 is :
 (1) 111.4 $\text{S cm}^2 \text{mol}^{-1}$ (2) 105.2 $\text{S cm}^2 \text{mol}^{-1}$
 (3) 130.6 $\text{S cm}^2 \text{mol}^{-1}$ (4) 150.2 $\text{S cm}^2 \text{mol}^{-1}$
22. The conductivity of a saturated solution of BaSO_4 is $3.06 \times 10^{-6} \text{ohm}^{-1} \text{cm}^{-1}$ and its molar conductance is 1.53 $\text{ohm}^{-1} \text{cm}^{-1} \text{mol}^{-1}$. The K_{sp} of BaSO_4 will be
 (1) 4×10^{-12} (2) 2.5×10^{-9}
 (3) 2.5×10^{-13} (4) 4×10^{-6}

GALVANIC CELL

23. In the galvanic cell
 $\text{Cu(s)} \mid \text{Cu}^{2+} (1 \text{ M}) \parallel \text{Ag}^+ (1 \text{ M}) \mid \text{Ag(s)}$
 the electrons will travel in the external circuit :
 (1) from Ag to Cu
 (2) from Cu to Ag
 (3) electrons do not travel in the external circuit
 (4) in any direction
24. The passage of electricity in the Daniell cell when Zn and Cu electrodes are connected is :
 (1) from Cu to Zn in the cell
 (2) from Cu to Zn out side the cell
 (3) from Zn to Cu outside the cell
 (4) in any direction in the cell
25. The equation representing the process by which standard reduction potential of zinc can be defined is
 (1) $\text{Zn}^{2+} (\text{s}) + 2\text{e}^- \longrightarrow \text{Zn} (\text{s})$
 (2) $\text{Zn} (\text{g}) \longrightarrow \text{Zn}^{2+} (\text{g}) + 2\text{e}^-$
 (3) $\text{Zn}^{2+} (\text{g}) + 2\text{e}^- \longrightarrow \text{Zn} (\text{s})$
 (4) $\text{Zn}^{2+} (\text{aq.}) + 2\text{e}^- \longrightarrow \text{Zn} (\text{s})$
26. A standard hydrogen electrode has zero electrode potential because :
 (1) Hydrogen is easiest to oxidize.
 (2) This electrode potential is assumed to be zero.
 (3) Hydrogen atom has only one electron.
 (4) Hydrogen is the lightest element.
27. Which is not true for a standard hydrogen electrode ?
 (1) The hydrogen ion concentration is 1 M.
 (2) Temperature is 25°C.
 (3) Pressure of hydrogen is 1 bar.
 (4) It contains a metallic conductor which does not adsorb hydrogen.
28. E° for the half cell
 $\text{Zn}^{2+} \mid \text{Zn}$ is -0.76 V. E.m.f. of the cell
 $\text{Zn} \mid \text{Zn}^{2+} (1\text{M}) \parallel 2\text{H}^+ (1\text{M}) \mid \text{H}_2 (1 \text{ atm})$ is :
 (1) -0.76 V (2) +0.76 V
 (3) -0.38 V (4) + 0.38 V
29. The chemical reaction,
 $2\text{AgCl(s)} + \text{H}_2(\text{g}) \longrightarrow 2\text{HCl(aq)} + 2\text{Ag(s)}$
 taking place in a galvanic cell is represented by the notation.
 (1) $\text{Pt(s)} \mid \text{H}_2(\text{g}).1\text{bar} \mid 1\text{MKCl(aq)} \mid \text{AgCl(s)} \mid \text{Ag(s)}$.
 (2) $\text{Pt(s)} \mid \text{H}_2(\text{g}).1\text{bar} \mid 1\text{MHCl(aq)} \mid 1\text{M Ag(aq)} \mid \text{Ag(s)}$.
 (3) $\text{Pt(s)} \mid \text{H}_2(\text{g}).1\text{bar} \mid 1\text{MHCl(aq)} \mid \text{AgCl(s)} \mid \text{Ag(s)}$.
 (4) $\text{Pt(s)} \mid \text{H}_2(\text{g}).1\text{bar} \mid 1\text{MHCl(aq)} \mid \text{Ag(s)} \mid \text{AgCl(s)}$.

30. $\text{Cu(s)} \mid \text{Cu}^{2+}(1 \text{ M}) \parallel \text{Zn}^{2+}(1 \text{ M}) \mid \text{Zn(s)}$
A cell represented above should have emf.
(1) Positive
(2) Negative
(3) Zero
(4) Cannot be predicted
31. Given electrode potentials :
 $\text{Fe}^{3+} + e \longrightarrow \text{Fe}^{2+}$; $E^\circ = 0.771 \text{ V}$
 $\text{I}_2 + 2e \longrightarrow 2\text{I}^-$; $E^\circ = 0.536 \text{ V}$
 E° cell for the cell reaction
 $2\text{Fe}^{3+} + 2\text{I}^- \rightarrow 2\text{Fe}^{2+} + \text{I}_2$ is -
(1) $(2 \times 0.771 - 0.536) = 1.006 \text{ V}$
(2) $(0.771 - 0.5 \times 0.536) = 0.503 \text{ V}$
(3) $0.771 - 0.536 = 0.235 \text{ V}$
(4) $0.536 - 0.771 = -0.235 \text{ V}$
32. Which of the following is not an anodic reaction-
(1) $\text{Ag}^+ \rightarrow \text{Ag} - e^-$
(2) $\text{Cu} \rightarrow \text{Cu}^{2+} + 2e^-$
(3) $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + e^-$
(4) $4\text{OH}^- \rightarrow 2\text{H}_2\text{O} + \text{O}_2 + 4e^-$
33. Which of the following statements is correct :-
(1) Oxidation occur at anode in both galvanic and electrolytic cell.
(2) Reduction occurs at anode in both galvanic and electrolytic cell,
(3) Reduction occur at anode in electrolytic cell where as oxidation occur at cathode in galvanic cell,
(4) Oxidation occur at anode in electrolytic cell where as reduction occur at anode in a galvanic cell,
34. Other things being equal, the life of a Daniell cell may be increased by :-
(1) Keeping low temperature
(2) Using large copper electrode
(3) Decreasing concentration of copper ions
(4) Using large zinc electrodes

ELECTROCHEMICAL SERIES

35. Zn can not displace following ions from their aqueous solution :
(1) Ag^+ (2) Cu^{2+}
(3) Fe^{2+} (4) Na^+
36. The standard reduction potentials at 25°C for the following half reactions are given against each :
 $\text{Zn}^{2+}(\text{aq}) + 2e^- \rightleftharpoons \text{Zn(s)}$, -0.762 V
 $\text{Cr}^{3+}(\text{aq}) + 3e^- \rightleftharpoons \text{Cr(s)}$, -0.740 V
 $2\text{H}^+ + 2e^- \rightleftharpoons \text{H}_2(\text{g})$, 0.00 V
 $\text{Fe}^{3+} + 2e^- \rightleftharpoons \text{Fe}^{2+}$, 0.77 V
Which is the strongest reducing agent ?
(1) Zn (2) Cr
(3) $\text{H}_2(\text{g})$ (4) $\text{Fe}^{2+}(\text{aq})$

37. Red hot carbon will remove oxygen from the oxide XO and YO but not from ZO. Y will remove oxygen from XO. Use this evidence to deduce the order of activity of the three metals X, Y and Z putting the most active first.
(1) XYZ (2) ZYX
(3) YXZ (4) ZXY
38. Adding powdered Pb and Fe to a solution containing 1.0 M in each of Pb^{2+} and Fe^{2+} ions, would result into the formation of :-
(1) More of Fe and Pb^{2+} ions
(2) More of Fe^{2+} and Pb^{2+} ions
(3) More of Pb and Fe^{2+} ions
(4) More of Fe and Pb
39. If a spoon of copper metal is placed in a solution of ferrous sulphate :
(1) Cu will precipitate out
(2) Iron will precipitate
(3) Cu and Fe will precipitate
(4) No reaction will take place
40. Which one will liberate Br_2 from KBr ?
(1) HI (2) I_2
(3) Cl_2 (4) SO_2
41. Using the standard electrode potential values given below, decide which of the statements, I, II, III and IV are correct. Choose the right answer from (1), (2), (3) and (4).
 $\text{Fe}^{2+} + 2e^- \rightleftharpoons \text{Fe}$; $E^\circ = -0.44 \text{ V}$
 $\text{Cu}^{2+} + 2e^- \rightleftharpoons \text{Cu}$; $E^\circ = +0.34 \text{ V}$
 $\text{Ag}^+ + e^- \rightleftharpoons \text{Ag}$; $E^\circ = +0.80 \text{ V}$
I. Copper can displace iron from FeSO_4 solution.
II. Iron can displace copper from CuSO_4 solution.
III. Silver can displace copper from CuSO_4 solution.
IV. Iron can displace silver from AgNO_3 solution.
(1) I and II (2) II and III
(3) II and IV (4) I and IV
42. The standard electrode potential value of the elements A, B and C are 0.68 , -2.50 and 0.50 V respectively. The order of their reducing power is :
(1) $A > B > C$ (2) $A > C > B$
(3) $C > B > A$ (4) $B > C > A$

- 43.** The oxidation potential of Zn, Cu, Ag, H₂ and Ni are 0.76 V, -0.34 V, -0.80 V, 0 V, 0.55 V respectively. Which of the following reaction will provide maximum voltage ?
 (1) $\text{Zn} + \text{Cu}^{2+} \longrightarrow \text{Cu} + \text{Zn}^{2+}$
 (2) $\text{Zn} + 2\text{Ag}^+ \longrightarrow 2\text{Ag} + \text{Zn}^{2+}$
 (3) $\text{H}_2 + \text{Cu}^{2+} \longrightarrow 2\text{H}^+ + \text{Cu}$
 (4) $\text{H}_2 + \text{Ni}^{2+} \longrightarrow 2\text{H}^+ + \text{Ni}$
- 44.** The following facts are available :-
 $2\text{X}^- + \text{Y}_2 \rightarrow 2\text{Y}^- + \text{X}_2$
 $2\text{W}^- + \text{Y}_2 \rightarrow \text{no reaction}$
 $2\text{Z}^- + \text{X}_2 \rightarrow 2\text{X}^- + \text{Z}_2$
 Which of the following statements is correct :-
 (1) $E^\circ_{\text{W}^-/\text{W}_2} > E^\circ_{\text{Y}^-/\text{Y}_2} > E^\circ_{\text{X}^-/\text{X}_2} > E^\circ_{\text{Z}^-/\text{Z}_2}$
 (2) $E^\circ_{\text{W}^-/\text{W}_2} < E^\circ_{\text{Y}^-/\text{Y}_2} < E^\circ_{\text{X}^-/\text{X}_2} < E^\circ_{\text{Z}^-/\text{Z}_2}$
 (3) $E^\circ_{\text{W}^-/\text{W}_2} < E^\circ_{\text{Y}^-/\text{Y}_2} > E^\circ_{\text{X}^-/\text{X}_2} > E^\circ_{\text{Z}^-/\text{Z}_2}$
 (4) $E^\circ_{\text{W}^-/\text{W}_2} > E^\circ_{\text{Y}^-/\text{Y}_2} < E^\circ_{\text{X}^-/\text{X}_2} < E^\circ_{\text{Z}^-/\text{Z}_2}$
- 45.** The standard reduction potential at 25 °C of Li⁺ / Li, Ba²⁺ / Ba, Na⁺ / Na and Mg²⁺ / Mg are -3.05 V, -2.73 V, -2.71 V and -2.37 V respectively. Which one of the following is the strongest oxidising agent ?
 (1) Na⁺ (2) Li⁺
 (3) Ba²⁺ (4) Mg²⁺
- 46.** A gas X at 1 atm is bubbled through a solution containing a mixture of 1 M Y⁻ and 1 M Z⁻ at 25 °C. If the reduction potential of Z > Y > X then :
 (1) Y will oxidise X and not Z
 (2) Y will oxidise Z and not X
 (3) Y will oxidise both X and Z
 (4) Y will reduce both X and Z
- 47.** The standard electrode potential of Zn, Ag and Cu are -0.76 V, 0.80 V and 0.34 V respectively, then :
 (1) Ag can oxidise Zn and Cu
 (2) Ag can reduce Zn²⁺ and Cu²⁺
 (3) Zn can reduce Ag⁺ and Cu²⁺
 (4) Cu can oxidise Zn and Ag
- 48.** Each of the three metals x, y and z were put in turn into aqueous solution of the other two. x + salt of y (or z) → y (or z) + salt of x
 Which one of the following observation is incorrect?
 (1) y + salt of x → no action observed
 (2) y + salt of z → z + salt of y
 (3) z + salt of x → x + salt of z
 (4) z + salt of y → no action observed
- 49.** A standard reduction electrode potentials of four elements are
 A = - 0.250 V , B = - 0.140 V
 C = - 0.126 V , D = - 0.402 V
 The element that displaces A from its compounds aqueous solution is :-
 (1) B
 (2) C
 (3) D
 (4) None of the above
- 50.** The following four colourless salt solutions are placed in separate test tubes and a strip of Cu is placed in each solution which finally turns blue :-
 (1) Zn(NO₃)₂ (2) Mg(NO₃)₂
 (3) KNO₃ (4) AgNO₃
- 51.** Which of the following displacement does not occur
 (1) $\text{Zn} + 2\text{H}^+ \rightarrow \text{Zn}^{2+} + \text{H}_2 \uparrow$
 (2) $\text{Fe} + 2\text{Ag}^+ \rightarrow \text{Fe}^{2+} + \text{Ag} \downarrow$
 (3) $\text{Cu} + \text{Fe}^{2+} \rightarrow \text{Cu}^{2+} + \text{Fe} \downarrow$
 (4) $\text{Zn} + \text{Pb}^{2+} \rightarrow \text{Zn}^{2+} + \text{Pb} \downarrow$

NERNST EQUATION

- 52.** $E^\circ(\text{Ni}^{2+}/\text{Ni}) = -0.25 \text{ V}$
 $E^\circ(\text{Au}^{3+}/\text{Au}) = 1.50 \text{ V}$
 The emf of the voltaic cell.
 $\text{Ni} / \text{Ni}^{2+} (1.0 \text{ M}) \parallel \text{Au}^{3+} (1.0 \text{ M}) \mid \text{Au}$ is :
 (1) 1.25 V (2) -1.75 V
 (3) 1.75 V (4) 4.0 V
- 53.** The emf of the cell in which the following reaction
 $\text{Zn(s)} + \text{Ni}^{2+} (a = 0.1) \rightleftharpoons \text{Zn}^{2+} (a = 1.0) + \text{Ni(s)}$
 occurs, is found to be 0.5105 V at 298 K. The standard e.m.f. of the cell is :-
 (1) -0.5105 V (2) 0.5400 V
 (3) 0.4810 V (4) 0.5696 V
- 54.** The emf of the cell
 $\text{Ti(s)} \mid \text{Ti}^+ (0.0001 \text{ M}) \parallel \text{Cu}^{2+} (0.01\text{M}) \mid \text{Cu(s)}$ is 0.83 V
 The emf of this cell will be increased by :-
 (1) Increasing the concentration of Cu²⁺ ions
 (2) Decreasing the concentration of Ti⁺
 (3) Increasing the concentration of both
 (4) (1) & (2) both
- 55.** The potential of hydrogen electrode
 ($P_{\text{H}_2} = 1 \text{ atm}$; $C_{\text{H}^+} = 0.1 \text{ M}$) at 25°C will be -
 (1) 0.00 V (2) -0.059 V
 (3) 0.118 V (4) 0.059 V

56. Which of the following represents the potential of silver wire dipped into 0.1 M AgNO_3 solution at 25°C ?
 (1) E°_{red} (2) $(E^\circ_{\text{red}} + 0.059)$
 (3) $(E^\circ_{\text{ox}} - 0.059)$ (4) $(E^\circ_{\text{red}} - 0.059)$
57. The potential of a hydrogen electrode at $\text{pH} = 1$ is
 (1) 0.059 V (2) 0.00 V
 (3) -0.059 V (4) 0.59 V
58. Consider the reaction
 $\text{Cl}_2(\text{g}) + 2\text{Br}^-(\text{aq}) \longrightarrow 2\text{Cl}^-(\text{aq}) + \text{Br}_2$
 The emf of the cell when
 $[\text{Cl}^-] = [\text{Br}_2] = [\text{Br}^-] = 0.01 \text{ M}$ and Cl_2 gas at 1 atm pressure will be (E° for the above reaction is = 0.29 V)
 (1) 0.54 V (2) 0.35 V
 (3) 0.24 V (4) -0.29 V
59. The standard emf for the cell reaction
 $\text{Zn} + \text{Cu}^{2+} \longrightarrow \text{Zn}^{2+} + \text{Cu}$ is 1.10 V at 25°C . The emf for the cell reaction when 0.1 M Cu^{2+} and 0.1 M Zn^{2+} solution are used at 25°C is :
 (1) 1.10 V (2) 0.110 V
 (3) -1.10 V (4) -0.110 V
60. E° for $\text{F}_2 + 2\text{e}^- \rightarrow 2\text{F}^-$ is 2.8 V,
 E° for $\frac{1}{2}\text{F}_2 + \text{e}^- \rightarrow \text{F}^-$ is ?
 (1) 2.8 V (2) 1.4 V (3) -2.8 V (4) -1.4 V
61. ΔG° of the cell reaction
 $\text{AgCl}(\text{s}) + \frac{1}{2}\text{H}_2(\text{g}) \rightarrow \text{Ag}(\text{s}) + \text{H}^+ + \text{Cl}^-$ is -21.52 kJ
 ΔG° of $2\text{AgCl}(\text{s}) + \text{H}_2(\text{g}) \rightarrow 2\text{Ag}(\text{s}) + 2\text{H}^+ + 2\text{Cl}^-$ is :
 (1) -21.52 kJ (2) -10.76 kJ
 (3) -43.04 kJ (4) 43.04 kJ
62. How much will the potential of $\text{Zn} / \text{Zn}^{2+}$ change if the solution of Zn^{2+} is diluted 10 times
 (1) increases by 0.03 V
 (2) decreases by 0.03 V
 (3) increases by 0.059 V
 (4) decreases by 0.059 V
63. How much will the potential of a hydrogen electrode change when its solution initially at $\text{pH} = 0$ is neutralised to $\text{pH} = 7$?
 (1) increases by 0.059 V
 (2) decreases by 0.059 V
 (3) increases by 0.41 V
 (4) decreases by 0.41 V
64. Which of the following will increase the voltage of the cell with following cell reaction
 $\text{Sn}_{(\text{s})} + 2\text{Ag}^+_{(\text{aq})} \rightarrow \text{Sn}^{2+}_{(\text{aq})} + 2\text{Ag}_{(\text{s})}$
 (1) Increase in the size of silver rod
 (2) Increase in the concentration of Sn^{2+} ions
 (3) Increase in the concentration of Ag^+ ions
 (4) Decrease in the concentration of Ag^+ ions
65. E° for the reaction $\text{Fe} + \text{Zn}^{2+} \rightarrow \text{Zn} + \text{Fe}^{2+}$ is -0.35 V. The given cell reaction is :
 (1) feasible (2) not feasible
 (3) in equilibrium (4) can't say anything
66. For a reaction - $\text{A}(\text{s}) + 2\text{B}^+ \rightarrow \text{A}^{2+} + 2\text{B}(\text{s})$
 K_c has been found to be 10^{12} . The E°_{cell} is:
 (1) 0.354 V (2) 0.708 V
 (3) 0.0098 V (4) 1.36 V
67. The standard electrode potential (E°) for OCl^-/Cl^- and $\text{Cl}^- / \frac{1}{2} \text{Cl}_2$ respectively are 0.94 V and -1.36 V. The E° value of $\text{OCl}^- / \frac{1}{2} \text{Cl}_2$ will be :
 (1) -2.20 V (2) -0.42 V
 (3) 0.52 V (4) 1.04 V
68. The standard reduction potentials of Cu^{2+}/Cu and $\text{Cu}^{2+}/\text{Cu}^+$ are 0.337 V and 0.153 V respectively. The standard electrode potential of Cu^+/Cu half cell is :
 (1) 0.184 V (2) 0.827 V
 (3) 0.521 V (4) 0.490 V
69. The hydrogen electrode is dipped in a solution of $\text{pH} = 3$ at 25°C . The potential of the cell would be :
 (1) 0.177 V (2) - 0.177 V
 (3) 0.087 V (4) 0.059 V
70. What is the potential of the cell containing two hydrogen electrodes as represented below
 $\text{Pt}; \text{H}_2(\text{g}) | \text{H}^+(10^{-8}\text{M}) || \text{H}^+(0.001 \text{ M}) | \text{H}_2(\text{g}); \text{Pt}$;
 (1) - 0.295 V (2) - 0.0591 V
 (3) 0.295 V (4) 0.0591 V
71. Consider the cell $\text{Cu}/\text{Cu}^{2+} || \text{Ag}^+/\text{Ag}$. If the concentration of Cu^{2+} and Ag^+ ions becomes ten times then the emf of the cell will :-
 (1) Becomes 10 times
 (2) Remains same
 (3) Increases by 0.0295 V
 (4) Decreases by 0.0295 V

72. The emf of the cell
 $\text{Ni} | \text{Ni}^{+2} (1.0 \text{ M}) || \text{Au}^{+3} (0.1 \text{ M}) | \text{Au}$
 $[E^\circ \text{ for } \text{Ni}^{+2}/\text{Ni} = -0.25 \text{ V}, E^\circ \text{ for } \text{Au}^{+3}/\text{Au} = 1.50 \text{ V}]$
 is given as:-
 (1) 1.25 V (2) -1.75 V
 (3) 1.75 V (4) 1.73 V

ELECTROLYSIS

73. When an electric current is passed through acidified water, 112 mL of hydrogen gas at STP collects at the cathode in 965 s. The current passed, in ampere is :
 (1) 1.0 (2) 0.5
 (3) 0.1 (4) 2.0
74. Two electrolytic cells one containing acidified ferrous chloride and another acidified ferric chloride are connected in series. The ratio of iron deposited at cathodes in the two cells when electricity is passed through the cells will be :
 (1) 3 : 1 (2) 2 : 1 (3) 1 : 1 (4) 3 : 2
75. A current of 9.65 A flowing for 10 minute deposits 3.0 g of a metal. The equivalent weight of the metal is :
 (1) 10 (2) 30 (3) 50 (4) 96.5
76. The same amount of electricity was passed through two separate electrolytic cells containing solutions of nickel nitrate $[\text{Ni}(\text{NO}_3)_2]$ and chromium nitrate $[\text{Cr}(\text{NO}_3)_3]$ respectively. If 0.3 g of nickel was deposited in the first cell, the amount of chromium deposited in the other cell is :-
 (at. wt. of Ni = 59, at. wt. of Cr = 52)
 (1) 0.1 g (2) 0.17 g
 (3) 0.3 g (4) 0.6 g
77. How many coulombs of electricity are required for the oxidation of 1 mol of H_2O to O_2 ?
 (1) $9.65 \times 10^4 \text{ C}$ (2) $4.825 \times 10^5 \text{ C}$
 (3) $1.93 \times 10^5 \text{ C}$ (4) $1.93 \times 10^4 \text{ C}$
78. When a quantity of electricity equal to that required to liberate 2.24 L of hydrogen at STP from 0.1 M aqueous H_2SO_4 is passed (At. mass of Cu = 63.5) then the mass of copper that will be deposited at cathode in electrolysis of 0.2 M solution of copper sulphate will be :
 (1) 1.59 g (2) 3.18 g
 (3) 6.35 g (4) 12.70 g
79. 10800 C of electricity through the electrolyte deposited 2.977 g of metal with atomic mass 106.4 g mol⁻¹. The charge on the metal cation is -
 (1) +4 (2) +3 (3) +2 (4) +1
80. On passing electricity through dilute H_2SO_4 solution the amount of substance liberated at the cathode and anode are in the ratio :
 (1) 1 : 8 (2) 8 : 1
 (3) 16 : 1 (4) 1 : 16
81. During electrolysis of fused calcium hydride, the hydrogen is produced at :
 (1) Cathode
 (2) Anode
 (3) Hydrogen is not liberated at all
 (4) H_2 produced reacts with oxygen to form water
82. A silver cup is plated with silver by passing 965 A current for one second, the mass of Ag deposited is :-(At. wt. of Ag = 107.87)
 (1) 9.89 g. (2) 107.87 g.
 (3) 1.0787 g. (4) 100.2 g.
83. When electricity is passed through a solution of AlCl_3 , 13.5 g Al is deposited. The number of Faradays must be :-
 (1) 5.0 (2) 1.0 (3) 1.5 (4) 3.0
84. A solution of sodium sulphate in water is electrolysed using inert electrodes. The product at the cathode and anode are respectively :-
 (1) H_2 , SO_2 (2) O_2 , H_2
 (3) O_2 , Na (4) H_2 , O_2
85. One Faraday of electricity will liberate one mole of the metal from the solution of
 (1) Auric chloride (2) Silver nitrate
 (3) Calcium chloride (4) Copper sulphate
86. When 96500 C of electricity are passed through barium chloride solution, the amount of barium deposited will be :-
 (1) 0.5 mol (2) 1.0 mol
 (3) 1.5 mol (4) 2.0 mol
87. A factory produces 40 kg of calcium in two hours by electrolysis. How much aluminium can be produced by the same current in two hours :-
 (At wt. of Ca = 40, Al = 27)
 (1) 22 kg (2) 18 kg
 (3) 9 kg (4) 27 kg

88. What would be the ratio of moles each of Ag^+ , Cu^{+2} , Fe^{+3} ions would be deposited by passage of same quantity of electricity through solutions of their salts :-

(1) 1 : 1 : 1

(2) $1 : \frac{1}{2} : \frac{1}{3}$

(3) $\frac{1}{3} : \frac{1}{2} : 1$

(4) 1 : 2 : 3

89. Electrolysis of aq. CuSO_4 causes :-

(1) An increase in pH

(2) A decrease in pH

(3) Either decrease or increase

(4) None

90. The passage of current liberates H_2 at cathode and Cl_2 at anode the solution is :-

(1) CuSO_4 (aq)

(2) CuCl_2 (aq.)

(3) NaCl (aq.)

(4) Water

COMMERCIAL CELLS

91. When lead accumulator is charged it is :

(1) an electrolytic cell

(2) a galvanic cell

(3) a Daniel cell

(4) none of the above

92. When a lead storage battery is charged :

(1) PbO_2 dissolves

(2) The lead electrode becomes coated with lead sulphate

(3) Sulphuric acid is regenerated

(4) The amount of acid decreases

EXERCISE-I (Conceptual Questions)

ANSWER KEY

Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	3	2	3	1	4	3	3	3	4	2	2	1	2	1	1
Que.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Ans.	1	2	3	3	1	2	4	2	2	4	2	4	2	3	2
Que.	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45
Ans.	3	1	1	4	4	1	2	3	4	3	3	4	2	2	4
Que.	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Ans.	1	3	3	3	4	3	3	2	4	2	4	3	2	1	1
Que.	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75
Ans.	3	1	4	3	2	1	3	3	2	3	3	4	1	4	3
Que.	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90
Ans.	2	3	3	1	1	2	3	3	4	2	1	2	2	2	3
Que.	91	92													
Ans.	1	3													

EXERCISE-II (Previous Year Questions)

AIPMT/NEET & AIIMS (2006-2018)

AIPMT 2006

1. If $E_{\text{Fe}^{+2}/\text{Fe}}^0 = -0.441\text{V}$ and $E_{\text{Fe}^{+3}/\text{Fe}^{+2}}^0 = 0.771\text{V}$ the standard EMF of the reaction $\text{Fe} + 2\text{Fe}^{+3} \rightarrow 3\text{Fe}^{+2}$ will be :
- (1) 0.330 V
 - (2) 1.653 V
 - (3) 1.212 V
 - (4) 0.111 V

2. A hypothetical electrochemical cell is shown



The emf measured is +0.20 V

The cell reaction is :

- (1) $\text{A}^+ + \text{B} \longrightarrow \text{A} + \text{B}^+$
- (2) $\text{A}^+ + \text{e}^- \longrightarrow \text{A}$, $\text{B}^+ + \text{e}^- \longrightarrow \text{B}$
- (3) The cell reaction cannot be predicted.
- (4) $\text{A} + \text{B}^+ \longrightarrow \text{A}^+ + \text{B}$

AIIMS 2006

3. The charge required for the reduction of 1 mol of MnO_4^- to MnO_2 is :
- (1) 1F
 - (2) 3F
 - (3) 5F
 - (4) 7F
4. The products formed when an aqueous solution of NaBr is electrolyzed in a cell having inert electrodes are :-
- (1) Na and Br_2
 - (2) Na and O_2
 - (3) H_2 , Br_2 and NaOH
 - (4) H_2 and O_2

AIPMT 2007

5. The equilibrium constant of the reaction : $\text{Cu(s)} + 2\text{Ag}^+(\text{aq.}) \longrightarrow \text{Cu}^{2+}(\text{aq.}) + 2\text{Ag(s)}$ $E^\circ = 0.46\text{V}$ at 298 K is :-
- (1) 2.0×10^{10}
 - (2) 4.0×10^{10}
 - (3) 4.0×10^{15}
 - (4) 2.4×10^{10}

AIPMT 2008

6. On the basis of the following E° values, the strongest oxidizing agent is :-
- $$[\text{Fe}(\text{CN})_6]^{4-} \rightarrow [\text{Fe}(\text{CN})_6]^{3-} + \text{e}^-, E^\circ = -0.35\text{V}$$
- $$\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-, E^\circ = -0.77\text{V}$$
- (1) Fe^{3+}
 - (2) $[\text{Fe}(\text{CN})_6]^{3-}$
 - (3) $[\text{Fe}(\text{CN})_6]^{4-}$
 - (4) Fe^{2+}

7. Kohlrausch's law states that at :-
- (1) Infinite dilution, each ion makes definite contribution to conductance of an electrolyte whatever be the nature of the other ion of the electrolyte.
 - (2) Infinite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte whatever be the nature of the other ion of the electrolyte.
 - (3) Finite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte whatever be the nature of the other ion of the electrolyte.
 - (4) Infinite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte depending on the nature of the other ion of electrolyte.

AIPMT 2009

8. Given :
- (i) $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}$, $E^\circ = 0.337\text{V}$
 - (ii) $\text{Cu}^{2+} + \text{e}^- \rightarrow \text{Cu}^+$, $E^\circ = 0.153\text{V}$
- Electrode potential, E° for the reaction, $\text{Cu}^+ + \text{e}^- \rightarrow \text{Cu}$, will be :-
- (1) 0.38 V
 - (2) 0.52 V
 - (3) 0.90 V
 - (4) 0.30 V
9. The equivalent conductance of $\frac{M}{32}$ solution of a weak monobasic acid is $8.0\text{ mho cm}^2 \text{eq}^{-1}$ and at infinite dilution is $400\text{ mho cm}^2 \text{eq}^{-1}$. The dissociation constant of this acid is :-
- (1) 1.25×10^{-4}
 - (2) 1.25×10^{-5}
 - (3) 1.25×10^{-6}
 - (4) 6.25×10^{-4}
10. Al_2O_3 is reduced by electrolysis at low potential and high current. If $4.0 \times 10^4\text{A}$ of current is passed through molten Al_2O_3 for 6 hours, what mass of aluminium is produced ? (Assume 100% current efficiency, At. mass of Al = 27 g mol^{-1})
- (1) $1.3 \times 10^4\text{g}$
 - (2) $9.0 \times 10^3\text{g}$
 - (3) $8.1 \times 10^4\text{g}$
 - (4) $2.4 \times 10^5\text{g}$

AIPMT 2010

11. An increase in equivalent conductance of a strong electrolyte with dilution is mainly due to:-
- (1) Increase in number of ions.
 - (2) Increase in ionic mobility of ions.
 - (3) 100% ionisation of electrolyte at normal dilution.
 - (4) Increase in both i.e. number of ions and ionic mobility of ions.

12. Consider the following relations for emf of a electrochemical cell :
- emf of cell = (Oxidation potential of anode) – (Reduction potential of cathode)
 - emf of cell = (Oxidation potential of anode) + (Reduction potential of cathode)
 - emf of cell = (Reduction potential of anode) + (Reduction potential of cathode)
 - emf of cell = (Oxidation potential of anode) – (Oxidation potential of cathode)

Which of the above relations are correct :

- (a) and (b)
 - (c) and (d)
 - (b) and (d)
 - (c) and (a)
13. Which of the following expressions correctly represents the equivalent conductance at infinite dilution of $\text{Al}_2(\text{SO}_4)_3$. Given that $\Lambda^\circ_{\text{Al}^{3+}}$ and $\Lambda^\circ_{\text{SO}_4^{2-}}$ are the equivalent conductances at infinite dilution of the respective ions :-

- $\Lambda^\circ_{\text{Al}^{3+}} + \Lambda^\circ_{\text{SO}_4^{2-}}$
- $(\Lambda^\circ_{\text{Al}^{3+}} + \Lambda^\circ_{\text{SO}_4^{2-}}) \times 6$
- $\frac{1}{3} \Lambda^\circ_{\text{Al}^{3+}} + \frac{1}{2} \Lambda^\circ_{\text{SO}_4^{2-}}$
- $2\Lambda^\circ_{\text{Al}^{3+}} + 3\Lambda^\circ_{\text{SO}_4^{2-}}$

14. For the reduction of silver ions with copper metals, the standard cell potential was found to be +0.46 V at 25 °C. The value of standard Gibbs energy, ΔG° will be ($F = 96500 \text{ C mol}^{-1}$)
- 98.0 kJ
 - 89.0 kJ
 - 89.0 J
 - 44.5 kJ

AIIMS 2010



15. Limiting molar conductance of H^+ and CH_3COO^- ions are $344 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ and $40 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ respectively. Molar conductance of 0.008 M CH_3COOH is 48. What will be the value of K_a for CH_3COOH :-
- 1.4×10^{-5}
 - 1.2×10^{-5}
 - 1.4×10^{-4}
 - 1×10^{-5}

16. E value for the cell reaction $\text{Cu} | \text{Cu}^{2+} (0.001 \text{ M}) || \text{Cu}^{2+} (0.1 \text{ M}) | \text{Cu}$ is :-
- $-\frac{RT}{F} \ln(0.01)$
 - $-\frac{RT}{F} \ln(0.1)$
 - $+\frac{RT}{F} \ln(0.01)$
 - $+\frac{RT}{F} \ln(0.1)$

AIPMT Pre. 2011

17. Standard electrode potential of three metals X, Y and Z are -1.2 V, +0.5 V and -3.0 V respectively. The reducing power of these metals will be :-
- $Y > Z > X$
 - $Y > X > Z$
 - $Z > X > Y$
 - $X > Y > Z$

18. The electrode potentials for



are +0.15 V and +0.50 V respectively. The value of $E^\circ_{\text{Cu}^{2+}/\text{Cu}}$ will be :-

- 0.500 V
- 0.325 V
- 0.650 V
- 0.150 V

19. Standard electrode potential for $\text{Sn}^{4+}/\text{Sn}^{2+}$ couple is +0.15 V and that for the Cr^{3+}/Cr couples is -0.74 V. These two couples in their standard state are connected to make a cell. The cell potential will be :-

- +1.19 V
- +0.89 V
- +0.18 V
- +1.83 V

20. If the E°_{cell} for a given reaction has a negative value, then which of the following gives the correct relationship for the values of ΔG° and K_{eq} ?

- $\Delta G^\circ > 0$; $K_{\text{eq}} > 1$
- $\Delta G^\circ < 0$; $K_{\text{eq}} > 1$
- $\Delta G^\circ < 0$; $K_{\text{eq}} < 1$
- $\Delta G^\circ > 0$; $K_{\text{eq}} < 1$

AIPMT Mains 2011

21. A solution contains Fe^{2+} , Fe^{3+} and I^- ions. This solution was treated with iodine at 35 °C. E° for $\text{Fe}^{3+} / \text{Fe}^{2+}$ is +0.77 V and E° for $\text{I}_2 / 2\text{I}^- = 0.536 \text{ V}$. The favourable redox reaction is :-

- Fe^{2+} will be oxidised to Fe^{3+}
- I_2 will be reduced to I^-
- There will be no redox reaction
- I^- will be oxidised to I_2

AIPMT Pre. 2012

22. Limiting molar conductivity of NH_4OH

(i.e. $\Lambda^\circ_{\text{m}(\text{NH}_4\text{OH})}$) is equal to:-

- $\Lambda^\circ_{\text{m}(\text{NH}_4\text{OH})} + \Lambda^\circ_{\text{m}(\text{NH}_4\text{Cl})} - \Lambda^\circ_{\text{m}(\text{HCl})}$
- $\Lambda^\circ_{\text{m}(\text{NH}_4\text{Cl})} + \Lambda^\circ_{\text{m}(\text{NaOH})} - \Lambda^\circ_{\text{m}(\text{NaCl})}$
- $\Lambda^\circ_{\text{m}(\text{NH}_4\text{Cl})} + \Lambda^\circ_{\text{m}(\text{NaCl})} - \Lambda^\circ_{\text{m}(\text{NaOH})}$
- $\Lambda^\circ_{\text{m}(\text{NaOH})} + \Lambda^\circ_{\text{m}(\text{NaCl})} - \Lambda^\circ_{\text{m}(\text{NH}_4\text{Cl})}$

AIPMT Mains 2012

23. Molar conductivities (Λ_m°) at infinite dilution of NaCl, HCl and CH_3COONa are 126.4, 425.9 and $91.0 \text{ S cm}^2 \text{ mol}^{-1}$ respectively. Λ_m° for CH_3COOH will be :-
 (1) $290.8 \text{ S cm}^2 \text{ mol}^{-1}$ (2) $390.5 \text{ S cm}^2 \text{ mol}^{-1}$
 (3) $425.5 \text{ S cm}^2 \text{ mol}^{-1}$ (4) $180.5 \text{ S cm}^2 \text{ mol}^{-1}$

AIIMS 2012

24. For $\text{Zn}^{2+} | \text{Zn} \text{ E}^\circ = -0.76 \text{ V}$ then EMF of this cell $\text{Zn} | \text{Zn}^{2+}(1\text{M}) || 2\text{H}^+(1\text{M}) | \text{H}_2(1 \text{ atm})$ will be :-
 (1) -0.76 V (2) 0.76 V
 (3) 0.38 V (4) -0.38 V

NEET-UG 2013

25. At 25°C molar conductance of 0.1 molar aqueous solution of ammonium hydroxide is $9.54 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ and at infinite dilution its molar conductance is $238 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$. The degree of ionisation of ammonium hydroxide at the same concentration and temperature is :-
 (1) 40.800% (2) 2.080%
 (3) 20.800% (4) 4.008%
26. A hydrogen gas electrode is made by dipping platinum wire in a solution of HCl of $\text{pH} = 10$ and by passing hydrogen gas around the platinum wire at 1 atm pressure. The oxidation potential of electrode would be ?
 (1) 1.18 V (2) 0.059 V
 (3) 0.59 V (4) 0.118 V

27. A button cell used in watches function as following
 $\text{Zn(s)} + \text{Ag}_2\text{O(s)} + \text{H}_2\text{O(l)} \rightleftharpoons 2\text{Ag(s)} + \text{Zn}^{2+}(\text{aq}) + 2\text{OH}^-(\text{aq})$
 If half cell potentials are
 $\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn(s)}; \text{E}^\circ = -0.76 \text{ V}$
 $\text{Ag}_2\text{O(s)} + \text{H}_2\text{O(l)} + 2\text{e}^- \rightarrow 2\text{Ag(s)} + 2\text{OH}^-(\text{aq}); \text{E}^\circ = 0.34 \text{ V}$
 The cell potential will be :-
 (1) 1.34 V (2) 1.10 V
 (3) 0.42 V (4) 0.84 V

AIIMS 2013

28. Electrode potential of hydrogen electrode is 18 mV then $[\text{H}^+]$ is :-
 (1) 0.2 (2) 1 (3) 2 (4) 5

AIPMT 2014

29. When $0.1 \text{ mol } \text{MnO}_4^{2-}$ is oxidised the quantity of electricity required to completely oxidise MnO_4^{2-} to MnO_4^- is :-
 (1) 96500 C (2) $2 \times 96500 \text{ C}$
 (3) 9650 C (4) 96.50 C
30. The weight of silver (at wt. = 108) displaced by a quantity of electricity which displaces 5600 mL of O_2 at STP will be :-
 (1) 5.4 g (2) 10.8 g
 (3) 54.0 g (4) 108.0 g

AIIMS 2014

31. Amount of charge is required to convert $17 \text{ g } \text{H}_2\text{O}_2$ into O_2 :-
 (1) 1F (2) 2F
 (3) 6F (4) None of these

AIPMT 2015

32. A device that converts energy of combustion of fuels like hydrogen and methane, directly into electrical energy is known as :-
 (1) Electrolytic cell (2) Dynamo
 (3) Ni-Cd cell (4) Fuel Cell

AIIMS 2015

33. For the cell reaction
 $\text{Mg} + 2\text{Ag}^+(0.0001\text{M}) \rightarrow \text{Mg}^{2+}(0.01\text{M}) + 2\text{Ag}$
 $\text{E}_{\text{cell}}^\circ = 3.177 \text{ V}$ then what will be the value of E_{cell} for the above reaction ?
 (1) 2.5 (2) 3.0 (3) 3.2 (4) 3.8
34. If electrode potential of $0.01 \text{ M } \text{Cu}^{2+} / \text{Cu}$ is 0.32 V then what will be its $\text{E}_{\text{RP}}^\circ$ -
 (1) 0.18 V (2) 0.28 V
 (3) 0.74 V (4) 0.38 V

NEET-I 2016

35. The pressure of H_2 required to make the potential of H_2 -electrode zero in pure water at 298 K is :-
 (1) 10^{-14} atm (2) 10^{-12} atm
 (3) 10^{-10} atm (4) 10^{-4} atm

NEET-II 2016

36. The molar conductivity of a 0.5 mol/dm^3 solution of AgNO_3 with electrolytic conductivity of $5.76 \times 10^{-3} \text{ S cm}^{-1}$ at 298 K is
 (1) $0.086 \text{ S cm}^2/\text{mol}$ (2) $28.8 \text{ S cm}^2/\text{mol}$
 (3) $2.88 \text{ S cm}^2/\text{mol}$ (4) $11.52 \text{ S cm}^2/\text{mol}$

37. During the electrolysis of molten sodium chloride, the time required to produce 0.10 mol of chlorine gas using a current of 3 A is
 (1) 220 minutes (2) 330 minutes
 (3) 55 minutes (4) 110 minutes

38. If the E_{cell}° for a given reaction has a negative value, which of the following gives the **correct** relationships for the values of ΔG° and K_{eq} ?
 (1) $\Delta G^{\circ} < 0$; $K_{\text{eq}} > 1$ (2) $\Delta G^{\circ} < 0$; $K_{\text{eq}} < 1$
 (3) $\Delta G^{\circ} > 0$; $K_{\text{eq}} < 1$ (4) $\Delta G^{\circ} > 0$; $K_{\text{eq}} > 1$

39. The number of electrons delivered at the cathode during electrolysis by a current of 1 A in 60 s is (charge on electron = 1.60×10^{-19} C)
 (1) 3.75×10^{20} (2) 7.48×10^{23}
 (3) 6×10^{23} (4) 6×10^{20}

AIIMS 2016

40. How many faraday charge is required for $\text{KMnO}_4/\text{Mn}^{2+}$:-
 (1) 3F (2) 2F (3) 5F (4) 1F

NEET(UG) 2017

41. In the electrochemical cell :-
 $\text{Zn} | \text{ZnSO}_4(0.01\text{M}) || \text{CuSO}_4(1.0\text{M}) | \text{Cu}$, the emf of this Daniel cell is E_1 . When the concentration of ZnSO_4 is changed to 1.0M and that of CuSO_4 changed to 0.01M, the emf changes to E_2 . Which one of the relationship is correct between E_1 and E_2 ?

(Given, $\frac{RT}{F} = 0.059$)

- (1) $E_1 < E_2$ (2) $E_1 > E_2$
 (3) $E_2 = 0 \neq E_1$ (4) $E_1 = E_2$

AIIMS 2017


42. If limiting molar conductances of Ca^{+2} and Cl^- ions are 119 and $76.3 \text{ Scm}^2 \text{ mol}^{-1}$ respectively. The limiting molar conductivity of CaCl_2 is :-
 (1) $347 \text{ S cm}^2 \text{ mol}^{-1}$ (2) $271.6 \text{ S cm}^2 \text{ mol}^{-1}$
 (3) $42.7 \text{ S cm}^2 \text{ mol}^{-1}$ (4) $195.3 \text{ S cm}^2 \text{ mol}^{-1}$

EXERCISE-II (Previous Year Questions)

Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	3	4	2	3	3	1	2	2	2	3	2	3	1	2	3
Que.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Ans.	2	3	2	2	4	4	2	2	2	4	3	2	3	3	4
Que.	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45
Ans.	1	4	2	4	1	4	4	3	1	3	2	2	4	2	2
Que.	46	47													
Ans.	4	3													

NEET(UG) 2018

43. Consider the change in oxidation state of Bromine corresponding to different EMF values as shown in the diagram below:




Then the species undergoing disproportionation is:-

- (1) BrO_3^- (2) BrO_4^-
 (3) Br_2 (4) HBrO

AIIMS 2018

44. For given reaction



If K_{eq} of reaction is $10^{15.2}$

and $E^{\circ}(\text{A}^{+2}/\text{A}) = 0.34\text{V}$


Then calculate $E^{\circ}(\text{B}^{+}/\text{B})$ will be :-

- (1) - 0.80 (2) 0.80
 (3) 1.27 (4) 1.60

45. For an electrolytic solution having concentration 0.1M and resistance is $1\text{K}\Omega$. If distance between electrode is 100 cm and area is 1 cm^2 then calculate λ_m in $\text{S cm}^2 \text{ mol}^{-1}$:-

- (1) 100 (2) 1000 (3) 500 (4) 50

46. What will be value of E_{cell} for reaction



Given E°_{cell} is 3.17 V

- (1) 2.67 (2) 2.13
 (3) 3.27 (4) 3.041

47. 36 g H_2O is electrolysed by passing 3 ampere current. What will be time required for complete decomposition (in hour) :-

- (1) 17.87 hour (2) 15.84 hour
 (3) 35.74 hour (4) 25.35 hour

ANSWER KEY

EXERCISE-III (Analytical Questions)

Check Your Understanding

- Specific conductance of 0.1 M Nitric acid is $6.3 \times 10^{-2} \text{ ohm}^{-1} \text{ cm}^{-1}$. The molar conductance of the solution is :
 (1) $630 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 (2) $315 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 (3) $100 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 (4) $6.300 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 (5) $63.0 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
- The highest electrical conductivity of the following aqueous solution is of
 (1) 0.1 M fluoroacetic acid
 (2) 0.1 M difluoroacetic acid
 (3) 0.1 M acetic acid
 (4) 0.1 M chloroacetic acid
- Resistance of a conductivity cell filled with a solution of an electrolyte of concentration 0.1 M is 100Ω . The conductivity of this solution is 1.29 S m^{-1} . Resistance of the same cell when filled with 0.02 M of the same solution is 520Ω . The molar conductivity of 0.02 M solution of the electrolyte will be.
 (1) $124 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 (2) $1240 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 (3) $1.24 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 (4) $12.4 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
- Resistance of 0.2 M solution of an electrolyte is 50Ω . The specific conductance of the solution is 1.4 S m^{-1} . The resistance of 0.5 M solution of the same electrolyte is 280Ω . The molar conductivity of 0.5 M solution of the electrolyte in $\text{S m}^2 \text{ mol}^{-1}$ is :
 (1) 5×10^3 (2) 5×10^2
 (3) 5×10^{-4} (4) 5×10^{-3}
- Resistance of 0.2 M solution of an electrolyte is 50Ω . The specific conductance of the solution is 1.3 S m^{-1} . If resistance of the 0.4 M solution of the same electrolyte is 260Ω then its molar conductivity is:-
 (1) $6250 \text{ S m}^2 \text{ mol}^{-1}$
 (2) $6.25 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 (3) $625 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 (4) $62.5 \text{ S m}^2 \text{ mol}^{-1}$
- The equivalent conductance of NaCl at concentration C and at infinite dilution are λ_C and λ_∞ , respectively. The correct relationship between λ_C and λ_∞ is given by :
 (where the constant B is positive)
 (1) $\lambda_C = \lambda_\infty - (B) \sqrt{C}$ (2) $\lambda_C = \lambda_\infty + (B) \sqrt{C}$
 (3) $\lambda_C = \lambda_\infty + (B) C$ (4) $\lambda_C = \lambda_\infty - (B) C$

Electrolyte	KCl	KNO ₃	HCl	NaOAc	NaCl
$\Lambda^\infty (\text{S cm}^2 \text{ mol}^{-1})$	149.9	145.0	426.2	91.0	126.5

- Calculate $\Lambda_{\text{HOAc}}^\infty$ using appropriate molar conductances of the electrolytes listed above at infinite dilution in H_2O at 25°C
 (1) 390.7 (2) 217.5 (3) 517.2 (4) 552.7
- Which of the following statements is TRUE for the electrochemical Daniel cell :
 (1) Electrons flow from copper electrode to zinc electrode.
 (2) Current flows from zinc electrode to copper electrode.
 (3) Cations move toward copper electrode.
 (4) Cations move toward zinc electrode.
- Aluminium displaces hydrogen from dilute HCl whereas silver does not, the E.M.F. of a cell prepared by combining Al/Al^{3+} and Ag/Ag^+ is 2.46 V. The reduction potential of silver electrode is +0.80 V. The reduction potential of aluminium electrode is :
 (1) - 3.26 V (2) +1.66 V
 (3) -1.66 V (4) 3.26 V
- Standard reduction electrode potential of three metals A, B and C are respectively + 0.5 V, - 3.0 V and -1.2 V. The reducing power of these metals are :
 (1) $C > B > A$ (2) $A > C > B$
 (3) $B > C > A$ (4) $A > B > C$
- Consider the following E° values
 $E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^\circ = +0.77 \text{ V}$
 $E_{\text{Sn}^{2+}/\text{Sn}}^\circ = -0.14 \text{ V}$
 Under standard conditions the potential for the reaction
 $\text{Sn(s)} + 2\text{Fe}^{3+}(\text{aq}) \longrightarrow 2\text{Fe}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq})$ is
 (1) 0.91 V (2) 1.40 V
 (3) 1.68 V (4) 0.63 V
- For the redox reaction :
 $\text{Zn(s)} + \text{Cu}^{2+} (0.1\text{M}) \rightarrow \text{Zn}^{2+} (1\text{M}) + \text{Cu(s)}$ taking place in a cell,
 E_{Cell}° is 1.10 V E_{Cell} for the cell will be
 $\left(2.303 \frac{RT}{F} = 0.0591 \right)$
 (1) 1.07 V (2) 0.82 V
 (3) 2.14 V (4) 1.80 V

- 13.** In a cell that utilises the reaction
- $$\text{Zn(s)} + 2\text{H}^+(\text{aq.}) \rightleftharpoons \text{Zn}^{2+}(\text{aq}) + \text{H}_2(\text{g})$$
- addition of H_2SO_4 to cathode compartment, will
- (1) increase the E_{cell} and shift equilibrium to the right
 - (2) lower the E_{cell} and shift equilibrium to the right
 - (3) lower the E_{cell} and shift equilibrium to the left
 - (4) increase the E_{cell} and shift equilibrium to the left
- 14.** The e.m.f. of a Daniell cell at 298 K is E_1 .
 $\text{Zn} | \text{ZnSO}_4 | | \text{CuSO}_4 | \text{Cu}$
 (0.01M) (1.0M)
 When the concentration of ZnSO_4 is 1.0 M and that of CuSO_4 is 0.01 M, the e.m.f. changed to E_2 . What is the relationship between E_1 and E_2
- (1) $E_1 > E_2$
 - (2) $E_1 < E_2$
 - (3) $E_1 = E_2$
 - (4) $E_2 = 0 \neq E_1$
- 15.** The reduction potential of hydrogen half-cell will be negative if :-
- (1) $p(\text{H}_2) = 2 \text{ atm}$ and $[\text{H}^+] = 1.0 \text{ M}$
 - (2) $p(\text{H}_2) = 2 \text{ atm}$ and $[\text{H}^+] = 2.0 \text{ M}$
 - (3) $p(\text{H}_2) = 1 \text{ atm}$ and $[\text{H}^+] = 2.0 \text{ M}$
 - (4) $p(\text{H}_2) = 1 \text{ atm}$ and $[\text{H}^+] = 1.0 \text{ M}$
- 16.** For a cell reaction involving two-electron change, the standard e.m.f. of the cell is found to be 0.295 V at 25° C. The equilibrium constant of the reaction at 25 °C will be :
- (1) 10
 - (2) 1×10^{10}
 - (3) 1×10^{-10}
 - (4) 29.5×10^{-2}
- 17.** For a spontaneous reaction the ΔG , equilibrium constant (K) and E_{Cell}^0 will be respectively
- (1) -ve, < 1, -ve
 - (2) -ve, > 1, -ve
 - (3) -ve, > 1, +ve
 - (4) +ve, > 1, -ve
- 18.** On the basis of the information available from the reaction
- $$\frac{4}{3}\text{Al} + \text{O}_2 \rightarrow \frac{2}{3}\text{Al}_2\text{O}_3,$$
- $\Delta G = -827 \text{ KJ mol}^{-1}$ of O_2 , the minimum e.m.f. required to carry out an electrolysis of Al_2O_3 is :-
 (F = 96500 C mol⁻¹)
- (1) 2.14 V
 - (2) 4.28 V
 - (3) 6.42 V
 - (4) 8.56 V

- 19.** Given below are the half-cell reactions :-
- $$\text{Mn}^{2+} + 2\text{e}^- \rightarrow \text{Mn}; \quad E^\circ = -1.18 \text{ V}$$
- $$2(\text{Mn}^{3+} + \text{e}^- \rightarrow \text{Mn}^{2+}); \quad E^\circ = +1.51 \text{ V}$$
- The E° for $3\text{Mn}^{2+} \rightarrow \text{Mn} + 2\text{Mn}^{3+}$ will be :
- (1) -0.33 V; the reaction will not occur
 - (2) -0.33 V; the reaction will occur
 - (3) -2.69 V; the reaction will not occur
 - (4) -2.69 V; the reaction will occur
- 20.** Given the data at 25°C,
- $$\text{Ag} + \text{I}^- \rightarrow \text{AgI} + \text{e}^-, \quad E^\circ = 0.152 \text{ V}$$
- $$\text{Ag} \rightarrow \text{Ag}^+ + \text{e}^-, \quad E^\circ = -0.800 \text{ V}$$
- What is the value of $\log K_{\text{sp}}$ for AgI ?
- $$\left(2.303 \frac{RT}{F} = 0.059 \text{ V} \right)$$
- (1) -8.12
 - (2) +8.612
 - (3) -37.83
 - (4) -16.13
- 21.** The metal that cannot be obtained by electrolysis of an aqueous solution of its salts is:
- (1) Cu
 - (2) Cr
 - (3) Ag
 - (4) Ca
- 22.** When during electrolysis of a solution of AgNO_3 9650 C of charge pass through the electroplating bath, the mass of silver deposited on the cathode will be :
- (1) 21.6 g
 - (2) 108 g
 - (3) 1.08 g
 - (4) 10.8 g
- 23.** Aluminium oxide may be electrolysed at 1000 °C to furnish aluminium metal (At. Mass = 27 amu; 1 Faraday = 96500 C). The cathode reaction is $\text{Al}^{3+} + 3\text{e}^- \rightarrow \text{Al}$
 To prepare 5.12 kg of aluminium metal by this method would require.
- (1) $5.49 \times 10^4 \text{ C}$ of electric charge
 - (2) $5.49 \times 10^1 \text{ C}$ of electric charge
 - (3) $5.49 \times 10^7 \text{ C}$ of electric charge
 - (4) $1.83 \times 10^7 \text{ C}$ of electric charge
- 24.** A current of 'i' A was passed for 't' s. through three cells P, Q and R connected in a series. These contain respectively silver nitrate, mercuric nitrate and mercurous nitrate. At the cathode of the cell P, 0.216 g of Ag was deposited. The weights of mercury deposited in the cathode of Q and R respectively are : (at. wt. of Hg = 200.59)
- (1) 0.4012 g and 0.8024 g
 - (2) 0.4012 g and 0.2006 g
 - (3) 0.2006 g and 0.4012 g
 - (4) 0.1003 g and 0.2006 g

- 25.** A galvanic cell is set up from a zinc bar weighing 100 g and 1.0 L of 1.0 M CuSO_4 solution. How long would the cell will run if it is assumed to deliver a steady current of 1.0 A. (Atomic mass of Zn = 65)
- (1) 1.1 hr. (2) 46 hr.
(3) 53.6 hr. (4) 24.00 hr.
- 26.** The electrochemical equivalent of silver is 0.0011180 g. When an electric current of 0.5 A is passed through an aqueous silver nitrate solution for 200 s, the amount of silver deposited is :
- (1) 1.1180 g (2) 0.11180 g
(3) 5.590 g (4) 0.5590 g
- 27.** A current of 9.65 A is passed through an aqueous solution of NaCl using suitable electrodes for 1000 s. Given that 1 faraday equals 96500 C the amount of NaOH (mol wt. 40.00) formed on electrolysis is :
- (1) 2.0 g (2) 8.0 g
(3) 4.0 g (4) 1.0 g.
- 28.** An electric current is passed through silver voltameter connected to a water voltameter. The cathode of the silver voltameter weighed 0.108 g more at the end of the electrolysis. The volume of oxygen evolved at STP is :
- (1) 56 cm^3 (2) 550 cm^3
(3) 5.6 cm^3 (4) 11.2 cm^3
(5) 22.4 cm^3
- 29.** 4.5 g of aluminium (at. mass 27 amu) is deposited at cathode from Al^{3+} solution by a certain quantity of electric charge. The volume of hydrogen produced at STP from H^+ ions in solution by the same quantity of electric charge will be –
- (1) 44.8 L (2) 11.2 L
(3) 22.4 L (4) 5.6 L
- 30.** The thermodynamic efficiency of cell is given by -
- (1) $\frac{\Delta H}{\Delta G}$ (2) $\frac{nFE}{\Delta G}$
(3) $-\frac{nFE}{\Delta H}$ (4) Zero

EXERCISE-III (Analytical Questions)

ANSWER KEY

Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	1	2	1	3	2	1	1	3	3	3	1	1	1	1	1
Que.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Ans.	2	3	1	3	4	4	4	3	3	3	2	3	3	4	3

EXERCISE-IV (Assertion & Reason)

Target AIIMS

Directions for Assertion & Reason questions

These questions consist of two statements each, printed as Assertion and Reason. While answering these Questions you are required to choose any one of the following four responses.

- (A) If both Assertion & Reason are True & the Reason is a correct explanation of the Assertion.
 (B) If both Assertion & Reason are True but Reason is not a correct explanation of the Assertion.
 (C) If Assertion is True but the Reason is False.
 (D) If both Assertion & Reason are false.

- Assertion :-** Electrolysis of molten calcium hydride produces hydrogen gas at anode.
Reason :- Hydrogen in calcium hydride is present as H^- ion.
 (1) A (2) B (3) C (4) D
- Assertion :-** Molar conductance of 0.1 M NH_4OH solution is less than that of 0.001M NH_4OH solution.
Reason :- Dilution increases the degree of ionisation of NH_4OH .
 (1) A (2) B (3) C (4) D
- Assertion :-** Salt bridge is generally used in the electrochemical cells.
Reason :- The ions of the electrolyte used in the salt bridge should have nearly same transport number.
 (1) A (2) B (3) C (4) D
- Assertion :-** The electrode potential of SHE is zero.
Reason :- SHE is a standard reference electrode.
 (1) A (2) B (3) C (4) D
- Assertion :-** Copper sulphate solution cannot be stored in iron-vessel.
Reason :- Copper atoms are oxidised by iron ions.
 (1) A (2) B (3) C (4) D
- Assertion :-** Absolute value of E_{red}^0 of an electrode cannot be determined.
Reason :- Neither oxidation nor reduction can take place alone.
 (1) A (2) B (3) C (4) D
- Assertion :-** A dry cell became dead after long time even if it has not been used.
Reason :- The NH_4Cl slowly and gradually corrodes the zinc container.
 (1) A (2) B (3) C (4) D
- Assertion :-** The molar conductance of weak electrolytes is low as compared to that of strong electrolytes at moderate concentrations.
Reason :- Weak electrolytes at moderate concentration dissociates to a much greater extent as compared to strong electrolytes.
 (1) A (2) B (3) C (4) D
- Assertion :-** Galvanised iron does not rust.
Reason :- Zinc has a more negative electrode potential than iron.
 (1) A (2) B (3) C (4) D
- Assertion :-** Increase in the concentration of copper half cell in Daniel cell increases the emf of the cell.
Reason :- According to the nernst equation

$$emf \text{ of Daniel cell} = E_{cell}^0 + \frac{0.059}{2} \log \frac{[Cu^{+2}]}{[Zn^{+2}]}$$
 (1) A (2) B (3) C (4) D
- Assertion :-** Sodium ions are discharged at the Hg electrode during electrolysis in preference to H^+ ions.
Reason :- The nature of electrode also affects the order of discharge of cations.
 (1) A (2) B (3) C (4) D
- Assertion :-** The cell constant of a cell depends upon the nature of the material of the electrodes.
Reason :- The observed conductance depends upon the nature of the electrolyte and the concentration of the solution.
 (1) A (2) B (3) C (4) D
- Assertion :-** At the end of electrolysis using platinum electrodes, an aqueous solution of copper sulphate turns colourless.
Reason :- Copper in $CuSO_4$ is converted to $Cu(OH)_2$ during the electrolysis.
 (1) A (2) B (3) C (4) D

- 14. Assertion :-** In electrolysis, the quantity of electricity needed for depositing 1 mol of silver is different from that required for 1 mol of copper.
Reason :- The atomic weight of silver and copper are different.
 (1) A (2) B (3) C (4) D
- 15. Assertion :-** pH of solution increases during electrolysis of aqueous solution of NaCl.
Reason :- Electrolysis is non-spontaneous process.
 (1) A (2) B (3) C (4) D
- 16. Assertion :-** A reaction is spontaneous if $E_{\text{cell}} = +ve$.
Reason :- For $E_{\text{cell}} = +ve$, (ΔG) is always $-ve$.
 (1) A (2) B (3) C (4) D
- 17. Assertion :-** Cu is stronger reducing agent than H_2 .
Reason :- E° of Cu^{+2}/Cu is negative.
 (1) A (2) B (3) C (4) D
- 18. Assertion :-** For cell reaction; at 298 K
 $Zn(s) + Cu^{+2}(1M) \rightleftharpoons Zn^{+2}(1M) + Cu(s)$
 $E_{\text{cell}}^0 = 0$.
Reason :- At standard condition ; cell potential is always zero.
 (1) A (2) B (3) C (4) D
- 19. Assertion :-** Molar conductivity of an electrolyte solution increases on dilution.
Reason :- Decrease in conductivity is overcome by increase in volume.
 (1) A (2) B (3) C (4) D
- 20. Assertion :-** In electrolytic refining, electrolytic cell consists of pure and impure Cu electrodes dipped in $CuSO_4(aq)$.
Reason :- Pure Cu is obtained at anode.
 (1) A (2) B (3) C (4) D
- 21. Assertion :-** Molar conductivity of CH_3COOH increases with dilution.
Reason :- The graph between λ_m v/s $(C)^{1/2}$ is a straight line for CH_3COOH .
 (1) A (2) B (3) C (4) D
- 22. Assertion :-** On electrolysis of aq. NaCl, Na is obtained.
Reason :- Na^+ is reduced at cathode.
 (1) A (2) B (3) C (4) D

EXERCISE-IV (Assertion & Reason)

ANSWER KEY

Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	1	1	2	2	3	1	1	3	1	1	1	4	3	2	2
Que.	16	17	18	19	20	21	22								
Ans.	1	4	4	1	3	3	4								

[illegible]

This image shows a full page of white paper with evenly spaced horizontal black lines, typical of notebook or primary writing paper. There are no margins, text, or other markings present.