

CHAPTER – 7

COORDINATE GEOMETRY

DISTANCE FORMULA

The distance between any two points $A(x_1, y_1)$ and $B(x_2, y_2)$ is given by

$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

or $AB = \sqrt{(\text{difference of abscissae})^2 + (\text{difference of ordinates})^2}$

Distance of a point from origin

The distance of a point $P(x, y)$ from origin O is given by $OP = \sqrt{x^2 + y^2}$

Problems based on geometrical figure

To show that a given figure is a

- ☞ Parallelogram – prove that the opposite sides are equal
- ☞ Rectangle – prove that the opposite sides are equal and the diagonals are equal.
- ☞ Parallelogram but not rectangle – prove that the opposite sides are equal and the diagonals are not equal.
- ☞ Rhombus – prove that the four sides are equal
- ☞ Square – prove that the four sides are equal and the diagonals are equal.
- ☞ Rhombus but not square – prove that the four sides are equal and the diagonals are not equal.
- ☞ Isosceles triangle – prove any two sides are equal.
- ☞ Equilateral triangle – prove that all three sides are equal.
- ☞ Right triangle – prove that sides of triangle satisfies Pythagoras theorem.

IMPORTANT QUESTIONS

Show that the points (1, 7), (4, 2), (-1, -1) and (-4, 4) are the vertices of a square.

Solution : Let $A(1, 7)$, $B(4, 2)$, $C(-1, -1)$ and $D(-4, 4)$ be the given points.

$$AB = \sqrt{(1-4)^2 + (7-2)^2} = \sqrt{9+25} = \sqrt{34}$$

$$BC = \sqrt{(4+1)^2 + (2+1)^2} = \sqrt{25+9} = \sqrt{34}$$

$$CD = \sqrt{(-1+4)^2 + (-1-4)^2} = \sqrt{9+25} = \sqrt{34}$$

$$DA = \sqrt{(1+4)^2 + (7-4)^2} = \sqrt{25+9} = \sqrt{34}$$

$$AC = \sqrt{(1+1)^2 + (7+1)^2} = \sqrt{4+64} = \sqrt{68}$$

$$BD = \sqrt{(4+4)^2 + (2-4)^2} = \sqrt{64+4} = \sqrt{68}$$

Since, $AB = BC = CD = DA$ and $AC = BD$, all the four sides of the quadrilateral $ABCD$ are equal and its diagonals AC and BD are also equal. Therefore, $ABCD$ is a square.

Find a point on the y-axis which is equidistant from the points $A(6, 5)$ and $B(-4, 3)$.

Solution : We know that a point on the y-axis is of the form $(0, y)$. So, let the point $P(0, y)$ be equidistant from A and B . Then $AP^2 = BP^2$

$$\Rightarrow (6-0)^2 + (5-y)^2 = (-4-0)^2 + (3-y)^2$$

$$\Rightarrow 36 + 25 + y^2 - 10y = 16 + 9 + y^2 - 6y \Rightarrow 4y = 36 \Rightarrow y = 9$$

So, the required point is $(0, 9)$.

Questions for practice

1. Show that the points A(1, 2), B(5, 4), C(3, 8) and D(-1, 6) are vertices of a square.
2. Show that the points A(5, 6), B(1, 5), C(2, 1) and D(6, 2) are vertices of a square.
3. Show that the points A(1, -3), B(13, 9), C(10, 12) and D(-2, 0) are vertices of a rectangle.
4. Show that the points A(1, 0), B(5, 3), C(2, 7) and D(-2, 4) are vertices of a rhombus.
5. Prove that the points A(-2, -1), B(1, 0), C(4, 3) and D(1, 2) are vertices of a parallelogram.
6. Find the point on x-axis which is equidistant from (7, 6) and (-3, 4).
7. Find the point on the x-axis which is equidistant from (2, -5) and (-2, 9).
8. Find a point on the y-axis which is equidistant from the points A(5, 2) and B(-4, 3).
9. Find a point on the y-axis which is equidistant from the points A(5, -2) and B(-3, 2).
10. Find the values of y for which the distance between the points P(2, -3) and Q(10, y) is 10 units.
11. Find the value of a, if the distance between the points A(-3, -14) and B(a, -5) is 9 units.
12. If the point A(2, -4) is equidistant from P(3, 8) and Q(-10, y), find the values of y. Also find distance PQ.

Section formula

The coordinates of the point P(x, y) which divides the line segment joining the points A(x₁, y₁) and B(x₂, y₂), internally, in the ratio m₁ : m₂ are

$$\left(\frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \right)$$

Mid-point formula

The coordinates of the point P(x, y) which is the midpoint of the line segment joining the points

A(x₁, y₁) and B(x₂, y₂), are $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

IMPORTANT QUESTIONS

Find the coordinates of the point which divides the line segment joining the points (4, -3) and (8, 5) in the ratio 3 : 1 internally.

Solution : Let P(x, y) be the required point.

Using the section formula, $x = \frac{m_2 x_1 + m_1 x_2}{m_1 + m_2}$, $y = \frac{m_2 y_1 + m_1 y_2}{m_1 + m_2}$ we get

$$x = \frac{3(8) + 1(4)}{3 + 1} = 7, y = \frac{3(5) + 1(-3)}{3 + 1} = 3$$

Therefore, (7, 3) is the required point.

In what ratio does the point (-4, 6) divide the line segment joining the points A(-6, 10) and B(3, -8)?

Solution : Let (-4, 6) divide AB internally in the ratio k : 1.

Using the section formula, $x = \frac{m_2 x_1 + m_1 x_2}{m_1 + m_2}$, $y = \frac{m_2 y_1 + m_1 y_2}{m_1 + m_2}$ we get

$$y = \frac{k(-8) + 1(10)}{k + 1} = 6$$

$$\Rightarrow -8k + 10 = 6k + 6 \Rightarrow -8k - 6k = 6 - 10$$

$$\Rightarrow -14k = -4 \Rightarrow k = \frac{4}{14} = \frac{2}{7}$$

Therefore, the point (-4, 6) divides the line segment joining the points A(-6, 10) and B(3, -8) in the ratio 2 : 7.

Questions for practice

1. Find the coordinates of the point which divides the join of $(-1, 7)$ and $(4, -3)$ in the ratio $2 : 3$.
2. Find the coordinates of the points of trisection of the line segment joining $(4, -1)$ and $(-2, -3)$.
3. Find the coordinates of the points of trisection (i.e., points dividing in three equal parts) of the line segment joining the points $A(2, -2)$ and $B(-7, 4)$.
4. Find the ratio in which the y-axis divides the line segment joining the points $(5, -6)$ and $(-1, -4)$. Also find the point of intersection.
5. Find the ratio in which the line segment joining the points $(-3, 10)$ and $(6, -8)$ is divided by $(-1, 6)$.
6. Find the ratio in which the line segment joining $A(1, -5)$ and $B(-4, 5)$ is divided by the x-axis. Also find the coordinates of the point of division.
7. Find the coordinates of the points which divide the line segment joining $A(-2, 2)$ and $B(2, 8)$ into four equal parts.
8. If the points $A(6, 1)$, $B(8, 2)$, $C(9, 4)$ and $D(p, 3)$ are the vertices of a parallelogram, taken in order, find the value of p .
9. If $(1, 2)$, $(4, y)$, $(x, 6)$ and $(3, 5)$ are the vertices of a parallelogram taken in order, find x and y .
10. In what ratio does the x-axis divide the line segment joining the points $(-4, -6)$ and $(-1, 7)$? Find the coordinates of the point of division.
11. If $P(9a - 2, -b)$ divides line segment joining $A(3a + 1, -3)$ and $B(8a, 5)$ in the ratio $3 : 1$, find the values of a and b .
12. If (a, b) is the mid-point of the line segment joining the points $A(10, -6)$ and $B(k, 4)$ and $a - 2b = 18$, find the value of k and the distance AB .
13. The centre of a circle is $(2a, a - 7)$. Find the values of a if the circle passes through the point $(11, -9)$ and has diameter $10\sqrt{2}$ units.
14. The line segment joining the points $A(3, 2)$ and $B(5, 1)$ is divided at the point P in the ratio $1:2$ and it lies on the line $3x - 18y + k = 0$. Find the value of k .
15. Find the coordinates of the point R on the line segment joining the points $P(-1, 3)$ and $Q(2, 5)$ such that $PR = \frac{3}{5}PQ$.
16. Find the values of k if the points $A(k + 1, 2k)$, $B(3k, 2k + 3)$ and $C(5k - 1, 5k)$ are collinear.
17. Find the ratio in which the line $2x + 3y - 5 = 0$ divides the line segment joining the points $(8, -9)$ and $(2, 1)$. Also find the coordinates of the point of division.
18. The mid-points D, E, F of the sides of a triangle ABC are $(3, 4)$, $(8, 9)$ and $(6, 7)$. Find the coordinates of the vertices of the triangle.

MCQ (1 MARK)

1. The distance of the point $P(4, -3)$ from the origin is
(a) 1 unit (b) 7 units (c) 5 units (d) 3 units
2. The distance between the points $A(2, -3)$ and $B(2, 2)$ is
(a) 2 units (b) 4 units (c) 5 units (d) 3 units
3. What is the midpoint of a line with endpoints $(-3, 4)$ and $(10, -5)$?
(a) $(-13, -9)$ (b) $(-6.5, -4.5)$ (c) $(3.5, -0.5)$ (d) none of these
4. If the distance between the points $(8, p)$ and $(4, 3)$ is 5 then value of p is
(a) 6 (b) 0 (c) both (a) and (b) (d) none of these
5. If the origin is the mid-point of the line segment joined by the points $(2, 3)$ and (x, y) , then the value of (x, y) is
(a) $(2, -3)$ (b) $(2, 3)$ (c) $(-2, 3)$ (d) $(-2, -3)$

6. The distance of the point $P(2, 3)$ from the x-axis is:
(a) 2 (b) 3 (c) 1 (d) 5
7. The distance between the points $A(0, 6)$ and $B(0, -2)$ is:
(a) 2 (b) 6 (c) 4 (d) 8
8. The distance of the point $P(-6, 8)$ from the origin is:
(a) 8 (b) 27 (c) 10 (d) 6
9. The distance between the points $(0, 5)$ and $(-5, 0)$ is:
(a) 5 (b) 52 (c) 25 (d) 10
10. AOBC is a rectangle whose three vertices are $A(0, 3)$, $O(0, 0)$ and $B(5, 0)$. The length of its diagonal is:
(a) 5 (b) 3 (c) 34 (d) 4
11. The perimeter of a triangle with vertices $(0, 4)$, $(0, 0)$ and $(3, 0)$ is:
(a) 5 (b) 12 (c) 11 (d) $7 + 5$
12. The points $(-4, 0)$, $(4, 0)$, $(0, 3)$ are the vertices of a :
(a) Right triangle (b) Isosceles triangle (c) Equilateral triangle (d) Scalene triangle
13. Point on x – axis has coordinates:
(a) $(a, 0)$ (b) $(0, a)$ (c) $(-a, a)$ (d) $(a, -a)$
14. Point on y – axis has coordinates:
(a) $(-a, b)$ (b) $(a, 0)$ (c) $(0, b)$ (d) $(-a, -b)$